

Pacific Journal of Mathematics

INTRINSIC EXTENSIONS OF RINGS

JOHN JOSEPH HUTCHINSON

INTRINSIC EXTENSIONS OF RINGS

JOHN J. HUTCHINSON

Faith posed the problem of characterizing the left intrinsic extensions of left quotient semisimple (simple) rings. In this paper a characterization is given for the left strongly intrinsic extensions of left quotient semisimple rings.

Section 1 consists of several definitions and known preliminary results. In §2 we define essential subdirect sums and develop several of their elementary properties. The results of §2 enable us to state and prove the main characterization theorem which appears in §3. In the last section it is shown that in the class of left quotient semisimple rings, the left strongly intrinsic extensions are exactly the left intrinsic extensions.

1. Preliminaries. Let R and S be nonzero associative rings (not necessarily with identities or commutative) where $S \subseteq R$. S is *left quotient simple*, *left quotient semisimple*, a *left Ore domain* if S has a left classical (and maximal) quotient ring which is respectively simple Artinian, semisimple Artinian, a division ring. The left classical quotient ring of S will be denoted \bar{S} , and left quotient semisimple (left quotient simple) will be written lqss (lqs). R is a *left intrinsic extension* of S if every nonzero left ideal of R has nonzero intersection with S . A left S -module M (denoted ${}_sM$) is an *essential extension* of a submodule N if every nonzero submodule of M has nonzero intersection with N (we also say N is essential in M). R is a *left essential extension* of S if ${}_sR$ is an essential extension of ${}_sS$. It is clear that every left essential extension of S is left intrinsic, but the converse is not always true (for instance when R is a proper field extension of a field S). A left ideal A of S is *closed* if S contains no proper left essential extensions of A (as left S -modules). The symbol $L(S)$ will denote the set of closed left ideals of S . R is a *left strongly intrinsic extension* of S if R is a left intrinsic extension of S , and for all $A \in L(S)$ there exists a left ideal B of R such that $B \cap S = A$. In any left S -module M , we denote by $Z({}_sM)$ the set of elements in M whose annihilator in S is an essential left ideal. Clearly $Z({}_sM)$ is a submodule of M .

THEOREM 1.1. *If $Z({}_sS) = 0$, then ${}_sS$ has a (unique up to isomorphism) maximal essential extension Q (called the maximal quotient ring of S) which has a ring structure compatible with the module structure; and Q is a regular, left self-injective ring such that*

$Z(Q) = 0$. Moreover $L(S)$ and $L(Q)$ are lattices, and $L(Q) \cong L(S)$ under contraction.

Proof. Follows from [1, Theorem 1, p 69] and [3, Corollary 2.6] and their proofs.

The following two lemmas appear in [2].

LEMMA 1.2. *If R is a left strongly intrinsic extension of S , then the following are equivalent: (i) $Z(S) = 0$, (ii) $Z({}_S R) = 0$, (iii) $Z({}_R R) = 0$.*

LEMMA 1.3. *If $Z({}_S S) = 0$ and R is a left strongly intrinsic extension of S , then $L(R) \cong L(S)$ under contraction.*

2. Essential subdirect sums. If R is a subdirect sum of rings $\{R_\alpha \mid \alpha \in A\}$ and $S = \sum_{\alpha \in A}^c R_\alpha$ is the complete direct sum of the R_α , then the subdirect sum is essential if R (identifying R and its canonical isomorphic image in S) is an essential left R -submodule of S .

Clearly an essential subdirect sum of nonzero rings is irredundant [5], and in the case of a finite number of factors, is essentially irredundant in the sense of [1, p 114]. It is an easily verified property of subdirect sums that if $B_i (i \in D)$ are disjoint subsets of A such that $A = \bigcup_{i \in D} B_i$ and $R_{B_i} = \{a \in \sum_{\alpha \in B_i}^c R_\alpha \mid \text{for some } b \in R \ a(\alpha) = b(\alpha) \text{ for all } \alpha \in B_i\}$, then for each $i \in D$, the R_{B_i} are rings which are subdirect sums in a natural way of the rings $\{R_\alpha \mid \alpha \in B_i\}$, and R is a subdirect sum in a natural way of the rings $\{R_{B_i} \mid i \in D\}$. If, in addition, each R_α is a subdirect sum of rings $\{T_{\alpha\gamma} \mid \gamma \in A_\alpha\}$, then R is a subdirect sum in a natural way of the rings $\{T_{\alpha\gamma} \mid \gamma \in A_\alpha \text{ whenever } \alpha \in A\}$. Each of these constructed subdirect sums will be referred to as the *induced subdirect sum*, and whenever we say "the subdirect sum" we are referring either to the original fixed subdirect sum or one of its various induced subdirect sums. Loosely speaking, we may think of the preceding remarks as saying that subdirect sums satisfy a generalized associative law. The results in this section will show that finite essential subdirect sums also have this nice property (finite irredundant subdirect sums do not).

LEMMA 2.1. *Let R be a subdirect sum of nonzero rings R_1, \dots, R_n . The subdirect sum is essential if and only if $R \cap R_i$ is an essential left R -submodule of R_i for $i = 1, 2, \dots, n$.*

Proof. If the subdirect sum is essential and W_1 is a nonzero R -submodule of R_1 , then W_1 is also a nonzero left R -submodule of $\bigoplus \sum_{i=1}^n R_i$, and so $W_1 \cap R \neq 0$. Since $W_1 \cap (R \cap R_1) = W_1 \cap R$ the

result follows. Conversely suppose $R \cap R_i$ is an essential R -submodule of R_i for each i . If $n = 1$ the result is trivial, so suppose $n = 2$. Let $0 \neq x_1 + x_2 \in R_1 \oplus R_2$ ($x_i \in R_i$) and assume $x_1 \neq 0$. Since $R \cap R_1$ is essential in R_1 , there exists an integer n and an $r \in R$ such that $0 \neq rx_1 + nx_1 = (r + n)x_1 \in R \cap R_1$. If $(r + n)x_2 = 0$, then $0 \neq (r + n)(x_1 + x_2) \in R$. If $(r + n)x_2 \neq 0$, then since $R \cap R_2$ is essential in R_2 there exists an integer m and an $s \in R$ such that $0 \neq (s + m)(r + n)x_2 \in R \cap R_2$. Then clearly $0 \neq (s + m)(r + n)(x_1 + x_2) \in R$, and R is essential in $R_1 \oplus R_2$. The result now follows by a simple induction.

PROPOSITION 2.2. *Let Q be a ring which is an essential subdirect sum of nonzero rings Q_1, \dots, Q_n . If each Q_i is an essential subdirect sum of nonzero rings $Q_{i,1}, \dots, Q_{i,k_i}$, then the induced subdirect sum of Q (of the $Q_{i,j}$) is essential. Also, if n_1, n_2, \dots, n_{k+1} are integers such that $1 = n_1 < n_2 < \dots < n_{k+1} = n + 1$, and Q'_i ($i = 1, 2, \dots, k$) are the induced subdirect sums of $Q_{n_i}, \dots, Q_{n_{i+1}-1}$, then Q is the essential subdirect sum of Q'_1, \dots, Q'_k and each Q'_i is the essential subdirect sum of $Q_{n_i}, \dots, Q_{n_{i+1}-1}$.*

Proof. The result follows by a fairly straightforward application of Lemma 2.1.

The following theorem is a modification of a theorem of Levy, [5, Theorem 6.1].

THEOREM 2.3. *R is lqss if and only if it is an essential subdirect sum of a finite number of lqs rings R_1, \dots, R_n for some n . In this case, we have $\bar{R} = \bigoplus \sum_{i=1}^n \bar{R}_i$.*

Proof. If R is lqss, then by [5, Theorem 6.1], R is an irredundant subdirect sum of lqs rings R_1, \dots, R_n for some n , and $R \subseteq R_1 \oplus \dots \oplus R_n \subseteq \bar{R}_1 \oplus \dots \oplus \bar{R}_n \subseteq \bar{R}$. Since \bar{R} is an essential extension of R , it follows that R is essential in $R_1 \oplus \dots \oplus R_n$, and R is then an essential subdirect sum of R_1, \dots, R_n . Conversely, if R is a finite essential (and so irredundant) subdirect sum of lqs rings R_1, \dots, R_n , then R is lqss by [5, Theorem 6.1].

THEOREM 2.4. *If R (S) is an essential subdirect sum of prime rings R_1, \dots, R_n (S_1, \dots, S_n), $S \subseteq R$, and each R_i is a left intrinsic extension of the corresponding S_i , then R is a left intrinsic extension of S .*

Proof. If $0 \neq x \in R$, then $x = x_1 + \dots + x_n$ ($x_i \in R_i$). We may

assume $x_1 \neq 0$, and so $0 \neq R_1 x_1 = R_1 x$ since R_1 is prime. Thus $R_1 x$ is a nonzero left ideal of R_1 , so $R_1 x \cap S_1 \neq 0$. $R_1 x \cap S_1$ is a nonzero left S -submodule of $S_1 \oplus \cdots \oplus S_n$, so $S \cap R_1 x \cap S_1 \neq 0$. Since S_1 is prime, $(S \cap R_1 x \cap S_1)^2 \neq 0$. Hence there exists $s, s' \in S \cap R_1 x \cap S_1$ such that $0 \neq ss' \in S$. If $s' = r_1 x$, then there exists $r \in R$ such that $r = r_1 + \cdots + r_n$. Hence $srx = sr_1 x = ss' \in R x \cap S$, so R is a left intrinsic extension of S .

3. The main theorem.

LEMMA 3.1. *If rings R and S are direct sums of division rings R_1, \dots, R_n and S_1, \dots, S_m respectively, and R is a left intrinsic extension of S such that their identities coincide, then $m = n$ and $R_1 \cap S = S_i$ for a suitable arrangement of the R_i .*

Proof. Since R_1 is a nonzero ideal of R , it follows that $R_1 \cap S$ is a nonzero ideal of S . The ideals of S are direct sums of some of the S_i , so $R_1 \cap S = \bigoplus \sum_{i=1}^k S_i$ for some rearrangement of the S_i . But $k = 1$, otherwise R_1 has nonzero zero divisors. Similarly, each R_i contains exactly one S_i , so $n \leq m$ and $R_i \cap S = S_i$ for $i = 1, 2, \dots, n$. Each S_i^* ($i = 1, 2, \dots, n$) is a multiplicative subgroup of R_i^* , so their identities coincide. Equating identities leads to a contradiction if $n < m$, so we must have $n = m$.

PROPOSITION 3.2. *If S is a left Ore domain, then R is a left intrinsic extension of S if and only if R is a left strongly intrinsic extension of S .*

Proof. Let R be a left intrinsic extension of S . By Theorem 1.1, $L(S) \cong L(\bar{S}) = \{0, \bar{S}\}$, so $L(S) = \{0, S\}$. The zero ideal of R and R itself contract to the elements of $L(S)$, so R is a left strongly intrinsic extension of S . Note that by Lemma 1.3, $L(R) = \{0, R\}$.

PROPOSITION 3.3. *If R is a left intrinsic extension of a left Ore domain S , then R is a left Ore domain.*

Proof. By Proposition 3.2, R is a left strongly intrinsic extension of S , and $\{0, R\} = L(R)$ which clearly satisfies the maximum condition. By Lemma 1.2, $Z({}_R R) = Z({}_S S) = 0$. If A is a nilpotent left ideal of R , then $A \cap S$ is a nilpotent left ideal of S . Thus $A \cap S = 0$, and $A = 0$. Hence R is semiprime, and by [3, Theorem 4.4], R is lqss. Thus $\{0, R\} = L(R) \cong L(\bar{R}) = \{0, \bar{R}\}$. By [1, Proposition 5, p. 71], $L(\bar{R})$ consists of the annihilator left ideals of \bar{R} . Since \bar{R} has an identity, \bar{R} is a domain. It follows that R is a left Ore domain.

THEOREM 3.4. *If S is lqss, then R is a left strongly intrinsic extension of S if and only if $S \subseteq R$ and one of the following is true:*

- (i) $\bar{S} = \bar{R}$ is semisimple Artinian,
- (ii) S (and R) is an essential subdirect sum of left Ore domains S'_1, \dots, S'_n (R'_1, \dots, R'_n) where R'_i is a left intrinsic extension of the corresponding S'_i ,
- (iii) S (and R) is an essential subdirect sum of nonzero rings S_1 and S_2 (R_1 and R_2) where $S_i \subseteq R_i$ for $i = 1, 2$ and such that (i) holds for S_2 and R_2 and (ii) holds for S_1 and R_1 .

Proof. By [3, Theorem 4.4], $L(S)$ satisfies the maximum condition, so by Lemma 1.3, so does $L(R)$. By Lemma 1.2, $Z({}_R R) = Z({}_S S) = 0$; and as in Proposition 3.3, R is semiprime. Thus by [3, Theorem 4.4], R is lqss. By Theorem 1.1, \bar{R} is a regular, semisimple, left self-injective ring. The lattice of principal left ideals of \bar{R} is complete by [6, Theorem 1], so by [6, Corollary to Theorem 4], \bar{R} can be decomposed into the direct sum of two ideal Q_1 and Q_2 in such a way that Q_1 is strongly regular and Q_2 does not contain any nonzero strongly regular ideals. By [2, Theorem 2.5], there is a subring T of Q_1 with the properties that:

- (a) T contains every idempotent of Q_1 ,
- (b) T is a strongly regular self-injective ring,
- (c) $\bar{S} = T \oplus Q_2$.

Since \bar{S} (\bar{R}) is semisimple Artinian, $\bar{S} = \bigoplus \sum_{i=1}^m F_i$ ($\bar{R} = \bigoplus \sum_{i=1}^{m'} D_i$) where each F_i (D_i) is simple Artinian. Since $\bar{S} = T \oplus Q_2$ ($\bar{R} = Q_1 \oplus Q_2$), we have $T = \bigoplus \sum_{i=1}^n F_i$ ($Q_1 = \bigoplus \sum_{i=1}^{n'} D_i$) where $0 \leq n \leq m$ ($0 \leq n' \leq m'$), and the F_i (D_i) are suitably arranged. Since strongly regular rings have no nonzero nilpotent elements, it follows that $F_1, \dots, F_n, D_1, \dots, D_{n'}$ are division rings (if $n \neq 0 \neq n'$). It is clear that Q_1 is a left intrinsic extension of T (so $T = 0$ if and only if $Q_1 = 0$). By property (a) the identities of T and Q_1 coincide, so by Lemma 3.1, $n = n'$ and $D_i \cap T = F_i$ for $i = 1, 2, \dots, n$.

Let e_1 be the identity of Q_1 (and T) and e_2 the identity of Q_2 . If $T \neq 0$, let d_1, \dots, d_n be the identities of D_1, \dots, D_n (and of F_1, \dots, F_n). Let $R_i = Re_i$ and $S_i = Se_i$ for $i = 1, 2$. Clearly $S_i \subseteq R_i \subseteq Q_i$; $Q_i = 0$ if and only if $R_i = 0$ if and only if $S_i = 0$; and $S(R)$ is a subdirect sum of S_1 and S_2 (R_1 and R_2).

We claim that if $Q_1 \neq 0$, then $\bar{R}_1 = Q_1$ and $\bar{S}_1 = T$; and if $Q_2 \neq 0$, then $\bar{S}_2 = \bar{R}_2 = Q_2$. Suppose $Q_1 \neq 0$ and r is regular in R . Clearly $re_1 \in R_1$ is regular in R_1 , so R_1 has regular elements. If r_1 is any regular element in R_1 and $q_1 r_1 = 0$ where $q_1 \in Q_1$, then $q_1 = c^{-1}b$ ($c, b \in R$),

so $0 = br_1 = (be_1)r_1$. Since r_1 is regular in R_1 , it follows that $be_1 = 0$, and so $q_1 = q_1e_1 = c^{-1}be_1 = 0$. Hence r_1 is not a zero divisor in Q_1 , so by [1, Corollary 4, p. 70], r_1 is invertible in Q_1 . If q_1 is given, then $q_1 = d^{-1}b$ ($d, b \in R$) and $q_1 = q_1e_1 = (d^{-1}b)e_1 = (de_1)^{-1}(be_1)$. Hence $Q_1 = \bar{R}_1$, and exactly the same argument gives that $T = \bar{S}_1$ and (if $Q_2 \neq 0$) that $Q_2 = \bar{R}_2 = \bar{S}_2$.

Since $S \subseteq S_1 \oplus S_2 \subseteq T \oplus Q_2 = \bar{S}$ and $R \subseteq R_1 \oplus R_2 \subseteq Q_1 \oplus Q_2 = \bar{R}$, it follows that $S(R)$ is an essential subdirect sum of S_1 and S_2 (R_1 and R_2).

If $R_1 = 0$, then $S_1 = 0$; so $S = S_2 \subseteq R_2 = R$ and $\bar{S} = \bar{S}_2 = Q_2 = \bar{R}_2 = \bar{R}$. This is condition (i).

If $R_1 \neq 0$, then $e_1 = d_1 + \cdots + d_n$, $S_1 = Se_1 \subseteq Sd_1 + \cdots + Sd_n$, and $R_1 = Re_1 \subseteq Rd_1 + \cdots + Rd_n$. If $S'_i = S_i d_i = Sd_i$ ($R'_i = R_i d_i = Rd_i$) for $i = 1, 2, \dots, n$, it follows as before that $S_1(R_1)$ is a subdirect sum of S'_1, \dots, S'_n (R'_1, \dots, R'_n). In exactly the same way as we proved that $\bar{R}_1 = Q_1$, we get that $\bar{S}'_i = F_i$ and $\bar{R}'_i = D_i$ for $i = 1, 2, \dots, n$. Also, as before, the subdirect sums are essential.

We next show that R'_1 is a left intrinsic extension of S'_1 (and similarly for R'_2, \dots, R'_n). Let $0 \neq x = rd_1 \in Rd_1 = R'_1$ ($r \in R$). Then $R'_1 x = Rx \neq 0$, so $Rx \cap R \neq 0$ (since $Rx \subseteq \bar{R}$). Since R is a left intrinsic extension of S , we have $Rx \cap R \cap S = Rx \cap S \neq 0$. Thus if $0 \neq s = r'x \in Rx \cap S$ ($r' \in R$), we have $0 \neq s = sd_1 \in Sd_1$. Hence $0 \neq s \in (Rd_1)x \cap Sd_1 = R'_1 x \cap S'_1$, and R'_1 is a left intrinsic extension of S'_1 .

If $R_2 = 0$, then $S_2 = 0$; so $S = S_1$ and $R = R_1$ which gives condition (ii). If R_1 and R_2 are not zero, then condition (iii) is satisfied.

Conversely, suppose condition (i) is true. Hence $S \subseteq R \subseteq \bar{S}$, and \bar{R} exists. Thus by [3, Corollary 2.6], $L(S) \cong L(\bar{S}) = L(\bar{R}) \cong L(R)$ under contraction, so R is a left strongly intrinsic extension of S .

In condition (ii), we have by Theorem 2.3 that $\bar{S} = \bigoplus \sum_{i=1}^n \bar{S}'_i$, and $\bar{R} = \bigoplus \sum_{i=1}^n \bar{R}'_i$, where \bar{S}'_i and \bar{R}'_i are division rings. Clearly $L(\bar{S}) \cong L(\bar{R})$ under contraction, and since $L(S) \cong L(\bar{S})$ and $L(R) \cong L(\bar{R})$, it follows that $L(S) \cong L(R)$ under contraction. By Theorem 2.4, R is a left intrinsic extension of S , so R is a left strongly intrinsic extension of S .

In condition (iii), $\bar{S}_2 = \bar{R}_2$ are semisimple Artinian, so S_2 and R_2 are lqss. Let $\bar{S}_2 = \bar{R}_2 = \bigoplus \sum_{i=1}^m F_i$, where F_i are simple Artinian rings with identities e_i ($i = 1, 2, \dots, m$). Let $S'_{n+i} = S_2 e_i$ and $R'_{n+i} = R_2 e_i$ for $i = 1, 2, \dots, m$. By Theorem 2.3 and the proof of [5, Theorem 6.1], we have that $S_2(R_2)$ is an essential subdirect sum of the lqs rings S'_i (R'_i) ($i = n+1, \dots, n+m$), and $\bar{S}'_i = \bar{R}'_i = F_{i-n}$ for $i = n+1, \dots, n+m$. Since $S'_i \subseteq R'_i \subseteq \bar{S}'_i$, (for $i = n+1, \dots, n+m$), it follows that \bar{R}'_i is a left intrinsic extension of S'_i for $i = 1, \dots, n+m$. Thus by Proposition 2.2 and Theorem 2.4, R is a left intrinsic extension of S . Also $\bar{S} = \bigoplus \sum_{i=1}^{n+m} \bar{S}_i = \bar{S}_1 \oplus \bar{S}_2$, and $\bar{R} = \bigoplus \sum_{i=1}^{n+m} \bar{R}_i = \bar{R}_1 + \bar{R}_2$, and as in the proof of case (ii), $L(\bar{S}_1) \cong L(\bar{R}_1)$ under contraction. Since

$L(\bar{S}_2) = L(\bar{R}_2)$, it follows that $L(\bar{S}) \cong L(\bar{R})$ under contraction. Again $L(S) \cong L(R)$ under contraction, so R is a left strongly intrinsic extension of S .

COROLLARY 3.5. *R is a left strongly intrinsic extension of a lqs ring S if and only if either $S \subseteq R \subseteq \bar{S}$ or S and R are left Ore domains such that R is a left intrinsic extension of S .*

Proof. If $S \subseteq R \subseteq \bar{S}$, then R is a left strongly intrinsic extension of S by case (i). If R and S are left Ore domains such that R is a left intrinsic extension of S , then the result follows from Proposition 3.2.

Conversely, since $\bar{S} = T \oplus Q_2$, we have either $T = 0$ or $Q_2 = 0$. If $Q_2 = 0$, then $\bar{S} = T = F_1$ and $\bar{R} = Q_1 = D_1$, so R and S are left Ore domains and R is a left intrinsic extension of S . If $T = 0$; then $S_1 = R_1 = 0$, $S = S_2$, and $R = R_2$ which is case (i).

4. Left intrinsic extensions. In this section, it is shown that, in the case of lqss rings, every left intrinsic extension is left strongly intrinsic.

LEMMA 4.1. *If R is a left intrinsic extension of S , then $Z_{(S)}S \subseteq Z_{(S)}R \subseteq Z_{(R)}R$.*

Proof. The first containment is clear. If $x \in R$ and $x \notin Z_{(R)}R$, then the left annihilator in R of x (denoted $l_R(x)$) is not an essential left ideal of R . Thus there exists a nonzero left ideal A of R such that $l_R(x) \cap A = 0$. Thus $0 = l_R(x) \cap A \cap S = l_S(x) \cap (A \cap S)$, and $A \cap S \neq 0$. Hence $x \notin Z_{(S)}R$, and so $Z_{(S)}R \subseteq Z_{(R)}R$.

LEMMA 4.2. *Let S have a left classical quotient ring. If $Z_{(R)}R = 0$ and R is a left intrinsic extension of S , then every regular element of S is a regular element of R .*

Proof. Let s be a regular element of S , and $r \in R$. If $rs = 0$, then $r \in l_R(s)$. Clearly, $l_S(s) = l_R(s) \cap S = 0$, so $l_R(s) = 0$ and $r = 0$. If $sr = 0$, then $(Ss)r = 0$ and $r \in Z_{(S)}R$. By Lemma 4.1, $Z_{(S)}R \subseteq Z_{(R)}R = 0$, so $r = 0$. Thus s is regular in R .

LEMMA 4.3. *Let S have a classical left quotient ring \bar{S} . If R is a left intrinsic extension of S where $Z_{(R)}R = 0$, then $\bar{S} \subseteq Q$ where Q is the maximal left quotient ring of R .*

Proof. Let M be the injective hull of R as a left R -module. By [1, Theorem 1, p. 69], $Q = \text{Hom}_R(M, M) \cong M$. If d is a regular

elements of S , define the map $\bar{f}: Rd \rightarrow R$ by $\bar{f}(rd) = r$ for all $r \in R$. The map is well defined by Lemma 4.2, and by the injectivity of M , there exists $f \in \text{Hom}_R(M, M)$ such that $f|_{Rd} = \bar{f}$. By [1, Theorem 1, p. 69], the canonical isomorphic image of d in Q is the unique $g \in \text{Hom}_R(M, M)$ such that $g(r) = rd$ for all $r \in R$. If 1 denotes the identity of Q , it follows that $R \subseteq \ker(1 - gf)$ and $Rd \subseteq \ker(1 - gf)$. R is left essential in Q , and it is easy to verify that Rd is also essential in Q . By [1, Theorem 1, p. 44], $1 - gf$ and $1 - fg$ are in the Jacobson radical of the semisimple ring Q . Hence $gf = fg = 1$. By the canonical injection of R into Q , we can consider R to be a subring of Q , and so d has a two-sided inverse f (henceforth denoted d^{-1}) in Q . Hence every regular element of S has a two-sided inverse in Q . If $T = \{a^{-1}b \mid b \in S, a \text{ regular in } S\}$, then $T \subseteq Q$ and T is a ring by Ore's condition for S , [see 4, p. 109]. Hence $\bar{S} = T \subset Q$.

LEMMA 4.4. *If S is a left self-injective ring and $Z({}_S S) = 0$, then every left intrinsic extension of S is a left strongly intrinsic extension of S .*

Proof. Let R be a left intrinsic extension of S . By [1, Theorem 1, p. 69], S is its own maximal left quotient ring and is a regular ring. If $A \in L(S)$, then by [1, Theorem 4, p. 70], $A = Se$ where $e^2 = e \in S$. Hence $Re \cap S = Se$, and R is a left strongly intrinsic extension of S .

THEOREM 4.5. *If R is an extension of a lqss ring S and $Z({}_R R) = 0$, then R is a left intrinsic extension of S if and only if R is a left strongly intrinsic extension of S .*

Proof. Let R be a left intrinsic extension of S and Q the maximal left quotient ring of R . By Lemma 4.3, $\bar{S} \subseteq Q$, and clearly Q is a left intrinsic extension of S . Also \bar{S} is left self-injective and $Z({}_S \bar{S}) = 0$, so by Lemma 4.4, Q is a left strongly intrinsic extension of the lqss ring \bar{S} . By Lemma 1.3, $L(Q) \cong L(\bar{S})$ under contraction, and since $L(Q) \cong L(R)$ and $L(S) \cong L(\bar{S})$ under contraction, it follows that $L(S) \cong L(R)$ under contraction. Hence R is a left strongly intrinsic extension of S .

THEOREM 4.6. *If R is a left intrinsic extension of a lqss ring S , then the following are equivalent:*

- (i) $Z({}_R R) = 0$,
- (ii) R is a left strongly intrinsic extension of S ,
- (iii) R is lqss.

Proof. (i) \Rightarrow (ii) by Theorem 4.5. (ii) \Rightarrow (iii) by the proof of Theorem 3.4. (iii) \Rightarrow (i) follows from [3, Theorem 4.4].

REFERENCES

1. C. Faith, *Lectures on injective modules and quotient rings*, Springer Verlag, New York, 1967.
2. C. Faith, and Y. Utumi, *Intrinsic extensions of rings*, Pacific J. Math. **14** (1964), 505-512.
3. R. E. Johnson, *Quotient rings with zero singular ideal*, Pacific J. Math. **11** (1961), 1385-1392.
4. J. Lambeck, *Lectures on rings and modules*, Blaisdell Publishing Company, Waltham, Mass., 1967.
5. L. Levy, *Unique subdirect sums of prime rings*, Trans. Amer. Math. Soc. **106** (1963), 67-76.
6. Y. Utumi, *On continuous regular rings and semisimple self-injective rings*, Canadian J. Math. **12** (1960), 597-605.

Received July 19, 1968. This paper is a portion of a dissertation written at The University of Kansas, under the direction of Professor P. J. McCarthy.

WASHINGTON STATE UNIVERSITY
PULLMAN, WASHINGTON 99163

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Willard Ellis Baxter, <i>Topological rings with property (Y)</i>	563
Sterling K. Berberian, <i>Note on some spectral inequalities of C. R. Putnam</i>	573
David Theodore Brown, <i>Galois theory for Banach algebras</i>	577
Dennis K. Burke and R. A. Stoltenberg, <i>A note on p-spaces and Moore spaces</i>	601
Rafael Van Severen Chacon and Stephen Allan McGrath, <i>Estimates of positive contractions</i>	609
Rene Felix Dennemeyer, <i>Conjugate surfaces for multiple integral problems in the calculus of variations</i>	621
Edwin O. Elliott, <i>Measures on countable product spaces</i>	639
John Moss Grover, <i>Covering groups of groups of Lie type</i>	645
Charles Lemuel Hagopian, <i>Concerning semi-local-connectedness and cutting in nonlocally connected continua</i>	657
Velmer B. Headley, <i>A monotonicity principle for eigenvalues</i>	663
John Joseph Hutchinson, <i>Intrinsic extensions of rings</i>	669
Harold H. Johnson, <i>Determination of hyperbolicity by partial prolongations</i>	679
Tilla Weinstein, <i>Holomorphic quadratic differentials on surfaces in E^3</i>	697
R. C. Lacher, <i>Cell-like mappings. I</i>	717
Roger McCann, <i>A classification of centers</i>	733
Curtis L. Outlaw, <i>Mean value iteration of nonexpansive mappings in a Banach space</i>	747
Allan C. Peterson, <i>Distribution of zeros of solutions of a fourth order differential equation</i>	751
Bhalchandra B. Phadke, <i>Polyhedron inequality and strict convexity</i>	765
Jack Wyndall Rogers Jr., <i>On universal tree-like continua</i>	771
Edgar Andrews Rutter, <i>Two characterizations of quasi-Frobenius rings</i>	777
G. Sankaranarayanan and C. Suyambulingom, <i>Some renewal theorems concerning a sequence of correlated random variables</i>	785
Joel E. Schneider, <i>A note on the theory of primes</i>	805
Richard Peter Stanley, <i>Zero square rings</i>	811
Edward D. Tymchatyn, <i>The 2-cell as a partially ordered space</i>	825
Craig A. Wood, <i>On general Z.P.I.-rings</i>	837