

Pacific Journal of Mathematics

A NOTE ON THE SIMILARITY OF MATRIX AND ITS
CONJUGATE TRANSPOSE

JOHN WALTER DUKE

A NOTE ON THE SIMILARITY OF A MATRIX AND ITS CONJUGATE TRANPOSE

JOHN W. DUKE

It is well-known that each square matrix A over a field is similar to its transpose A^T and there exists a nonsingular symmetric matrix P for which $PA^T = AP$. The purpose of this note is to show that if A is similar to its conjugate transpose A^* then, under certain conditions, there exists a nonsingular Hermitian matrix Q for which $QA^* = AQ$.

Let f be an automorphism of order 2 on a field F and let K be the fixed field of f . For each x in F , we denote $f(x)$ by \bar{x} . If $A = (a_{ij})$ is a matrix over F , let $A^* = (b_{ij})$ where $b_{ij} = \overline{a_{ji}}$. A matrix M is called Hermitian (skew-Hermitian) provided $M^* = M$ ($M^* = -M$).

Taussky and Zassenhaus [1] have shown that for each square matrix over a field, there exists a nonsingular symmetric matrix which transforms the given matrix into its transpose. Our main result is

THEOREM 1. *Suppose F is an infinite field whose characteristic is different from 2. If a matrix A over F is similar to A^* , there exists a nonsingular Hermitian matrix Q over F for which $QA^* = AQ$.*

We shall utilize the following lemmas in both of which z denotes an element of F which is not in K .

LEMMA 1. *Every element of F can be expressed uniquely in the form $a + bz$ where both a and b lie in K .*

Proof. If c belongs to F , it is clear that

$$c = c - (c - \bar{c})(z - \bar{z})^{-1}z + (c - \bar{c})(z - \bar{z})^{-1}z$$

since $z \neq \bar{z}$. This is the required form since both

$$a = c - (c - \bar{c})(z - \bar{z})^{-1}z$$

and

$$b = (c - \bar{c})(z - \bar{z})^{-1}$$

lie in K . The uniqueness of the expression follows from the fact that z does not belong to K .

LEMMA 2. *If $c = r + sz$ and $d = t + uz$ with r, s, t , and u in K and $c/\bar{c} = d/\bar{d}$, then $ru = st$.*

Lemma 2 implies that there exists a one-to-one correspondence between K and the set of all elements c/\bar{c} where $c = r + z$ and r ranges over K . If F is infinite, Lemma 1 implies that K is infinite.

Proof of Theorem 1. Suppose $PA^* = AP$ with P nonsingular. Since the characteristic of F is not 2, the matrix P can be expressed as the sum of an Hermitian matrix H and a skew-Hermitian matrix S . Hence $HA^* = AH$ and it remains to show that H may be chosen nonsingular.

Since $(cP)A^* = A(cP)$ for all c in F , we want to choose c so that $M = cP + \bar{c}P^*$ is nonsingular. The matrix M is nonsingular if and only if $-c/\bar{c}$ is distinct from all of the eigenvalues of $P^{-1}P^*$. Since there exist infinitely many values of $-c/\bar{c}$, an element c can be properly chosen and the proof is complete.

In regard to finite fields, we have

THEOREM 2. *Suppose A is a square matrix of order n over a field F whose characteristic is different from 2 and $PA^* = AP$ with P nonsingular. If there exists an element y in F such that y^m does not belong to K for $1 \leq m \leq n+1$, there exists a nonsingular Hermitian matrix Q for which $QA^* = AQ$.*

Proof. Utilizing the same decomposition of P as in the proof of Theorem 1, it is sufficient to show there exists an element c in F such that $cP + \bar{c}P^*$ is nonsingular. For c nonzero, $cP + \bar{c}P^*$ is nonsingular if and only if $-\bar{c}/c$ is not an eigenvalue of $P(P^*)^{-1}$. Hence let k_1, k_2, \dots, k_t be the distinct eigenvalues of $P(P^*)^{-1}$ in F and let

$$W = \{1, -k_1, -k_2, \dots, -k_t\}.$$

If for each nonzero x in F there exists k in W such that $\bar{x} = kx$, then k^r belongs to W for all positive integers r since $\bar{x}^r = k^r x^r$. In particular, for the element y mentioned in the hypothesis of the theorem, $\bar{y} = dy$ for some d in W and hence the elements d^i , for $1 \leq i \leq n+2$, all belong to W . Since W contains only $t+1$ elements and $0 \leq t \leq n$, it follows that $d^i = d^j$ for some integers i and j , $i < j$, between 1 and $n+2$, inclusively. Hence $j-i \leq n+1$ and $d^{j-i} = 1$ since $d \neq 0$. Therefore

$$f(y^{j-i}) = d^{j-i}y^{j-i} = y^{j-i}$$

implies y^{j-i} belongs to K . This contradiction shows the existence of

some c in F such that $\bar{c} \neq kc$ for all k in W : hence c does not belong to K and $cP + \bar{c}P^*$ is nonsingular as required.

As a simple application of Theorem 2, suppose $F = GF(p^{2s})$ with $p \neq 2$ and let $f(x) = x^{p^s}$ for all x in F . By considering a generator of the multiplicative group of F , one may verify the result for matrices over F of order less than p^s .

REFERENCE

1. Olga Taussky and Hans Zassenhaus, *On the similarity transformation between a matrix and its transpose*, Pacific J. Math. **9** (1959), 893-896.

Received January 3, 1969. This note is included in the author's doctoral dissertation prepared under the direction of Professor Burton W. Jones at the University of Colorado.

ANGELO STATE UNIVERSITY
SAN ANGELO, TEXAS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. 36, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 108 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Kokusai Bunko Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 31, No. 2

December, 1969

Efraim Pacillas Armendariz, <i>Quasi-injective modules and stable torsion classes</i>	277
J. Adrian (John) Bondy, <i>On Ulam's conjecture for separable graphs</i>	281
Vasily Cateforis and Francis Louis Sandomierski, <i>On commutative rings over which the singular submodule is a direct summand for every module</i>	289
Rafael Van Severen Chacon, <i>Approximation of transformations with continuous spectrum</i>	293
Raymond Frank Dickman and Alan Zame, <i>Functionally compact spaces</i>	303
Ronald George Douglas and Walter Rudin, <i>Approximation by inner functions</i>	313
John Walter Duke, <i>A note on the similarity of matrix and its conjugate transpose</i>	321
Micheal Neal Dyer and Allan John Sieradski, <i>Coverings of mapping spaces</i>	325
Donald Campbell Dykes, <i>Weakly hypercentral subgroups of finite groups</i>	337
Nancy Dykes, <i>Mappings and realcompact spaces</i>	347
Edmund H. Feller and Richard Laham Gantos, <i>Completely injective semigroups</i>	359
Irving Leonard Glicksberg, <i>Semi-square-summable Fourier-Stieltjes transforms</i>	367
Samuel Irving Goldberg and Kentaro Yano, <i>Integrability of almost cosymplectic structures</i>	373
Seymour Haber and Charles Freeman Osgood, <i>On the sum $\sum (n\alpha)^{-t}$ and numerical integration</i>	383
Sav Roman Harasymiv, <i>Dilations of rapidly decreasing functions</i>	395
William Leonard Harkness and R. Shantaram, <i>Convergence of a sequence of transformations of distribution functions</i>	403
Herbert Frederick Kreimer, Jr., <i>A note on the outer Galois theory of rings</i>	417
James Donald Kuelbs, <i>Abstract Wiener spaces and applications to analysis</i>	433
Roland Edwin Larson, <i>Minimal T_0-spaces and minimal T_D-spaces</i>	451
A. Meir and Ambikeshwar Sharma, <i>On Ilyeff's conjecture</i>	459
Isaac Namioka and Robert Ralph Phelps, <i>Tensor products of compact convex sets</i>	469
James L. Rovnyak, <i>On the theory of unbounded Toeplitz operators</i>	481
Benjamin L. Schwartz, <i>Infinite self-interchange graphs</i>	497
George Szeto, <i>On the Brauer splitting theorem</i>	505
Takayuki Tamura, <i>Semigroups satisfying identity $xy = f(x, y)$</i>	513
Kenneth Tolo, <i>Factorizable semigroups</i>	523
Mineko Watanabe, <i>On a boundary property of principal functions</i>	537
James Juei-Chin Yeh, <i>Singularity of Gaussian measures in function spaces with factorable covariance functions</i>	547