

Pacific Journal of Mathematics

RINGS OF ANALYTIC FUNCTIONS

JULIANNE SOUCHEK

RINGS OF ANALYTIC FUNCTIONS

JULIANNE SOUCHEK

If F is an open Riemann surface and $A(F)$ is the set of all analytic functions on F , then $A(F)$ is a ring under pointwise addition and multiplication. This paper is concerned with proper subrings R of $A(F)$ which are isomorphic images of $A(G)$, the ring of all analytic functions on an open Riemann surface G , under a homomorphism Φ which maps constant functions onto themselves. The ring R has the form $\{g \circ \phi: g \in A(G), \phi \text{ an analytic map from } F \text{ into } G\}$, and will be denoted R_ϕ . Relations between ϕ , R_ϕ and the spectrum of R_ϕ are given as necessary and sufficient conditions for the existence of a Riemann surface G such that R is isomorphic to $A(G)$.

Open Riemann surfaces will be denoted by F and G , the rings of all analytic functions on F and G with pointwise addition and multiplication will be denoted by $A(F)$ and $A(G)$, and Φ will denote a homomorphism from $A(G)$ into $A(F)$ which maps constant functions onto themselves. Let Φ be such a homomorphism. In [5, pp. 272–273] H. L. Royden shows there is an analytic mapping ϕ of F into G such that $\Phi(g) = g \circ \phi$, and that if Φ is an isomorphism onto $A(F)$ then ϕ is a one-to-one, onto analytic mapping. If ϕ is an analytic mapping of F into G , then Φ defined by $\Phi(g) = g \circ \phi$, $g \in A(G)$, is a homomorphism from $A(G)$ into $A(F)$ which preserves constant functions. When ϕ is one-to-one and onto, Φ is an isomorphism.

The image of $A(G)$ under Φ is the set $\{g \circ \phi: g \in A(G), \phi \text{ is an analytic map of } F \text{ into } G\}$ denoted by R_ϕ . R_ϕ is a subring of $A(F)$ and contains the constant functions, since $\Phi(\lambda) = \lambda$ for λ a constant function. The following conditions are equivalent: R_ϕ properly contains the constant functions, Φ is an isomorphism, ϕ is not a constant function. Theorems 1 and 2 give other relations between ϕ and R_ϕ .

THEOREM 1. *If R_ϕ properly contains the constant functions, then R_ϕ contains $1/f$ whenever $f \in R_\phi$, $f(z) \neq 0$ on F , if and only if ϕ maps F onto G .*

Proof. Let ϕ map F onto G , $f \in R_\phi$, $f(z) \neq 0$ on F . Then $f = \Phi h$ for some $h \in A(G)$ and $1/h \in A(G)$ if $h(y) \neq 0$ for $y \in G$. Suppose $h(a) = 0$. Since $a = \phi(z)$ for some $z \in F$, $0 = h(a) = h(\phi(z)) = \Phi h(z) = f(z)$. This contradicts $f(z) \neq 0$ on F . Thus $h(a) \neq 0$ for $a \in G$, $1/h \in A(G)$, and $1/f = \Phi(1/h) \in R_\phi$.

Suppose R_ϕ contains $1/f$ when $f \in R_\phi$, $f(z) \neq 0$ on F . Let $a \in G$.

There is $g \in A(G)$ such that $g(a) = 0$ and $g(w) \neq 0$ for $w \neq a$ [1, pp. 591-592]. The function $\Phi g \in R_\phi$. If $\Phi g(z) = g \circ \phi(z) \neq 0$ for $z \in F$, then there is $h \in R_\phi$ such that $(\Phi g)(h) = 1$. There is $k \in A(G)$ such that $h = \Phi k$. Then $(\Phi g)(\Phi k) = 1$ and $\Phi(gk) = 1$ but Φ is an isomorphism implies $gk = 1$ and $g(a)k(a) = 1$. This contradicts $g(a) = 0$. Therefore $g(\phi(z)) = 0$ and $\phi(z) = a$ for some $z \in F$.

A straightforward argument shows

THEOREM 2. *If R_ϕ properly contains the constant functions, then R_ϕ separates the points of F if and only if ϕ is one-to-one.*

Let R be a ring of analytic functions defined on F . The spectrum of R , ΣR , is the set of nonzero homomorphisms π from R into the complex numbers such that $\pi(\lambda) = \lambda$ for λ a constant function. For $x \in F$ the point evaluation mapping $\pi_x = \{(f, f(x)): f \in R\}$ is a homomorphism from R into the complex numbers, and $\pi_x(\lambda) = \lambda$ for λ a constant function. Therefore ΣR always contains the point evaluation mappings defined on R . In [5, p. 272] H. L. Royden shows that the spectrum of $A(F)$ is the set of point evaluation mappings π_x defined on $A(F)$, $x \in F$. For $f \in R$ let $\hat{f} = \{(\pi, \pi f): \pi \in \Sigma R\}$; \hat{f} is a function from ΣR into the complex numbers. Let \hat{R} denote $\{\hat{f}: f \in R\}$. With pointwise addition and multiplication \hat{R} is a ring containing the constant functions and is isomorphic to R under $f \rightarrow \hat{f}$.

For $y \in G$, let ψ_y denote an element of $\Sigma A(G)$. The mapping $P = \{(y, \psi_y): y \in G\}$ is a one-to-one function from G onto $\Sigma A(G)$. If $R = \Phi(A(G))$ and Φ is an isomorphism, $L = \{(\pi, \pi \circ \Phi): \pi \in \Sigma R\}$ is a one-to-one function from ΣR onto $\Sigma A(G)$. The mapping $\pi \rightarrow \pi \circ \Phi = \psi_y \rightarrow y$ which is $P^{-1} \circ L$ defines a one-to-one correspondence between ΣR and G when Φ is an isomorphism.

THEOREM 3. *Let $R_\phi = \Phi(A(G))$, Φ be an isomorphism from $A(G)$ into $A(F)$ which preserves constant functions. Let M be the function from $\Sigma A(F)$ into ΣR_ϕ defined by $M(\pi_x) = \pi_x|_{R_\phi}$. Then M is onto if and only if ϕ is onto, and M is one-to-one if and only if ϕ is one-to-one.*

Proof. The proof that M is one-to-one if and only if ϕ is one-to-one follows from Theorem 2 and the fact that $A(F)$ separates the points of F .

Let $\pi \in \Sigma R_\phi$. Then $\pi \circ \Phi \in \Sigma A(G)$ implies there is $y \in G$ such that $\pi \circ \Phi = \psi_y$, where $\psi_y(g) = g(y)$ for $g \in A(G)$. There are two cases: $y \in \phi(F)$, $y \notin \phi(F)$. If $y \in \phi(F)$, then $y = \phi(x)$ for some $x \in F$ and $\pi(\Phi g) = g(y) = g(\phi(x)) = \Phi g(x)$ for every $g \in A(G)$, $\pi(\Phi g) = \Phi g(x)$ for

every $f = \Phi g \in R_\phi$. This implies $\pi = M(\pi_x)$. If $y \notin \phi(F)$, then $y \neq \phi(x)$ for $x \in F$, and it may be shown that for every $x \in F$ there is $f \in R_\phi$ such that $\pi(f) \neq f(x)$. Let $x \in F$. Then $\phi(x) \in G$. $y \in G$, $y \neq \phi(x)$, and $A(G)$ separates the points of G implies there is a $g \in A(G)$ such that $g(y) \neq g(\phi(x))$. From $\Phi(g) \in R_\phi$ and $\pi(\Phi g) = g(y) \neq g(\phi(x)) = \Phi g(x)$ it follows that $\pi \neq M(\pi_x) = \pi_x|_{R_\phi}$.

For $\pi \in \Sigma R_\phi$, $\pi \circ \Phi = \psi_y \in \Sigma A(G)$, and it has been shown $\pi \in M(\Sigma A(F))$ if and only if $y \in \phi(F)$.

From Theorem 3 and since ΣR_ϕ and G are in one-to-one correspondence, it follows that the point evaluation maps in ΣR_ϕ are in one-to-one correspondence with the points $\phi(x) \in \phi(F)$, and the elements of ΣR_ϕ which are not point evaluation maps are in one-to-one correspondence with the points in $G - \phi(F)$.

Theorem 4 contains a necessary condition which a subring R of $A(F)$ must satisfy if R is to be $\Phi(A(G))$, the isomorphic image of $A(G)$ under Φ for some open Riemann surface G . The corollary to Theorem 5 gives a set of sufficient conditions on R in order that R be $\Phi(A(G))$ when $\Phi g = g \circ \phi$ and $\phi: F \rightarrow G$ is an onto mapping.

Suppose F is an open Riemann surface, $p \in F$, f is analytic at p and τ is a local uniformizer which maps a neighborhood of p onto $\{z: |z| < \rho\}$ for some $\rho > 0$, $\tau(p) = 0$. There is a number $r > 0$ such that $f \circ \tau^{-1}(z) = \sum_{i=0}^{\infty} a_i z^i$ for $|z| < r$. The multiplicity of f at p is defined as $\inf \{k: k \neq 0 \text{ and } a_k \neq 0\}$, denoted $n(p; f)$. The multiplicity $n(p; f)$ of f at p does not depend on τ . If R contains functions other than constants, $m = \inf \{n(p; f): f \in R\}$ is defined, and $n(p; f) = m$ for some $f \in R$.

THEOREM 4. *Let $p \in F$, R_ϕ contain functions other than constants and let $m = \{\inf n(p; f): f \in R_\phi\}$. There is a local uniformizer τ at p with the properties: $\tau(0) = p$, for some $\rho > 0$, τ maps $\{z: |z| < \rho\}$ onto a neighborhood of p , and if $f \in R_\phi$, $f \circ \tau(z) = \sum_{i=0}^{\infty} a_i (z^m)^i$ for $|z| < \rho$.*

The proof of Theorem 4 is based on two lemmas:

LEMMA 1. *If $p \in F$, $m = \inf \{n(p; f): f \in R_\phi\}$ and $f \in R_\phi$, then $n(p; f) = km$, where k is a positive integer.*

LEMMA 2. *Given $\sum_{i=m}^{\infty} c_i z^i$ convergent for $|z| < \rho$, $c_m \neq 0$, $m \neq 0$, there is $\sum_{i=1}^{\infty} b_i z^i$ convergent for $|z| < \rho$, $b_1 \neq 0$, such that $(\sum_{i=1}^{\infty} b_i z^i)^m = \sum_{i=m}^{\infty} c_i z^i$.*

Lemma 1 follows from the two relations: For $f \in R_\phi$, $f = g \circ \phi$ for

some $g \in A(G)$, which implies $n(p; f) = (n(p; \phi))(n(\phi(p); g))$, and if $m = \inf \{n(p; f): f \in R_\phi\}$ then $n(p; \phi) = m$. Lemma 2 is proved by defining W a subset of the natural numbers N as $W = \{n \in N: b_1, b_2, \dots, b_n \text{ can be defined in such a way that the coefficients of } z^i \text{ for } 1 \leq m \leq i \leq m + n - 1 \text{ of } (\sum_{i=1}^{\infty} b_i z^i)^m \text{ and } \sum_{i=m}^{\infty} c_i z^i \text{ are equal}\}$ and using induction to show $W = N$.

Proof of Theorem 4. Let τ_p be a local uniformizer about p such that $\tau_p(0) = p$. If $m = \inf \{n(p; f): f \in R_\phi\}$, there is $f_p \in R_\phi$ and $\rho > 0$ such that $f_p \circ \tau_p(z) = \sum_{i=m}^{\infty} c_i z^i$ for $|z| < \rho$, $c_m \neq 0$, and the range of $\sum_{i=m}^{\infty} c_i z^i$ contains $|z| < \rho^m$.

There is a power series $\sum_{i=1}^{\infty} b_i z^i$, $b_1^m = c_m$, such that $\sum_{i=m}^{\infty} c_i z^i = (\sum_{i=1}^{\infty} b_i z^i)^m$ for $|z| < \rho$ as stated in Lemma 2. $k(z) = \sum_{i=1}^{\infty} b_i z^i$ is defined for $|z| < \rho$, is one-to-one, and its range contains $|z| < \rho$. Thus $k^{-1}(y)$ is defined for $|y| < \rho$ and $f_p \circ \tau_p \circ k^{-1}(z) = (\sum_{i=1}^{\infty} b_i (k^{-1}(z))^i)^m = z^m$ for $|z| < \rho$, $\tau_p \circ k^{-1}(0) = p$. The function $\tau = \tau_p \circ k^{-1}$ is a local uniformizer about p and there is $f_p \in R_\phi$ such that $f_p \circ \tau(z) = z^m$ for $|z| < \rho$.

Let $f \in R_\phi$, f not a constant function. Then $f \circ \tau(z) = \sum_{i=0}^{\infty} a_i z^i$ for $|z| < \rho$. Let N denote the natural numbers and define $W = \{n \in N: f \circ \tau(z) = \sum_{i=0}^n a_{m j_i} z^{m j_i} + z^{m j_n} h_n(z), \text{ where } h_n(z) = \sum_{i=1}^{\infty} b_{n, i} z^i \text{ and } j_i \text{ are nonnegative integers, } 0 = j_0 < j_1 < \dots < j_n\}$.

It follows from Lemma 1 that for $|z| < \rho$, $f \circ \tau(z) = \sum_{i=0}^{\infty} a_i z^i = a_0 + a_{m j_1} z^{m j_1} + z^{m j_1} h_1(z)$, where $h_1(0) = 0$. If $k \in W$, then $f \circ \tau(z) = \sum_{i=0}^k a_{m j_i} z^{m j_i} + z^{m j_k} h_k(z)$, $h_k(0) = 0$. Since $f \in R_\phi$, $z^m \in R_\phi$ and constants are contained in R_ϕ , $z^{m j_k} h_k(z) = f(z) - \sum_{i=0}^k a_{m j_i} z^{m j_i} \in R_\phi$. If $h_k \neq 0$, $n(p; z^{m j_k} h_k) = m j_{k+1}$ and $f \circ \tau(z) = \sum_{i=0}^{k+1} a_{m j_i} z^{m j_i} + z^{m j_{k+1}} h_{k+1}(z)$, where $h_{k+1}(z) = \sum_{i=1}^{\infty} b_{k+1, i} z^i$ on $|z| < \rho$ and $j_{k+1} > j_k$. If $h_k = 0$, then the above statement is true with $a_{m j_{k+1}} = 0$, $h_{k+1} = 0$. By induction $W = N$ and $f \circ \tau(z) = \sum_{i=0}^{\infty} a_{m i} z^{m i}$ on $|z| < \rho$.

If R , a subring of $A(F)$, has the property that for every $a \in F$, $f \in R$, for some local uniformizer τ about a , $f \circ \tau(z) = \sum_{i=0}^{\infty} a_i (z^{m(a)})^i$ for $m(a) = \inf \{n(a; f): f \in R\}$, then R has property (ξ) . If R contains functions other than constants and has property (ξ) , then for $a \in F$, $m(a) = \inf \{n(a; f): f \in R\} = 1$ if R separates the points of F .

THEOREM 5. *If R is a subring of $A(F)$ which contains functions other than constants and has property (ξ) , then there is an open Riemann surface G , an analytic mapping ϕ of F onto G , and a separating subring S of $A(G)$ such that S is isomorphic to R under $\hat{f} \mapsto \hat{f} \circ \phi$, $\hat{f} \in S$.¹*

Proof. Let $G = \{\pi_p: p \in F\}$ where $\pi_p = \{(f, f(p)): f \in R\}$ and $\phi =$

$\{(p, \pi_p): p \in F\}$. The topology on G will be that which makes ϕ continuous and open. If N_p is an open neighborhood of $p \in F$, then $N_{\pi_p} = \{\pi_q: q \in N_p\}$ is an open neighborhood of π_p . The set G with this topology is a connected Hausdorff space.

Let $p \in F$, $\pi_p \in G$ and $m = \inf \{n(p; f): f \in R\}$. By the same argument used in the beginning of the proof of Theorem 4, there is a function $f_p \in R$ and a local uniformizer τ about p such that $\tau(0) = p$ and $f_p \circ \tau(z) = z^m$ for $|z| < \rho^{1/m}$ for some $\rho > 0$. Then for $f \in R$, $f \circ \tau(z) = \sum_{i=0}^{\infty} a_i(z^m)^i = g_f(z^m)$ for $|z| < \rho^{1/m}$, g_f analytic on $|z| < \rho$.

It will be shown that $\sigma_\tau = \{(z^m, \pi_{\tau(z)}): |z| < \rho^{1/m}\}$ is a local uniformizer about π_p . If $z_1^m = z_2^m$, then $f \circ \tau(z_1) = g_f(z_1^m) = g_f(z_2^m) = f \circ \tau(z_2)$, for $f \in R$ implies $\pi_{\tau(z_1)} = \pi_{\tau(z_2)}$, which implies that σ_τ is a function. If $\pi_{\tau(z_1)} = \pi_{\tau(z_2)}$ then in particular $f_p \circ \tau(z_1) = f_p \circ \tau(z_2)$, which implies $z_1^m = z_2^m$, and σ_τ is one-to-one. Since the relations $z^m \rightarrow z \rightarrow \tau(z) \rightarrow \phi(\tau(z)) = \pi_{\tau(z)}$ are open and continuous, σ_τ is open and continuous. Thus σ_τ is a homeomorphism from $\{w: |w| < \rho\}$ onto $\phi \circ \tau(\{z: |z| < \rho^{1/m}\}) = N_{\pi_p}$.

If $\pi \in W = \sigma_{\tau_2}(|z| < \rho_2) \cap \sigma_{\tau_1}(|z| < \rho_1)$, there are points z_1, z_2 such that $\pi_{\tau_1}(z_1) = \pi_{\tau_2}(z_2)$. Then $f \circ \tau_1(z_1) = f \circ \tau_2(z_2)$ for every $f \in R$, and $z_1^{m_1} = f_1(\tau_1(z_1)) = f_1(\tau_2(z_2)) = g_{f_1}(z_2^{m_2})$, so g_{f_1} is analytic on $\{w: |w| < \rho_2\}$, which contains $\sigma_{\tau_2}^{-1}(W)$. This shows that $z_1^{m_1} = \sigma_{\tau_1}^{-1} \circ \sigma_{\tau_2}(z_2^{m_2})$ is analytic on $\sigma_{\tau_2}^{-1}(W)$ to $\sigma_{\tau_1}^{-1}(W)$. The function σ_τ is a local uniformizer of a neighborhood of π_p , and G is a Riemann surface.

For $f \in R$, let $\hat{f} = \{(\pi_p, f(p)): p \in F\}$, $S = \{\hat{f}: f \in R\}$. Since f is continuous and ϕ is open, \hat{f} is continuous. The function \hat{f} is analytic at π_p , because if $|w| < \rho$, $w = z^m$, then $\hat{f} \circ \sigma_\tau(w) = \hat{f}(\pi_{\tau(z)}) = f(\tau(z)) = \sum_{i=0}^{\infty} a_i(z^m)^i = \sum_{i=0}^{\infty} a_i w^i$. The mapping ϕ is analytic at p , because $\sigma_\tau^{-1} \circ \phi \circ \tau(z) = \sigma_\tau^{-1}(\pi_{\tau(z)}) = z^m$ for $|z| < \rho^{1/m}$. With pointwise addition and multiplication, S is a ring and is isomorphic to R under the mapping $\hat{f} \rightarrow \hat{f} \circ \phi = f$. The ring S separates the points of G . Since S contains functions which are not constant and are analytic on G , G is an open Riemann surface.

If S is to be $A(G)$, then by Theorem 3 the mapping $M(\pi_p) = \pi_p|_R$ from $\Sigma A(F)$ to ΣR must be onto, since ϕ is an onto mapping of F to G . Thus ΣR may contain only point evaluation mappings and $\Sigma R = G$.

COROLLARY TO THEOREM 5. *If R is a subring of $A(F)$ which properly contains the constant functions and has property (ξ) , if ΣR contains only point evaluation mappings, and R contains all $f \in A(F)$ such that $f \circ \tau_p(z) = \sum_{i=0}^{\infty} a_i(z^m)^i$ for $|z| < \rho^{1/m}$, $p \in F$, $m = \inf \{n(p; f): f \in R\}$, then $\Sigma R = G$ is an open Riemann surface, and R is isomorphic to $S = A(G)$.*

¹ This result and proof are similar to one given by M. Heins for a subfield of the field of all meromorphic functions on a Riemann surface [2, pp. 268-269].

Proof. Everything except $S = A(G)$ was shown in the proof of Theorem 5. The function $\hat{f} \in A(G)$ if and only if for every $\pi_p \in G$, $\hat{f} \circ \sigma_{\tau_p}(w) = \sum_{i=0}^{\infty} a_i w^i$ for $|w| < \rho$. Let $\hat{f} \in A(G)$, $p \in F$, $\pi_p \in G$, and $f = \hat{f} \circ \phi$. Then $f \in A(F)$ and $f \in R$, because for $|z| < \rho^{1/m}$, $f \circ \tau_p(z) = \hat{f} \circ \phi(\tau_p(z)) = \hat{f}(\pi_{\tau_p(z)}) = \hat{f} \circ \sigma_{\tau_p}(z^m) = \sum_{i=0}^{\infty} a_i (z^m)^i$.

If $R = \{\hat{f} \circ \phi: \hat{f} \in S\}$ and S separates the points of G , then R separates the points of F if and only if ϕ is a one-to-one function. If S separates the points of G , and $S = A(G)$, then R may not separate the points of F , because if it did ϕ would be a one-to-one, onto analytic function from F to G , and $R = A(F)$. If $S \neq A(G)$ there may be a surface H , a mapping ϕ_1 and a separating subring T of $A(H)$ such that ϕ_1 is analytic and one-to-one but not onto, and $T = A(H)$.

In this part of the paper it is noted that if $R = \Phi(A(G))$, then ΣR with the Gelfand topology is an open Riemann surface, and \hat{R} which is isomorphic to R , is the ring of all analytic functions on ΣR . Theorem 8 gives sufficient conditions on a subring R of $A(F)$ and on \hat{R} in order that ΣR be an open Riemann surface and \hat{R} be a ring of analytic functions on ΣR . In conclusion sufficient conditions for \hat{R} to be $A(\Sigma R)$ are given.

If R is a ring of complex valued functions on F , then the Gelfand topology on ΣR is the weakest topology on ΣR which makes each element of \hat{R} continuous, where $\hat{R} = \{\hat{f}: f \in R\}$, $\hat{f} = \{(\pi, \pi f): \pi \in \Sigma R\}$. Let $\pi_0 \in \Sigma R$, K be a finite subset of \hat{R} , $\varepsilon > 0$. An open neighborhood of π_0 will be $\{\pi \in \Sigma R: |\hat{f}(\pi) - \hat{f}(\pi_0)| < \varepsilon \text{ for } \hat{f} \in K\}$. If $R = \Phi(A(G))$ and Φ is an isomorphism, then ΣR and $\Sigma A(G)$ with the Gelfand topology are homeomorphic under the mapping $L(\pi) = \pi \circ \Phi$ from ΣR onto $\Sigma A(G)$. The mapping $P(y) = \psi_y$ from G onto $\Sigma A(G)$ with the Gelfand topology is one-to-one, onto and continuous. The mapping P is also open. As Royden observes [4, pp. 287-288], this is a consequence of a theorem of Remmert that an open Riemann surface can be mapped one-to-one and holomorphically into C^3 [3, p. 118]. Thus $P^{-1} \circ L$ is a homeomorphism from ΣR with the Gelfand topology onto G .

THEOREM 6. *If R is a subring of $A(F)$ such that $R = \Phi(A(G))$, and if Φ is an isomorphism which preserves constant functions, then ΣR with the Gelfand topology is an open Riemann surface, and \hat{R} is the ring of all analytic functions on ΣR . Moreover \hat{R} is isomorphic to R .*

Proof. The spectrum of R with the Gelfand topology is a Hausdorff space. It is homeomorphic to G under the mapping $L^{-1} \circ P$,

and is connected. Let $\pi_q \in \Sigma R$ where $q \in G$, $\psi_q \in \Sigma A(G)$, and $L^{-1} \circ P$ maps $q \rightarrow \psi_q \rightarrow \pi_q$. If N_q is a neighborhood of q then $N_{\pi_q} = L^{-1} \circ P(N_q)$ is a neighborhood of π_q . There exists $h_q \in A(G)$ which has a simple zero at q [1, pp. 591-592]. h_q is a local uniformizer on a neighborhood of q , $N_q = h_q^{-1}(|z| < \rho)$ for some $\rho > 0$. If $\sigma_q = h_q|_{N_q}$, then $h_q \circ \sigma_q^{-1}(z) = z$ for $|z| < \rho$. For $h \in A(G)$, $y \in N_q$, $h(y) = \sum_{i=0}^{\infty} a_i(h_q(y))^i$.

If $f_q = \phi h_q$ then \hat{f}_q is a local uniformizer on $N_{\pi_q} = L^{-1} \circ P(N_q)$. From $\hat{f}_q(\pi_y) = h_q(y)$ follows $\hat{f}_q(\pi_y) = h_q \circ P^{-1} \circ L(\pi_y)$, $\pi_y \in N_{\pi_q}$, which implies \hat{f}_q is a homeomorphism of N_{π_q} onto $|z| < \rho$. If $\pi_y \in N_{\pi_{q_1}} \cap N_{\pi_{q_2}}$, then $\hat{f}_{q_1}(\pi_y) = h_{q_1}(y) = \sum_{i=0}^{\infty} a_i(h_{q_2}(y))^i = \sum_{i=0}^{\infty} a_i(\hat{f}_{q_2}(\pi_y))^i$ since $\pi_y \in N_{\pi_{q_2}}$ or $y \in N_{q_2}$. The function \hat{f}_q is a local uniformizer on N_{π_q} and ΣR is a Riemann surface.

The ring \hat{R} is contained in $A(\Sigma R)$, because if $\hat{f} \in \hat{R}$, $\pi_y \in N_{\pi_q}$, $z = \hat{f}_q(\pi_y)$, then $\hat{f} \circ \hat{f}_q^{-1}(z) = \hat{f}(\pi_y) = h(y) = \sum_{i=0}^{\infty} a_i(h_q(y))^i = \sum_{i=0}^{\infty} a_i(\hat{f}_q(\pi_y))^i = \sum_{i=0}^{\infty} a_i z^i$. The function $T(q) = \pi_q$ is an analytic map of G onto ΣR . If θ is analytic on ΣR , then $\theta \circ T \in A(G)$ and $\theta \in \hat{R}$ because $\theta(\pi_q) = \theta \circ T(q) = \psi_q(\theta \circ T) = \pi_q(f)$ for $f = \Phi(\theta \circ T)$. This implies $\theta = \hat{f}$. Thus $\hat{R} = A(\Sigma R)$. Since \hat{R} contains functions which are analytic and are not constant on ΣR , ΣR is an open Riemann surface.

THEOREM 7. *Let $R = \Phi(A(G))$. If $\hat{\pi} \in \Sigma R$, then $\hat{\pi}^{-1}(0)$ is a principal maximal ideal of R , and every principal maximal ideal of R is the kernel of $\pi \in \Sigma R$. If $\hat{\pi}^{-1}(0)$ is generated by f , then \hat{f} is a local homeomorphism on a neighborhood $N_{\hat{\pi}}$ of $\hat{\pi}$ and if $\pi \in N_{\hat{\pi}}$, $\hat{k} \in \hat{R}$, then $\hat{k}(\pi) = \sum_{i=0}^{\infty} a_i(\hat{f}(\pi))^i$.*

Proof. If $\hat{\pi} \in \Sigma R$, then $\hat{\pi} \circ \Phi = \psi_q \in \Sigma A(G)$ and $\hat{\pi}^{-1}(0) = \Phi(\psi_q^{-1}(0))$. The kernel of ψ_q , $M_q = \psi_q^{-1}(0)$, is a principal maximal ideal of $A(G)$, and every principal maximal ideal of $A(G)$ is a kernel of $\psi \in \Sigma A(G)$ [5, pp. 271-272]. If h generates M_q , then h has a single zero and it is a simple zero at q [5]. Thus h is a homeomorphism on a neighborhood of q , N_q . If $f = \phi h$, then $\hat{\pi}^{-1}(0)$ is the ideal generated by f . Also \hat{f} is a uniformizer on $N_{\hat{\pi}} = L^{-1} \circ P(N_q)$, and if $\pi \in N_{\hat{\pi}}$, $\hat{k} \in \hat{R}$, then $\hat{k}(\pi) = \sum_{i=0}^{\infty} a_i(\hat{f}(\pi))^i$ as shown in the proof of Theorem 6.

LEMMA. *Let S be a ring of continuous functions on X with identity. Then X is not connected if and only if S is contained in a ring Q of continuous functions on X , where $Q = I_1 + I_2$, I_1, I_2 proper ideals of Q , $I_1 \cap I_2 = \{0\}$.*

THEOREM 8. *Let R be a subring of $A(F)$ which properly contains the constant functions, and suppose \hat{R} is not contained in a ring Q of continuous functions on ΣR where $Q = I_1 + I_2$, I_1, I_2 proper ideals of Q , $I_1 \cap I_2 = \{0\}$. If for $\hat{\pi} \in \Sigma R$, $\hat{\pi}^{-1}(0)$ is a principal ideal of R*

generated by f and \hat{f} , the function in \hat{R} which corresponds to f in R , is a homeomorphism on a neighborhood of $\hat{\pi}$, and for π in this neighborhood, $g \in R$, $\pi g = \sum_{i=0}^{\infty} a_i(\pi f)^i$, then ΣR is an open Riemann surface and \hat{R} is a ring of analytic functions on ΣR .

Proof. The spectrum of R with the Gelfand topology is a Hausdorff space. By the lemma ΣR is connected. Let $\hat{\pi} \in \Sigma R$. There is \hat{f} a homeomorphism of $N_{\hat{\pi}}$ onto $|z| < \rho$ for some $\rho > 0$. If $\pi \in N_{\hat{\pi}}$, $g \in R$, then $\hat{g}(\pi) = \sum_{i=0}^{\infty} a_i(\hat{f}(\pi))^i$. If $\pi \in N_{\pi_1} \cap N_{\pi_2} = W$ then $\hat{f}_1 \circ \hat{f}_2^{-1}(\hat{f}_2(\pi)) = \hat{f}_1(\pi) = \sum_{i=0}^{\infty} a_i(\hat{f}_2(\pi))^i$ implies $\hat{f}_1 \circ \hat{f}_2^{-1}$ is analytic on $\hat{f}_2(W)$. $\{(N_{\pi}, \hat{f}_{\pi}): \pi \in \Sigma R\}$ defines an analytic structure on ΣR . It is immediate that $\hat{R} \subset A(\Sigma R)$. Since \hat{R} contains functions which are not constant and are analytic on ΣR , ΣR is an open Riemann surface.

If $\{R_n\}$ is a sequence of subrings of $A(F)$ such that R_n satisfies the conditions of Theorem 8, $\Sigma R_n|_{R_1} = \Sigma R_1$, $R_{n-1} \subset R_n$, then the chain has a maximal element, $\{\hat{f} \circ \phi: \hat{f} \in A(\Sigma R_1) \text{ and } \phi(x) = \pi_x, x \in F\}$. Let $\hat{\pi} \in \Sigma R_1$ and \hat{f} be a local homeomorphism at $\hat{\pi}$. If R_1 satisfies the conditions of Theorem 8 and contains all functions g in $A(F)$ such that $\hat{g}(\pi) = \sum_{i=0}^{\infty} a_i(\hat{f}(\pi))^i$ for $\pi \in N_{\hat{\pi}}$, π and $\hat{\pi}$ elements of ΣR_1 , then $\hat{R}_1 = A(\Sigma R_1)$, because if $\hat{g} \notin \hat{R}_1$, then there is $\hat{\pi} \in \Sigma R_1$ such that $\hat{g} \circ \hat{f}^{-1}$ is not analytic on $\{z: |z| < \rho\}$ which implies $\hat{g} \notin A(\Sigma R_1)$.

The author wishes to thank Professors H. B. Curtis and H. E. Lacey for their suggestions and encouragement.

REFERENCES

1. H. Behnke, and F. Sommer, *Theorie der analytischen Funktionen einer komplexen Veränderlichen*, Springer-Verlag, Berlin, 1962.
2. M. Heins, *Algebraic structure and conformal mapping*, Trans. Amer. Math. Soc. **89** (1958), 267-276.
3. R. Remmert, *Sur les espaces analytiques holomorphiquement separables et holomorphiquement convexes*, C. R. Acad. Sci. Paris **243** (1956), 118-121.
4. H. Royden, *Function algebras*, Bull. Amer. Math. Soc. **69** (1963), 281-298.
5. ———, *Rings of analytic and meromorphic functions*, Trans. Amer. Math. Soc. **83** (1956), 269-276.

Received July 24, 1968, and in revised form October 14, 1969. This paper is part of the author's Ph. D. thesis submitted to the University of Texas at Austin.

SAM HOUSTON STATE UNIVERSITY
HUNTSVILLE, TEXAS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mir Maswood Ali, <i>On some extremal simplexes</i>	1
Silvio Aurora, <i>On normed rings with monotone multiplication</i>	15
Silvio Aurora, <i>Normed fields which extend normed rings of integers</i>	21
John Kelly Beem, <i>Indefinite Minkowski spaces</i>	29
T. F. Bridgland, <i>Trajectory integrals of set valued functions</i>	43
Robert Jay Buck, <i>A generalized Hausdorff dimension for functions and sets</i>	69
Vlastimil B. Dlab, <i>A characterization of perfect rings</i>	79
Edward Richard Fadell, <i>Some examples in fixed point theory</i>	89
Michael Benton Freeman, <i>Tangential Cauchy-Riemann equations and uniform approximation</i>	101
Barry J. Gardner, <i>Torsion classes and pure subgroups</i>	109
Vinod B. Goyal, <i>Bounds for the solution of a certain class of nonlinear partial differential equations</i>	117
Fu Cheng Hsiang, <i>On $C, 1$ summability factors of Fourier series at a given point</i>	139
Lawrence Stanislaus Husch, Jr., <i>Homotopy groups of PL-embedding spaces</i>	149
Daniel Ralph Lewis, <i>Integration with respect to vector measures</i>	157
Marion-Josephine Lim, <i>$\mathcal{L} - 2$ subspaces of Grassmann product spaces</i>	167
Stephen J. Pierce, <i>Orthogonal groups of positive definite multilinear functionals</i>	183
W. J. Pugh and S. M. Shah, <i>On the growth of entire functions of bounded index</i>	191
Siddani Bhaskara Rao and Ayyagari Ramachandra Rao, <i>Existence of triconnected graphs with prescribed degrees</i>	203
Ralph Tyrrell Rockafellar, <i>On the maximal monotonicity of subdifferential mappings</i>	209
R. Shantaram, <i>Convergence of a sequence of transformations of distribution functions. II</i>	217
Julianne Soucek, <i>Rings of analytic functions</i>	233
Ted Joe Suffridge, <i>The principle of subordination applied to functions of several variables</i>	241
Wei-lung Ting, <i>On secondary characteristic classes in cobordism theory</i>	249
Pak-Ken Wong, <i>Continuous complementors on B^*-algebras</i>	255
Miyuki Yamada, <i>On a regular semigroup in which the idempotents form a band</i>	261