

# Pacific Journal of Mathematics

**APPROXIMATE UNITS AND MAXIMAL ABELIAN  
 $C^*$ -SUBALGEBRAS**

CHARLES A. AKEMANN

# APPROXIMATE UNITS AND MAXIMAL ABELIAN $C^*$ -SUBALGEBRAS

CHARLES A. AKEMANN

In recent years it has become clear that the study of  $C^*$ -algebras without a unit element is more than just a mildly interesting extension of the "typical" case of a  $C^*$ -algebra with unit. A number of important examples of  $C^*$ -algebras rarely have a unit, for example the group  $C^*$ -algebras and algebras of the form  $I \cap I^*$  where  $I$  is a closed left ideal of a  $C^*$ -algebra. J. Dixmier's book, *Les  $C^*$ -algebras et leurs representations*, carries through all the basic theory of  $C^*$ -algebras for the no-unit case, and his main tool is the approximate identity which such algebras have. Many  $C^*$ -algebra questions can be answered for a  $C^*$ -algebra without unit by embedding such an algebra in a  $C^*$ -algebra with unit. Some problems, especially those which involve approximate units, are not susceptible to this approach. This paper will study some problems of this type.

Theorem 1.1 states that if  $\mathcal{U}$  is a norm separable  $C^*$ -algebra and  $\{f_1, \dots, f_n\}$  is a finite set of orthogonal pure states of  $\mathcal{U}$  (i.e.,  $\|f_i - f_j\| = 2$  if  $i \neq j$ ), then there exists a maximal abelian  $C^*$ -subalgebra  $A$  of  $\mathcal{U}$  such that  $f_k|_A$  is pure ( $k = 1, \dots, n$ ) and  $f_k|_A$  has unique pure state extension to  $\mathcal{U}$  ( $k = 1, \dots, n$ ). This extends the prototype result of Aarnes and Kadison by (a) allowing a finite number of pure states instead of just one, (b) dropping the assumption that  $1 \in \mathcal{U}$ , and (c) proving uniqueness of the pure state extension. In §2 two examples are constructed which show that the uniqueness assertion of Theorem 1.1 cannot be extended to the nonseparable case, and that even in the separable case the subalgebra  $A$  must be carefully chosen to insure uniqueness of pure state extension. Theorem 1.2 and Example 2.3 show that a very desirable majorization property of approximate units does not quite carry over from the abelian case to the general case. (If it did, several important problems, including the Stone-Weierstrass problem would have been solved.) Theorem 1.3 extends the author's characterization of approximate units of  $C^*$ -algebras to approximate right units for left ideals of  $C^*$ -algebras.

## 1. Theorems.

**THEOREM 1.1** *Let  $\mathcal{U}$  be a norm separable  $C^*$ -algebra and  $\{f_1, \dots, f_n\}$  a finite set of orthogonal pure states (i.e.,  $\|f_j - f_i\| = 2$  if  $i \neq j$ ) of  $\mathcal{U}$ . Then there exists a maximal abelian  $C^*$ -subalgebra*

$A \subset \mathcal{U}$  such that  $f_k|_A$  is pure ( $k = 1, \dots, n$ ), and  $f_k$  is the unique pure state of  $\mathcal{U}$  which extends  $f_k|_A$  ( $k = 1, \dots, n$ ).

*Proof.* We shall consider  $\mathcal{U}$  acting on  $H$  under its universal representation (the direct sum of all the cyclic representations due to positive linear functionals on  $\mathcal{U}$  [5, p. 43]). Recall [5, p. 236] that the weak closure of  $\mathcal{U}$  can be identified with the second dual  $\mathcal{U}^{**}$  of  $\mathcal{U}$ . Let  $x_k \in H_{f_k} \subset H$  be a unit vector such that for any  $a \in \mathcal{U}$  we have  $\langle ax_k, x_k \rangle = f_k(a)$ . By [6, Corollary 7] we may choose a positive operator  $b \in \mathcal{U}$  with  $\|b\| \leq 1$  and  $bx_k = (k/n)x_k$  for  $k = 1, \dots, n$ . Choose real-valued functions  $\{\varphi_k\}_{k=1}^n$  of a real variable with the following properties: (1)  $\varphi_k(t)\varphi_j(t) = 0$  if  $k \neq j$ , (2)  $\varphi_k(k/n) = 1$ , (3)  $0 \leq \varphi_k(t) \leq 1$ , (4)  $\varphi_k$  is continuous; where  $t$  is any real number and  $k, j = 1, \dots, n$ .

Define  $b_k = \varphi_k(b)$  by the spectral theorem. Let  $p_k$  be the range projection of  $b_k$ , and let  $p$  be the orthogonal projection on the subspace spanned by  $\{x_1, \dots, x_n\}$ . Using the terminology of [3] we see that  $(1 - p)$  and  $\{p_k\}_{k=1}^n$  are open projections for  $\mathcal{U}$ , and hence for  $\tilde{\mathcal{U}}$ , the  $C^*$ -algebra formed from  $\mathcal{U}$  by adjoining the unit 1. Note that  $p$  commutes with  $b$  and hence with each  $p_k$ . Set

$$I_k = \{a \in \mathcal{U} : p_k(1 - p)a(1 - p)p_k = a\}$$

for each  $k = 1, \dots, n$ . Since  $\mathcal{U}$  is norm separable, we may choose a strictly positive element  $a_k \in I_k$  with  $\|a_k\| = 1$  by [1]. (I.e., for any positive linear functional  $h$  on the  $C^*$ -algebra  $I_k$ ,  $h(a_k) = 0$  implies  $h = 0$ .) By [3, II. 7]  $p_k(1 - p)$  is open, hence  $1 - p_k(1 - p)$  is the null projection of  $a_k$  for each  $k = 1, \dots, n$ .

Now we define  $c_k = b_k - b_k a_k b_k$  for  $k = 1, \dots, n$ . Each  $c_k$  is self-adjoint and  $c_k x_j = \delta_{kj} x_j$  (Kronecker Delta). Also  $c_k c_j = 0$  if  $j \neq k$ . Define  $I_0 = \{a \in \mathcal{U} : (1 - p)a(1 - p) = a\}$ . Then let  $\Gamma_0 = \{a \in I_0 : c_k a = a c_k \text{ for all } k = 1, \dots, n\}$ . Let  $\Gamma_1$  be the  $C^*$ -algebra generated by  $\{c_k\}_{k=1}^n$  and  $\Gamma_0$ , and let  $A$  be a maximal abelian  $C^*$ -subalgebra of  $\Gamma_1$  containing  $\{c_k\}_{k=1}^n$ . It is clear that  $f_k|_A$  is multiplicative, hence pure, on  $A$ , since  $f_k|_{I_0} = 0$  for each  $k = 1, \dots, n$ . We shall show that  $A$  is a maximal abelian  $C^*$ -subalgebra of  $\mathcal{U}$ .

Suppose  $c \in \mathcal{U}$  with  $ca = ac$  for all  $a \in A$ . We may suppose  $c \geq 0$  since  $\{d \in \mathcal{U} : da = ad \text{ for all } a \in A\}$  is a  $C^*$ -algebra, hence generated by its positive elements. For each  $k = 1, \dots, n$ , we define a scalar  $\lambda_k$  as follows. If  $cx_k = 0$ , let  $\lambda_k = 0$ . If  $cx_k \neq 0$ , let  $y = cx_k / \|cx_k\|$ . Since  $cc_k = c_k c$ , we get  $c_k(cx_k) = c(c_k x_k) = cx_k$ , so  $c_k y = y$ . Thus  $1 = \langle c_k y, y \rangle = \langle b_k y, y \rangle - \langle a_k b_k y, b_k y \rangle$ . Since  $b_k \geq 0$ ,  $a_k \geq 0$ ,  $\|b_k\| \leq 1$ , this means  $b_k y = y$  and  $a_k b_k y = 0 = a_k y$ . By the above, this means  $(1 - p_k(1 - p))y = y$ , since  $(1 - p_k(1 - p))$  is the null projection of  $a_k$ .

But  $b_k y = y$  implies  $y = p_k y$ . Hence  $y = p_k p y$ . This means that  $y = \lambda_k x_k$ , since  $p_k p$  is the projection on the one dimensional subspace spanned by  $x_k$ . Since the foregoing was valid for each  $k = 1, \dots, n$ , we can define  $d = c - \sum_{k=1}^n \lambda_k c_k$ , and note that  $(1 - p)d(1 - p) = d$ , so that  $d \in I_0$ . But  $da = ad$  for all  $a \in A$ , so  $d \in I'$ , hence  $d \in A$  by maximality of  $A$ . This proves that  $A$  is a maximal abelian  $C^*$ -subalgebra of  $\mathcal{U}$ .

Lastly, we shall prove the uniqueness assertion. Suppose  $g$  is a pure state of  $\mathcal{U}$  with  $g|_A = f_k|_A$ . Then there exists a unit vector  $y \in H_g \subset H$  such that  $\langle ay, y \rangle = g(y)$  for all  $a \in \mathcal{U}$ . Thus  $\langle c_k y, y \rangle = \langle c_k x_k, x_k \rangle = 1 = \langle b_k y, y \rangle - \langle a_k b_k y, y \rangle$ . Using the same argument as above, we get  $b_k y = y$  and  $a_k y = 0$ . As before, this implies  $y = \lambda x_k$ , so  $g = f_k$ .

One is tempted to ask if the hypothesis of separability can be dropped. Example 2.1 shows that we cannot hope for unique state extensions in the (very) nonseparable case. However, the question of the existence of the maximal abelian  $C^*$ -subalgebra is still open. For the separable case one might ask if a pure state of a maximal abelian  $C^*$ -subalgebra always has unique pure state extension to the whole algebra. Example 2.2 shows that this is not the case.

We now turn to a quite different problem, that of majorizing an element of a  $C^*$ -algebra by an element of a subalgebra. If  $\mathcal{U}$  is an abelian  $C^*$ -algebra and  $B \subset \mathcal{U}$  a  $C^*$ -subalgebra which contains a positive increasing approximate unit for all of  $\mathcal{U}$ , then for every  $a \geq 0$  in  $\mathcal{U}$  there is  $b \in B$  with  $b \geq a$ . In fact, with a slightly more refined argument one can choose  $b$  so that  $\|b\| = \|a\|$ . Example 2.3 shows that this last assertion is (in general) false if the hypothesis of commutativity is dropped. The following result shows how close we can come to the abelian case.

**THEOREM 1.2** *Let  $A$  be a  $C^*$ -algebra and  $A_0$  a  $C^*$ -subalgebra of  $A$  which contains a positive, increasing approximate unit  $\{a_\alpha\}_{\alpha \in I}$  for  $A$ . Then given  $a_0 \geq 0$  in  $A$  and  $\varepsilon > 0$ , there exists  $b \geq 0$  in  $A_0$  with  $b \geq a_0$  and  $\|b\| \leq \|a_0\| + \varepsilon$ .*

*Proof.* First let us note that if the theorem is true for all  $a_0 \in A$  with  $\|a_0\| = 1$ , then it is true for all  $a_0 \in A$ . Thus we may assume  $\|a_0\| = 1$ . Given  $\varepsilon > 0$  we may choose  $\alpha_1 \in I$  such that  $\|a_0 - a_{\alpha_1} a_0 a_{\alpha_1}\| < \varepsilon/2$ . Since  $\{a_\alpha\}$  is an increasing positive approximate identity and  $\|a_0\| = 1$ ,  $a_{\alpha_1} \geq a_{\alpha_1} a_0 a_{\alpha_1}$ . Set  $a_1 = a_0 - a_{\alpha_1} a_0 a_{\alpha_1}$ . Then find  $\alpha_2 \in I$  such that  $\|a_1 - a_{\alpha_2} a_1 a_{\alpha_2}\| < \varepsilon/4$ . Now  $\|(2/\varepsilon)a_1\| \leq 1$ , so as above

$$a_{\alpha_2} \geq a_{\alpha_2}^2 \geq a_{\alpha_2} \left( \frac{2}{\varepsilon} a_1 \right) a_{\alpha_2}.$$

Thus  $(\varepsilon/2)a_{\alpha_2} \geq a_{\alpha_2}a_1a_{\alpha_2}$ . Set  $a_2 = a_1 - a_{\alpha_2}a_1a_{\alpha_2}$ , and continue by induction to get sequences  $\{\alpha_n\}_{n=1}^\infty \subset I$  and  $\{a_n\}_{n=0}^\infty \subset A$  so that  $a_n = a_{n-1} - a_{\alpha_n}a_{n-1}a_{\alpha_n}$  for  $n > 0$  ( $a_0$  as given),  $\|a_n\| \leq \varepsilon/2^n$  for  $n > 0$ , and

$$\left( \frac{\varepsilon}{2^{n-1}} \right) a_{\alpha_n} \geq a_{\alpha_n} a_{n-1} a_{\alpha_n}.$$

Thus the series  $\sum_{n=1}^\infty a_{\alpha_n} a_{n-1} a_{\alpha_n}$  is absolutely convergent to  $a_0$ . Also

$$a_0 = \sum_{n=1}^\infty a_{\alpha_n} a_{n-1} a_{\alpha_n} \leq a_{\alpha_1} + \sum_{n=2}^\infty \left( \frac{\varepsilon}{2^{n-1}} \right) a_{\alpha_n},$$

the right hand side also converging absolutely to an element  $b \in A_0$ . Clearly

$$\|b\| \leq \|a_{\alpha_1}\| + \sum_{n=2}^\infty \left\| \left( \frac{\varepsilon}{2^{n-1}} \right) a_{\alpha_n} \right\| \leq 1 + \varepsilon,$$

so we have the theorem.

In order to state the last result of this section we introduce a definition.

**DEFINITION.** We say a pure state  $f$  of  $\mathcal{U}$  is *pure on a closed left ideal  $I$*  if  $f$  is pure on the  $C^*$ -algebra  $I \cap I^*$ .

**THEOREM 1.3** *Let  $I$  be a closed left ideal of a  $C^*$ -algebra  $\mathcal{U}$ . An increasing directed net of positive operators  $\{a_\alpha\} \subset I$  is an approximate right unit for  $I$  if  $f(a_\alpha) \rightarrow 1$  for every pure state  $f$  of  $\mathcal{U}$  which is pure on  $I$ .*

*Proof.* Let  $\{a_\alpha\} \subset I$  be a positive increasing directed net in  $I$  with  $f(a_\alpha) \rightarrow 1$  for every pure state  $f$  of  $\mathcal{U}$  which is pure on  $I$ . Now set  $I_0 = I \cap I^*$ , and note that  $\{a_\alpha\} \subset I_0$ . By [4, 5.1]  $\{a_\alpha\}$  is an approximate unit for  $I_0$ , since every pure state of  $I_0$  has (by [7]) a pure state extension to  $\mathcal{U}$ . Let  $b \in I$ . Then

$$\begin{aligned} \|b - ba_\alpha\|^2 &= \|(b^* - a_\alpha b^*)(b - ba_\alpha)\| \\ &= \|b^*b - b^*ba_\alpha - a_\alpha b^*b + a_\alpha b^*ba_\alpha\|_\alpha \rightarrow 0 \end{aligned}$$

since  $b^*b \in I_0$  and  $\{a_\alpha\}$  is an approximate unit for  $I_0$ . Thus  $\|b - ba_\alpha\|_\alpha \rightarrow 0$ .

2. **Examples.** This first example will be a nonseparable  $C^*$ -algebra  $A$  which has no positive increasing approximate unit  $\{a_\alpha\}$  consisting of pairwise commuting elements. That is, no maximal abelian  $C^*$ -subalgebra of  $A$  contains an approximate unit for  $A$ . By considering  $\tilde{A}$  the  $C^*$ -algebra obtained from  $A$  by adjoining a unit, we get a pure state  $f$  on  $\tilde{A}$  with  $f|_A = 0$  (considering  $A \subset \tilde{A}$ ) which cannot be the unique pure state extension of any pure state of any maximal abelian  $C^*$ -subalgebra of  $\tilde{A}$ .

**EXAMPLE 2.1** Let  $\Gamma$  be an index set of cardinality  $2^c$ , where  $c$  is the cardinality of the set of real numbers. Let  $H$  be a Hilbert space with an orthonormal basis of cardinality  $2^{2^c}$ . Choose a family  $\{H_\gamma\}_{\gamma \in \Gamma}$  of orthogonal subspaces of  $H$  with  $\dim(H_\gamma) = \dim(H)$  for all  $\gamma \in \Gamma$ . For each fixed  $\gamma \in \Gamma$ , choose a family  $\{H_\gamma^\alpha\}_{\alpha \in \Gamma}$  of subspace of  $H_\gamma$  which are orthogonal and such that  $\dim(H_\gamma^\alpha) = \dim(H)$  for all  $\gamma, \alpha \in \Gamma$ . Let  $H_0 = \Sigma \oplus \{H_\gamma : \gamma \in \Gamma\}$ , and we shall work in  $B(H_0)$ , the algebra of all bounded operators on  $H_0$ . For each pair  $(\alpha, \gamma) \in \Gamma \times \Gamma$  with  $\alpha \neq \gamma$ , choose projections  $p_{\alpha\gamma}$  and  $q_{\alpha\gamma}$  on  $H_\gamma^\alpha$  such that  $p_{\alpha\gamma}q_{\alpha\gamma} \neq q_{\alpha\gamma}p_{\alpha\gamma}$ . Define for each  $\beta \in \Gamma$  a projection  $p_\beta \in B(H_0)$  by defining  $p_\beta$  on each subspace  $H_\gamma^\alpha$  as follows:  $p_\beta|H_\gamma^\alpha = 0$  if  $\alpha \neq \beta$  and  $\gamma \neq \beta$ ,  $p_\beta|H_\gamma^\alpha = 1_{\beta\beta}$  if  $\alpha = \beta = \gamma$ ,  $p_\beta|H_\beta^\alpha = p_{\alpha\beta}$  if  $\alpha \neq \beta$ , and  $p_\beta|H_\alpha^\beta = q_{\beta\alpha}$  if  $\beta \neq \alpha$ .

Now we let  $A$  be the  $C^*$ -algebra generated by  $\{p_\gamma\}_{\gamma \in \Gamma}$ . We note the following facts.

- (1)  $p_\gamma p_\alpha \neq p_\alpha p_\gamma$  unless  $\gamma = \alpha$ .
- (2) If  $1_{\alpha\gamma}$  denotes the projection on  $H_\gamma^\alpha$ , then  $1_\gamma^\alpha \in A'$ , the commutant of  $A$  in  $B(H_0)$ , and  $1_\gamma^\alpha \cdot 1_\rho^\beta = 0$  unless  $\alpha = \beta$  and  $\gamma = \rho$ .
- (3)  $p_\gamma p_\alpha p_\beta = 0$  if  $\gamma \neq \alpha \neq \beta \neq \gamma$ .

These are immediate from the definition of the  $\{p_\gamma\}$ .

Now define the  $*$ -homomorphism  $\varphi_{\alpha\gamma} : A \rightarrow B(H_0)$  by  $\varphi_{\alpha\gamma}(a) = 1_{\alpha\gamma}a$ , and let  $A_{\alpha\gamma}$  be the kernel of this homomorphism for each pair  $(\alpha, \gamma) \in \Gamma \times \Gamma$ . We note that for each  $\gamma \in \Gamma$ ,  $A = A_{\gamma\gamma} + \{\lambda p_\gamma : \lambda \text{ a scalar}\}$ . Now suppose  $A_1 \subset A$  is a maximal abelian  $C^*$ -subalgebra of  $A$  and that  $A_0$  contains an approximate identity for all of  $A$ . Then for each  $\gamma \in \Gamma$ , surely  $\varphi_{\gamma\gamma}$  is nonzero on  $A_0$ , so that  $A_0$  contains a positive element of the form  $p_\gamma + a_\gamma$  where  $a_\gamma \in A_{\gamma\gamma}$ . We shall prove that this implies that  $A_0$  is not abelian.

For each  $\gamma \in \Gamma$ , we are assuming  $A_0$  contains a positive (fixed) element of the form  $p_\gamma + a_\gamma$  with  $a_\gamma \in A_{\gamma\gamma}$ . Now for each fixed  $\gamma \in \Gamma$ ,  $a_\gamma$  can be written as the sum of a series of products of the  $\{p_\alpha\}_{\alpha \in \Gamma}$  with suitable scalar coefficients. This series can be chosen so that every term of the series lies in  $A_{\gamma\gamma}$ . We shall fix such a series for each  $\gamma$ , and define  $F_\gamma = \{\alpha \in \Gamma : p_\alpha \text{ appears as a factor in one of the}$

terms of the series for  $a_\gamma$ . Since each  $F_\gamma$  is countable and  $\text{card}(\Gamma) = 2^c$ , we may choose a collection  $\{F_\gamma: \gamma \in K\}$ , where  $K$  is an uncountable subset of  $\Gamma$  (any subset  $K$  with  $\text{card}(K) = c$  will work), and  $\bigcup \{F_\gamma: \gamma \in K\} \neq \Gamma$ . Then for fixed  $\beta \in \Gamma \sim \bigcup \{F_\gamma: \gamma \in K\}$ ,  $p_\beta$  does not appear as a factor in any term of the series for  $a_\gamma$  for any  $\gamma \in K$ . Since  $(p_\beta + a_\beta) \in A_0$  for some  $a_\beta \in A_{\beta\beta}$  and  $K$  is uncountable (and the series for  $a_\beta$  is countable), there exists some  $\gamma \in K$  (which we now fix) with  $p_\gamma a_\beta 1_{\gamma\beta} = 0 = 1_{\gamma\beta} a_\beta p_\gamma$ .

We now show that  $(p_\gamma + a_\gamma)$  and  $(p_\beta + a_\beta)$  do not commute. We need only check that  $1_{\gamma\beta}(p_\alpha + a_\gamma)(p_\beta + a_\beta) \neq 1_{\gamma\beta}(p_\beta + a_\beta)(p_\gamma + a_\gamma)$ . Now

$$\begin{aligned} 1_{\gamma\beta}(p_\gamma + a_\gamma)(p_\beta + a_\beta) &= 1_{\gamma\beta}(p_\gamma p_\beta + p_\gamma a_\beta + a_\gamma p_\beta + a_\gamma a_\beta) = 1_{\gamma\beta}(p_\gamma p_\beta) \\ &= q_{\gamma\beta} p_{\gamma\beta} \neq p_{\gamma\beta} q_{\gamma\beta} = 1_{\gamma\beta}(p_\beta p_\gamma) \\ &= 1_{\gamma\beta}(p_\beta + a_\beta)(p_\gamma + a_\gamma). \end{aligned}$$

This is seen as follows.  $1_{\gamma\beta} p_\gamma a_\beta = 0 = 1_{\gamma\beta} a_\beta p_\gamma$  by the choice of  $\gamma$  above. Also  $1_{\gamma\beta} a_\gamma a_\beta = 1_{\gamma\beta} a_\beta a_\gamma = 0$  because no term of the series for  $a_\gamma$  contains  $p_\beta$  as a factor, and each term lies in  $A_{\gamma\gamma}$ . Finally  $1_{\gamma\beta} p_\beta a_\gamma = 1_{\gamma\beta} a_\gamma p_\beta = 0$  by the choice of  $\beta \in \Gamma \sim \bigcup \{F_\alpha: \alpha \in K\}$ . We have proved that  $A_0$  is not abelian, so this contradiction establishes the result that  $A$  has no maximal abelian  $C^*$ -subalgebra which contains an approximate unit for all of  $A$ .

Now if  $\tilde{A}$  is the  $C^*$ -algebra consisting of  $A$  with the identity adjoined, define the pure state  $f$  on  $\tilde{A}$  by  $f(a + \lambda 1) = \lambda$ , where  $a \in A$  (every element of  $\tilde{A}$  can be written uniquely in the form  $a + \lambda 1$  with  $a \in A$ ). If  $\tilde{A}_0$  is a maximal abelian  $C^*$ -subalgebra of  $\tilde{A}$ , then  $\tilde{A}_0 = A_0 + \{\lambda 1\}$ , where  $A_0$  is a maximal abelian  $C^*$ -subalgebra of  $A$ , and  $A_0$  is the kernel of  $f|_{\tilde{A}_0}$ . If  $\{a_\alpha\}_{\alpha \in I} \subset A_0$  is a positive increasing approximate unit for  $A_0$ , the only pure state of  $\tilde{A}_0$  vanishing on all the  $a_\alpha$  ( $\alpha \in I$ ) is  $f|_{\tilde{A}_0}$ . However, it follows immediately from [4, p. 531] and [3, Th. II. 17] that there are at least two pure states of  $A$  vanishing on  $\{a_\alpha\}$  or else  $\{a_\alpha\}$  would be an approximate unit for all of  $A$ —contradicting the conclusion above.

In the next example we construct a separable  $C^*$ -algebra  $A$  with unit and a maximal abelian  $C^*$ -subalgebra  $A_0$  of  $A$  and two pure states  $f, g$  of  $A$  with  $f = g$  and  $f|_{A_0} = g|_{A_0}$ .

**EXAMPLE 2.2** For each  $n = 1, 2, \dots$  let  $H_n$  be a two-dimensional Hilbert space with fixed orthonormal basis  $\{e_n, e'_n\}_{n=1}^\infty$ . Let  $H = \sum_{n=1}^\infty \oplus H_n$ . Let  $C$  be the  $C^*$ -algebra of all operators  $b$  on  $H$  such that  $b(H_n) \subset H_n$  and  $\lim_{n \rightarrow \infty} \|b|_{H_n}\| = 0$ . ( $C$  is the  $C^*(\infty)$  direct sum of the algebras  $B(H_n)$ .) Let  $p$  be a projection on  $H$  defined on

each  $H_n$  by the matrix  $\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$  with respect to the given basis. Finally let  $A$  be the  $C^*$ -algebra generated by  $C$ ,  $p$ , and  $1$  (the identity operator on  $H$ ). Let  $A_0$  be the maximal abelian  $C^*$ -subalgebra of  $A$  consisting of all operators  $b$  in  $A$  with  $be_n = \lambda_n e_n$ ,  $be'_n = \lambda'_n e'_n$  for all  $n = 1, 2, \dots$ . ( $A_0$  is the algebra of operators in  $A$  diagonalized by the given basis.) Now  $A/A \cap C$  is two-dimensional so there are two pure states  $f$  and  $g$  on  $A$  with  $f \neq g$  and  $\ker(f) \supset A \cap C$ ,  $\ker(g) \supset A \cap C$ . Since  $A_0/A_0 \cap C$  is one dimensional,  $f|_{A_0} = g|_{A_0}$ .

Our last result is an example which proves that one cannot take  $\varepsilon = 0$  in Theorem 1.2.

**EXAMPLE 2.3** Let  $H$  be a separable Hilbert space and  $\{e_n\}_{n=1}^\infty$  an orthonormal basis for  $H$ . Let  $A$  be the algebra of all compact operators on  $H$  and  $A_0 = \{a \in A: \text{each } e_i \text{ is an eigenvector of } a\}$ . That means  $A_0$  is the algebra of compact diagonal operators for the basis  $\{e_n\}$ . For each  $n = 1, 2, \dots$  define  $q_n$  to be the orthogonal projection on the subspace spanned by  $\{e_1, \dots, e_n\}$ . It is known that  $\{q_n\}$  is a positive, increasing approximate identity for  $A$ . Also  $A_0$  is the  $C^*$ -algebra generated by  $\{q_n\}$ . Thus the theorem applies.

Now let  $p$  be the orthogonal projection on the subspace spanned by any vector  $x = \sum_{i=1}^\infty x_i e_i$ , where we assume  $x_i \neq 0$  for all  $i$  and  $\|x\| = 1$ . To see that the  $\varepsilon$  condition of the theorem is necessary, suppose there was some  $b \in A_0$ ,  $\|b\| = 1$ ,  $b \geq p$ . Thus

$$\langle bx, x \rangle = \sum_{n=1}^\infty \langle be_n, e_n \rangle |x_n|^2 \geq \langle px, x \rangle = \langle x, x \rangle = \sum_{n=1}^\infty |x_n|^2.$$

This would mean  $\langle be_n, e_n \rangle = 1$  for all  $n$ , contradicting the compactness of the operator  $b$ .

## BIBLIOGRAPHY

1. J. Aarnes and R. V. Kadison, *Pure states and approximate identities*, Proc. Amer. Math. Soc. **21** (1969), 749-752.
2. C. Akemann, *The dual space of an operator algebra*, Trans. Amer. Math. Soc. (2) **126** (1967), 286-302.
3. ———, *The general Stone-Weierstrass problem*, J. Functional Analysis **4** (1969), 277-294.
4. ———, *Interpolation in  $W^*$ -algebras*, Duke Math. J. **35** (1968), 525-533.
5. J. Dixmier, *Les  $C^*$ -Algebres et leurs representations*, Gauthier-Villars, Paris, 1964.
6. J. Glimm and R. V. Kadison, *Unitary operators in  $C^*$ -algebras*, Pacific J. Math. **10** (1960), 547-566.
7. R. V. Kadison and I. M. Singer, *Extensions of pure states*, Amer. J. Math. **81** (1959), 383-400.
8. S. Sakai, *The theory of  $W^*$ -algebras*, mimeographed notes, Yale University, New Haven, 1962.
9. M. Takesaki, *On the conjugate space of an operator algebra*, Tohoku Math. J. **10** (1958), 194-203.

10. ———, *Singularity of positive functionals*, Proc. Japan Acad. **35** (1959), 365-366.

Received April 10, 1969.

UNIVERSITY OF CALIFORNIA  
SANTA BARBARA, CALIFORNIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Charles A. Akemann, <i>Approximate units and maximal abelian <math>C^*</math>-subalgebras</i> .....	543
Gail Atneosen, <i>Wild points of cellular arcs in 2-complexes in <math>E^3</math> and cellular hulls</i> .....	551
John Logan Bryant and De Witt Sumners, <i>On embeddings of 1-dimensional compacta in a hyperplane in <math>E^4</math></i> .....	555
H. P. Dikshit, <i>On a class of Nörlund means and Fourier series</i> .....	559
Nancy Dykes, <i>Generalizations of realcompact spaces</i> .....	571
Hector O. Fattorini, <i>Extension and behavior at infinity of solutions of certain linear operational differential equations</i> .....	583
Neal David Glassman, <i>Cohomology of nonassociative algebras</i> .....	617
Neal Hart, <i>Ulm's theorem for Abelian groups modulo bounded groups</i> .....	635
Don Barker Hinton, <i>Continuous spectra of second-order differential operators</i> .....	641
Donald Gordon James, <i>On Witt's theorem for unimodular quadratic forms. II</i> .....	645
Melvin F. Janowitz, <i>Principal multiplicative lattices</i> .....	653
James Edgar Keesling, <i>On the equivalence of normality and compactness in hyperspaces</i> .....	657
Adalbert Kerber, <i>Zu einer Arbeit von J. L. Berggren über ambivalente Gruppen</i> .....	669
Keizō Kikuchi, <i>Various <math>m</math>-representative domains in several complex variables</i> .....	677
Jack W. Macki and James Stephen Muldowney, <i>The asymptotic behaviour of solutions to linear systems of ordinary differential equations</i> .....	693
Andy R. Magid, <i>Locally Galois algebras</i> .....	707
T. S. Ravisankar, <i>On differentially simple algebras</i> .....	725
Joseph Gail Stampfli, <i>The norm of a derivation</i> .....	737
Francis C.Y. Tang, <i>On uniqueness of central decompositions of groups</i> .....	749
Robert Charles Thompson, <i>Some matrix factorization theorems. I</i> .....	763
Robert Charles Thompson, <i>Some matrix factorization theorems. II</i> .....	811