

Pacific Journal of Mathematics

CELL-LIKE MAPPINGS. II

R. C. LACHER

CELL-LIKE MAPPINGS, II

R. C. LACHER

This paper is an addendum to the previous paper, Cell-like Mappings, I. Therein, the category of cell-like maps between ENR's was established, homotopy-theoretic characterizations of cell-like maps were given, and the image of a cell-like map on an ENR was studied. In the present paper, three related topics are considered: the relationship between (sometimes global) properties of a map and local properties of its mapping cylinder; limits of cell-like maps; and preservation of tameness properties under cell-like maps. Loose descriptions of some of the results follow.

If an onto map between metric spaces has its image locally collared in its mapping cylinder, then the two spaces are stably homeomorphic. If a proper, onto map between ENR's has its mapping cylinder locally k -connected mod its image for all k , then the map is cell-like (hence a proper homotopy equivalence).

The limit of a sequence of cell-like maps between ENR's is cell-like. Likewise, if a proper map between ENR's is "concordantly" approximated by cell-like maps, it is cell-like.

The property of having ULC^1 complements (for compact sets in ENR's) is preserved under monotone maps.

In an appendix, the nonexistence of two types of isolated singularities is proved.

Since this paper is a continuation of [17], none of the definitions from [17] will be restated. All conventions, notation, etc., from [17] carry over. When referencing [17], we will use (I. i. j) to mean "result i.j. of [17]."

As usual, R is the real line, $I = [0, 1]$, R^n is euclidean n -space, B^n is the unit ball in R^n , and $S^n = \partial B^{n+1}$.

1. **Locally trivial mapping cylinders.** If $f: X \rightarrow Y$ is a map, the mapping cylinder Z_f of f is the quotient space of $(X \times I) \cup (Y \times 2)$ obtained by identifying $(x, 1)$ with $(f(x), 2)$ for each $x \in X$. X is identified with the image of $X \times 0$ in Z_f and Y is identified with the image of $Y \times 2$ in Z_f . We have the natural projection $p: Z_f \rightarrow Y$ (a cell-like map when f is proper) and the map $q: X \times I \rightarrow Z_f$, the quotient map restricted to $X \times I$. (q is cell-like if and only if f is cell-like.)

Local collars. Let Y be a closed subset of the space Z , $y \in Y$. A local collar of Y in Z at y is an embedding $\gamma: V \times [0, 1) \rightarrow Z$ such that $\gamma(V \times [0, 1))$ is open in Z and $\gamma(v, 0) = y$ for all $v \in V$, where V

is some neighborhood of y in Y . (See [4].)

We say Y is *locally collared* in Z if it is locally collared in Z at each point of Y .

The following theorem serves to illustrate the power of certain types of hypotheses concerning the way the range of a map is attached to its mapping cylinder.

THEOREM 1.1. *Let X and Y be metric spaces, $f: X \rightarrow Y$ an onto map. If Y is locally collared in Z_f (e.g., if Z_f is a topological manifold with boundary $X \cup Y$) then $X \times R \approx Y \times R$.*

Proof. Y is closed in Z_f , so by [4] Y is collared in Z_f . That is, there is an embedding γ of $Y \times [0, 1)$ onto an open subset of Z_f such that $\gamma(y, 0) = y$ for all $y \in Y$. Note that, by the definition of Z_f , X is also collared in Z_f . In fact, if we let $\lambda = q|X \times [0, 1)$, then $\lambda(x, 0) = x$ for all $x \in X$ and $\lambda(X \times [0, 1)) = Z_f - Y$.

SUBLEMMA. *For any $0 < \eta < 1$, there is a homeomorphism $h: Z_f \approx Z_f$ such that*

- (1) $h|(X \cup Y) = \text{identity}$, and
- (2) $h\gamma(Y \times [0, 1)) \supset Y \cup \lambda(X \times [\eta, 1))$.

Proof of sublemma. Since X is paracompact, we can find open covers $\{U_\alpha\}$ and $\{V_\alpha\}$ of X such that $\bar{V}_\alpha \subset U_\alpha$, $\{U_\alpha\}$ being locally finite, and such that there exists a number $0 < \phi_\alpha < 1$ with

$$\lambda(x, t) \in \gamma(Y \times [0, 1))$$

for all $x \in V_\alpha$ and $\phi_\alpha \leq t < 1$. We extend ϕ_α to a continuous function $\bar{\phi}_\alpha: X \rightarrow [0, \phi_\alpha]$ which is zero outside U_α . Now let

$$\phi(x) = \max \bar{\phi}_\alpha(x).$$

Clearly ϕ is continuous and maps X into $[0, 1)$. Moreover,

$$\lambda(x, t) \in \gamma(Y \times [0, 1))$$

for all $x \in X$ and $\phi(x) \leq t < 1$.

We can easily find a homeomorphism g of $X \times [0, 1]$ onto itself such that

$$\begin{aligned} g|X \times 0 &= \text{identity}, \\ g|x \times [(\phi(x) + 1)/2, 1] &= \text{identity}, \text{ and} \\ g(x \times [\phi(x), 1]) &= x \times [\eta, 1] \end{aligned}$$

hold for all $x \in X$. (g is defined linearly on each interval $x \times [0, \eta]$, $x \times [\eta, \phi(x)]$, $x \times [\phi(x), (\phi(x) + 1)/2]$, and $x \times [(\phi(x) + 1)/2, 1]$. Continuity of ϕ implies g is a homeomorphism.) Define h by

$$h = \begin{cases} \lambda g \lambda^{-1} & \text{on } Z_f - Y \\ \text{identity} & \text{on } Y \end{cases}.$$

This completes the proof of the sublemma.

The completion of the proof of (1.1) uses a typical “monotone union” argument to show that $(Z_f - X, Y) \approx (Y \times (0, 1], Y \times 1)$ and hence that $X \times (0, 1) \approx Z_f - X - Y \approx Y \times (0, 1)$. (See [3] or [15].)

Locally shrinkable maps. Let $f: X \rightarrow Y$ be an onto map. We say f is locally shrinkable (by pseudo-isotopies) if and only if, for each $y \in Y$, there exist a neighborhood V of y in Y and a proper homotopy $h: f^{-1}(V) \times I \rightarrow f^{-1}(V)$ such that h_t is an onto homeomorphism for $0 \leq t < 1$ and $\{h_t^{-1}(x) | x \in f^{-1}(V)\} = \{f^{-1}(y) | y \in V\}$.

COROLLARY 1.2. *Let X and Y be locally compact metric spaces, $f: X \rightarrow Y$ a proper, onto map. If f is locally shrinkable, then $X \times R \approx Y \times R$.*

Proof. For $y \in Y$, let V and h be given as in the definition of locally shrinkable, $U = f^{-1}(V)$. The conditions imply that $qH^{-1}: U \times I \rightarrow Z_f$ is an embedding, where $H(x, t) = h(x, t) \times t$. (See [10].) Also, the image of $U \times I$ under qH^{-1} is $p^{-1}(V)$, a neighborhood of y in Z_f . Finally, note that $h_1 f^{-1}: V \rightarrow U$ is a homeomorphism of V onto U , so $\gamma = qH^{-1}(h_1 f^{-1} \times T): V \times I \rightarrow Z_f$ is a local collar at y , where $T(t) = 1 - t$. Hence, Y is locally collared in Z_f .

2. Local connectivity for maps.

UV-properties. Let A be a subset of the space X , and let \mathcal{F} be a homotopy invariant covariant functor from a category containing inclusion maps of open sets (respectively based open sets) of X to the category of based sets. We say that $A \subset X$ has property $UV(\mathcal{F})$ if and only if, for any neighborhood U of A in X , there exists a neighborhood V of A in U such that the inclusion-induced map $\mathcal{F}(V) \rightarrow \mathcal{F}(U)$ (respectively $\mathcal{F}(V, v) \rightarrow \mathcal{F}(U, v)$) is zero (for all $v \in V$). Property $UV(\mathcal{F})$ is a topological property of A in the following sense:

THEOREM 2.1. *Let A be a compact set in the ANR X . If there*

exists an embedding $f: A \rightarrow Y$, where Y is an ANR and $f(A) \subset Y$ has property $UV(\mathcal{F})$, then $A \subset X$ has property $UV(\mathcal{F})$. (See the argument that (a) \Rightarrow (d) in [16].)

Relative local properties. Let Z be a space, Y a closed subset of Z , and let \mathcal{F} be a homotopy invariant covariant functor from a category containing inclusion maps of open subsets of Z to the category of based sets. For $y \in Y$, we say Z is $LC(\mathcal{F}) \bmod Y$ at y if and only if, for any neighborhood U of y in Z , there is a neighborhood V of y in U such that the inclusion-induced map $\mathcal{F}(V - Y) \rightarrow \mathcal{F}(U - Y)$ is zero. If Z is $LC(\mathcal{F}) \bmod Y$ for all $y \in Y$, we say that Z is $LC(\mathcal{F}) \bmod Y$.

Local properties of maps. Let $f: X \rightarrow Y$ be a map. We say that f is $LC(\mathcal{F})$ if and only if Z_f is $LC(\mathcal{F}) \bmod Y$.

Standard examples. There are special notations for $UV(\mathcal{F})$ and $LC(\mathcal{F})$ for certain \mathcal{F} . When $\mathcal{F} = \tilde{H}_k(-; G)$ (reduced singular homology with coefficients in the group G) we get $k - uv(G)$ and $k - lc(G)$, respectively. When $\mathcal{F} = \tilde{H}_0(-; G) \oplus \cdots \oplus \tilde{H}_k(-; G)$, we get $uv^k(G)$ and $lc^k(G)$, respectively. When $\mathcal{F} = [S^k, -]$ (based homotopy classes of maps $S^k \rightarrow -$) we get $k - UV$ and $k - LC$, respectively. And when $\mathcal{F} = [S^0, -] \oplus \cdots \oplus [S^k, -]$, we get UV^k and LC^k , respectively.

For an appropriate space K , $\mathcal{F} = [K, -]$ also yields UV^∞ and LC^∞ , respectively. When we are dealing with ANR's, $K =$ disjoint union of all homotopy types of ANR's, would do. When dealing with separable metric spaces, $K =$ disjoint union of all separable metric spaces works. And other examples can be worked up.

THEOREM 2.2. *Let X and Y be locally compact metric spaces, let $f: X \rightarrow Y$ be a proper, onto map, and let \mathcal{F} be a homotopy invariant covariant functor from a category containing inclusion maps of open sets of X and $X \times I$ to the category of based sets. Then the following are equivalent:*

- (a) *Each inclusion $f^{-1}(y) \subset X$ has property $UV(\mathcal{F})$, $y \in Y$.*
- (b) *f is $LC(\mathcal{F})$.*

REMARKS. Theorem 2.2 relates the (recently defined) UV -properties to the (classical) local connectivity properties. For results on the latter, see [11], [22] and [26], among others. The results from [22] can be translated, using (2.2) above, into an "Hurewicz" theorem: Let $f: X \rightarrow Y$ be a proper, onto map between locally compact ANR's; if each $f^{-1}(y)$ has property UV^k then each has $uv^k(G)$ for all G ; and, if each $f^{-1}(y)$ has UV^1

and $uv^k(Z)$, where $Z = \text{integers}$, then each has UV^k . (Compare with Theorems 4.1 and 4.2 of [18].) Theorems 2 and 3 of [11] can be translated, again using (2.2), to yield (I.3.1). (See the remark on page 617.)

Proof. First assume that each inclusion $f^{-1}(y) \subset X$ has $UV(\mathcal{F})$. Let $y \in Y$, and suppose that U is a given neighborhood of y in Z_f . Find a neighborhood U_1 of y in Y and a number t , $0 < t < 1$, such that $p^{-1}(U_1) \cap q(X \times (t, 1])$ is contained in U . Then, we can find a neighborhood V_1 of y in U_1 such that the inclusion-induced map

$$\mathcal{F}(f^{-1}(V_1)) \rightarrow \mathcal{F}(f^{-1}(U_1))$$

is zero. Define $V = p^{-1}(V_1) \cap q(X \times (t, 1])$. Since \mathcal{F} is homotopy invariant, the inclusion-induced

$$\text{map } \mathcal{F}(f^{-1}(V_1) \times (t, 1)) \rightarrow \mathcal{F}(f^{-1}(U_1) \times (t, 1))$$

is zero. But the restriction of q to $f^{-1}(U_1) \times (t, 1)$ takes the pair

$$(f^{-1}(U_1) \times (t, 1), f^{-1}(V_1) \times (t, 1))$$

homeomorphically into $(U - Y, V - Y)$, with $f^{-1}(V_1) \times (t, 1)$ mapping onto $V - Y$. Thus it is clear that $\mathcal{F}(V - Y) \rightarrow \mathcal{F}(U - Y)$ is zero, so Z_f is $LC(\mathcal{F}) \bmod Y$ at y .

Now suppose Z_f is $LC(\mathcal{F}) \bmod Y$. Let $y \in Y$, and let U_1 be a neighborhood of y in Y , $U = p^{-1}(U_1)$. By assumption, there is a neighborhood V_1 of y in Y such that $\mathcal{F}(V - Y) \rightarrow \mathcal{F}(U - Y)$ is zero, where $V = p^{-1}(V_1) \cap q(X \times (t, 1])$ for some $t < 1$. Let $U_0 = f^{-1}(U_1)$ and $V_0 = f^{-1}(V_1)$. Then q maps $(U_0 \times [0, 1], V_0 \times (t, 1))$ homeomorphically onto $(U - Y, V - Y)$, so it is clear that $\mathcal{F}(V_0) \rightarrow \mathcal{F}(U_0)$ is zero. Since $f^{-1}(y)$ has arbitrarily small neighborhood pairs of the form (U_0, V_0) , we see that $f^{-1}(y) \subset X$ has property $UV(\mathcal{F})$.

REMARK. If one were to desire a definition of “locally homotopically collared” pertaining to $Y \subset Z$, “ $LC^\infty \bmod Y$ ” might be a good choice, at least when Y is locally contractible. This interpretation, combined with known facts, yields some interesting analogues (as well as conjectures) about cell-like maps and locally shrinkable maps.

COROLLARY 2.3. *Let X and Y be locally compact metric spaces, $f: X \rightarrow Y$ a proper, onto map. If f is LC^k then*

$$f_*: \pi_q(X, x) \rightarrow \pi_q(Y, f(x))$$

is an isomorphism for $0 \leq q \leq k$ (and an epimorphism for $q = k + 1$, provided Y is an ANR).

Proof. Apply (I.3.1). If Y is an ANR, use the technique in the proof of (I.2.4) to see that f_* is epic when $q = k + 1$.

COROLLARY 2.4. *Let X and Y be ENR's, $f: X \rightarrow Y$ a proper, onto map. f is cell-like if and only if it is LC^∞ .*

3. Limits of cell-like maps. The following was obtained independently by R. Finney. (See [12].)

THEOREM 3.1. *Let X and Y be ENR's and let $f: X \rightarrow Y$ be a proper, onto map. If there exists a sequence of proper, cell-like maps of X onto Y which converges to f (in the compact-open topology) then f is cell-like.*

Before proving (3.1), we need the following lemma on monotone maps. (A map $f: X \rightarrow Y$ is *monotone* if $f^{-1}(y)$ is compact and connected for each $y \in Y$.)

LEMMA 3.2. *Let X and Y be locally compact metric spaces, f a proper map of X onto Y . Let $\{f_n\}$ be a sequence of proper monotone maps of X onto Y which converges to f . Suppose that $y \in V \subset \bar{V} \subset U$, where V and U are open sets in Y , V is connected, and U has compact closure. Then there exists an integer m such that*

$$f^{-1}(y) \subset f_n^{-1}(V) \subset f^{-1}(U)$$

for $n \geq m$.

Proof. Suppose that $f^{-1}(y) \not\subset f_n^{-1}(V)$ for infinitely many n . Then, there is an infinite sequence $\{x_i\}$, with $x_i \in f^{-1}(y) - f_{n_i}^{-1}(V)$. Since $f^{-1}(y)$ is compact, we may assume that $\{x_i\}$ converges to some $x \in f^{-1}(y)$. Thus $f_{n_i}(x_i)$ converges to $f(x) = y$, so almost all of $f_{n_i}(x_i)$ must lie in V , contrary to the choice of $x_i \notin f_{n_i}^{-1}(V)$. Hence there is an integer k such that $f^{-1}(y) \subset f_n^{-1}(V)$ for $n \geq k$.

Now suppose that $f_n^{-1}(\bar{V}) \not\subset f^{-1}(U)$ for infinitely many n . Since $f_n^{-1}(\bar{V}) \cap \overline{f^{-1}(U)} \neq \emptyset$ for all $n \geq k$, and each $f_n^{-1}(\bar{V})$ is compact and connected, we can find an infinite sequence $\{z_i\}$, with $z_i \in f_{n_i}^{-1}(\bar{V}) \cap B$, where $B = \overline{f^{-1}(U)} \cap (X - f^{-1}(U))$. B is compact, so we may assume that $\{z_i\}$ converges to some $z \in B$. Since $\{f_{n_i}(z_i)\}$ converges to $f(z)$, and $z_i \in f_{n_i}^{-1}(\bar{V})$, we see that $f(z) \in \bar{V} \subset U$, contradicting the fact that $z \notin f^{-1}(U)$. We conclude that there is an integer $m \geq k$ such that

$$f_n^{-1}(V) \subset f^{-1}(U)$$

for $n \geq m$.

Proof of (3.1). We want to show that each inclusion $f^{-1}(y) \subset X$ has property UV^∞ . The result will then follow from (I.1.1). Let U, V , and W be connected neighborhoods of $y \in Y$, chosen so that $\bar{W} \subset V \subset \bar{V} \subset U$ and \bar{U} is compact. Let $\{f_n\}$ be a sequence of proper, cell-like maps which converges to f . Choose W so that W is contractible in V . By (3.2), we can find an integer m such that

$$f^{-1}(y) \subset f_m^{-1}(W) \subset f_m^{-1}(V) \subset f^{-1}(U) .$$

By (I.1.2), $f_m^{-1}(W)$ is contractible in $f_m^{-1}(V)$. Since $f^{-1}(y)$ has arbitrarily small neighborhoods of the form $f^{-1}(U)$, $f^{-1}(y)$ has property UV^∞ .

THEOREM 3.3. *Let $f: X \rightarrow Y$ be an onto map between ENR's. Suppose $f \times 0$ extends to a proper map $f': X \times [0, 1] \rightarrow Y \times [0, 1]$ such that $f'|X \times (0, 1)$ is a cell-like map of $X \times (0, 1)$ onto $Y \times (0, 1)$. Then f is cell-like.*

Proof. When $\alpha: X \rightarrow (0, 1)$ is continuous, let ϕ_α be the map

$$\phi_\alpha(x, t) = q\left(x, 1 - \frac{t}{\alpha(x)}\right), x \in X, 0 \leq t \leq \alpha(x) .$$

Let $g_\alpha = f' \phi_\alpha^{-1}$. Then g_α is a cell-like map of Z_f onto a subset of $Y \times [0, 1]$. (g_α is a homeomorphism on Y and is f' "twisted" on $x \times [0, 1]$.) Moreover, $g_\alpha(y) = y \times 0$, and $g_\alpha(Z_f - Y) \subset Y \times (0, 1)$.

Let α be chosen so that

$$Y \times [0, \tfrac{1}{2}] \subset \bigcup_{x \in X} f'(x \times [0, \alpha(x)]) = g_\alpha(Z_f) .$$

Then Y is collared in $g_\alpha(Z_f)$. Also, g_α is cell-like, $g_\alpha^{-1}(g_\alpha(Z_f) - Y) = Z_f - Y$, and $g_\alpha|Y$ is a homeomorphism, so it follows as in the proof of (4.2) below that Z_f is $LC^\infty \bmod Y$. By (2.4), f is cell-like.

4. Preservation of tameness properties. The tameness properties referred to involve the LC^k properties. We think of the compact set A in the ANR X as being "locally k -tame" in X whenever X is $LC^k \bmod A$; clearly, this is a topological property of the pair (X, A) .

It should be noted that, when A is closed in the compact ANR X , X is $LC^k \bmod A$ if and only if $X - A$ is ULC^k (uniformly locally k -connected).

THEOREM 4.1. *Let X and Y be locally compact ANR's, and let A and B be compact sets in X and Y , respectively. Suppose that $f: X \rightarrow Y$ is a proper, onto UV^k -map such that $f^{-1}(B) = A$ and $f|A$ is one-to-one. If X is $LC^{k+1} \bmod A$ then Y is $LC^{k+1} \bmod B$.*

THEOREM 4.2. *Let X, Y, A, B , and f be as in the hypothesis of Theorem 4.1. If Y is $LC^k \bmod B$ then X is $LC^k \bmod A$.*

Proof of (4.1.). We will show that Y is $LC^{k+1} \bmod B$ at each point of B . Let U be a neighborhood of $y \in B$ in Y , and let V be a neighborhood of y in U such that any map $S^q \rightarrow f^{-1}(V - B)$ extends to a map $B^{q+1} \rightarrow f^{-1}(U - B)$, $0 \leq q \leq k + 1$, using the fact X is $LC^{k+1} \bmod A$ at $f^{-1}(x)$.

Since f induces epimorphisms on π_q when restricted to any inverse open set, it follows that any map $S^q \rightarrow (V - B)$ extends to a map $B^{q+1} \rightarrow (U - B)$, $0 \leq q \leq k + 1$, and Y is $LC^{k+1} \bmod B$ at y .

Proof of (4.2.). We want to show that X is $LC^k \bmod A$ at each point of A . Let U be a neighborhood of $x \in A$ in X . Since $f^{-1}f(x) = x$, there is a neighborhood W of $f(x)$ in Y such that $f^{-1}(W) \subset U$. Now, Y is $LC^k \bmod B$ at $f(x)$, so there is a neighborhood W' of $f(x)$ in Y such that any map $S^q \rightarrow (W' - B)$ extends to a map $B^{q+1} \rightarrow (W - B)$, $0 \leq q \leq k$. Let $V = f^{-1}(W')$. Again, since f induces isomorphisms on π_q when restricted to any inverse open set in $X - A$, we see that any map $S^q \rightarrow (V - A)$ extends to a map $B^{q+1} \rightarrow (U - A)$, $0 \leq q \leq k$, and X is $LC^k \bmod A$ at x .

COROLLARY 4.3. *Let M^m and N^n be unbounded PL manifolds with $m \geq 5$ and $n \geq 5$. Further, let P^p and Q^q be compact polyhedra topologically embedded in M and N , respectively, $m - p \geq 3$, $n - q \geq 3$. Finally, let $f: M \rightarrow N$ be a proper, onto monotone map such that $f^{-1}(Q) = P$ and $f|P$ is one-to-one.*

(1) *If P is tame in M then Q is tame in N .*

(2) *If each $f^{-1}(y)$ has property 1-UV and Q is tame in N then P is tame in M .*

Proof. When $n - p \geq 3$, M is $LC^1 \bmod P$ if and only if P is tame in M . The difficult part of this statement is due to Bryant and Seebeck [5].

5. Maps on Euclidean space. In [8], Cohen pointed out that, if $f: R^3 \rightarrow R^3$ is the example described by Bing in [2] of a proper, monotone noncell-like map, there is no proper, onto map $f': R^4 \rightarrow R^4$ whose nondegenerate point-inverses are the same as those of $f \times 0$. A higher-dimensional analogue of this result can be obtained.

THEOREM 5.1. *Let $f: R^n \rightarrow R^n$ be a proper, onto UV^1 -map. If there exists a proper, onto map $f': R^{n+1} \rightarrow R^{n+1}$ whose nondegenerate point-inverses are the same as those of $f \times 0$, then f is cell-like (and hence cellular if $n \neq 4$).*

Thus, for example, f could not be any of the generalizations of Bing's example described in § 6 of [18], with $k \geq 2$ and $l \geq 2$.

Proof. Note that the condition implies that Z_f embeds in R^{n+1} with the crinkled end being $f'(R^n \times 0)$. Thus Z_f is $lc^k \text{ rel } f(R^n)$ for all k , by Theorem II. 5. 35 [26], and hence f is UV^1 -trivial and uv^k -trivial for all k by Theorem 2.2. Using the "Hurewicz" theorem mentioned in the remark following (2.2), we see that f is UV^k -trivial for all k , hence cell-like.

For $n \leq 3$, it is not necessary to assume that f is UV^1 -trivial in the above proposition. (See [23] and Theorem 2 of [19].)

THEOREM 5.2. (*Lacher and Wright.*) *Let $f: R^n \rightarrow R^n$ be a proper, onto map, $n \leq 3$. If there exists a proper, onto map $f': R^{n+1} \rightarrow R^{n+1}$ whose nondegenerate point-inverses are the same as those of $f \times 0$, then f is shrinkable by a pseudo-isotopy.*

THEOREM 5.3. *Let $f: R^{n+1} \rightarrow R^{n+1}$ be a proper, onto map,*

$$S_f = \{x \in R^{n+1} | f^{-1}f(x) \neq x\}.$$

If $S_f \subset R^n \times 0$, $f(R^n \times 0) \approx R^n$, and $\overline{f(S_f)}$ is a polyhedron of dimension $\leq n - 3$, tame in both $f(R^n \times 0)$ and R^{n+1} , then f is cell-like.

Proof. By Theorem 6.1 of [7], $f(R^n)$ is locally flat in R^{n+1} . Assuming $n \geq 3$, and applying [6], we obtain a homeomorphism h of R^{n+1} onto itself such that $hf(R^n) = R^n$. By Theorem 3.3, $hf|R^n$ is cell-like, hence $f|R^n$ is cell-like. Since $S_f \subset R^n$, f is cell-like.

REMARKS. 1. D. R. McMillan has recently proved the following: If $f: R^{n+1} \rightarrow R^{n+1}$ is a proper, onto map, and $S(f) \subset R^n \times 0$, then each $f^{-1}(y)$ has property $uv^\infty(Z)$.

2. The results of this section should be compared with [28].

Appendix: The nonexistence of two types of isolated singularities. Let M be a manifold (without boundary), Y a Hausdorff space, and $f: M \rightarrow Y$ a proper, onto map. Define two "singular sets" as follows:

$$C_f = \{y \in Y | f^{-1}(y) \text{ is not cellular in } M\},$$

and

$$E_f = \{y \in Y | Y \text{ is not locally euclidean at } y\}.$$

We show below that $C_f - E_f$ has no isolated points (assuming $\dim M \geq 5$, f is strongly acyclic, and M is simply connected), and that $E_f - \bar{C}_f$ has no isolated points (assuming $\dim M \neq 4, 5$). (Note: E_f is

a closed subset of Y . \bar{C}_f denotes the closure of C_f in Y .)

A1. *Noncellular points of a strongly acyclic map.* A map $f: X \rightarrow Y$ is *strongly acyclic* (cf. [20]) if and only if each inclusion $f^{-1}(y) \subset X$, $y \in Y$, has property $UV(\mathcal{S})$, where $\mathcal{S} = \tilde{H}_*(-; \mathbb{Z})$. (I.e., \mathcal{S} is reduced singular homology with integral coefficients.) According to Corollary 3.3 of [18], if X is an ENR then a proper map $f: X \rightarrow Y$ is strongly acyclic if and only if $\tilde{H}^*(f^{-1}(y)) = 0$ for all $y \in Y$. (\tilde{H}^* is reduced Čech cohomology.) McMillan [20] and [21] and Wright [27] have studied strongly acyclic maps on 3-manifolds. In particular, Wright has shown that if $f: M^3 \rightarrow N^3$ is a proper, onto, strongly acyclic map between (open or closed) 3-manifolds, then C_f is a closed, locally finite set in N , and, hence, if $M = S^3$ or R^3 , $C_f = \emptyset$. Theorem 1 below could be considered a weak analogue of this last statement.

We recall two methods of producing strongly acyclic maps.

EXAMPLE 1. Let H be a topological k -manifold which is a homology k -sphere, and let A be the closure of the complement of a locally flat k -cell in H . By taking the quotient map cross the identity map, we obtain a strongly acyclic map $H \times S^{n-k} \rightarrow S^k \times S^{n-k}$ between n -manifolds with $\dim C_f = n - k$, $k \geq 3$. In particular, one can have C_f a finite set when the domain is not simply connected.

EXAMPLE 2. Let $f: R^n \rightarrow R^n$ be a proper, onto map, and let $S_f = \{x \in R^n \mid f^{-1}f(x) \neq x\}$. If $f(S_f)$ is compact and 0-dimensional, then f is strongly acyclic. (See [25].)

Finally, we should point out that there is a strongly acyclic map of S^n onto itself, $n \geq 4$, with $\dim \bar{C}_f = 1$. (See [21].)

THEOREM 1. *Let M and N be (open or closed) topological n -manifolds, $n \geq 5$, and let $f: M \rightarrow N$ be a proper, onto, strongly acyclic map. If M is simply connected then C_f has no isolated points.*

Proof. Suppose that y_0 is a point of N which has a neighborhood V such that $f^{-1}(y)$ is cellular in M for all $y \neq y_0$ in V . Since

$$\check{H}^* f^{-1}(y_0) = 0,$$

$f^{-1}(y_0)$ has a *PL* neighborhood, according to Theorem 2 of [13]. We assume, therefore, that V is an open n -cell and that $f^{-1}(V) = U$ is a *PL* manifold.

Let W be a closed neighborhood of $f^{-1}(y_0)$ in U , chosen to be a compact *PL* manifold. Let $W_0 = W - f^{-1}(y_0)$, $U_0 = U - f^{-1}(y_0)$, and $V_0 = V - \{y_0\}$. Since $f|_{U_0}: U_0 \rightarrow V_0$ is cellular, we see that W_0 is 1-connected at infinity. Applying Theorem 3.10 of [24], we can replace

W by another compact manifold W' whose boundary is connected and simply connected. Applying Van-Kampen's Theorem, we see that $0 = \pi_1(\overline{M - W'}) * \pi_1(W')$, so W' is simply connected. Therefore $f^{-1}(y_0)$ has property UV^1 .

It follows from Theorem 4.2 of [18] that $f^{-1}(y_0)$ has property UV^∞ and hence is cellular in M .

A2. Noneuclidean points of a cellular map. Let $f: S^n \rightarrow Y$ be an onto map whose only nondegenerate point-inverse is A . Then A is cellular if and only if Y is a manifold. I.e., $C_f = \emptyset$ if and only if $E_f = \emptyset$. Such a statement is not true about maps in general, but Theorem 2 below is a partial converse to Theorem 1 in this sense.

THEOREM 2. *Let M be a topological n -manifold (open or closed), $n \neq 4, 5$, and let Y be a Hausdorff space. If $f: M \rightarrow Y$ is a proper, cellular map, then E_f has no isolated points.*

Proof. Let $y_0 \in Y$, and suppose that y_0 has a neighborhood V such that $V - \{y_0\}$ is an open topological manifold. Then $\dim(V - \{y_0\}) = n$, so V is an ENR by Corollary I.3.3.

Let $W = f^{-1}(V - \{y_0\})$. Since $f|_W$ is a proper homotopy equivalence when restricted to any inverse open set (see Theorem I. 1.2). and since $f^{-1}(y_0)$ is cellular in M , it is clear that y_0 has a neighborhood $V' \subset V$ such that the inclusion-induced map $H^*(V - \{y_0\}; Z_2) \rightarrow H^*(V' - \{y_0\}; Z_2)$ is zero. It follows from the Kirby-Siebenmann triangulation theorem [14] that y_0 has a neighborhood $V'' \subset V'$ such that $V'' - \{y_0\}$ has a PL structure. (See the remark following Theorem 2 of [13]).

We may as well assume, then, that $V - \{y_0\}$ has a PL structure. Let ε be the end of V determined by y_0 . Then $\{f^{-1}(U) | U \in \varepsilon\}$ determines an end $f^{-1}(\varepsilon)$ of W , the same one determined by $f^{-1}(y_0)$. Moreover, $f^{-1}(\varepsilon)$ is $(n - 2)$ -connected, so ε is $(n - 2)$ -connected. It follows immediately from [24] that, in case $n \geq 6$, ε has a collar neighborhood (which must be $S^{n-1} \times [0, \infty)$ by [9]). Thus V is a topological (in fact, PL) manifold, and $y_0 \notin E_f$.

If $n \leq 3$, $f|_W$ is properly homotopic to a homeomorphism: $W \approx V - \{y_0\}$, so the result is easy in that case. (See [1].)

REFERENCES

1. S. Armentrout, *Cellular decompositions of 3-manifolds that yield 3-manifolds*, Bull. Amer. Math. Soc. **75** (1969), 453-455.
2. R. H. Bing, "Decompositions of E^3 " *Topology of 3-manifolds*, M. K. Fort, Jr. (Editor), Prentice-Hall, Englewood Cliffs, N. J., 1962.
3. M. Brown, *The monotone union of open n -cells in an open n -cell*, Proc. Amer. Math.

Soc. **12** (1961), 812-814.

4. ———, *Locally flat imbeddings of topological manifolds*, Ann. of Math. (2) **75** (1962), 331-341.
5. J. L. Bryant and C. L. Seebeck, III, *Locally nice embeddings in codimension three*, Bull. Amer. Math. Soc. **74** (1968), 378-380.
6. J. C. Cantrell, *Almost locally flat embeddings of S^{n-1} in S^n* , Bull. Amer. Math. Soc. **69** (1963), 716-718.
7. J. C. Cantrell and R. C. Lacher, *Local flattening of a submanifold*, Quart. J. Math. (Oxford Second Series) **20** (1969), 1-10.
8. M. Cohen, *Simplicial structures and transverse cellularity*, Ann. of Math. (2) **85** (1967), 218-245.
9. E. H. Connell, *A topological h -cobordism theorem for $n \geq 5$* , Illinois J. Math. **11** (1967), 300-309.
10. R. H. Crowell, *Invertible isotopies*, Proc. Amer. Math. Soc. **14** (1963), 658-664.
11. S. Eilenberg and R. L. Wilder, *Uniform local connectedness and contractibility*, Amer. J. Math. **64** (1942), 613-622.
12. R. L. Finney, *Uniform limits of compact cell-like maps*, Notices Amer. Math. Soc. **15** (1968), 942.
13. J. Hollingsworth and R. B. Sher, *Triangulating neighborhoods in topological manifolds*.
14. R. C. Kirby and L. C. Siebenmann, *On the triangulation of manifolds and the Hauptvermutung*, Bull. Amer. Math. Soc. **75** (1969), 742-49.
15. K. W. Kwun, *Uniqueness of the open cone neighborhood*, Proc. Amer. Math. Soc. **15** (1964), 476-479.
16. R. C. Lacher, *Cell-like spaces*, Proc. Amer. Math. Soc. **20** (1969), 598-602.
17. ———, *Cell-like mappings, I*, Pacific J. Math. **30** (1969), 717-731.
18. ———, *Cellularity criteria for maps*, Michigan Math. J. **17** (1970), 385-396.
19. R. C. Lacher and A. H. Wright, *Mapping cylinders and 4-manifolds*, Proc. 1969 Ga. Top. Cont.
20. D. R. McMillan, *Compact, acyclic subsets of three-manifolds*, Michigan Math. J. **16** (1969), 129-136.
21. ———, *Acyclicity in three-manifolds*, Bull. Amer. Math. Soc. **76** (1970), 942-964.
22. M. H. A. Newman, *Local connection in locally compact spaces*, Proc. Amer. Math. Soc. **1** (1950), 44-53.
23. V. Nicholson, *Mapping cylinder neighborhoods*, Trans. Amer. Math. Soc. **143** (1969), 259-268.
24. L. C. Siebenmann, *The obstruction to finding a boundary for a manifold of dimension greater than five*, Princeton University Ph. D. Thesis (University Microfilms #66-5012), 1965.
25. K. A. Sitnikov, *On continuous mappings of open sets of a Euclidean space*, Mat. Sbornik N. S. **31** (1952), 439-458; (Russian) (MR#14,489).
26. R. L. Wilder, *Topology of manifolds*, A. M. S. Colloquium Publication 32, New York, 1949.
27. A. Wright, *Monotone mappings of compact 3-manifolds*, Ph. D. Thesis, University of Wisconsin, 1969.
28. J. Zaks, *Trivially extending decompositions of E^n* , Pacific J. Math. **29** (1969), 727-729.

Received June 13, 1969, and in revised form February 2, 1970. Research supported in part by NSF grant GP-11943.

THE FLORIDA STATE UNIVERSITY
TALLAHASSEE, FLORIDA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

John D. Arrison and Michael Rich, <i>On nearly commutative degree one algebras</i>	533
Bruce Alan Barnes, <i>Algebras with minimal left ideals which are Hilbert spaces</i>	537
Robert F. Brown, <i>An elementary proof of the uniqueness of the fixed point index</i>	549
Ronn L. Carpenter, <i>Principal ideals in F-algebras</i>	559
Chen Chung Chang and Yiannis (John) Nicolas Moschovakis, <i>The Suslin-Kleene theorem for V_κ with cofinality $(\kappa) = \omega$</i>	565
Theodore Seio Chihara, <i>The derived set of the spectrum of a distribution function</i>	571
Tae Geun Cho, <i>On the Choquet boundary for a nonclosed subspace of $C(S)$</i>	575
Richard Brian Darst, <i>The Lebesgue decomposition, Radon-Nikodym derivative, conditional expectation, and martingale convergence for lattices of sets</i>	581
David E. Fields, <i>Dimension theory in power series rings</i>	601
Michael Lawrence Fredman, <i>Congruence formulas obtained by counting irreducibles</i>	613
John Eric Gilbert, <i>On the ideal structure of some algebras of analytic functions</i>	625
G. Goss and Giovanni Viglino, <i>Some topological properties weaker than compactness</i>	635
George Grätzer and J. Sichler, <i>On the endomorphism semigroup (and category) of bounded lattices</i>	639
R. C. Lacher, <i>Cell-like mappings. II</i>	649
Shiva Narain Lal, <i>On a theorem of M. Izumi and S. Izumi</i>	661
Howard Barrow Lambert, <i>Differential mappings on a vector space</i>	669
Richard G. Levin and Takayuki Tamura, <i>Notes on commutative power joined semigroups</i>	673
Robert Edward Lewand and Kevin Mor McCrimmon, <i>Macdonald's theorem for quadratic Jordan algebras</i>	681
J. A. Marti, <i>On some types of completeness in topological vector spaces</i>	707
Walter J. Meyer, <i>Characterization of the Steiner point</i>	717
Saad H. Mohamed, <i>Rings whose homomorphic images are q-rings</i>	727
Thomas V. O'Brien and William Lawrence Reddy, <i>Each compact orientable surface of positive genus admits an expansive homeomorphism</i>	737
Robert James Plemmons and M. T. West, <i>On the semigroup of binary relations</i>	743
Calvin R. Putnam, <i>Unbounded inverses of hyponormal operators</i>	755
William T. Reid, <i>Some remarks on special disconjugacy criteria for differential systems</i>	763
C. Ambrose Rogers, <i>The convex generation of convex Borel sets in euclidean space</i>	773
S. Saran, <i>A general theorem for bilinear generating functions</i>	783
S. W. Smith, <i>Cone relationships of biorthogonal systems</i>	787
Wolmer Vasconcelos, <i>On commutative endomorphism rings</i>	795
Vernon Emil Zander, <i>Products of finitely additive set functions from Orlicz spaces</i>	799
G. Sankaranarayanan and C. Suyambulingom, <i>Correction to: "Some renewal theorems concerning a sequence of correlated random variables"</i>	805
Joseph Zaks, <i>Correction to: "Trivially extending decompositions of E^n"</i>	805
Dong Hoon Lee, <i>Correction to: "The adjoint group of Lie groups"</i>	805
James Edward Ward, <i>Correction to: "Two-groups and Jordan algebras"</i>	806