

Pacific Journal of Mathematics

**ON A SIX-DIMENSIONAL PROJECTIVE REPRESENTATION
OF $PSU_4(3)$**

JOHN HATHWAY LINDSEY, II

ON A SIX DIMENSIONAL PROJECTIVE REPRESENTATION OF $PSU_4(3)$

J. H. LINDSEY, II

In the course of an investigation of six-dimensional complex linear groups, it was discovered that a central extension of Z_6 by $PSU_4(3)$ has a representation of degree six. In fact, this representation has as its image the unimodular subgroup $X(G)$ of index 2 of the following 6-dimensional matrix group: \langle all 6 by 6 permutation matrices; all unimodular diagonal matrices of order 3; $I_6 - Q/3$ where Q has all its entries equal to one \rangle . This matrix group leaves the following lattice invariant: $\{(a_1, \dots, a_6) \mid a_i \in Z(\omega) \text{ where throughout this paper } \omega \text{ is a primitive third root of unity; } a_i - a_j \in \sqrt{-3}Z(\omega) \text{ for all } i, j; \sum_{i=1}^6 a_i \in 3Z(\omega)\}$. The generators of the matrix group are similar to the following generators for an 8-dimensional complex linear group with Jordan-Holder constituents Z_2 , the non-trivial simple constituent of $O_8(2)$, Z_2 : \langle all 8 by 8 permutation matrices, all unimodular diagonal matrices of order 2, $I_8 - P/4$ where P has all entries equal to 1 \rangle .

The projective representation of $PSU_4(3)$ can be used to construct a 12-dimensional representation $Y(H)$, a central extension of Z_6 by the Suzuki group, which leads to the known 24-dimensional projective representation of the Conway group. In fact, H has a subgroup K isomorphic to a central extension of $\langle Z_6 \times Z_3 \rangle$ by $PSU_4(3)$. Also, $Y \mid H$ has two six-dimensional constituents coming from the above matrix group where the constituents are related by an outer automorphism of $PSU_4(3)$ which does not lift to the central extension of Z_6 by $PSU_4(3)$ with the six-dimensional representation. We obtain two commuting automorphisms, α and β respectively, of G from $I_6 - Q/3$ and complex conjugation. For $PSU_4(3)$, the outer automorphism group is dihedral of order eight with its center corresponding to complex conjugation of $X(G)$. The entire automorphism group lifts to K . We may take the center of K to be $\langle a, b, c \rangle$ with a and b of order 3 and c of order 2, with $G \cong K/b$, and with $\alpha(a) = a, \alpha(b) = b^{-1}, \beta(a) = a^{-1}, \beta(b) = b^{-1}$. We can also find an automorphism γ of K with $\gamma(a) = b$ and $\gamma(b) = a$. We give the character table of K giving only one representative of each family of algebraically conjugate characters and classes. Irrational characters and classes are underlined. Only one class in each coset of $Z(K)$ is represented by the character tables. The characters in the table $\widetilde{U}_4(3)$ give the characters with $Z(K)$ in the kernel. The succeeding five character tables in order give the following linear characters, respectively, on $Z(K)$: $\theta(a) = \theta(b) = 1, \theta(c) = -1$; $\theta(a) = \omega, \theta(b) = \theta(c) = 1$; $\theta(a) = \omega^{-1}, \theta(b) = 1, \theta(c) = -1$; $\theta(a) =$

$\theta(b) = \omega, \theta(c) = 1; \theta(a) = \theta(b) = \omega, \theta(c) = -1$. The characters with other actions are obtained by applying elements of the outer automorphism group. The automorphism α transposes π_7 with π_7^{-1} ; and N_1 with N_1^{-1} in the character tables. The automorphism β transposes N_1 with N_1^{-1} ; and N_2 with N_2^{-1} . The automorphism γ transposes T_1 with $T_2; JT_1$ with $JT_2; N_1$ with $N_2; N_1^{-1}$ with N_2^{-1} ; and possibly π_7 with π_7^{-1} . As $SU_4(3)/\Omega(ZSU_4(3))$ has the centralizer of some central involution isomorphic to the centralizer of some central involution J in G , presumably $SU_4(3)/\Omega_1(ZSU_4(3)) \cong G/0_3(Z(G))$.

The first four character tables give the characters of the central extension of $\langle d \rangle = Z_6$ by $LF(3, 4)$ with a six dimensional, complex representation. Respectively, they give the following linear characters on $\langle a \rangle$: $\theta(a) = 1, \theta(a) = \omega, \theta(a) = -1, \theta(a) = -\omega$. The characters with $\theta(a) = \omega^{-1}$ or $\theta(a) = -\omega^{-1}$ come from complex conjugation of the second and fourth table respectively.

We let $\widetilde{U}_4(3) = PSU_4(3)$ and let S_p be a p -Sylow subgroup of whatever group is in question. The term ‘‘Blichfeldt’’ refers to the theorem in [1] that no primitive complex linear group contains an element with some eigenvalue within 60 degrees of all the other eigenvalues of the element. Where clear, we use χ_n to refer to the previously discussed character of G of degree n . Finally, $a(X, Y, Z)$ is the coefficient of the conjugacy class containing Z in the product of the classes containing X and Y .

This paper fills a gap in [9] concerning groups G with a faithful unimodular representation X with character χ of degree six and \bar{G} simple of order $2^7 3^6 35$ where $Z = Z(G)$ and $\bar{G} = G/Z$. We also know by [9, § 8], that $C(S_5) = S_5 Z, C(S_7) = S_7 Z, 4/t_5 = [N(S_5) : C(S_5)] = 4$, and $6/t_7 = [N(S_7) : C(S_7)] = 3$. Also, the principal 7-block $B_6(7)$ has degree equation $1 + 729 = 640 + 90$. Finally, by [9, § 8], $\chi(G) \cong Q(\omega), 3 \parallel Z$, and we may take $X(S_3)$ to be

$$\left\langle \text{diag}(1, 1, \omega, 1, 1, \omega^{-1}), \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \oplus I_3, I_3 \oplus \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \right\rangle.$$

I learned from the referee that this representation was discovered earlier by Mitchell, [10]. Mitchell also showed that this linear group and the first orthogonal group on six indices with modulus three have isomorphic nonsolvable Jordan-Holder constituents. Hammill, [6] and Todd, [12] also worked on this linear group with the latter constructing the character table of $\widetilde{U}_4(3)$.

2. The character table. By the above, $|Z| = 6$ since χ_{20} , the character of the skew-symmetric tensors of $X \otimes X \otimes X$, does not have a constituent of degree 90 or 640. There is a character χ_{640} , completing the 2-block of χ_{640} in G/Z_3 . Since $\chi_{640'}$ is the 7-exceptional character in the block $B_1(7)$ with characters whose kernel is Z_3 , and G/Z_3 does not have a character of degree 6, χ_{20} is irreducible. Degrees divisible exactly by 2 or 4 and $\equiv \pm 1 \pmod{7}$ and $\equiv 0$ or $\pm 1 \pmod{5}$ are 6, 36, 90, 20, and 540. The possibilities are $660 - 36 = 624$, $660 + 90 = 750$, $660 + 20 = 680$, and $660 - 540 = 120$. The degree equation is $20 + 640 = 120 + 540$. By [5], 3-7 block separation in G/Z_3 , these characters are in the same 3-block of G/Z_3 . Let $T \in Z(S_3)$, $\chi(T) = -3$. Then $(\text{mod } 3) |G|\chi_{640}(T)/(540)|C(T)| \equiv |G|\chi_{20}(T)/(20)|C(T)| \equiv (-7)|G|/(20)|C(T)| =$ some 3-unit, so $\chi_{640}(T)$ is divisible exactly by 27 and $|C(T)|/|Z| > 3^6$.

A 7-block whose characters have kernel Z_2 contains χ_{15} from the skew-symmetric tensors (irreducible since G/Z_2 has no representation of degree 6) and $\chi_{729'}$ completing a 3-block of defect 1. There is another degree divisible exactly by 3 which must be 384, 24, 15, 60, 120, 480, or 960. The degree 24 is impossible since

$$\chi_{24}(\pi_7)\overline{\chi_{24}(\pi_7)} = 2,$$

but $\chi_{24}\overline{\chi_{24}}$ cannot fit $\chi_0 + \chi_{729}$ in $B_0(7)$ inside. The possibilities are $729 + 15 - 384 = 360$, $744 + 15 = 759$, $744 + 60 = 804$, $744 + 120 = 864$, $744 + 480 = 1224$, and $744 + 960 = 1704$. Since G/Z_2 has no representation of degree 6, χ_{24} corresponding to the symmetric tensors of $X \otimes X$, is irreducible. In the case of 864 there is a 5-block with degree equation $864 + 864 + 729 = 21 + \dots$ and the fifth degree is too large. Therefore, the 7-block has degree equation $15 + 729 = 384 + 360$. Suppose that G has an element J with $X(J)$ having eigenvalues $i, i, i, -i, -i, -i$. Then $\chi_{15}(J) = (0^2 - (-6))/2 = 3$. Also χ_{384} has 2-defect 0 and $\chi_{384}(J) = 0$. Since $t_7 = 2$, $a_{J, J, \pi_7} = 0$ in G/Z and G/Z_2 , so

$$3^2/15 + \chi_{729'}(J)^2/729 - \chi_{360}(J)^2/360 = 0$$

and $3 + \chi_{729'}(J) = \chi_{360}(J)$. Then $9|\chi_{729'}(J)|, 3|\chi_{360}(J)|, 27|\chi_{729'}(J)|$, and $4|\chi_{360}(J)|$; so $\chi_{729'}(J) \equiv -27 \pmod{108}$. Then $\chi_{729'}(J) = -27$, otherwise $|\chi_{729'}(J)| > 80$ and the sum is negative. Then in $B_0(7)$,

$$\chi_0(J) = 1, \chi_{729}(J) = -27, \chi_{640}(J) = 0, \chi_{90}(J) = 1 - 27 = -26,$$

and $1^2/1 + 27^2/729 - 26^2/90 \neq 0$, a contradiction. Therefore, J cannot exist. We have a character $\chi_{384'}$ faithful on Z completing a 2-block containing χ_{384} . Then a 5-block faithful on Z contains characters of degree 6 and 384. Now $1 = (\chi_{15}, \chi_6\chi_6) = (\overline{\chi}_6\chi_{15}, \chi_6)$ so $\overline{\chi}_6\chi_{15}$ contains χ_6 as a constituent. Also $\overline{\chi}_6\chi_{15} - \chi_6$ has an irreducible constituent of degree $\equiv -1 \pmod{5}$ and divisible by 6: 84 or 24. By the previous

$\chi_{24}\bar{\chi}_{24}(\pi_7)$ argument, 24 is impossible and the 5-block contains the degree 6, 384, and 84. We have another degree divisible exactly by 2: 6, 486, 126, or 1134. The possibilities are

$$384 + 84 - 6 - 6 = 456, 486 - 462 = 24$$

already shown to be an impossible degree,

$$462 - 126 = 336, \text{ and } 462 + 1134 = 1596 .$$

The degree equation is $6 + 126 + 336 = 84 + 384$. As with 84, $\bar{\chi}_6\chi_{21} - \chi_6$ is a character. Since $(\bar{\chi}_6\chi_{21}, \chi_6) = 1$, $\bar{\chi}_6\chi_{21} - \chi_6$ has no constituent of degree 6. Therefore, from the 5-block, all its constituents have degrees divisible by 30, and must be 120, 90, or 60. The degree 90 would imply the impossible degree 30. If 60, then a 7-block has degree equation $6 + 384 = 60 + 330$, impossible. Therefore, it is irreducible, and the 7-block is $6 + 384 = 120 + 270$. If J gives an involution in G/Z , then possibly replacing J by $-J$, $X(J)$ has eigenvalues 1, 1, 1, 1, -1, -1 as $\chi(G) \cong Q(\omega)$ and eigenvalues $i, i, i, -i, -i, -i$ are impossible. In $\bar{G} = G/Z, \langle \pi_5 \rangle$ is self-centralizing and $a_{J,J,\pi_5} = 0$ or 5. Now $|C_G(J)| = |C_{\bar{G}}(\bar{J})| |Z|$ and $a_{J,J,\pi_5} = 0$ or 5 in $G/Z, G/Z_2, G/Z_3$, and G . Then looking successively at $G, G/Z_2, G/Z_3$, and G we see that $\sum \chi_i(J)^2 \chi_i(\pi_5) / \chi_i(1)$ over each 5-block is 0 or 5 $|C_{\bar{G}}(\bar{J})|^2 / |\bar{G}|$. By 2-block orthogonality on (I, J) , $\chi_{384}(J) = 0$. Also $\chi_6(J) = 2, \chi_{84}(J) = 2(4 - 6)/2 - 2 = -4$. Then $\chi_{126}(J) + \chi_{336}(J) = -4 - 2 = -6$. Let $a = \chi_{336}(J)$. We may find some J in $Z(S_2)$ with $2^7 | \sum = 4/6 + a^2/336 + (6 + a)^2/126 - 16/84$. Then $4|a$ and we may let $a = 4b$. Multiply the sum by 63:

$$2^7 | 42 + 3b^2 + 8b^2 + 24b + 18 - 12 = 11b^2 + 24b + 48 .$$

Then $4|b$ and if $c = b/4$, then $8 | 11c^2 + 6c + 3$. Then c is odd. Since $|6 + 16c| < 126$, we have $c = \pm 1, \pm 3, \pm 5$, or ± 7 . Also $11c^2 \equiv 11 \equiv 3 \pmod{8}$, so $6c \equiv 2 \pmod{8}$ and $c \equiv 3 \pmod{4}$. The possibilities are $11 - 6 + 3 = 8, 99 + 18 + 3 = 120$ impossible by the factor 5 since $5 \nmid |C_{\bar{G}}(\bar{J})|, 275 - 30 + 3 = 248$ divisible by 31 and impossible, $539 + 42 + 3 = 584$ divisible by 73. Therefore,

$$c = -1, 5 | C_{\bar{G}}(\bar{J})|^2 / |\bar{G}| = (8)(4)(4)/63 ,$$

and $|C_{\bar{G}}(\bar{J})| = 2^9$. Then J inverts a 5-element and there is only one such class of such $J \pmod{Z}$. If another involution J_1 does not invert a 5-element, then $2^7 | 0 = \sum \chi_i(J_1)^2 \chi_i(\pi_5) / \chi_i(1)$, and the above leads to a contradiction. Therefore, G/Z has a unique class of involutions. Suppose that there is an element F with $X(F)$ having eigenvalues 1, 1, 1, 1, $i, -i$. Then

$$\chi_{15}(F) = (4^2 - 2)/2 = 7, \chi_{21}(F) = (4^2 + 2)/2 = 9, \chi_{84}(F) = 28 - 4 = 24 ,$$

and $\chi_{120}(F) = 36 - 4 = 32$. However, $32^2 + 24^2 > 2^9 = |C_{\bar{c}}(\bar{F}^2)| \geq |C_{\bar{c}}(\bar{F})|$, a contradiction.

3. The centralizer of an involution. Let J be an involution with $X(J) = I_4 \oplus -I_2$. Then $X|C(J) = U \oplus V$ and $\chi|C(J) = \theta + \phi$ where θ corresponds to U and $\theta(J) = 4$. If α is a field automorphism fixing ω , then $\theta^\alpha + \phi^\alpha = \theta + \phi$, $\theta^\alpha = \theta$, and $\phi^\alpha = \phi$ since θ^α and θ are the sums of irreducible characters of $\chi|C(J)$ with J in the kernel. Therefore, $\theta(C(J))$ and $\phi(C(J))$ are contained in $Q(\omega)$. Let K be the subgroup of $C(J)$ of elements k such that $(\det V(k))^{2^m} = 1$ for some m . Then $|K| = 2^9|Z|/3 = 2^8 \cdot 9$. Suppose $x \in \ker U$. Then x is a 2-element, otherwise, some power y of v has order 3 with $\theta(y) = 4$, $\phi(y) = -1$, and Jy contradicts Blichfeldt. If x has order 4, then $X(x)$ has eigenvalues $1, 1, 1, 1, i, -i$; already shown impossible. Therefore, $\ker U = \langle J \rangle$ and $|U(K)| = 2^9$.

Suppose U has 2-dimensional spaces S and T as spaces of imprimitivity or invariant spaces. Then H of index 1 or 2 in $U(K)$ has $\theta|H = \mu + \nu$ corresponding to the 2-dimensional spaces S and T . Let L be a 2-Sylow subgroup of $U(K)$. Unless $[U(K):H] = 2$ and $\mu|L \cap H$ and $\nu|L \cap H$ are irreducible, H has an abelian subgroup A of order 2^5 , impossible (if A has an element of order 8, the linear characters of $\theta|A$ are algebraic conjugates and faithful, so $|A| = 8$. Therefore, irrational characters of $\theta|A$ occur in pairs and have image of order 4. Rational characters have image of order 2. Therefore, $|A| \leq 16$). Therefore, μ and ν are irreducible and a 2-element $x \in C(J)$ transposes S and T . If $\mu \notin Q(\omega)$, then μ and ν are algebraic conjugates, μ is faithful on H , and $H \cap L$ has an abelian subgroup of index 2 and order at least 2^5 , impossible. Therefore, $\mu, \nu \in Q(\omega)$ and $\mu|L \cap H, \nu|L \cap H$ are rational. Then

$$|\mu(L \cap H)|, |\nu(L \cap H)| \leq [2/(2 - 1)] + [2/2] + \dots = 3.$$

Since $|L \cap H| = 2^6, L \cap H = \ker \nu \times \ker \mu$. In 2 by 2 matrix blocks let $U(x) = \begin{pmatrix} 0 & W \\ Y & 0 \end{pmatrix}$. Then $U(x^2) = \begin{pmatrix} WY & 0 \\ 0 & YW \end{pmatrix}$ is contained in a conjugate in H of $\text{Ker } \nu \times \text{Ker } \mu = L \cap H$, a 2-Sylow subgroup of H . Therefore, $\begin{pmatrix} WY & 0 \\ 0 & I_2 \end{pmatrix} = U(y)$ is contained in H . Now $U(y^{-1}x) = \begin{pmatrix} 0 & Y^{-1} \\ Y & 0 \end{pmatrix}$. Changing coordinates by conjugation with $\begin{pmatrix} I_1 & 0 \\ 0 & Y \end{pmatrix}$ and replacing x by $y^{-1}x$, we may take $U(x) = \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}$. Since $\mu|L \cap H$ is irreducible and $L \cap H = \text{Ker } \nu \times \text{Ker } \mu$, there is a 2-element y with $U(y) = -I_2 \oplus I_2$. Then $U((xy)^2) = -I_4$, so $V((xy)^2) \neq -I_2$. However, ϕ is rational and $1 = \det U(xy) = \det V(xy)$. Therefore, $\phi(xy) \pm 2$. If $\text{Ker } \nu$ has an element T of order 3, then $\mu(T) = -1, \nu(T) = 2$, and $X(J(xy)^{-1}TxyT^{-1})$ has eigenvalues $\omega, \bar{\omega}, \omega, \bar{\omega}, -1, -1$; contrary to

Blichfeldt. Therefore, the representation corresponding to μ has image of order 72. Then $\mu \subseteq Q(\omega)$ implies that there is a 3-element g with $\mu(g) = 2\omega$. Then $\nu(g) = 1 + \omega$, otherwise, $\nu(g) = 2\bar{\omega}$ and $X(J(xy)^{-1}gxyg^{-1})$ contradicts Blichfeldt. Now $\phi(g) = \omega + \bar{\omega}$, otherwise, $\phi(g) = 2$ and $X(J(xy)^{-1}gxyg)$ has eigenvalues $\omega, \bar{\omega}, \omega, \bar{\omega}, -1, -1$ and contradicts Blichfeldt. There exists a 2-element z with $\mu(z) = i + (-i)$, $\mu(z^2) = -2$, and $\nu(z) = 2$. Then $\phi(z) = i + (-i)$ and $\phi(z^2) = -2$, otherwise, $X(z)$ or $X(zJ)$ has eigenvalues $i, -i, 1, 1, 1, 1$. Then $\theta(z^{-1}g^{-1}zg) = 4$ implies that $z^{-1}g^{-1}zg \in \langle J \rangle$. As $Jz^{-1}g^{-1}z$ has order 6, it cannot equal g^{-1} , and $z^{-1}g^{-1}zg$ is the identity in G . Then $V(z)$ with eigenvalues $i, -i$ commutes with $V(g)$ with eigenvalues $\omega, \bar{\omega}$ contrary to $\phi \subseteq Q(\omega)$.

Now suppose that U is monomial, but not imprimitive on 2-dimensional subspaces. Then there exists a 3-element g corresponding to a permutation of order 3. As before, $U(K)$ has no abelian subgroup of order 32, so the image of $U(K)$ under ρ , the natural permutation representation on four letters has order eight and must be S_4 . Then $U(K)$ has an element T of order 3 in $\text{Ker } \rho$ and conjugates of some commutator of T with a transposition show that $U(K)$ contains all diagonal matrices of order 3 and determinant 1. Then $27 \mid |U(K)|$, a contradiction.

Now by Blichfeldt's classification of groups of degree 4, $U(K)$ modulo $Z(U(K))$ has a subgroup N of the tensor product of 2-dimensional representations W of $M = GL(2, 3)$. Also, N has index 2 or 1 in $U(K)$. Now $Z(U(K)) \subseteq \langle -I_4 \rangle$ since $\det U(k)$ for $k \in K$ is a 2^m -th root of 1 and $\theta \subseteq Q(\omega)$. Let $U|N = A \otimes B$. Now $W(M) \otimes I_2$ does not appear as a subgroup modulo scalars of $U(K)$ since eigenvalues $\gamma, \gamma, \gamma^{-1}, \gamma^{-1}$ with $\gamma^2 = i$ or $i, i, 1, 1$ contradict 2-rationality of θ . Therefore, the image under A of $\text{Ker } B$ in $M/Z(M)$ has order at most 12. The image of N under B in $M/Z(M)$ has order at most 24. This gives $|N| \leq |Z(N)|(12)(24) \leq 2^9$. We must have equality. Then an element x takes $A \otimes B$ to $B \otimes A$. Therefore, $N \supset W(SL(2, 3)) \otimes I_2$, $I_2 \otimes W(SL(2, 3))$ after elements of $W(SL(2, 3))$ are changed by scalar multiplication. Also, the quaternions $Q = SL(2, 3)'$ can have $W(Q)$ taken as the matrices in [1, § 57]. Since $\det U$ is a 2^m -th root of 1 we may also use the matrix in § 57 for a 3-element S in $W(SL(2, 3))$. Let g be a 3-element with $U(g) = S \otimes I_2$. Then $V(g)$ has eigenvalues $\omega, \bar{\omega}$; otherwise $\phi(g) = 2$ and gJ has eigenvalues $\omega, \bar{\omega}, \omega, \bar{\omega}, -1, -1$; contrary to Blichfeldt. If h is a 3-element with $U(h) = I_2 \otimes S$, then, similarly, $\phi(h) = -1$. Also $U(g)$ and $U(h)$ commute, $V(g)$ and $V(h)$ commute modulo $\langle J \rangle$, and $V(g)$ and $V(h)$ commute. Both may be taken as diagonal. There exists $E \in W(M)$ with $E^{-1}SE = S^{-1}$. Let $V(g) = \omega \oplus \bar{\omega}$. If necessary, we may replace h with h^{-1} and change coordinates of U by conjugation with $I_2 \otimes E$ to take $V(h) = \omega \oplus \bar{\omega}$. If $x \in C(J)$ with $U(x) \in W(Q) \otimes I_2$ and $U(x)$ of order 4, then $U(x)$ has

eigenvalues $i, i, -i, -i$ and $V(x)$ cannot have eigenvalues $i, -i$. Possibly replacing x by Jx , we may take $\phi(x) = 2$. Because of equality in $|N| \leq 2^8 9$, $U(K)$ contains a tensor product of elements in

$$W(GL(2, 3)) - W(SL(2, 3)) .$$

By [1, § 57], we may take this element $U(y)$ as $\alpha((\gamma \oplus \gamma^{-1}) \otimes (\gamma \oplus \gamma^{-1}))$ where $\gamma^2 = i$. Then $U(y)$ has eigenvalues $\alpha i, \alpha, \alpha, -\alpha i$. By 2-rationality, $\alpha = \pm 1$ and $U(y)$ is determined. The action of $U(y)$ on the group of order 3: $W(SL(2, 3)) \otimes W(SL(2, 3)) / \langle W(Q) \otimes W(Q), S \otimes S^{-1} \rangle$ is non-trivial. Therefore,

$$V(y)^{-1} V(g) V(y) = V(y)^{-1} V(h) V(y) = V(g)^{-1}$$

(since $-V(g)^{-1}$ is not a 3-element). Since $1 = \det U(y) = \det V(y)$, we may choose coordinates so that $V(y) = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$. The element x flipping $W(SL(2, 3)) \otimes I_2$ to $I_2 \otimes W(SL(2, 3))$ is determined modulo $W(M) \otimes W(M) / \langle U(y), W(SL(2, 3)) \otimes W(SL(2, 3)) \rangle$ and modulo scalars to be $1 \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus 1$. We may take x as $\alpha(1 \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus 1)$ or $\alpha(1 \oplus \begin{pmatrix} 0 & 1 \\ i & 0 \end{pmatrix} \oplus i)$. As θ is rational on 2-elements, 2α or $\alpha(1 + i)$ is rational. Therefore, $\alpha = \pm 1$, and we are in the first case, so $U(x)$ is determined. Then $-1 = \det U(x) = \det V(x)$ and $V(x)$ has eigenvalues $1, -1$. Since the action of $U(x)$ on $W(SL(2, 3)) \otimes W(SL(2, 3)) / \langle W(Q) \otimes W(Q), S \otimes S^{-1} \rangle$ is trivial, $V(x)$ and $V(g)$ commute. Possibly replacing x by xJ we may take $V(x) = 1 \oplus -1$. Therefore, $C(J)$ and $X(C(J))$ are completely determined. In fact $C(J)/Z$ is isomorphic to $\widetilde{C(I_2 \oplus -I_2)}$ in $\widetilde{U_4(3)}$: $(W(SL(2, 3)) \otimes I_2) \oplus \dots \rightarrow SL(2, 3) \oplus I_2; (I_2 \otimes W(SL(2, 3))) \oplus \dots \rightarrow I_2 \oplus SL(2, 3); \left(\begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix} \right) \oplus \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \rightarrow \begin{pmatrix} i & 0 \\ 0 & 1 \end{pmatrix} \oplus \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix}$ here both elements have the same action on the central product of $SL(2, 3)$ with itself, the square of the left element is $\left(\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right) \oplus -I_2 \approx \left(\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \otimes \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \right) \oplus I_2$. The square of the right element is $-i \left(\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \oplus \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \right); 1 \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus 1 \oplus 1 \oplus -1 \rightarrow \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}$. Here both elements have order 2. Both elements have identical action on the central product of $SL(2, 3)$ with itself. The commutator of $X(x)$ with $X(y)$ is $I_4 \oplus -I_2$. The corresponding commutator in $\widetilde{U_4(3)}$ is $i \oplus i \oplus -i \oplus -i$. This shows that $C(J)/Z$ is isomorphic to the centralizer of an involution in $PSU_4(3)$. By Phan's characterization of $PSU_4(3)$, $PSU_4(3) \cong G/Z$.

4. The normalizer of $Z(S_3)$. Earlier, for

$$T = \text{diag} (\omega, \omega, \omega, \bar{\omega}, \bar{\omega}, \bar{\omega}) ,$$

we showed that $|C(T)/Z| > 3^8$ and T is centralized by an involution in $\bar{G} = G/Z$. We may take T in $C(J)$ and \bar{J} in the center of a Sylow-2-subgroup of $C(T)/Z$. As $\chi(T) = -3$, $U(T) = S^{\pm 1} \otimes I_2$ or $I_2 \otimes S^{\pm 1}$, say the former. Then

$$U(C(TJ)) = \langle U(T), U(Z), I_2 \otimes SL(2, 3) \rangle, |C(T)| = 3^8 |Z|,$$

and T is conjugate to T^{-1} . As the constituents of $X|C(T)$ are not algebraically conjugate, $X(C(T)) = \langle -I_6 \rangle \times H$ where H = the subgroup of $X(C(T))$ whose action on the homogeneous ω -space of $X(T)$ has determinant = to a third root of 1. A Sylow-2-subgroup of H is Q , the quaternions. Let -1 have order 2 in $Z(G)$. Now $\langle \pm J \rangle = Z(Q)$ is represented faithfully in the ω or the $\bar{\omega}$ space of H , say the ω space with $\zeta =$ the corresponding constituent of $X|H$. If ζ is monomial, then $\pm J$, being a square in H , is diagonal and conjugating

with $\begin{pmatrix} 0 & a & 0 \\ 0 & 0 & b \\ c & 0 & 0 \end{pmatrix} \oplus I_3$ (the first component is taken to correspond to ζ),

we have $C(T)/Z$ contains an elementary abelian subgroup of order 4, a contradiction. Therefore, the representation corresponding to ζ is the Hessian group in [1, § 79], except that $\omega \oplus 1 \oplus 1$ has been changed by a scalar. As an element inverting T flips the constituents of $X|C(T)$, taking $H \supset S_3$ with $X(S_3)$ in the normal form given at the start of this chapter, $X(C(T)) \subset \{M_1 \oplus M_2 | M_i$ appears in the Hessian group in [1], except that $\text{diag}(1, 1, \omega)$ replaces $\omega^{-1/3} \text{diag}(1, 1, \omega)\}$. As the normal subgroup K of order 27 of the Hessian group appears independently in each component, we may examine the components of $X(H)$ modulo

K . Let i be the image of $\begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \bar{\omega} \\ 1 & \bar{\omega} & \omega \end{pmatrix} / (\omega - \bar{\omega})$ in this homomorphism.

Since Q is represented faithfully in the top component and some element in $X(C(T))$ flips the components, Q is represented faithfully in the bottom component. By changing coordinates by conjugating with a power of $\text{diag}(1, 1, 1, \omega, 1, 1)$, we may assume that $X(C(T))$ contains $i \oplus \pm \bar{i}$ (i stands for a coset of 3 by 3 matrices and \bar{i} is obtained by complex conjugation of the entries) where

$$j = (\text{diag } 1, 1, \bar{\omega})i(\text{diag } (1, 1, \omega)), -1 = i^2,$$

and $k = ij$. If $X(C(T))$ contains $i \oplus -\bar{i}$, then, conjugating with $T_1 = \text{diag}(1, 1, \omega, 1, 1, \bar{\omega}) \in S_3$, we have $j \oplus -\bar{j}$ and $k \oplus -\bar{k} \in X(C(T))$ and

$$(i \oplus -\bar{i})(j \oplus -\bar{j})(k \oplus -\bar{k}) = -1 \oplus 1 \in X(C(T)),$$

contrary to $8 || |H|$. Since $\text{diag}(1, 1, \omega)$, i , and K generate the Hessian group, $H = \langle K \oplus I_3, I_3 \oplus K, M \oplus \bar{M}$ where M is any matrix in the Hessian group changed as shown by scalars.

$X(N(\langle T \rangle))$ is obtained from $X(C(T))$ by addition of a 2-element

$X(x) = \begin{pmatrix} 0 & E \\ F & 0 \end{pmatrix}$ where E and F are 3 by 3 matrices normalizing the Hessian group, and, hence, in the Hessian group modulo scalar multiplication. By multiplication with an element in $C(T)$ we may take E as scalar and, changing coordinates by conjugation with a direct sum of 3 by 3 scalar matrices, we may take $E = I_3$. Again, we are only interested in F modulo K . If F is scalar, then by determinant, $F = -I_3$ and $X(x^2) = -I_6$, impossible. The other possibilities are $F =$ some scalar times $-1, \pm i, \pm j$, or $\pm k$ in the notation of the previous paragraph. If not -1 , then replace x by $T_1^a x T_1^a$ to take $F =$ some scalar times $\pm i$. The scalar is $-I_3$ by determinant = 1. Then

$$(-I_6)X(x^2) = \begin{pmatrix} \pm i & 0 \\ 0 & \pm i \end{pmatrix}.$$

Possibly replacing this by its third power, we have $\begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & \frac{0}{i} \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$, contrary to $8 || |H|$. Therefore, $F =$ some scalar times -1 and the scalar is -1 by determinant = 1. This completely determines $X(N(\langle T \rangle))$.

5. The correlation between $X(C(J))$ and $X(N(\langle T \rangle))$ for $T \in C(J)$. Take $X(T) = (S \otimes I_2) \oplus \omega \oplus \omega^{-1}$ in our normal form for $X(C(J))$. Let $GL(2, 3)$ and $SL(2, 3)$ be the 2-dimensional matrix groups in [1, § 57] and ϕ be an isomorphism from $SL(2, 3)$ to $SL(2, 3)/0_2(SL(2, 3)) \cong Z_3$ with $\phi(S) = 1$ and $0_2(SL(2, 3))$ isomorphic to the quaternions. Then $X(N(\langle JT \rangle)) = \langle X(JT) = (S \otimes I_2) \oplus -\omega \oplus -\omega^{-1}; (I_2 \otimes u) \oplus (\omega \oplus \omega^{-1})^{\phi(u)}$ for $u \in SL(2, 3)$; $Y = \left(y \otimes \begin{pmatrix} \gamma & 0 \\ 0 & \gamma^{-1} \end{pmatrix} \right) \oplus \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ for some

$$y \in \begin{pmatrix} \gamma & 0 \\ 0 & \gamma^{-1} \end{pmatrix} 0_2(SL(2, 3))$$

with $y^{-1}Sy = S^{-1}; -\omega I_2 \rangle$. We get a subgroup of order at least $2^7 3^8$ of $X(G)$ generated by our normal form for $N(\langle T \rangle)$ and the image under conjugation by a matrix R of our normal form for $X(C(J))$ where R conjugates $X(JT)$ and $X(N(\langle JT \rangle))$, in our normal form for $X(C(J))$, to $X(JT)$ and $X(N(\langle JT \rangle))$, respectively, in our normal form for $X(N(\langle T \rangle))$. Therefore, R is determined modulo multiplication on the left by a matrix P fixing $X(JT)$ and $X(N(\langle JT \rangle))$ in the normal form for $X(C(J))$. As we are only interested in the image of $X(C(J))$ under conjugation by R . We are only interested in P modulo multiplication on the left by a matrix fixing $X(JT)$, $X(N(\langle JT \rangle))$, and $X(C(J))$. As $0_2(0^2(X(N(\langle JT \rangle)))) = \langle (I_2 \otimes u) \oplus I_2 \text{ such that } u \in 0_2(SL(2, 3)) \rangle$, by [7, Satz 3] and [1], $P = (A \otimes B) \oplus C$ where $B \in GL(2, 3)$, $A \in C_{GL(2, C)}(S)$, and $C \in GL(2, C)$ where C is the complex number field. If $B \notin SL(2, 3)$,

then P conjugates $(S \otimes S^{-1}) \oplus I_2$ to $(S \otimes Sv) \oplus I_2$ for some

$$v \in O_2(SL(2, 3)) ,$$

a contradiction, since the former, but not the latter is in $X(N(\langle JT \rangle))$. Therefore, multiplying P by an element in $X(N(\langle JT \rangle))$, we may take $B = I_2$. Also,

$$\begin{aligned} (A^{-1}yA)^{-1}S(A^{-1}yA) &= (A^{-1}yA)^{-1}(A^{-1}SA)(A^{-1}yA) \\ &= A^{-1}y^{-1}SyA = A^{-1}S^{-1}A = S^{-1} . \end{aligned}$$

Therefore, $A^{-1}yA \in N_{GL(2,3)}(\langle S \rangle) - C_{GL(2,3)}(\langle S \rangle)$ where

$$N_{GL(2,3)}(\langle S \rangle) = \langle y, S, ZGL(2, 3) \rangle .$$

Multiplying P on the left by a power of $X(T)$, we may take $A^{-1}yA$ in $\langle y, ZGL(2, 3) \rangle$ of order 4 and $A^{-1}yA \in yZGL(2, 3) = y\langle -I_2 \rangle$. Let $Q \in GL(6, C)$ be the matrix which acts as I_3 on the space where $X(T)$ acts as ωI_3 , and acts as $-I_3$ on the space where $X(T)$ acts as $\omega^{-1}I_3$. Then for $W \in N_{GL(6,C)}(X(\langle T \rangle))$, $W^{-1}(X(T))W = X(T)^a$ and $Q^{-1}W^{-1}QW = (-1)^{[(a-1)/2]}I_6$ with a equal to either 1 or -1 . Therefore, Q normalizes $X(N(\langle T \rangle))$ and $X(N(\langle JT \rangle))$. Also, $Q \in C(J), C(T)$, and

$$C((I_2 \otimes O_2(SL(2, 3))) \oplus I_2) ,$$

and $Q^{-1}Y^{-1}QY = -I_6$. If we are allowed the possibility of replacing P by QP , then we may take $A^{-1}yA = y$. Then, as $\langle y, S \rangle$ is an irreducible two dimensional group on which A acts trivially, A and $A \otimes I_2$ are scalar. As the homomorphism $C(J) \rightarrow U(C(J))$ has kernel J , and $A \otimes I_2$ centralizes $U(N(\langle JT \rangle))$, C centralizes $V(N(\langle JT \rangle))/\langle -I_2 \rangle$. Then C centralizes $V(T) = w \oplus w^{-1}$, and C is diagonal. Let

$$F = 1 \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus 1 \oplus 1 \oplus -1 .$$

Then $V(F)$ is centralized by C . As $V(C(J)) = \langle V(N(\langle JT \rangle)), V(F) \rangle$, C centralizes $V(C(J))/\langle -I_2 \rangle$, and P normalizes $X(C(J))$.

Therefore, $X(JT)$ and $X(N(\langle JT \rangle))$ determine $X(C(J))$ except possibly for conjugation of $X(C(J))$ by a matrix U which is $\pm I_3$ on the homogeneous spaces of $X(T)$. Now $\langle C(J), N(\langle T \rangle) \rangle$ has index in G dividing 35. As $B_0(7)$ has only $\bar{\chi}_0$ with degree < 35 ,

$$G = \langle C(J), N(\langle T \rangle) \rangle .$$

We put $X(N(\langle T \rangle))$ in our normal form. Then $X(JT)$ and $X(\langle NJT \rangle)$ determine $X(C(J))$ within conjugation by U . However,

$$\begin{aligned} U^{-1}\langle X(C(J)), X(N(\langle T \rangle)) \rangle U &= \langle U^{-1}X(C(J))U, U^{-1}X(N(\langle T \rangle))U \rangle \\ &= \langle U^{-1}X(C(J))U, X(N(\langle T \rangle)) \rangle \end{aligned}$$

so the similarity class of the representation is not affected by replacing $X(C(J))$ by $U^{-1}X(C(J))U$. Therefore, there, is at most one unimodular, 6-dimensional, complex, linear group projectively representing a simple group of order $2^7 3^8 35$.

6. Existence of $X(G)$. We shall show that $G_1 = \langle x, D, P \rangle$, where $x = V \oplus \bar{V}$ and $V = \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \bar{\omega} \\ 1 & \bar{\omega} & \omega \end{pmatrix} / (\omega - \bar{\omega})$, $D = \langle \text{all diagonal matrices of order 3 and determinant 1} \rangle$, and $P = \langle \text{all permutation matrices} \rangle$ has a central extension of Z_6 by $U_4(3)$ as a subgroup of index 2. First we show it is finite. In fact, $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus I_4$ has a total of 126 conjugates, $C_1 \cup C_2$, where C_1 consists of 45 monomial matrices and C_2 of 81 conjugates of $z = I_6 - Q/3$ where $Q = (q_{i,j})$ and $q_{i,j} = 1$. $\langle C_1 \rangle$ has no invariant subspaces, so only scalars commute with all conjugates. If S_i are sets of matrices, define $S_1^{-1}S_2S_1 = \{y \mid y = s_1^{-1}s_2s_1 \text{ for } s_i \in S_i\}$. Then $C_2 = D^{-1}zD$. Let $M = DP = PD$. Now $M^{-1}C_1M = C_1$ and $M^{-1}C_2M = M^{-1}(D^{-1}zD)M = M^{-1}zM = D^{-1}(P^{-1}zP)D = D^{-1}zD = C_2$. It only remains show that $x^{-1}(C_1 \cup C_2)x = C_1 \cup C_2$. Let $\{U_i\}$ be the top 3 by 3 blocks of the 9 elements of C_1 whose bottom 3 by 3 block is I_3 . Then $\{U_i\} = -I_3$ {2-elements in the normal subgroup of order 54 of the Hessian group, [1, § 79]}. As the top left 3 by 3 block of x is contained in the Hessian group, conjugation by x permutes these 9 elements. We may reverse the roles of the top left and the bottom right to show that x permutes 9 more elements of C_1 . As $x^{-1}zx$ is a permutation matrix transposing 1 and 4, $x^{-1}zx$ has eigenvalues $-1, 1, 1, 1, 1, 1$. Suppose that $d = \text{diag}(d_1, \dots, d_6) \in D$ with $d_1d_2d_3 = 1$. Then in each row and column of $d^{-1}xd$, and nonzero entries are distinct and have sum 0, or are identical. Then $u_d = (d^{-1}xd)^{-1}z(d^{-1}xd) = I_6 - C_d$ where nonzero entries of C_d are sixth roots of 1. As z and u_d are unitary, u_d has entries 1 or 0 on the diagonal and third roots of 1 off the diagonal and is monomial. Then $u_d \in C_1$ since u_d has eigenvalues $-1, 1, 1, 1, 1, 1$. Therefore, $x^{-1}dxd^{-1}x \in dC_1d^{-1} = C_1$ where d runs through 27 cosets of $\langle wI_6 \rangle$. This gives the other $27 = 45 - 9 - 9$ elements in C_1 and $x^{-1}C_1 \cup C_2x \supset C_1$; $C_1 \cup C_2 \supset xC_1x^{-1} = x^{-1}(x^2C_1x^2)x = x^{-1}C_1x$ as $-I_6x^2 \in P$. It only remains to show that $x^{-1}dxd^{-1}x \in C_2$ where $d_1d_2d_3 = \omega$ or $\bar{\omega}$, say $\bar{\omega}$ without loss of generality. We may find e in $\langle D, \text{diag}(\omega, 1, 1, 1, 1, 1) \rangle$ with $(\omega - \bar{\omega})d^{-1}xde = (a_{i,j})$; $\{a_{1,j}, a_{2,j}, a_{3,j}\} = \{1, \bar{\omega}, \bar{\omega}\}$ counting multiplicity for $j = 1, 2, 3$; and $\{a_{4,j}, a_{5,j}, a_{6,j}\} = \{-1, -\omega, -\omega\}$ for $j = 4, 5, 6$. As $d^{-1}xde$ is unitary, the ± 1 's appear in different rows. Then the product of the nonzero entries in the first and the fourth rows is still -1 , and $e \in D$. Now $(\omega - \bar{\omega})d^{-1}xdeQ$ and $(\omega - \bar{\omega})Qd^{-1}xde$ have all their entries equal to $\bar{\omega} + \bar{\omega} + 1 = -\omega - \omega - 1$. Then $zd^{-1}xde = d^{-1}xdez$, $d^{-1}x^{-1}dxd^{-1}xd = eze^{-1}$, and $x^{-1}dxd^{-1}x = deze^{-1}d^{-1} \in DzD^{-1} = C_2$.

G_1 is primitive since D contains any proper normal reducible subgroup of M and x does not preserve the monomial form of M . Furthermore, G_1 may be made unimodular by replacing odd permutation matrices by their products with iI_6 . As $3^7 \nmid |G_1|$, by [9]'s classification of groups of degree 6, G_1 contains a central extension of Z_6 by $U_4(3)$ as normal subgroup, G . However, G_1 contains an element with eigenvalues $-1, 1, 1, 1, 1, 1$ and G contains no element with eigenvalues $-i, i, i, i, i, i$. By [8], $7^2 \nmid |G_1/Z|$. By [4], $3F, S_7$ is self-centralizing in G_1/Z , otherwise G_1 has a normal p -subgroup not contained in Z for some prime p , a contradiction. Since $[N_G(S_7); C_G(S_7)] = 3$ and $[N_{G_1}(S_7); C_{G_1}(S_7)] \leq 6$, $[G_1:G] \leq 2$ and $[G_1:G] = 2$. For any unimodular finite linear group normalizing $X(G)$, applying this argument to G_2 in place of G_1 shows that $[G_2: X(G)] = 2$, so G_1 is maximal among finite unimodular 6-dimensional complex linear groups normalizing $X(G)$.

7. $LF(3, 4)$. From [9] we may have a six-dimensional group $X(G)$ with $G/Z(G)$ simple of order $2^6 3^2 5$, $\chi(G) \subseteq Q(w)$, and $B_6(5)$ with degree equation: $1 + 63 = 64$. As S_5 is self-centralizing in $\bar{G} = G/Z$ and $B_6(5)$ does not contain the degree 6, $|Z| \neq 1$. If $|Z| = 2$, then $B_6(5)$ contains the degrees 6 and 64 from a 2-block of defect 1, impossible as $64 - 6 = 58$ cannot be a degree. If $|Z| = 3$, then $B_6(5)$ contains the degrees 6 and 63 from a 3-block of defect 1, impossible as $63 + 6 = 69$. Therefore, $|Z| = 6$. Let J be any involution in \bar{G} . Then 0 or $5 = \alpha_{J, J, \tau_5} = |\bar{G}|(1 + \chi_{63}(J)^2/63 - \chi_{64}(J)^2/64)/|C_{\bar{G}}(J)|^2$. Now, χ_{64} has 2-defect 0, so $\chi_{64}(J) = 0$ and $\chi_{63}(J) = 1 - \chi_{64}(J) = 1$. Then $5|C_{\bar{G}}(J)|^2 = 2^6 3^2 5(1 + 1/63) = 2^{12} 5$ and $|C_{\bar{G}}(J)| = 2^6$. Therefore, $C(J)$ has a normal 2-Sylow-subgroup, and by [11], $\bar{G} \approx LF(3, 4)$. As $U_4(3)$ has a subgroup isomorphic to $LF(3, 4)$ and $LF(3, 4)$ has no projective representation of degree ≤ 5 , by § 6, G exists with a representation of degree 6. By private communication with N. Burgoyne, G is unique, and the subgroup of the outer automorphism group with trivial action on Z has order 2. A group $G_1 \triangleright G$ with $[G_1:G] = 2$ comes from the product of a field and a graph automorphism.

APPENDIX.

$G = \text{Some Central Extension of } Z_8 \text{ by } LF(3, 4).$

Element	$\theta = (1 + \sqrt{5})/2$			G/Z		$\phi = (1 + \sqrt{-7})/2$		
	I	π_5	π_7	T	J	F_1	F_2	F_3
Order	1	5	7	3	2	4	4	4
$C(g)$	g	5	7	9	64	16	16	16
	1	1	1	1	1	1	1	1
<u>63</u>	θ	0	0	0	-1	-1	-1	-1
64	-1	1	1	1	0	0	0	0
20	0	-1	-1	2	4	0	0	0
<u>45</u>	0	$-\phi$	0	0	-3	1	1	1
35	0	0	0	-1	3	3	-1	-1
35	0	0	0	-1	3	-1	3	-1
35	0	0	0	-1	3	-1	-1	3

G/Z_2

21	1	0	0	0	5	1	1	1
<u>63</u>	θ	0	0	0	-1	-1	-1	-1
84	-1	0	0	0	4	0	0	0
15	0	1	0	0	-1	3	-1	-1
15	0	1	0	0	-1	-1	3	-1
15	0	1	0	0	-1	-1	-1	3
<u>45</u>	0	$-\phi$	0	0	-3	1	1	1

G/Z_3

I	π_5	π_7	T	J	F_1	F_2	F_3
36	1	1	0	-4	0	0	0
64	-1	1	1	0	0	0	0
<u>28</u>	θ	0	1	4	0	0	0
90	0	-1	0	-2	-2	0	0
<u>10</u>	0	$-\phi$	1	-2	2	0	0
70	0	0	-2	2	2	0	0

G

36	1	1	0	-4	0	0	0
<u>42</u>	$-\theta$	0	0	-2	2	0	0
90	0	-1	0	-2	-2	0	0
<u>60</u>	0	ϕ	0	4	0	0	0
6	1	-1	0	2	2	0	0

$\widehat{U}_4(3), G/Z, \omega^3 = 1, v = \omega - \bar{\omega}.$																	
Element	I	π_3	π_7	J	T	F	T_1	JT	FT	JT_1	JT_2	N_1	N_2	T_2	E	F_1	T_3
Order	1	5	7	2	3	4	3	6	12	6	6	9	9	3	8	4	3
$C(g)$	g	5	7	$2^7 9$	$2^3 6$	96	$2^3 8$	72	12	36	36	27	27	$2^3 3^5$	8	16	81
	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	90	0	-1	10	9	-2	9	1	1	1	1	0	0	9	0	2	0
	640	0	$(-1 + \sqrt{-7})/2$	0	-8	0	-8	0	0	0	0	1	1	-8	0	0	1
	729	-1	1	9	0	-3	0	0	0	0	0	0	0	0	-1	1	0
	35	0	0	3	8	3	8	0	0	0	3	-1	2	-1	-1	-1	-1
	189	-1	0	-3	27	5	0	3	-1	0	0	0	0	0	1	1	0
	896	1	0	0	32	0	-4	0	0	0	0	-1	-1	-4	0	0	-4
	21	1	0	5	-6	1	3	2	-2	-1	-1	0	0	3	-1	1	3
	<u>280</u>	0	0	-8	10	0	10	-2	0	-2	1	1	$2\bar{\omega} - \omega$	1	0	0	1
	35	0	0	3	8	3	-1	0	0	3	0	2	-1	8	-1	-1	-1
	140	0	0	12	5	4	-4	-3	1	0	0	-1	-1	-4	0	0	5
	<u>280</u>	0	0	-8	10	0	1	-2	0	1	-2	$2\bar{\omega} - \omega$	1	10	0	0	1
	560	0	0	-16	-34	0	2	2	0	2	2	-1	-1	2	0	0	2
	315	0	0	11	-9	-1	-9	-1	-1	-1	2	0	0	18	1	-1	0
	315	0	0	11	-9	-1	18	-1	-1	2	-1	0	0	-9	1	-1	0
	420	0	0	4	-39	4	6	1	1	-2	-2	0	0	6	0	0	-3
	210	0	0	2	21	-2	3	5	1	-1	-1	0	0	3	0	-2	3

$G/Z_3, \phi^3 = 1, \phi^2 = -1.$

I	π_5	π_7	J	T	F	T_1	JT	FT	JT_1	JT_2	\bar{N}_1	\bar{N}_2	T_2	E	T_3
20	0	-1	-4	-7	4	2	-1	1	2	2	-1	-1	2	0	2
640	0	$(-1 + \sqrt{-7})/2$	0	-8	0	-8	0	0	0	0	1	1	-8	0	1
120	0	1	8	12	0	-6	-4	0	2	2	0	0	-6	0	3
540	0	1	-12	-27	4	0	3	1	0	0	0	0	0	0	0
896	1	0	0	32	0	-4	0	0	0	0	-1	-1	-4	0	-4
56	1	0	8	2	0	11	2	0	-1	2	-1	2	2	0	2
70	0	0	2	-11	2	7	-1	-1	-1	2	1	$1 + 3\omega$	-2	0	-2
56	1	0	8	2	0	2	2	0	2	-1	2	-1	11	0	2
504	-1	0	8	18	0	-9	2	0	-1	2	0	0	18	0	0
504	-1	0	8	18	0	18	2	0	2	-1	0	0	-9	0	0
70	0	0	2	-11	2	-2	-1	-1	2	-1	$1 + 3\omega$	1	7	0	-2
70	0	0	2	16	2	7	-4	2	-1	-1	1	1	7	0	-2
210	0	0	-10	21	2	3	-1	-1	-1	-1	0	0	3	$2i$	3
630	0	0	-14	-18	-6	9	-2	0	1	1	0	0	9	0	0
560	0	0	16	-34	0	2	-2	0	-2	-2	-1	-1	2	0	2

$G/Z_8, \omega^8 = 1, v = \omega - \bar{\omega} = \sqrt{-3}$.													
I	π_5	π_7	J	T	F	T_1	JT	FT	JT_1	JT_2	N_2	E	F_1
15	0	1	-1	6	3	3	2	0	-1	2	$-v$	1	-1
21	1	0	5	3	1	6	-1	1	2	2	$\bar{\omega}v$	-1	1
729	-1	1	9	0	-3	0	0	0	0	0	0	-1	1
105	0	0	-7	15	5	3	-1	-1	-1	2	$-\bar{\omega}v$	-1	1
105	0	0	9	15	1	3	3	1	3	0	$-\bar{\omega}v$	1	1
384	-1	-1	0	24	0	12	0	0	0	0	$-v$	0	0
360	0		8	-18	0	-9	2	0	-1	2	0	0	0
756	1		-12	27	-4	0	3	-1	0	0	0	0	0
336	1		16	-6	0	6	-2	0	-2	-2	ωv	0	0
210	0		2	3	-2	15	-1	1	-1	2	v	0	-2
105	0		9	-12	1	12	0	-2	0	0	$-\omega v$	1	1
420	0		4	33	4	-6	1	1	-2	-2	$-\omega v$	0	0
945	0		-15	-27	1	0	-3	1	0	0	0	1	1
315	0		-5	-36	3	9	4	0	1	-2	0	-1	-1
630	0		6	9	2	-9	-3	-1	3	0	0	0	-2

$G, v = \omega - \bar{\omega} = \sqrt{-3}$.

I	π_5	π_7	J	T	F	T_1	JT	FT	JT_1	JT_2	N_2	E
6	1	-1	2	-3	2	3	-1	-1	-1	2	$-\omega v$	0
84	-1	0	-4	-15	4	6	-1	1	2	2	v	0
126	1	0	10	18	2	9	-2	2	1	-2	0	0
384	-1	-1	0	24	0	12	0	0	0	0	$-v$	0
336	1	0	-16	-6	0	6	2	0	2	2	ωv	0
120	0	1	8	-6	0	15	2	0	-1	2	$\bar{\omega} v$	0
270	0	$(1 + \sqrt{-7})/2$	-6	27	2	0	-3	-1	0	0	0	0
420	0	0	12	-21	-4	12	-3	-1	0	0	$-\omega v$	0
210	0	0	6	-24	6	-3	0	0	-3	0	ωv	0
840	0	0	-8	-42	0	-3	-2	0	1	-2	$\bar{\omega} v$	0
630	0	0	18	9	2	-9	3	-1	3	0	0	0
840	0	0	-8	12	0	6	4	0	-2	-2	v	0
630	0	0	2	9	-2	-9	-1	1	-1	2	0	$2i$

An Extension of Z_3 by $\widetilde{U}_4(3)$ (faithful characters).

I	π_5	π_7	J	T	F	JT	FT	JT_1	JT_2	E	F_1
36	1	1	4	9	4	1	1	-2	-2	0	0
720	0	-1	16	18	0	-2	0	-2	-2	0	0
729	-1	1	9	0	-3	0	0	0	0	-1	1
45	0	$(-1 + \sqrt{-7})/2$	-3	-9	1	3	1	0	0	-1	1
189	-1	0	-3	27	5	3	-1	0	0	1	1
126	1	0	14	-9	2	-1	-1	2	2	0	2
756	1	0	-12	27	-4	3	-1	0	0	0	0
315	0	0	-5	18	3	-2	0	4	-2	-1	-1
315	0	0	-5	18	3	-2	0	-2	4	-1	-1
630	0	0	6	-45	2	3	-1	0	0	0	-2
315	0	0	11	18	-1	2	2	2	2	1	-1
945	0	0	-15	-27	1	-3	1	0	0	1	1

An Extension of Z_6 by $\widetilde{U}_4(3)$ (faithful characters).

I	π_5	π_7	J	T	F	JT	FT	JT_1	JT_2	E
90	0	-1	-2	-18	6	-2	0	-2	-2	0
126	1	0	10	-9	2	1	-1	-2	4	0
126	1	0	10	-9	2	1	-1	4	-2	0
540	0	1	-12	-27	4	3	1	0	0	0
630	0	0	18	36	2	0	2	0	0	0
1260	0	0	4	-9	-4	1	-1	-2	-2	0
504	-1	0	8	-36	0	-4	0	2	2	0
720	0	-1	-16	18	0	2	0	2	2	0
270	0	$(1 + \sqrt{-7})/2$	-6	27	2	-3	-1	0	0	0
126	1	0	-6	-9	-2	-3	1	0	0	2i

BIBLIOGRAPHY

1. H. F. Blichfeldt, *Finite Collineation Groups*, University of Chicago Press, Chicago, 1917.
2. R. Brauer. *Investigations on group characters*, Ann. of Math. **42** (1941), 936-958.
3. ———, *On groups whose order contains a prime to the first power*, I, II, Amer. J. Math. **64** (1942), 401-440.
4. ———, *Über endlichen lineare Gruppen von Primzahlgrad*, Math. Ann. **169** (1967), 73-96.
5. R. Brauer and H. F. Tuan, *On simple groups of finite order*, I, Bull. Amer. Math. Soc. **51** (1945), 756-766.
6. C. M. Hamill, *On a finite group of order 6,531,840*, Proc. London Math. Soc. **52** (1951), 401-454.
7. B. Huppert, *Lineare auflösbare Gruppen*, Math. Zeit. **67** (1957), 479-518.
8. J. H. Lindsey, II, *A generalization of Feit's theorem*, Trans. Amer. Math. Soc. **155** (1970), 250-271.
9. ———, *Finite Linear groups of degree six* (to appear).
10. H. H. Mitchell, *Determination of all primitive collineation groups in more than four variables which contain homologies*, Amer. J. Math. **36** (1914), 1-12.
11. M. Suzuki, *Finite groups in which the centralizer of an element of order 2 is 2-closed*, Amer. J. Math. (2) **82** (1965), 191-212.
12. J. A. Todd, *The characters of a collineation group in five dimensions*, Proc. Royal Soc. of London (A) **200** (1950).

Received January 28, 1970.

HARVARD UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

George E. Andrews, <i>On a partition problem of H. L. Alder</i>	279
Thomas Craig Brown, <i>An interesting combinatorial method in the theory of locally finite semigroups</i>	285
Yuen-Kwok Chan, <i>A constructive proof of Sard's theorem</i>	291
Charles Vernon Coffman, <i>Spectral theory of monotone Hammerstein operators</i>	303
Edward Dewey Davis, <i>Regular sequences and minimal bases</i>	323
Israel (Yitzchak) Nathan Herstein and Lance W. Small, <i>Regular elements in P.I.-rings</i>	327
Marcel Herzog, <i>Intersections of nilpotent Hall subgroups</i>	331
W. N. Hudson, <i>Volterra transformations of the Wiener measure on the space of continuous functions of two variables</i>	335
J. H. V. Hunt, <i>An n-arc theorem for Peano spaces</i>	351
Arnold Joseph Insel, <i>A decomposition theorem for topological group extensions</i>	357
Caulton Lee Irwin, <i>Inverting operators for singular boundary value problems</i>	379
Abraham A. Klein, <i>Matrix rings of finite degree of nilpotency</i>	387
Wei-Eihn Kuan, <i>On the hyperplane section through a rational point of an algebraic variety</i>	393
John Hathway Lindsey, II, <i>On a six-dimensional projective representation of $PSU_4(3)$</i>	407
Jorge Martinez, <i>Approximation by archimedean lattice cones</i>	427
J. F. McClendon, <i>On stable fiber space obstructions</i>	439
Mitsuru Nakai and Leo Sario, <i>Behavior of Green lines at the Kuramochi boundary of a Riemann surface</i>	447
Donald Steven Passman, <i>Linear identities in group rings. I</i>	457
Donald Steven Passman, <i>Linear identities in group rings. II</i>	485
David S. Promislow, <i>The Kakutani theorem for tensor products of W^*-algebras</i>	507
Richard Lewis Roth, <i>On the conjugating representation of a finite group</i> ...	515
Bert Alan Taylor, <i>On weighted polynomial approximation of entire functions</i>	523
William Charles Waterhouse, <i>Divisor classes in pseudo Galois extensions</i>	541
Chi Song Wong, <i>Subadditive functions</i>	549
Ta-Sun Wu, <i>A note on the minimality of certain bitransformation groups</i>	553
Keith Yale, <i>Invariant subspaces and projective representations</i>	557