

Pacific Journal of Mathematics

INVARIANT SUBSPACES AND PROJECTIVE REPRESENTATIONS

KEITH YALE

INVARIANT SUBSPACES AND PROJECTIVE REPRESENTATIONS

KEITH YALE

Let Γ be a subgroup of the real line R with the discrete topology, and let G be its compact dual group. This paper shows the existence of a (nontrivial) simply invariant subspace of $L^2(G)$ which is not of the form $\varphi H^2(G)$ provided Γ contains at least two rationally independent elements. The proof relies heavily on the existence of a nontrivial local projective representation of the two-dimensional torus.

Helson and Lowdenslager [4] showed the existence of a simply invariant subspace not of the form $\varphi H^2(G)$ in case Γ contains an infinite set of rationally linearly independent elements. We use the correspondence introduced in [4] between simply invariant subspaces and cocycles but in contrast to [4] we use nontrivial local projective multipliers to show that the appropriate cohomology group is nontrivial.

The connection between invariant subspaces and cocycles is discussed in § 2 and in § 3 we will give a quotient group argument which allows us to reduce the general problem to its specialization on the two-dimensional torus. Sections 4 and 5 relate the notion of projective representation with a cocycle and it is shown that a nontrivial projective representation gives rise to a cocycle whose corresponding subspace is not of the form $\varphi H^2(G)$.

2. Preliminaries. Let G be an arbitrary locally compact Abelian group dual to Γ and let A be a continuous one-parameter subgroup of G which we also denote by $\{e_t | t \text{ in } R\}$. Haar measure in G will be denoted by dx and will be normalized to have total mass one in case G is compact. As usual, a.e. (x) means for all but a set of Haar measure zero. A (Borel) function φ on G is said to be *unitary* in case $\varphi(x)$ has modulus one a.e. (x) .

DEFINITION. A function A on $A \times G$ is said to be a *cocycle* on G in case:

$$(2.1) \quad A(e_t, \cdot) \text{ is a unitary function for each } e_t \text{ in } A,$$

$$(2.2) \quad A(e_t + e_u, x) = A(e_u, x)A(e_t, x - e_u) \text{ for all } e_t, e_u \text{ in } A$$

and a.e. (x) , and

$$(2.3) \quad A \text{ is strongly continuous in the sense that } A(e_t, \cdot)f$$

is a continuous function from R into $L^2 = L^2(G)$ for f in L^2 .

Cocycles of the form

$$(2.4) \quad A(e_i, x) = \varphi(x)/\varphi(x - e_i), \quad \text{all } e_i \text{ in } A, \text{ a.e. } (x)$$

for some unitary function φ are called *coboundaries*. We will frequently denote $A(e_i, x)$ by $A(t, x)$.

If λ is in Γ we let χ_λ be the character on G defined by $\chi_\lambda(x) = x(\lambda)$ for all x in G ; the corresponding unitary representation V_λ of Γ is given by

$$(2.5) \quad V_\lambda(\lambda)f(x) = \chi_\lambda(x)f(x)$$

for all f in L^2 . Any bounded operator on L^2 which commutes with all the $V_\lambda(\lambda)$ is necessarily a multiplication by a function in L^∞ . Let U_0 be the unitary representation of G defined by

$$(2.6) \quad U_0(x)f(y) = f(y - x)$$

for all f in L^2 .

For the remainder of this section we will let Γ be a subgroup of the real line R . Let G be the compact Abelian group dual to the discretely topologized Γ . A closed subspace \mathcal{M} of L^2 is said to be *simply invariant* in case $V_\lambda(\lambda)\mathcal{M} \subseteq \mathcal{M}$ if and only if $\lambda \geq 0$. The Hardy space H^2 consists of those functions f in L^2 whose Fourier transforms $\hat{f}(\lambda) = \int \chi_{-\lambda}(x)f(x)dx$ vanish for $\lambda < 0$. Subspaces of the form $\mathcal{M} = \varphi H^2 = \{\varphi f: f \text{ in } H^2\}$ where φ is a unitary function are simply invariant and in the case where G is a circle all simply invariant subspaces are of this form.

In order to avoid the rather special circle group we will henceforth suppose that Γ is dense in R . The characters e_i defined by $e_i(\lambda) = \exp(it\lambda)$ are distinct and provide a continuous one-parameter dense subgroup A of G . A correspondence is exhibited in [3, 4] between simply invariant subspaces \mathcal{M} (suitably normalized) and cocycles A in such a way that $\mathcal{M} = \varphi H^2$ if and only if A is the coboundary (2.4). We therefore wish to construct cocycles which are not coboundaries.

If A is a coboundary then A can be extended from $A \times G$ to $G \times G$ so that (2.4) remains valid with t replaced by an arbitrary y in G and conversely. Moreover, the multiplication operator $A(y, \cdot)$ is the strong operator limit of a sequence $A(t_n, \cdot)$ where e_{t_n} tends to y in G ; this observation will be useful later. Equivalently, A is a coboundary if and only if the unitary representation $U(t) = A(t, \cdot)U_0(t)$ can be extended from A to a (strongly continuous) unitary representation of G . A cocycle was constructed in [4] (in case Γ is suitably large) for which the unitary representation did not extend to G . However, it is conceivable that $U(t)$ might extend to a (local) projective representation of G ; this idea is turned around and will be used to extract cocycles

from projective representations.

There is a superficial answer to our problem in case Γ is not all of R for then there are trivial cocycles which are not coboundaries. For example, let $A(t, x) = \exp(-it\lambda)$ for some fixed real λ not in Γ . If λ were in Γ then A would be the coboundary with unitary function χ_λ but with λ not in Γ there is no unitary function φ such that $\exp(-it\lambda) = \varphi(x)/\varphi(x - e_t)$. Conversely, if A is a cocycle which is constant a.e. (x) for each t (the null set depending upon t), then $A(t, x) = \exp(-it\lambda)$, a.e. (x) for some fixed λ in R . We will call cocycles of this form *constant cocycles*. Consequently the nontrivial problem [3, p.149] is to find cocycles which are not products of constant cocycles and coboundaries.

The cocycles defined in [3] were measurable functions on $A \times G$ but we will have no need for cocycles to be product measurable. Anyway, one can pass from one version to another [3, p.145], [2]. Also we have departed from [3] by making an insignificant sign change in our definition of cocycle.

3. Reduction to the torus. Suppose that $\Gamma_0 \subseteq \Gamma$ are subgroups of the discrete real line and let G_0 and G be their compact dual groups. To each cocycle A_0 on G_0 we will associate a cocycle A on G in such a way that if A is the product of a constant cocycle and a coboundary then so is A_0 . Since the two-dimensional torus T^2 is dual to the group of lattice points Z^2 and Z^2 is isomorphic to a subgroup $\Gamma_0 \subseteq \Gamma$ of any group $\Gamma \subseteq R$ with at least two independent elements it will be sufficient to construct a cocycle on T^2 which is not the product of a constant cocycle and a coboundary.

Define a closed subgroup $H = \{x \text{ in } G \mid \chi_\lambda(x) = 1 \text{ for all } \lambda \text{ in } \Gamma_0\}$ of G so that G_0 can be identified with G/H . Let π be the usual quotient map from G onto G/H and let e_t and ε_t be the previously defined one-parameter groups A and A_0 in G and G_0 . One can verify $\pi(e_t) = \varepsilon_t$ by noting that ε_t is the restriction of e_t from A to A_0 .

If A_0 is a cocycle on G_0 we define a cocycle A on G by

$$(3.1) \quad A(e_t, x) = A_0(\varepsilon_t, \pi(x))$$

for all (e_t, x) in $A \times G$.

For each t in R the measurable function $A(e_t, \cdot)$ on G is certainly unitary because $\pi^{-1}(S)$ is a null set in G whenever S is a null set in G/H . The cocycle identity (2.2) is easy enough to verify with the aid of $\pi(e_t) = \varepsilon_t$ so all that remains is the strong continuity.

Let the Haar measures dx and dx_0 in G and G/H both be normalized to have total mass one. There is a normalization for the Haar measure $d\xi$ on H such that

$$(3.2) \quad \int_G f(x)dx = \int_{G/H} \left(\int_H f(x + \xi) d\xi \right) dx_0$$

for all f in $L^1(G)$.

Let f be in $L^2(G)$ and put

$$g(x_0) = \int_H |f(x + \xi)|^2 d\xi$$

where $x_0 = \pi(x)$. A straight-forward computation with (3.2) shows that $A(e_t, \cdot)f$ moves continuously in $L^2(G)$ as t varies because $A_0(e_t, \cdot)\sqrt{g}$ moves continuously in $L^2(G/H)$.

THEOREM. *If A is the product of a constant cocycle and a coboundary then so is A_0 .*

Proof. For some constant cocycle C and some unitary function φ on G we have

$$C(t)A(t, x) = \varphi(x)/\varphi(x - e_t)$$

for each real t and almost all x .

It is advantageous to normalize by choosing λ in R such that $\int \chi_\lambda(x)\varphi(x)dx$ does not vanish and putting $\psi = \chi_\lambda\varphi$. The cocycle $B = \chi_\lambda CA$ is really the coboundary.

$$(3.3) \quad B(t, x) = \psi(x)/\psi(x - e_t)$$

and we have $B_0 = \chi_\lambda CA_0$. Consequently it is sufficient to show that B_0 is a coboundary and we will do this by arguing that ψ must be constant on cosets of H .

Since $B(t, x) = B_0(t, \pi(x))$ it follows that $B(t, x) = B(t, x + h)$ for all real t and all (x, h) in $G \times H$. Now the coboundary B can be extended to $G \times G$ and, in fact, $B(y, \cdot)$ is a limit in $L^2(G)$ of a sequence $B(t_n, \cdot)$ where e_{t_n} goes to y in G . Therefore, passing to a subsequence if necessary, $B(t_n, x)$ tends to $B(y, x)$ for almost all x and we can conclude

$$(3.4) \quad B(y, x) = B(y, x + h)$$

for all y in G , h in H and almost all x in G .

From (3.3) (valid now for t replaced by any element in G) and (3.4) we have

$$(3.5) \quad \psi(x + \xi) = B(h, x)\psi(x + \xi - h)$$

for every ξ in H and almost all x in G . Integrating this last expression with respect to Haar measure $d\xi$ on H we find

$$\int_H \psi(x + \xi) d\xi = B(h, x) \int_H \psi(x + \xi) d\xi.$$

Now $\int_H \psi(x + \xi) d\xi$ does not vanish since $\int_G \psi(x) dx$ is not zero (consider (3.2)) and so we may conclude $B(h, x) = 1$ for all h in H and almost all x in G .

It follows from (3.5) that ψ is constant on cosets of H and so we can define a unitary function ψ_0 on G/H by $\psi_0(\pi(x)) = \psi(x)$. Clearly B_0 is a coboundary determined by ψ_0 . That completes the proof.

4. Projective representations and projective cocycles. Let G be a locally compact Abelian group. A strongly continuous function U from G into the unitary operators on some Hilbert space is said to be a *projective representation* if

$$(4.1) \quad U(x)U(y) = \omega(x, y)U(x + y)$$

for some function ω of modulus one and if $U(0) = 1$. We say that ω is the *multiplier* of the representation and it is not difficult to show that it satisfies the identity $\omega(x, y)\omega(x + y, z) = \omega(y, z)\omega(x, y + z)$ and the normalizing condition $\omega(x, 0) = \omega(0, x) = 1$. Moreover, ω is continuous on $G \times G$. Conversely, given a function ω with these properties one can construct a projective representation U_ω with multiplier ω . Indeed, define U_ω on L^2 by

$$(4.2) \quad U_\omega(x)f(y) = \omega(x, y - x)f(y - x).$$

The projective representation U_ω is of the form

$$(4.3) \quad U_\omega(x) = A_\omega(x, \cdot)U_0(x)$$

where $A_\omega(x, y) = \omega(x, y - x)$ is a function of modulus one on $G \times G$. The (projective) group property of U_ω implies that

$$(4.4) \quad \omega(x, y)A_\omega(x + y, z) = A_\omega(x, z)A_\omega(y, z - x)$$

and the strong continuity of U_ω implies that $A_\omega(x, \cdot)$ is a strongly continuous operator valued function in x .

Observe that A_ω differs from the ordinary cocycle (§ 2) in two respects; first, A_ω is a function on $G \times G$ instead of merely on $A \times G$, and, secondly, (4.4) replaces (2.2). We say that A_ω is a *projective cocycle*.

We say that ω is trivial if

$$(4.5) \quad \omega(x, y) = p(x)p(y)/p(x + y)$$

for some continuous function p of modulus one on G . In this case any projective representation U with multiplier ω can be made into an

ordinary representation merely by multiplying $U(x)$ by $p(x)$. The product of two multipliers is again a multiplier and two multipliers whose quotient is trivial are said to be *equivalent*.

If ω and σ are equivalent multipliers so that

$$(4.6) \quad \omega(x, y)/\sigma(x, y) = p(x)p(y)/p(x + y)$$

then a direct computation will give

$$(4.7) \quad A_\omega(x, y)/A_\sigma(x, y) = p(x)(\varphi(y)/\varphi(y - x))$$

where $\varphi(y) = 1/p(y)$. In particular if ω is trivial then A_ω is p times a coboundary and conversely.

Now suppose that G has a continuous one-parameter subgroup $A = \{e_t | t \in R\}$ and let A_ω be a projective cocycle on G with U_ω the corresponding projective representation as given by (4.3). We wish to extract an ordinary cocycle A from A_ω in such a way that A will not be the product of a constant cocycle and a coboundary if ω is a nontrivial multiplier.

Restrict U_ω to A so that it is a projective representation of the reals. It follows that (see the last paragraph of this section) U_ω is equivalent to an ordinary representation U given by

$$(4.8) \quad U(e_t) = p(e_t) U_\omega(e_t)$$

where

$$(4.9) \quad \omega(e_t, e_u) = p(e_t)p(e_u)/p(e_t + e_u)$$

for some continuous function p on A and for all $e_t, e_u \in A$. Observe that U satisfies the Weyl commutation relation

$$(4.10) \quad U(e_t) V_0(\lambda) = \chi_\lambda(-e_t) V_0(\lambda) U(e_t)$$

because U_ω does.

Consequently the operator $U(e_t) U_0(-e_t)$ commutes with all the $V_0(\lambda)$ so that

$$(4.11) \quad U(e_t) = A(e_t, \cdot) U_0(e_t), e_t \in A,$$

for some ordinary cocycle A .

From (4.8) and (4.11) we see that

$$(4.12) \quad A(e_t, x) = p(e_t) A_\omega(e_t, x)$$

for all $e_t \in A$ and a.e. (x) .

We say that A is the cocycle induced by A_ω ; it is uniquely determined up to a constant cocycle factor. If A is the product of a constant cocycle $e^{it\lambda}$ and a coboundary $\varphi(x)/\varphi(x - e_t)$ then (4.12) and (4.7) imply that ω is trivial.

This analysis will have to be refined to yield the desired result on the torus T^2 for T^2 has no nontrivial multipliers. However, there are $\frac{1}{2}n(n-1) + 1$ inequivalent *local* multipliers on T^n or R^n as shown by Bargmann [1] and local multipliers are sufficient for our purposes. Notice, in particular, that R has no nontrivial local projective representations.

5. Local multipliers and cocycles on T^2 . A *local projective* multiplier ω on the torus T^2 is a continuous function on some neighborhood $\mathcal{N} \times \mathcal{N}$ of the identity in $T^2 \times T^2$ which satisfies the same functional equation and normalizing condition as a multiplier whenever x, y and $x + y$ belong to \mathcal{N} . Unfortunately (4.3) cannot be used to define a local projective representation U_ω , or, equivalently, a local projective cocycle A_ω . We must resort to an *ad hoc* construction of U_ω starting from a specific nontrivial local projective multiplier ω . We can then extract a cocycle from U_ω in much the same manner as in § 4 and it is a matter of detail to prove that A is not the product of a constant cocycle and a coboundary.

Let T^2 be realized as the square $[-\pi, \pi] \times [-\pi, \pi]$ with the opposite edges identified and let \mathcal{N} be the open neighborhood $(-\pi, \pi) \times (-\pi, \pi)$ of the identity. For a one-parameter subgroup A we will take the familiar winding line with irrational slope α .

Define ω on $\mathcal{N} \times \mathcal{N}$ by

$$(5.1) \quad \begin{aligned} \omega(x, y) &= \exp i((x_2 - \alpha x_1)y_1 - (y_2 - \alpha y_1)x_1) \\ &= \exp i(x_2y_1 - y_2x_1) \end{aligned}$$

where $x = (x_1, x_2)$, $y = (y_1, y_2)$ with $-\pi < x_i, y_i < \pi$. This is the canonical example of a nontrivial local projective multiplier on T^2 [1].

Since the complement of \mathcal{N} is a null set we can regard $\omega(x, \cdot)$ as a unitary function on T^2 for each fixed $x \in \mathcal{N}$. Now put $A_\omega(x, y) = \omega(x, y - x)$ whenever $x \in \mathcal{N}$ and $y \in \mathcal{N} + x$. Then $A_\omega(x, \cdot)$ is a unitary function on T^2 for each fixed $x \in \mathcal{N}$ (the exceptional null set depends upon x). For $x \in \mathcal{N}$ we define the unitary operator $U_\omega(x)$ by

$$(5.2) \quad U_\omega(x) = A_\omega(x, \cdot) U_0(x).$$

It is easily verified that U_ω is a strongly continuous operator valued function on \mathcal{N} .

We will now extract a cocycle A from A_ω even though $A_\omega(x, \cdot)$ is *not* defined for all x . The discussion parallels that of § 4 and will only be given in outline.

Let A_1 denote the connected segment of $A \cap \mathcal{N}$ (relative to the ordinary real line topology on A) which contains the identity and choose a proper segment A_0 of A_1 such that $0 \in A_0 \subseteq A_0 + A_0 \subseteq A_1$.

For $x, y \in A_0$, U_ω satisfies (4.1) so that U_ω is a local projective representation of the reals. Consequently U_ω is equivalent to a local ordinary representation U ; this means that equations (4.8) and (4.9) hold for some continuous function p on A_1 (say) and for all $e_i, e_u \in A_0$. The local representation U can be extended to a representation U (keeping the same notation) of A [5, Th. 63] which must satisfy the Weyl commutation relation (4.10). Exactly as before we have $U(e_i) = A(e_i, \cdot) U_0(e_i)$, $e_i \in A$, for some ordinary cocycle A . We say that A is the cocycle induced by A_ω ; notice that

$$(5.3) \quad A(e_i, x) = p(e_i) A_\omega(e_i, x)$$

holds only for $e_i \in A_0$, a.e. (x) .

We will now show that A is not the product of a constant cocycle C and a coboundary. If, on the contrary, A is such a product, then

$$(5.4) \quad A_\omega(e_i, x) = \bar{p}(e_i) (\varphi(x) / \varphi(x - e_i))$$

holds for all $e_i \in A_0$, a.e. (x) where we have relabeled the continuous function C/p on A_1 by \bar{p} . In terms of the unitary operators U_ω and $U(y) = (\varphi(\cdot) / \varphi(\cdot - y)) U_0(y)$, $y \in T^2$, equation (5.4) becomes $U_\omega(e_i) = \bar{p}(e_i) U(e_i)$ for all $e_i \in A_0$.

We wish to extend p from A_0 to $A \cap \mathcal{N}$ in such a way that (5.4) remains valid. A continuity argument will then enable us to extend p from $A \cap \mathcal{N}$ to \mathcal{N} and this will imply that ω is trivial.

To extend p from A_0 to $A \cap \mathcal{N}$ let $y \in A \cap \mathcal{N}$ so that $y \in MA_0 = \{Me_i | e_i \in A_0\}$ for some integer $M > 0$. Thus $e_i = yM \in A_0$ and suppose, for the moment, that $ne_i \in \mathcal{N}$ for all $n \leq M$. Then

$$\begin{aligned} U(y) &= U(Me_i) = (U(e_i))^M \\ &= (p(e_i) U_\omega(e_i))^M \\ &= [(p(e_i))^M \prod_{k=1}^{M-1} \omega(e_i, (M-k)e_i)] U_\omega(y) \end{aligned}$$

and we can define $p(y)$ to be the value of the expression in the brackets which obviously is independent of the representation $y = Me_i$.

This definition of $p(y)$ is valid whenever $(M-k)e_i$ is in the domain of $\omega(e_i, \cdot)$, i.e., whenever $ne_i \in \mathcal{N}$ for all $0 \leq n \leq M$. For each M there are only finitely many $y \in MA_0$ such that $ne_i \notin \mathcal{N}$ for some $0 \leq n \leq M$. For these exceptional values we can define $p(y)$ by continuity (relative to the usual real line topology on A) so that

$$(5.5) \quad U(y) = p(y) U_\omega(y)$$

holds for all $y \in A \cap \mathcal{N}$, or, equivalently, so that (5.4) holds for all e_i in $A \cap \mathcal{N}$.

To extend p from $A \cap \mathcal{N}$ to \mathcal{N} we need only note that $A \cap \mathcal{N}$ is dense in \mathcal{N} . Let $y \in \mathcal{N}$ and choose a sequence $y_n \in A \cap \mathcal{N}$ which

converges to y . Hence $p(y_n)I = U(y_n)U_\omega(-y_n)$ tends strongly to

$$U(y)U_\omega(-y)$$

and this limit must be of the form $p(y)I$. Alternately, $U(y)U_\omega(-y)$ is a multiple of the identity for each y in \mathcal{N} because it commutes with all bounded operators when y varies over a dense subset of $A \cap \mathcal{N}$. We have now constructed a continuous function p on \mathcal{N} such that (5.5) holds for all y in \mathcal{N} . Since U_ω is a nontrivial local projective representation of \mathcal{N} this is a contradiction. Hence the induced cocycle A cannot be the product of a constant cocycle and a coboundary. That completes the proof.

An interesting question remains. If A is a cocycle on T^2 can one find a local projective cocycle A_ω which induces A ? An affirmative answer should enable one to settle some of the open function theoretic questions on T^2 .

The author gratefully acknowledges useful conversations with Professors F. Forelli, J. E. Gilbert and H. Helson.

REFERENCES

1. V. Bargmann, *On unitary ray representations of continuous groups*, Ann. of Math. **59** (1954), 1-46.
2. T. W. Gamelin, *Uniform Algebras*, Prentice Hall, Englewood Cliffs. N. J., 1969.
3. H. Helson, *Compact groups with ordered duals*, Proc. London Math. Soc. (3) **14A** (1965), 144-156.
4. H. Helson and D. Lowdenslager, *Invariant subspaces*, Proc. Int. Symp. on Linear Spaces, Jerusalem (1961), 251-262.
5. L. Pontrajagin, *Topological Groups*, Princeton, 1939.

Received April 22, 1969.

UNIVERSITY OF MONTANA
MISSOULA, MONTANA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

George E. Andrews, <i>On a partition problem of H. L. Alder</i>	279
Thomas Craig Brown, <i>An interesting combinatorial method in the theory of locally finite semigroups</i>	285
Yuen-Kwok Chan, <i>A constructive proof of Sard's theorem</i>	291
Charles Vernon Coffman, <i>Spectral theory of monotone Hammerstein operators</i>	303
Edward Dewey Davis, <i>Regular sequences and minimal bases</i>	323
Israel (Yitzchak) Nathan Herstein and Lance W. Small, <i>Regular elements in P.I.-rings</i>	327
Marcel Herzog, <i>Intersections of nilpotent Hall subgroups</i>	331
W. N. Hudson, <i>Volterra transformations of the Wiener measure on the space of continuous functions of two variables</i>	335
J. H. V. Hunt, <i>An n-arc theorem for Peano spaces</i>	351
Arnold Joseph Insel, <i>A decomposition theorem for topological group extensions</i>	357
Caulton Lee Irwin, <i>Inverting operators for singular boundary value problems</i>	379
Abraham A. Klein, <i>Matrix rings of finite degree of nilpotency</i>	387
Wei-Eihn Kuan, <i>On the hyperplane section through a rational point of an algebraic variety</i>	393
John Hathway Lindsey, II, <i>On a six-dimensional projective representation of $PSU_4(3)$</i>	407
Jorge Martinez, <i>Approximation by archimedean lattice cones</i>	427
J. F. McClendon, <i>On stable fiber space obstructions</i>	439
Mitsuru Nakai and Leo Sario, <i>Behavior of Green lines at the Kuramochi boundary of a Riemann surface</i>	447
Donald Steven Passman, <i>Linear identities in group rings. I</i>	457
Donald Steven Passman, <i>Linear identities in group rings. II</i>	485
David S. Promislow, <i>The Kakutani theorem for tensor products of W^*-algebras</i>	507
Richard Lewis Roth, <i>On the conjugating representation of a finite group</i> ...	515
Bert Alan Taylor, <i>On weighted polynomial approximation of entire functions</i>	523
William Charles Waterhouse, <i>Divisor classes in pseudo Galois extensions</i>	541
Chi Song Wong, <i>Subadditive functions</i>	549
Ta-Sun Wu, <i>A note on the minimality of certain bitransformation groups</i>	553
Keith Yale, <i>Invariant subspaces and projective representations</i>	557