

Pacific Journal of Mathematics

AN ELEMENTARY DEFINITION OF SURFACE AREA IN E^{n+1} FOR SMOOTH SURFACES

LOUIS I. ALPERT AND L. V. TORALBALLA

AN ELEMENTARY DEFINITION OF SURFACE AREA IN E^{n+1} FOR SMOOTH SURFACES

LOUIS I. ALPERT AND L. V. TORALBALLA

The present paper concerns the difficulty which one encounters in text books of Advanced Calculus of giving a simple and elementary definition of area of a smooth non-parametric surface in E^{n+1} such that, within the same elementary framework, one can then prove that the area so defined is equal to the classical area integral.

The authors were first made aware of the considerable interest of such a task in 1955 with the publication of Angus Taylor's now classic textbook "Advanced Calculus". The following statement is taken from page 384 of this book:

"It is logically and aesthetically desirable to have a definition of surface area which is directly geometric, and which does not put too many restrictions on the surface. A good definition ought not to depend upon the method of representing the surface analytically, and should not be limited to smooth surfaces. The demand for such a definition poses a very difficult problem, however. It may surprise the student to know that the problem has occupied the attention of many able mathematicians over the last fifty years, and that the end of research on the question is not yet in sight."

In the present paper we present an idea which seems to answer the questions raised by Angus Taylor for surfaces $S: z=f(x_1, \dots, x_n)$, which are continuous with their first order partial derivatives. The idea is to develop a scheme for the construction of sequences of suitably chosen polyhedra inscribed within the given surface, such that the corresponding sequences of the polyhedral areas converge to the classical area integral for the surface, and hence to the Lebesgue area of S .

In previous papers [1], [7] we discussed our definition of area for surfaces $S: z=f(x_1, x_2)$. In [7] we took in consideration surfaces $z=f(x_1, x_2)$ with f continuous with its first order partial derivatives. In [1] we gave a necessary and sufficient condition in order that for a surface $z=f(x_1, x_2)$ there are sequences of inscribed polyhedra satisfying the requirements of our definitions (see [1]).

J. A. Serret [6] in 1868 proposed a geometric definition of area, but H. A. Schwartz [5] in 1882 proved that Serret's definition was incorrect. Other geometric definitions of area and constructions have been proposed, and we mention here for example the ones of S. Kempisty [3] for surfaces $S: z=f(x_1, x_2)$ with f absolutely continuous in the sense of Tonelli. For general expositions concerning area, in particular, Le-

2. **The n -hedra.** Given a set of $n + 1$ points in \mathcal{E}^{n+1} , if the matrix of the coordinates of these $n + 1$ points is of rank n , this set determines an n -hedron, or n -simplex. This is the closed convex subset of \mathcal{E}^{n+1} which is bounded by the $n + 1$ $(n - 1)$ -hyperplanes determined by the given set of $n + 1$ points. An n -hedron determines the n -hyperplane in which it lies.

Given two vectors U and V in \mathcal{E}^{n+1} , the angle $\alpha = (U, V)$ between U and V is determined from the equation $U \cdot V = (U)(V)\cos \alpha$. Given two n -hyperplanes in \mathcal{E}^{n+1} by their dihedral angle we shall mean the acute angle between their normals. An n -hedron n of whose faces ($n-1$ hedras) are at right angles is called a right n -hedron. Given an n -hedron T in \mathcal{E}^{n+1} , we define the area of T to be the n -dimensional volume of T in the standard manner.

3. **Projections.** We distinguish x_{n+1} and call it z . Given an n -hedron T in E^{n+1} , its projection on the (x_1, x_2, \dots, x_n) hyperplane need not be an n -hedron. This occurs, for instance, if T is orthogonal to the hyperplane. We assume here that T lies on an hyperplane $H: z = c + m_1x_1 + \dots + m_nx_n$. Then, the projection of T on the (x_1, x_2, \dots, x_n) hyperplane, or $\text{Proj } T$, is also an n -hedron. If α is the dihedral angle between the hyperplane H determined by T and the (x_1, x_2, \dots, x_n) hyperplane, A is the area of T , and A' is the area of $\text{Proj } T$, then $A = A' \sec \alpha$, where

$$\sec \alpha = (1 + m_1^2 + \dots + m_n^2)^{1/2}.$$

4. **Surfaces in \mathcal{E}^{n+1} .** Let E be an open and connected set in the (x_1, x_2, \dots, x_n) hyperplane such that its closure \bar{E} is capable of being decomposed as the union of n -hedra in the natural manner. We say that \bar{E} is polyhedral. Let f be a real-valued function defined and continuous on \bar{E} . The locus in \mathcal{E}^{n+1} of the points $(x_1, x_2, \dots, x_n, z)$, where $z = f(x_1, x_2, \dots, x_n)$, a function having \bar{E} for domain, is called an n surface in \mathcal{E}^{n+1} , or more briefly a surface. We wish to give a definition of the area of this surface in the case where f is continuously partially differentiable on \bar{E} . We refer to such a surface as a continuously partially differentiable surface.

Let $\Gamma: x_1 = F_1(t), \dots, x_n = F_n(t), a \leq t \leq b$, be any parametric curve in \bar{E} passing through (x_1^0, \dots, x_n^0) for $t = t_0$. Then its image on S , or

$$C: x_1 = F_1(t), \dots, x_n = F_n(t), z = f[F_1(t), \dots, F_n(t)],$$

$a \leq t \leq b$, is a curve on C passing through Q for $t = t_0$. Assuming that each dx_i/dt exists and is continuous on $[a, b]$, it follows that there exists a tangent vector v to Γ at (x_1^0, \dots, x_n^0) , and a tangent

Vector V to C at Q . If $\Gamma_1, \dots, \Gamma_n$ are n such curves in \bar{E} , if C_1, \dots, C_n are the corresponding curves on S , and if we have chosen the curves Γ in such a way that the n vectors v_1, \dots, v_n are linearly independent, then the corresponding n tangent vectors V_1, \dots, V_n determine an n -hyperplane H in E^{n+1} .

One shows that for all such sets of curves in S , the corresponding n -hyperplane is unique. We refer to its normal line as the normal to S at Q .

If T is an n -hedron all of whose vertices are in S , we say that T is inscribed in S . By $D(T)$, the *deviation* of S on T , we mean the supremum of the set of the acute angles between the normal to T and the normals to the portion of S which is subtended by T (i.e., the portion of S whose projection on the x_1, x_2, \dots, x_n hyperplane is identical to that of T).

Let $\{P_1, P_2, \dots, P_m\}$ be the vertices of a decomposition of \bar{E} into a finite set of n -hedra. For each i , let $Q_i = f(P_i)$. The set

$$\{Q_1, Q_2, \dots, Q_m\}$$

determines a polyhedron which is inscribed on S . This polyhedron is composed of a finite set of n -hedra which are inscribed on S . By the *norm* of such a polyhedron we mean the largest of the diameters of its faces. By the *deviation norm* of the polyhedron we mean the largest of the deviations on its faces. By the area of this polyhedron we mean the sum of the areas of the n -hedra which compose it. We refer to these n -hedra as the faces of the polyhedron.

5. **The geometric basis.** We make use of the following additional properties of \mathcal{E}^{n+1} .

(a) Let U_1, U_2, \dots, U_n be any n vectors in \mathcal{E}^{n+1} , such that $|\cos(U_i, U_j)| < k$, $0 < k < 1$, for every $i \neq j$. Then for each $\varepsilon > 0$ there exists $\delta > 0$ such that if U'_1, U'_2, \dots, U'_n are any n vector such that $|\sin(U_i, U'_i)| < \delta$, for each i , then

$$|\sin(U_1 \times U_2, \dots \times U_n, U'_1 \times U'_2 \times \dots \times U'_n)| < \varepsilon.$$

(b) Let $P \in \bar{E}$. Let U be any vector in the x_1, x_2, \dots, x_n plane. We define the directional derivative of f in the direction U in the standard manner.

Under the hypothesis that f is continuously partially differentiable on \bar{E} , then the directional derivative of f is uniformly continuous on \bar{E} , i.e., for each $\varepsilon > 0$ there exists $\delta > 0$ such that if $(x'_1, x'_2, \dots, x'_n)$ and $(x''_1, x''_2, \dots, x''_n)$ are in \bar{E} and $\rho((x'_1, x'_2, \dots, x'_n), (x''_1, x''_2, \dots, x''_n)) < \delta$,

then the absolute value of the difference between the directional derivatives at $(x'_1, x'_2, \dots, x'_n)$ and $(x''_1, x''_2, \dots, x''_n)$ in the direction of the vector from the first point to the latter, is less than ε .

The directional derivative is uniformly Lipschitzian on \bar{E} .

(c) There exist positive numbers k and δ , $k < 1$, such that if P, P_1 and P_2 are any three distinct points in \bar{E} such that

$$(1) \quad \rho(P, P_1) < \delta$$

$$(2) \quad \rho(P, P_2) < \delta \text{ and}$$

$$(3) \quad \cos \overrightarrow{(PP_1, PP_2)} = 0,$$

then $\cos \overrightarrow{(QQ_1, QQ_2)} < k$, where $Q = f(P)$, $Q_1 = f(P_1)$, $Q_2 = f(P_2)$.

(d) Let P_1 and P_2 be any two distinct points in \bar{E} , $Q_1 = f(P_1)$ and $Q_2 = f(P_2)$. Let $\overline{P_1P_2}$ be the linear interval determined by P_1 and P_2 and $\overline{Q_1Q_2}$ be the linear interval determined by Q_1 and Q_2 . Let the curve $C = f(\overline{P_1P_2})$. Then there exists a point R in C such that the tangent line to C at R is parallel to $\overline{Q_1Q_2}$.

(e) With the notation as in (d), let the deviation $D(P_1, P_2)$ denote the supremum of the acute angles ϕ between $\overline{Q_1Q_2}$ and any tangent line to C . Then for every $\varepsilon > 0$, there exists $\delta > 0$ such that if $0 < \rho(P_1, P_2) < \delta$, then $D(P_1, P_2) < \varepsilon$.

(f) For each $\varepsilon > 0$ there exists $\delta > 0$ such that if P_1 and P_2 are any two distinct points of \bar{E} such that $\rho(P_1, P_2) < \delta$, then $\psi < \varepsilon$, where ψ is the acute angle between the normals to S at $f(P_1)$ and $f(P_2)$.

(g) If \bar{E} is polyhedral, it can be decomposed into a set of n -hedra each of which is a right n -hedron. Moreover, for each real number r , there exists a decomposition of \bar{E} into a set of right n -hedra, the diameter of each of which is less than r .

We now proceed to the main theory.

We consider infinite sequences of polyhedra inscribed on S . A sequence (Π_1, Π_2, \dots) of such polyhedra is said to be a *proper* sequence if the corresponding sequence (N_1, N_2, \dots) of norms and the corresponding sequence (ϕ_1, ϕ_2, \dots) of deviation norms both converge to zero.

We now give our definition of the area of the surface $S = f(\bar{E})$, where f is continuously partially differentiable on \bar{E} . If to every proper sequence (Π_1, Π_2, \dots) of polyhedra inscribed on S the corresponding sequence (A_1, A_2, \dots) of the polyhedral areas converges, then we say that S is quadrable and refer to the necessarily unique limit of (A_1, A_2, \dots) as the area of the surface S .

THEOREM 1. *Let $f(x_1, x_2, \dots, x_n)$ be defined and be continuously partially differentiable on \bar{E} . Then there exists a proper sequence (Π_1, Π_2, \dots) of polyhedra inscribed on S .*

Proof. For every positive number r , there exists a decomposition of \bar{E} into a finite set of right n -hedra whose diameters are all less than r . Their vertices determine a finite set of points in S whose projection is precisely the set of these vertices. This set of points in S determines a polyhedron Π which is inscribed on S . We show that by making the norm of the decomposition of \bar{E} sufficiently small, we can make the deviation norm of Π arbitrarily small.

Let $\varepsilon > 0$ be given.

By property (g) there exists a decomposition of \bar{E} into a set of right n -hedra the diameter of each of which is arbitrarily small. By property (c) there exist real numbers k, δ_1 , $k < 1$, such that if $PP_1P_2 \cdots P_n$ is a right n -hedron in \bar{E} , (with P the right angled vertex) of diameter $< \delta_1$, then $|\cos(\overrightarrow{QQ_i}, \overrightarrow{QQ_j})| < k$ for $i \neq j$, where $Q_i = f(P_i)$ and $Q_j = f(P_j)$. Let the decomposition of \bar{E} be by right n -hedra each of diameter less than δ_1 .

By property (a), there exists a positive number θ such that if $\sin(\overrightarrow{QQ_i}, \overrightarrow{QQ_{i'}}) < \theta$ for each i , then the acute angle between

$$QQ_1 \times QQ_2 \times \cdots \times QQ_n$$

and $QQ'_1 \times QQ'_2 \times \cdots \times QQ'_n$ is less than $\varepsilon/3$.

By properties (d) and (e) there exists a positive number δ_2 such that if $PP_1P_2 \cdots P_n$ is a right n -hedron of diameter less than δ_2 , then, for each i , the acute angle between the chord $\overline{QQ_i}$ and the tangent line at Q to the curve in S subtended by $\overline{QQ_i}$ is less than θ . It follows that the acute angle between the normal to the polyhedral face $QQ_1Q_2 \cdots Q_n$ and the surface normal at Q is less than $\varepsilon/3$.

By property (f) there exists a positive number δ_3 such that if the diameter of the n -hedron $PP_1 \cdots P_n$ is less than δ_3 , then the angle between the surface normals at any two points of the portion of S which is subtended by the polyhedral face $QQ_1 \cdots Q_n$ is less than $\varepsilon/3$.

Let δ be the least of $\delta_1, \delta_2, \delta_3$. If D is any decomposition of \bar{E} into right n -hedra each of diameter less than δ , then if $QQ_1 \cdots Q_n$ is any of the polyhedral faces, the supremum of the angles between the normal to the n -hedron $QQ_1 \cdots Q_n$ and the portion of S which is subtended by $QQ_1 \cdots Q_n$ is less than ε .

Thus, corresponding to a sequence $(\varepsilon_1, \varepsilon_2, \dots)$ converging to zero, there exists a sequence of polyhedra with corresponding sequence of norms converging to zero and also with corresponding sequence of deviation norms converging to zero.

THEOREM 2. *Let $f(x_1, x_2, \dots, x_n)$ be defined and continuously*

partially differentiable on \bar{E} . Then, for every proper sequence (Π_1, Π_2, \dots) of n -hedra inscribed on $S = f(\bar{E})$, the corresponding sequence (A_1, A_2, \dots) of polyhedral areas converges to the multiple integral

$$\int_{\bar{E}} (1 + z_{x_1}^2 + \dots + z_{x_n}^2)^{1/2} dx_1 dx_2 \dots dx_n.$$

Proof. For the sake of notations, let z_{x_h} denote $\partial z / \partial x_h$, $h=1, \dots, n$. For each m , the projection of Π_m constitutes a decomposition D_m of \bar{E} into a finite set of n -hedra. Let the n -hedron $\Delta_{mr} = QQ_1 \dots Q_n$ be a face of Π_m and let $\Delta'_{mr} = \text{Proj}(QQ_1 \dots Q_n) = PP_1 \dots P_n$. Let β_{mr} be the acute angle between the normals to Δ_{mr} and to Δ'_{mr} . Let A_{mr} and A'_{mr} denote the respective areas of Δ_{mr} and Δ'_{mr} . Then $A_{mr} = A'_{mr} \sec \beta_{mr}$ where β_{mr} is the angle between the z -axis and the normal to Δ_{mr} . The area of A_m of Π_m is given by $\sum_r A'_{mr} \sec \beta_{mr}$.

Let P_{mr} be any point in Δ'_{mr} and let Q_{mr} be the point of S whose projection is P_{mr} . Let θ_{mr} denote the acute angle between the surface normal at Q_{mr} and the z -axis.

We associate to the sequence (Π_1, Π_2, \dots) certain related sequences:

$$\begin{aligned} &(\Pi_1, \Pi_2, \dots) \\ &(\phi_1, \phi_2, \dots) \\ &(\Sigma_1, \Sigma_2, \dots) \\ &(\Sigma'_1, \Sigma'_2, \dots). \end{aligned}$$

The sequence (ϕ_1, ϕ_2, \dots) is the corresponding sequence of deviation norms. The sequence $(\Sigma_1, \Sigma_2, \dots)$ is the corresponding sequence of polyhedral areas, $\Sigma_m = \sum_r A'_{mr} \sec \beta_{mr}$. In the fourth sequence, $\Sigma'_m = \sum_r A'_{mr} \sec \theta_{mr}$. Here, $\sec \theta_{mr}$ is the value of $(1 + z_{x_1}^2 + \dots + z_{x_n}^2)^{1/2}$ at some point of Δ'_{mr} . Thus the sequence $(\Sigma'_1, \Sigma'_2, \dots)$ is a sequence of Riemann sum of the function $(1 + z_{x_1}^2 + \dots + z_{x_n}^2)^{1/2}$ on \bar{E} with corresponding sequence of norms converging to zero. Since

$$(1 + z_{x_1}^2 + \dots + z_{x_n}^2)^{1/2}$$

is continuous on \bar{E} , the sequence $(\Sigma'_1, \Sigma'_2, \dots)$ converges to the multiple integral

$$\int_{\bar{E}} (1 + z_{x_1}^2 + \dots + z_{x_n}^2)^{1/2} dx_1 dx_2 \dots dx_n.$$

We now consider the sequence $(\Sigma_1, \Sigma_2, \dots)$. Let θ denote the acute angle between the surface normal at a point in S and the z -axis. $\sec \theta = (1 + z_{x_1}^2 + \dots + z_{x_n}^2)^{1/2}$ is bounded on \bar{E} . Thus there exists an angle $\theta^* > 0$ such that $\theta < \theta^*$ for all points of \bar{E} (i.e., for all points of S). Since $\sec \theta$ is uniformly continuous on the closed

interval $[\theta, \theta^*]$ for every $\eta > 0$ there exists $\tau > 0$ such that if $0 < \theta_1 < \theta^*$, $\theta < \theta_2 < \theta^*$ and $|\theta_1 - \theta_2| < \tau$, then $|\sec \theta_1 - \sec \theta_2| < \eta$.

We now compare the sequences $(\Sigma_1, \Sigma_2, \dots)$ and $(\Sigma'_1, \Sigma'_2, \dots)$.

Let $\varepsilon > 0$ be given. Take $\varepsilon/2V$ where V is the volume (area) of \bar{E} . There exists $\tau > 0$ such that if $|\theta_1 - \theta_2| < \tau$, then

$$|\sec \theta_1 - \sec \theta_2| < \frac{\varepsilon}{2V}.$$

Since (ϕ_1, ϕ_2, \dots) converges to zero, there exists a positive integer N_1 such that if $m > N_1$ then $\phi_m < \tau$. Thus if $m > N_1$, then

$$|\Sigma_m - \Sigma'_m| = |\Sigma_r A'_{mr} (\sec \beta_{mr} - \sec \theta_{mr})| < \frac{\varepsilon}{2V} \Sigma A'_{mr} = \frac{\varepsilon}{2}.$$

Since $(\Sigma'_1, \Sigma'_2, \dots)$ converges to \oint , there exists a positive integer N_2 such that if $m > N_2$ then $|\Sigma'_m - \oint| < \varepsilon/2$. Let N be the larger of N_1 and N_2 . If $m > N$, then

$$\begin{aligned} \left| \Sigma_m - \oint \right| &= \left| \Sigma_m - \Sigma'_m + \Sigma'_m - \oint \right| \\ &\leq |\Sigma_m - \Sigma'_m| + \left| \Sigma'_m - \oint \right| \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Thus, $(\Sigma_1, \Sigma_2, \dots)$ converges to \oint .

REFERENCES

1. L. V. Toralballa, *Piecewise flatness and surface area*, Annales Polonici Mathematici, **21** (1969), 223-230.
2. L. Cesari, *Surface Area*, Princeton University Press 1956.
3. S. Kempisty, *Sur la méthode triangulaire du calcul de l'aire d'une surface courbe*, Bull. Soc. Math. France, **64** (1936), 119-132.
4. T. Rado, *Length and area*, Amer. Math. Soc. Colloquium Publ., 1948.
5. H. A. Schwartz, *Sur une définition erronée de l'aire d'une surface courbe*, Ges. Math. Abhandl. **2** (1882), 309-311, 369-370.
6. J. A. Serret, *Cours de calcul différentiel et integral*, 2 vols., Paris 1868.
7. L. V. Toralballa, *Directional deviational norms and surface area*, L'Enseignement Mathématique (2), **13** (1967).

Received February 5, 1970.

MARIST COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Louis I. Alpert and L. V. Toralballa, <i>An elementary definition of surface area in E^{n+1} for smooth surfaces</i>	261
Eamon Boyd Barrett, <i>A three point condition for surfaces of constant mean curvature</i>	269
Jan-Erik Björk, <i>On the spectral radius formula in Banach algebras</i>	279
Peter Botta, <i>Matrix inequalities and kernels of linear transformations</i>	285
Bennett Eisenberg, <i>Baxter's theorem and Varberg's conjecture</i>	291
Heinrich W. Guggenheimer, <i>Approximation of curves</i>	301
A. Hedayat, <i>An algebraic property of the totally symmetric loops associated with Kirkman-Steiner triple systems</i>	305
Richard Howard Herman and Michael Charles Reed, <i>Covariant representations of infinite tensor product algebras</i>	311
Domingo Antonio Herrero, <i>Analytic continuation of inner function-operators</i>	327
Franklin Lowenthal, <i>Uniform finite generation of the affine group</i>	341
Stephen H. McCleary, <i>0-primitive ordered permutation groups</i>	349
Malcolm Jay Sherman, <i>Disjoint maximal invariant subspaces</i>	373
Mitsuru Nakai, <i>Radon-Nikodým densities and Jacobians</i>	375
Mitsuru Nakai, <i>Royden algebras and quasi-isometries of Riemannian manifolds</i>	397
Russell Daniel Rupp, Jr., <i>A new type of variational theory sufficiency theorem</i>	415
Helga Schirmer, <i>Fixed point and coincidence sets of biconnected multifunctions on trees</i>	445
Murray Silver, <i>On extremal figures admissible relative to rectangular lattices</i>	451
James DeWitt Stein, <i>The open mapping theorem for spaces with unique segments</i>	459
Arne Stray, <i>Approximation and interpolation</i>	463
Donald Curtis Taylor, <i>A general Phillips theorem for C^*-algebras and some applications</i>	477
Florian Vasilescu, <i>On the operator $M(Y) = TYS^{-1}$ in locally convex algebras</i>	489
Philip William Walker, <i>Asymptotics for a class of weighted eigenvalue problems</i>	501
Kenneth S. Williams, <i>Exponential sums over $GF(2^n)$</i>	511