

# Pacific Journal of Mathematics

**MINIMAL INJECTIVE COGENERATORS FOR THE CLASS OF  
MODULES OF ZERO SINGULAR SUBMODULE**

VASILY CATEFORIS

# MINIMAL INJECTIVE COGENERATORS FOR THE CLASS OF MODULES OF ZERO SINGULAR SUBMODULE

VASILY C. CATEFORIS

Let  $R$  be a ring (with 1) of zero singular right ideal and let  $Q$  be its maximal right quotient ring; let  $\mathcal{N}$  be the class of all (unitary) right  $R$ -modules of zero singular submodule. An element  $M$  of  $\mathcal{N}$  is said to be an injective cogenerator for  $\mathcal{N}$  if  $M$  is an injective module and every element of  $\mathcal{N}$  can be embedded in a direct product of copies of  $M$ ;  $M$  is said to be a minimal injective cogenerator for  $\mathcal{N}$  if  $M$  is the only direct summand of  $M$ , which is an injective cogenerator for  $\mathcal{N}$ . This paper deals with the question of existence and uniqueness of a minimal injective cogenerator for  $\mathcal{N}$  (and in  $\mathcal{N}$ ). If a minimal injective cogenerator for  $\mathcal{N}$  exists, then it is (a) isomorphic to a minimal faithful direct summand of  $Q$ , (b) isomorphic to a direct summand of every injective cogenerator for  $\mathcal{N}$  (and in  $\mathcal{N}$ ) and (c) unique (up to isomorphism). Whether  $Q$  is (or is not) a prime ring, affects the structure, though not the existence, of a minimal injective cogenerator for  $\mathcal{N}$ : a minimal injective cogenerator for  $\mathcal{N}$ , if it exists, is (up to isomorphism) a faithful minimal right ideal of  $Q$  iff  $Q$  is a prime ring and so in this case  $Q$  is a minimal injective cogenerator for  $\mathcal{N}$  iff  $Q$  is a division ring. On the other hand, if  $R_R$  is finite dimensional (Goldie) then a minimal injective cogenerator for  $\mathcal{N}$  exists; it is  $Q$  iff  $Q$  is (ring) isomorphic to a finite product of division rings.

We begin with a list of conventions, assumptions and well known facts:

(a) By a ring  $R$  it is meant an associative ring  $R$  with 1, whose singular right ideal [2] is zero.  $R_R$  is used when  $R$  is considered as a right  $R$ -module.

(b)  $Q$  denotes the maximal right quotient ring [2] of a ring  $R$  and so  $Q$  is a Von Neumann regular ring, i.e. a ring every principal right ideal of which is a direct summand, and also the injective hull of  $R_R$  [2].

Now for the rest of the list, let  $R$  be a given ring.

(c) By a module  $M$  it is meant a unitary right  $R$ -module  $M$ ;  $Z(M)$  denotes the singular submodule of  $M$  [2] and a module of zero singular submodule is called (for short) nonsingular.  $\mathcal{N}$  denotes the class of all nonsingular  $R$ -modules.

(d) For each module  $M$ ,  $E(M)$  denotes the injective hull [2] of  $M$ . If  $M$  and  $N$  are modules such that  $M \subseteq N$  we write  $M \subseteq' N$  to denote the fact that  $M$  is essential in  $N$  ( $N$  is an essential extension of  $M$

[2]).

(e) A module  $C$  is said to be  $M$ -torsionless, for a given module  $M$ , if  $C$  can be embedded in a direct product of copies of  $M$ , or, equivalently, if  $\cap \ker f = (0)$  where  $f$  ranges over  $\text{Hom}_R(C, M)$ .

(f) A module  $M$  is said to be a cogenerator for a class  $\mathcal{A}$  of modules, if every module in  $\mathcal{A}$  is  $M$ -torsionless, and an injective cogenerator if, also,  $M$  is injective.

(g) Whenever a cogenerator  $M$  for  $\mathcal{N}$  is considered it is assumed that  $M$  is also in  $\mathcal{N}$ . As a corollary to Gentile's [3, p. 427, Prop. 1] we have:

**PROPOSITION 0.1.**  *$Q$  is an injective cogenerator for  $\mathcal{N}$ .*

(h) Perhaps the most crucially, certainly the most often used result is the following consequence of [6, p. 119, Remark] and [7, p. 226, Lemma 2.3]:

**LEMMA 0.2.** *If  $A$  is an injective module and  $C$  is a nonsingular module, then any homomorphism  $f: A \rightarrow C$  splits (i.e.  $\ker f$  is a direct summand of  $A$ ).*

The following will also be of frequent use:

**LEMMA 0.3.** *If  $I$  is a right ideal of  $R$ , then  $E(I) = eQ$  for some idempotent in  $Q$ .*

(i) For a nonempty subset  $S$  of a module  $M$ ,  $r. \text{ann}_R S = \{r \in R / sr = 0, \text{ for all } s \in S\}$  and thus a module  $M$  is faithful if  $r. \text{ann}_R M = (0)$ ; a module  $M$  is said to be minimal faithful if  $M$  is faithful and no proper ( $\neq M$ ) direct summand of  $M$  is (faithful).

**1. Minimal injective cogenerators for  $\mathcal{N}$ .** Let  $R$  be a ring. We start with a generalization of a theorem of Armendariz [1, p. 568, Theorem 3].

**THEOREM 1.1.** *For a nonsingular module  $M$ , the following statements are equivalent:*

(a)  *$M$  is a cogenerator for  $\mathcal{N}$ .*

(b)  *$M$  contains a faithful submodule  $D$  such that  $D$  contains the injective hull of every one of its finitely generated submodules.*

*Proof.* (a) implies (b). By hypothesis  $\text{Hom}_R(Q, M) \neq 0$  and so, by Lemma 0.2,  $M$  contains nonzero injective submodules. Let  $D$  be the sum of all injective submodules of  $M$ ; if  $N_1, \dots, N_k$  are injective sub-

modules of  $M$  (finitely many) then  $N_1 + \cdots + N_k$  is also injective as it is a homomorphic image of the injective module  $N_1 \times \cdots \times N_k$  (Lemma 0.2 again). It follows that  $D$  contains the injective hull of each of its finitely generated submodules. Now observe that  $\text{Hom}_R(Q, M) = \text{Hom}_R(Q, D)$  and thus  $Q$  is  $D$ -torsionless. It follows that  $D$  is faithful; in fact we have shown that  $D$  is a cogenerator for  $\mathcal{N}$  as  $Q$  is (Proposition 0.1).

(b) *implies* (a). If  $a$  is a nonzero element of  $Q$ , then for some  $r \in R$ ,  $ar$  is a nonzero element of  $R$ . Since  $D ar \neq 0$ , there exists  $d \in D$  such that  $dar \neq 0$ , and thus a module map  $f: arR \rightarrow daR$  such that  $f(ar) \neq 0$ . Since  $E(\text{Im } f) \subseteq D$ , the map  $f$  has an extension  $f': Q \rightarrow D$ , and  $f'(a) \neq 0$ . Thus  $Q$  is  $D$ -torsionless and hence  $M$ -torsionless; by Proposition 0.1  $M$  is a cogenerator for  $\mathcal{N}$ .

The following quite obvious corollaries to the above theorem are singled out for later usage.

**COROLLARY 1.1.1.** *A right ideal  $A$  of  $Q$  is a cogenerator for  $\mathcal{N}$  if and only if  $A$  is a faithful  $R$  - (or  $Q$  -) module.*

**COROLLARY 1.1.2.** *An injective nonsingular module  $M$  is a cogenerator for  $\mathcal{N}$  if and only if  $M$  is faithful.*

The rest of this section is devoted to results about the existence and uniqueness of a minimal injective cogenerator for  $\mathcal{N}$ , this concept defined in the obvious manner as follows:

**DEFINITION 1.2.** A nonsingular module  $M$  is said to be a minimal injective cogenerator for  $\mathcal{N}$  if (a)  $M$  is an injective cogenerator for  $\mathcal{N}$  and (b) no direct summand of  $M$  different from  $M$  is a cogenerator for  $\mathcal{N}$ .

We have as a corollary to Theorem 1.1:

**PROPOSITION 1.3.** *An injective nonsingular module  $M$  is a minimal injective cogenerator for  $\mathcal{N}$  if and only if  $M$  is a minimal faithful module.*

The quotient field  $K$  of a commutative integral domain  $R$  is isomorphic to a submodule of every nonzero torsion-free injective  $R$ -module  $M$ . A similar result about a nonzero nonsingular injective module  $M$  and  $Q$  does not in general hold, even when  $M$  is assumed faithful. However some theory relating  $Q$  and the nonzero injective elements of  $\mathcal{N}$  is possible and essentially a consequence of the following result

(Theorem Z) contained in a theorem of J. Zelamowitz [8, Theorem 2]:

**THEOREM Z.** *If  $M$  is a nonsingular module then there exists a collection of right ideals of  $R$ ,  $\{I_\alpha: \alpha \in A\}$ , such that  $M$  is an essential extension of a submodule isomorphic to  $\bigoplus I_\alpha$  (external direct sum).*

In view of Lemma 0.3 we have immediately:

**COROLLARY Z.1.** *A nonsingular injective module  $M$  is (up to isomorphism) the injective hull of a direct sum  $\bigoplus e_\alpha Q$ , where  $\{e_\alpha: \alpha \in A\}$  is a set of (not necessarily orthogonal) idempotents of  $Q$ .*

We now take a closer look at the direct summands  $eQ$  ( $e^2 = e$ ) of  $Q$  on the way to establishing results on the existence and uniqueness of a minimal injective cogenerator for  $\mathcal{N}$ .

**LEMMA 1.4.** *If  $e$  and  $f$  are idempotents of  $Q$  other than 0 or 1, then  $\text{Hom}(fQ, eQ) = (0)$  if and only if  $eQ$  and  $fQ$  have no isomorphic nonzero direct summands.*

*Proof.* Clear. (Use Lemma 0.2 for the *if* part; use the injectivity of  $eQ$ , or  $fQ$ , for the *only if* part).

**REMARK.** If  $eQ$  and  $fQ$  have no isomorphic nonzero direct summands, then  $eQfQ = 0$  and  $fQeQ = 0$  because  $\text{Hom}_R(fQ, eQ) \cong eQf$  (as groups) and the fact that the condition on the direct summands is a symmetric one.

**DEFINITION 1.5.** We say that the modules  $M$  and  $N$  share a nonzero direct summand if  $M$  and  $N$  have isomorphic nonzero direct summands.

**LEMMA 1.6.** *If  $\{f_\alpha: \alpha \in A\}$  is a set of idempotents in  $Q$  such that  $N = E(\bigoplus_\alpha f_\alpha Q)$  is an injective cogenerator for  $\mathcal{N}$ , then for each nonzero idempotent  $e$  in  $Q$ ,  $eQ$  shares a nonzero direct summand with some  $f_\beta Q$ .*

*Proof.* Let  $f: eQ \rightarrow N$  be a homomorphism such that  $f(e) \neq 0$ . Since  $N$  is an essential extension of  $\bigoplus f_\alpha Q$ , there exists  $r \in R$  such that  $0 \neq f(e)r \in \bigoplus f_\alpha Q$  and so for some  $\beta \in A$   $\pi_\beta f(e)r \neq 0$  where  $\pi_\beta: \bigoplus f_\alpha Q \rightarrow f_\beta Q$  is the canonical projection. It follows now by Lemma 0.2 that  $\pi_\beta f(erQ)$  is a nonzero direct summand shared by  $eQ$  and  $f_\beta Q$ .

**COROLLARY.** *If in  $Q$  there exist nonzero idempotents  $e_1$  and  $e_2$*

such that  $e_1 Q e_2 Q = 0$  then there also exist nonzero idempotents  $f_1$  and  $f_2$  such that  $f_i Q$ ,  $i = 1, 2$ , is (isomorphic to) a submodule of  $N$  and  $f_1 Q f_2 Q = 0$  ( $N$  is an injective cogenerator for  $\mathcal{N}$ ).

DEFINITION 1.7. (a) If  $e$  and  $f$  are idempotents in  $Q$ , the summands  $eQ$  and  $fQ$  of  $Q$  are said to be orthogonal if  $eQfQ = (0)$ .

(b) A nonzero right ideal  $B$  of  $Q$  is said to be only orthogonally decomposable if whenever  $B = X \oplus Y$ , for right ideals  $X$  and  $Y$  of  $Q$ , then  $XY = (0)$ .

It is easy to see that  $Q$  need not have any orthogonal summands different from  $(0)$  and  $Q$ ; in fact we have:

LEMMA 1.8.  $Q$  is a prime ring if and only if  $Q$  has no orthogonal summands other than  $(0)$  and  $Q$ .

*Proof.* A prime ring is one in which, for example, the product of nonzero principal right ideals is nonzero. Every principal right ideal of  $Q$  is a direct summand of  $Q$ .

REMARK. If  $Q$  is not a prime ring and  $N$  is an injective cogenerator for  $\mathcal{N}$ , then  $N$  contains (isomorphic copies of) orthogonal nonzero summands of  $Q$  (Lemma 1.6).

Now we consider an existence theorem.

THEOREM 1.9. The following statements are equivalent:

- (a) There exists a minimal injective cogenerator  $M$  for  $\mathcal{N}$ .
- (b) There exists a maximal set  $\{e_\alpha Q : \alpha \in A\}$  of pairwise orthogonal summands of  $Q$  such that each  $e_\alpha Q$  is only orthogonally decomposable.
- (c) There exists a minimal faithful right ideal  $fQ$  for some idempotent  $f$  in  $Q$ .

In particular if  $M$  is a minimal injective cogenerator for  $\mathcal{N}$ , then  $M \cong fQ$  for some minimal faithful right ideal direct summand  $fQ$  of  $Q$ .

*Proof.* (a) implies (b). Let  $\{e_\alpha Q : \alpha \in A\}$  be a set of summands of  $Q$  such that  $M = E(\oplus e_\alpha Q)$  (given by Corollary Z.1). We show at once that the summands  $\{e_\alpha Q\}$  are pairwise orthogonal and each only orthogonally decomposable. To this end suppose  $e_\alpha Q$  and  $e_\beta Q$  ( $\alpha \neq \beta$ ) share a direct summand and so there exist module decompositions  $e_\alpha Q = A' \oplus A''$  and  $e_\beta Q = B' \oplus B''$  with  $A' \cong B'$  and both  $A'$  and  $B'$  nonzero. These (decompositions) induce a module decomposition  $M = A' \oplus B' \oplus C$ ; now since  $A' \cong B'$  and  $M$  is a cogenerator for  $\mathcal{N}$ , it follows by the definition of cogenerator that, for example,  $B' \oplus C$  is

also a cogenerator for  $\mathcal{N}$ , contrary to the minimality of  $M$ . It follows that  $e_\alpha Q$  and  $e_\beta Q$  are orthogonal whenever  $\alpha \neq \beta$  and, by the same argument, that each  $e_\alpha Q$  is only orthogonally decomposable. Finally, the set  $\{e_\alpha Q: \alpha \in A\}$  is a maximal set of pairwise orthogonal summands of  $Q$  by Lemma 1.6.

(b) *implies* (c). Let  $fQ = E(\bigoplus e_\alpha Q)$ , where  $\{e_\alpha Q: \alpha \in A\}$  is as given in (b) (and so in particular the direct sum  $\bigoplus e_\alpha Q$  is internal). To show that  $fQ$  is faithful, assume that, on the contrary, there exists a nonzero idempotent  $e$  in  $Q$  such that  $fQe = (0)$ ; it follows that  $e_\alpha QeQ = (0)$  for each  $\alpha \in A$  and so  $\{e_\alpha Q: \alpha \in A\} \cup \{eQ\}$  is a set of pairwise orthogonal summands of  $Q$ , properly containing  $\{e_\alpha Q: \alpha \in A\}$ , contrary to the latter's maximality. To show that  $fQ$  is minimal faithful, suppose that  $fQ = B \oplus C$  where  $B$  is faithful and so, by Corollary 1.1.2, an injective cogenerator for  $\mathcal{N}$ . It needs to be shown that  $C = (0)$ . If  $C$  is not zero, that there exists a nonzero idempotent  $e$  in  $Q$  such that  $eQ \subset C$  and since  $\bigoplus e_\alpha Q \cong' fQ$  it may be assumed that  $eQ \subset \bigoplus e_\alpha Q$ . Furthermore as  $eQ$  shares a direct summand with some  $e_\alpha Q$  (Lemma 1.6.) and as the  $e_\alpha Q$ 's are orthogonal, it may be assumed that  $eQ \subset e_\beta Q$ , for some  $\beta \in A$ . Now since  $B$  is an injective cogenerator for  $\mathcal{N}$ , it follows from Lemma 1.6 (and Corollary Z.1) that  $eQ$  shares a direct summand with some summand  $e'Q$  of  $B$ , for some nonzero idempotent  $e'$  in  $Q$ , and it may be assumed that  $e'Q$  is isomorphic to a summand of  $eQ$ . As in the case of  $eQ$ , it may be assumed that  $e'Q$  is contained in one of the summands  $e_\alpha Q$  and as they are orthogonal it must be that  $e'Q \subset e_\beta Q$ . Now since  $e'Q \subset B$  and  $eQ \subset C$ , it follows from  $B \cap C = (0)$ , that  $e'Q \cap eQ = (0)$  and thus  $e'QeQ = (0)$ , as  $e_\beta Q$  is only orthogonally decomposable; however  $e'QeQ \neq (0)$  (Lemma 1.4) and thus the assumption  $C \neq (0)$  has led to a contradiction.

(c) *implies* (a). Proposition 1.3.

REMARKS. (1) It should be clear from the preceding considerations, that if  $Q$  is a prime ring then a minimal injective cogenerator for  $\mathcal{N}$  exists if and only if  $Q$  has nonzero socle. The case in which  $Q$  (or  $R$ ) is prime (including this remark) will be considered in detail in the next section.

(2) Any injective cogenerator  $M$  for  $\mathcal{N}$  contains a submodule isomorphic to a faithful right ideal  $fQ$ , for some nonzero idempotent  $f$  of  $Q$ . In fact if  $\{f_j Q: j \in J\}$  is a maximal set of pairwise orthogonal summands of  $Q$  contained in  $M$  (such exist by the Corollary to Lemma 1.6 in case  $Q$  is not prime) then  $fQ$  can be chosen to be  $E(\bigoplus f_j Q)$ .

The following is a uniqueness theorem.

**THEOREM 1.10.** *If  $M$  is a minimal injective cogenerator for  $\mathcal{N}$ , then*

- (a)  *$M$  is unique up to isomorphism, and*
- (b)  *$M$  is isomorphic to a submodule of every injective cogenerator  $N$  for  $\mathcal{N}$ .*

*Proof.* In view of Theorem 1.9 and Remark (2) following it, for a proof of both (a) and (b) of this theorem, it is sufficient to show that if  $e$  and  $f$  are idempotents in  $Q$  such that  $eQ$  is minimal faithful and  $fQ$  is faithful, then  $eQ$  is isomorphic to a direct summand of  $fQ$ . We show this next:

If an ideal  $A$  of  $Q$  is such that  $eQfQA = (0)$  then  $fQA = (0)$  and so  $A = (0)$  as both  $eQ$  and  $fQ$  are faithful. It follows that  $eQfQ = eQ$ , as  $eQfQ$  is a faithful direct summand of  $eQ$ . Thus there exist elements  $p$  and  $q$  in  $Q$  such that  $e = epfq$  and so the homomorphism  $h: fQ \rightarrow eQ$  given by  $h(fx) = epfx$  is an epimorphism. It follows from Lemma 0.2 that  $eQ$  is isomorphic to a direct summand of  $fQ$ .

We do not know whether, in general, the property of being isomorphic to a submodule of every injective cogenerator for  $\mathcal{N}$ , characterizes the minimal injective cogenerator, among the injective cogenerators for  $\mathcal{N}$ .

A simple example to put the results of this section in some concrete form is the case when  $R$  is a commutative ring. It is easy to show that then  $Q$ , also, is a commutative ring and  $Q$  is only orthogonally decomposable.  $Q$  is the unique minimal injective cogenerator for  $\mathcal{N}$ .

**2. Nonsingular uniform modules; rings of finite Goldie dimension; prime rings.** The assumption that  $R$  is an associative ring with 1, of zero singular right ideal and that  $Q$  is its maximal right quotient ring, continues in force.

A module  $M$  is said to be finite dimensional (in the sense of Goldie) [4, p. 202] if it contains no infinite direct sum of nonzero submodules and we call  $R$  a finite dimensional ring if  $R_R$  is a finite dimensional module. A module  $U$  is said to be uniform if  $U \neq 0$  and  $U$  is an essential extension of every one of its nonzero submodules. A uniform right ideal of  $R$  is, then, a uniform submodule of  $R_R$ .

For each module  $M$ ,  $\text{Soc}(M)$  denotes the socle of  $M$ .

In this section we are primarily interested in nonsingular uniform modules, as when they exist "in abundance" (e.g. when the sum of uniform right ideals of  $R$  is essential in  $R_R$ ), then they determine the minimal injective cogenerator for  $\mathcal{N}$  in a simple way; in fact (they



determine it) quite in the manner in which the (nonisomorphic) simple modules and their injective hulls determine the minimal injective cogenerator of the category of all modules. Thus we proceed in the following with a sequence of facts about nonsingular uniform modules (when they exist), some of them, probably, well known.

**LEMMA 2.1.** *A homomorphism  $f: U \rightarrow A$  where  $U$  and  $A$  are nonsingular modules and  $U$  is uniform is either the zero map or a monomorphism.*

*Proof.*  $U/\ker f$  is a nonsingular module, since  $A$  is and so if  $\ker f \neq (0)$  then it must be that  $\ker f = U$ , since  $U/\ker f$  is, then, its own singular submodule as well.

**DEFINITION 2.2.** A uniform module  $U$  is said to be equivalent to a uniform module  $V$ , and then we write  $U \sim V$ , if  $E(U) \cong E(V)$  or, equivalently, if there exists monomorphism  $A \rightarrow V$  for some nonzero submodule  $A$  of  $U$ .

It is clear that this relation is an equivalence relation on uniform modules.

**LEMMA 2.3.** *The following statement about a uniform module  $U$  are true:*

- (a)  $Z(U) = U$  or  $Z(U) = (0)$
- (b) If  $Z(U) = (0)$ , then  $E(U)$  is isomorphic (as an  $R$ -module) to a minimal right ideal of  $Q$ .
- (c)  $Z(U) = (0)$  if and only if  $U$  is equivalent to a uniform right ideal of  $R$ .

*Proof.* (a) follows from the fact that  $Z(U/Z(U)) = (0)$  [5, p. 270, Proposition 2.3] and at the same time,  $Z(U/Z(U)) = U/Z(U)$ , if  $Z(U) \neq (0)$ .

(b) If  $Z(U) = (0)$  then there exists embedding (of  $R$ -modules)  $U \rightarrow Q$  (Proposition 0.1 and Lemma 2.1). We may thus assume that  $U$  is a uniform  $R$ -submodule of  $Q$  and further assume that  $U = qR$  for some  $0 \neq q \in Q$ , since  $qR \sim U$  for every  $0 \neq q \in U$ . Thus  $E(U) = E(qR) = qQ = eQ$  for some idempotent  $e$  in  $Q$ . Now  $eQ$  is a uniform  $R$ -submodule of  $Q_R$ , as  $U$  is, and so  $eQ$  is a uniform ideal of  $Q$ . Since  $Q$  is Von Neumann regular, it follows that  $eQ$  is a minimal right ideal of  $Q$ .

(c) Since injective hulls of nonsingular modules are nonsingular modules,  $Z(U) = (0)$  if  $U$  is equivalent to a uniform right ideal of  $R$ . On the other hand, using the notation of part (b) above if  $Z(U) = (0)$ , we have  $I = eQ \cap R$ , a uniform right ideal of  $R$  such that  $E(I) = eQ = E(U)$ .

REMARK. Nonsingular uniform modules need not exist: a ring  $R$  such that  $\text{Soc}(Q_Q) = (0)$  exists (see Example following the Corollary to Proposition 2.11).

DEFINITION 2.4 (Terminology). For each module  $M$ ,  $\mathcal{U}(M)$  denotes the (module) sum of all uniform submodules of  $M$ . If  $M$  has no uniform submodules, then we write:  $\mathcal{U}(M) = (0)$ .

Any finite dimensional ring  $R$  satisfies  $\mathcal{U}(R_R) \subseteq' R_R$  [4, p. 202, Theorem 1.1]. If, on the other hand,  $R$  is an infinite direct product of fields, then  $\mathcal{U}(R_R) \subseteq' R_R$  but  $R$  is not finite dimensional.

PROPOSITION 2.5. For a ring  $R$  the following statements are equivalent:

- (a)  $\mathcal{U}(R_R) \subseteq' R_R$
- (b)  $\mathcal{U}(M) \subseteq' M$  for every nonzero nonsingular module  $M$
- (c)  $\text{Soc}(Q_Q) \subseteq' Q_Q$ .

*Proof.* (a) implies (b). It is sufficient to show that  $\mathcal{U}(M) \neq (0)$  in case  $M$  is a nonzero, nonsingular cyclic module. If  $M$  is such, then there exists epimorphism  $f: R_R \rightarrow M$ ; now it cannot happen that  $f(U(R_R)) = (0)$  as  $Z(M) = (0)$ , and this would imply  $f(R) = (0)$ , though  $M \neq 0$ . Thus  $f(U) \neq 0$  for some uniform right ideal of  $R$  and it follows from Lemma 2.1 that  $U \cong f(U) \subset M$ .

(b) implies (c). Condition (b) in particular implies that  $\mathcal{U}(R_R) \subseteq' R_R$  and so (c) follows from part (b) of Lemma 2.3.

(c) implies (a). For each minimal right ideal  $A$  of  $Q$ ,  $A \cap R$  is a uniform right ideal of  $R$  and so if  $\text{Soc}(Q_Q) = \sum A_i$  where  $\{A_i\}$  are the minimal right ideal of  $Q$ , then  $\sum(A_i \cap R) \subseteq \mathcal{U}(R)$ . On the other hand  $\sum(A_i \cap R) \subseteq' (\sum A_i) \cap R \subseteq' R_R$ .

REMARK. Finite dimensional rings  $R$  have been characterized by F. L. Sandomierski [6, p. 115, Theorem 1.6] as those for which  $\text{Soc}(Q_Q) = Q_Q$ .

THEOREM 2.6. Let  $R$  be a ring such that  $\mathcal{U}(R_R) \subseteq' R_R$  and let  $\{e_\alpha Q: \alpha \in A\}$  be a complete set of non-isomorphic minimal right ideals of  $Q$ , where  $\{e_\alpha: \alpha \in A\}$  is an appropriate set of primitive, orthogonal idempotents of  $Q$ . The ideal  $eQ = E(\bigoplus e_\alpha Q)$  is, then, a minimal injective cogenerator for  $\mathcal{N}$ .

*Proof.* We apply Theorem 1.9 (b). The summands  $\{e_\alpha Q\}$  are pairwise orthogonal because they are (pairwise) non-isomorphic minimal right ideals of  $Q$  and each  $e_\alpha Q$  is, clearly, only orthogonally decomposable. It remains to show that the set  $\{e_\alpha Q: \alpha \in A\}$  is a maximal

set of pairwise orthogonal summands of  $Q$ ; now if  $eQ$  is a nonzero summand of  $Q$ , then  $\text{Soc}(eQ) \neq (0)$ , by Proposition 2.5 (c), and so  $eQ$  contains a nonzero summand isomorphic to some  $e_\alpha Q$ , as  $\{e_\alpha Q: \alpha \in A\}$  is a complete set of non-isomorphic minimal right ideals of  $Q$ . Thus  $e_\alpha QeQ \neq (0)$ , for some  $\alpha \in A$ .

In view of Theorem 1.10 (b) and Lemma 2.3 (b), the following corollary to Theorem 2.6 is immediate.

**COROLLARY.** *If  $R$  is a ring such that  $\mathcal{Z}(R_R) \subseteq' R_R$  and  $M$  is a nonsingular module, then  $M$  is a cogenerator for  $\mathcal{N}$  if, and only if  $M$  contains a copy of the injective hull of every uniform nonsingular module.*

**THEOREM 2.7.** *If  $R$  is finite dimensional, then the following statements are equivalent:*

- (a)  $Q$  is a minimal injective cogenerator for  $\mathcal{N}$ .
- (b)  $Q$  (as a ring) is isomorphic to a finite direct product  $\Delta_1 \oplus \cdots \oplus \Delta_n$  of division rings  $\Delta_i$ .

*Proof.* (a) implies (b).  $Q$  is artinian semi-simple in this case [6, p. 115, Theorem 1.6] and so there exist primitive orthogonal idempotents  $e_1, \dots, e_n$  such that  $Q = e_1 Q \oplus \cdots \oplus e_n Q$ . Condition (a) now implies that the minimal ideals  $e_i Q$  are (pairwise) non-isomorphic. It follows now from the structure theory of artinian semi-simple rings that each  $e_i Q$  is a division ring and the sum  $e_1 Q \oplus \cdots \oplus e_n Q$  is a ring direct sum.

(b) implies (a). Each  $\Delta_i$  is a minimal  $Q$ -ideal and they (the ideals  $\Delta_i$ ) are non-isomorphic. Now use Theorem 2.6.

Now we turn our attention to the case when  $R$  is a prime ring.

**LEMMA 2.8.** *For a ring  $R$  the following statements are equivalent:*

- (a)  $R$  is a prime ring.
- (b) Every nonzero, nonsingular module  $M$  is a faithful module.

*Proof.* (a) implies (b). Over a prime ring  $R$  a two-sided ideal of  $R$  is either an essential submodule of  $R_R$  or it is zero (e.g. [1, p. 570, Lemma 3]). Since  $Z(M) = 0$ , it follows that  $r.\text{ann}_R(M) = (0)$ .

(b) implies (a). A nonzero right ideal  $A$  of  $R$  is a nonzero, nonsingular module and so  $r.\text{ann}_R A = (0)$ .

**PROPOSITION 2.9.** *Let  $R$  be a prime ring and let  $M$  be a nonzero, nonsingular injective module. The following statements about  $M$  are, then, equivalent:*

- (a)  $M$  is a minimal injective cogenerator for  $\mathcal{N}$ .

- (b) *A nonzero homomorphism  $f: M \rightarrow C$ , where  $C$  is nonsingular, is a monomorphism.*  
 (c)  *$M$  is a uniform module.*

*Proof.* (a) *implies* (b). It follows from Lemma 0.2 that  $M \rightarrow \text{Im } f$  splits and so  $\ker f$  is a direct summand of  $M$ . Now if  $\ker f \neq 0$  then  $\ker f$  is faithful (Lemma 2.8) contrary to minimality of  $M$  (Proposition 1.3). Thus it must hold that  $\ker f = 0$ .

(b) *implies* (c). If  $M$  is not uniform then, since  $M$  is injective, it is possible to find nonzero submodules  $A$  and  $B$  of  $M$  such that  $M = A \oplus B$ . Such a decomposition, however, gives rise to a homomorphism, e.g. the projection  $M \rightarrow B$ , of the kind which is forbidden by condition (b).

(c) *implies* (a).  $M$  is faithful, by Lemma 2.8, and so  $M$  is an injective cogenerator by Corollary 1.1.2. Condition (c) implies that  $M$  has no direct summands other than  $(0)$  and  $M$  and so (a) follows.

**THEOREM 2.10.** *If  $R$  is a prime ring, then a minimal injective cogenerator for  $\mathcal{N}$  exists if and only if  $\mathcal{U}(R_R) \neq (0)$ . Furthermore if  $\mathcal{U}(R_R) \neq (0)$  then  $\mathcal{U}(R_R) \subseteq' R_R$  and there is only one (up to isomorphism) nonsingular simple  $Q$ -module.*

*Proof.* If  $\mathcal{U}(R_R) \neq (0)$  then there exists a uniform right ideal  $U$  of  $R$  and thus also a nonsingular injective uniform  $R$ -module, namely  $E(U) \subset Q$ . It follows from Proposition 2.9 that  $E(U)$  is a minimal injective cogenerator for  $\mathcal{N}$ . On the other hand if  $M$  is a minimal injective cogenerator for  $\mathcal{N}$  then  $M$  is uniform (Proposition 2.9) and so  $\mathcal{U}(R_R) \neq (0)$  (Lemma 2.3 (c)). Now for the second part of the theorem, assume  $\mathcal{U}(R_R) \neq (0)$ . It follows (Lemma 2.3 (b)) that  $Q$  has a minimal right ideal  $fQ$ , where  $f$  is some (primitive) idempotent of  $Q$ ; now if  $eQ$  is any nonzero summand of  $Q$  then  $fQeQ \neq (0)$ , as  $Q$  is prime, and so  $\text{Soc}(eQ) \neq (0)$  (by Lemma 1.4, for example). It follows that  $\text{Soc}(Q_Q) \subseteq' Q_Q$  and so  $\mathcal{U}(R_R) \subseteq' R_R$ . Finally if  $S$  is any nonsingular simple  $Q$ -module, then  $S \cong gQ$  for some primitive idempotent  $g$  of  $Q$  (by Lemma 0.2) and, as before,  $fQ \cong gQ$ .

**COROLLARY.** *If  $R$  is a prime ring then  $Q$  is a minimal injective cogenerator for  $\mathcal{N}$  if and only if  $Q$  is a division ring.*

*Proof.* If  $Q$  is a minimal injective cogenerator for  $\mathcal{N}$  then  $Q$  is a uniform  $R$ -module (Proposition 2.9) and thus  $Q_Q$  is a minimal right  $Q$ -ideal. A ring with  $1 \neq 0$  and no right ideals other than zero and itself is, of course, a division ring.

Now we obtain an example of a prime ring  $R$  such that  $\mathcal{Z}(R) = (0)$ .

By a simple ring  $R$  we mean a ring in which  $(0)$  and  $R$  are the only two-sided ideals.

**PROPOSITION 2.11.** *If  $R$  has no divisors of zero  $\neq 0$  then  $Q$  is a simple ring.*

*Proof.* Let  $A$  be a nonzero two-sided ideal of  $Q$  and thus consider  $a \in R \cap A$ ,  $a \neq 0$  ( $R \cap A \neq (0)$  as  $R_R \cong' Q_R$ ). Since  $R$  has no divisors of zero  $\neq 0$  we have  $r \cdot \text{ann}_R a = (0)$  and, hence,  $r \cdot \text{ann}_Q a = (0)$ . Now  $Qa = Qe$  for some idempotent  $e$  in  $Q$  and so  $a(1 - e) = 0$ . Thus  $1 - e \in r \cdot \text{ann}_Q a = (0)$  and so  $e = 1$ ; we have  $Qa = Q$ , for some  $a \in A$ , and so  $Q = A$ .

**COROLLARY.** *If  $R$  has no divisors of zero  $\neq 0$  then a nonsingular uniform module exists if and only if  $Q$  is a division ring.*

*Proof.* If  $\mathcal{Z}(R) \neq (0)$  then  $\text{Soc}(Q) \neq (0)$  and as  $\text{Soc}(Q)$  is a two-sided ideal, it follows that  $\text{Soc}(Q) = Q$  or that  $Q$  is simple semi-simple artinian. Now every element of  $R - \{0\}$  is invertible in  $Q$  and so every element  $q$  of  $Q$  has the form  $ad^{-1}$  for appropriate elements  $a$  and  $d$  of  $R$  (i.e.  $Q$  is also the classical quotient ring [2] of  $R$ ). It follows that if  $q \neq 0$  then  $q^{-1} = da^{-1}$  exists and  $Q$  is a division ring.

**AN EXAMPLE.** Since a finite dimensional ring  $R$  (of zero singular right ideal) has an artinian semi-simple maximal quotient ring  $Q$  [6, p. 115, Theorem 1.6] it follows from the proof of the above corollary that a ring  $R$  which has no divisors of zero  $\neq 0$  can be one of only two Goldie dimensions (as a right  $R$ -module): either of dimension one (i.e. either  $R_R$  is a uniform module) or of infinite dimension (i.e.  $\mathcal{Z}(R) = (0)$ ). A ring  $R$  of the latter kind is the ring  $R = K[x, y]$  where  $K$  is a field and  $x, y$  are non-commuting indeterminates (but  $x\alpha = \alpha x$  and  $\alpha y = y\alpha$  for all  $\alpha \in K$ ). Since  $xR \cap yR = (0)$ ,  $R_R$  is not uniform and so  $\mathcal{Z}(R_R) = (0)$ .

I wish to thank Professor E. Enochs for patiently listening to me explaining the ideas in this paper and offering corrective and constructive suggestions.

## REFERENCES

1. E. P. Armendariz, *On finite-dimensional torsion-free modules and rings*, Proc. Amer. Math. Soc., **24** (1970), 566-571.
2. C. Faith, *Lectures on injective modules and quotient rings*, Springer-Verlag, New York; 1967.

3. E. Gentile, *Singular submodule and injective hull*, Indag. Math., **24** (1962), 426-433.
4. A. W. Goldie, *Semi-prime rings with maximum condition*, Proc. London Math. Soc., **10** (1960), 201-220.
5. ———, *Torsion-free modules and rings*, J. of Algebra, **1** (1964), 268-287.
6. F. L. Sandomierski, *Semi-simple maximal quotient rings*, Trans. Amer. Math. Soc., **128** (1967), 112-120.
7. ———, *Nonsingular rings*, Proc. Amer. Math. Soc., **19** (1968), 225-230.
8. J. Zelmanowitz, *Injective hulls of torsion free modules*, Canad. J. of Math., Vol. XXII, No. 6 (1971), 1094-1101.

Received September 15, 1971 and in revised form November 11, 1971.

UNIVERSITY OF KENTUCKY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California 94305

C. R. HOBBY

University of Washington  
Seattle, Washington 98105

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index. to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.



Wazir Husan Abdi, <i>A quasi-Kummer function</i> .....	521
Vasily Cateforis, <i>Minimal injective cogenerators for the class of modules of zero singular submodule</i> .....	527
W. Wistar (William) Comfort and Anthony Wood Hager, <i>Cardinality of <math>k</math>-complete Boolean algebras</i> .....	541
Richard Brian Darst and Gene Allen DeBoth, <i>Norm convergence of martingales of Radon-Nikodym derivatives given a <math>\sigma</math>-lattice</i> .....	547
M. Edelstein and Anthony Charles Thompson, <i>Some results on nearest points and support properties of convex sets in <math>c_0</math></i> .....	553
Richard Goodrick, <i>Two bridge knots are alternating knots</i> .....	561
Jean-Pierre Gossez and Enrique José Lami Dozo, <i>Some geometric properties related to the fixed point theory for nonexpansive mappings</i> .....	565
Dang Xuan Hong, <i>Covering relations among lattice varieties</i> .....	575
Carl Groos Jockusch, Jr. and Robert Irving Soare, <i>Degrees of members of <math>\Pi_1^0</math> classes</i> .....	605
Leroy Milton Kelly and R. Rottenberg, <i>Simple points in pseudoline arrangements</i> .....	617
Joe Eckley Kirk, Jr., <i>The uniformizing function for a class of Riemann surfaces</i> .....	623
Glenn Richard Luecke, <i>Operators satisfying condition <math>(G_1)</math> locally</i> .....	629
T. S. Motzkin, <i>On <math>L(S)</math>-tuples and <math>l</math>-pairs of matrices</i> .....	639
Charles Estep Murley, <i>The classification of certain classes of torsion free Abelian groups</i> .....	647
Louis D. Nel, <i>Lattices of lower semi-continuous functions and associated topological spaces</i> .....	667
David Emroy Penney, II, <i>Establishing isomorphism between tame prime knots in <math>E^3</math></i> .....	675
Daniel Rider, <i>Functions which operate on <math>\mathbb{F}L_p(T)</math>, <math>1 &lt; p &lt; 2</math></i> .....	681
Thomas Stephen Shores, <i>Injective modules over duo rings</i> .....	695
Stephen Simons, <i>A convergence theorem with boundary</i> .....	703
Stephen Simons, <i>Maximinimax, minimax, and antiminimax theorems and a result of R. C. James</i> .....	709
Stephen Simons, <i>On Ptak's combinatorial lemma</i> .....	719
Stuart A. Steinberg, <i>Finitely-valued <math>f</math>-modules</i> .....	723
Pui-kei Wong, <i>Integral inequalities of Wirtinger-type and fourth-order elliptic differential inequalities</i> .....	739
Yen-Yi Wu, <i>Completions of Boolean algebras with partially additive operators</i> .....	753
Phillip Lee Zenor, <i>On spaces with regular <math>G_\delta</math>-diagonals</i> .....	759