

# Pacific Journal of Mathematics

**CONCERNING BANACH SPACES WHOSE DUALS ARE  
ABSTRACT  $L$ -SPACES**

JONNIE BEE BEDNAR AND HOWARD E. LACEY

## CONCERNING BANACH SPACES WHOSE DUALS ARE ABSTRACT $L$ -SPACES

J. BEE BEDNAR AND H. ELTON LACEY

**The purpose of this paper is to give general methods for constructing Banach spaces whose duals are linearly isometric to abstract  $L$  spaces. These methods are based on annihilating certain subspaces of the duals of certain Banach spaces and in the existence of "affine" maps from a compact Hausdorff space  $X$  to the space of regular Borel measures on  $X$ .**

1. Introduction. There have been several papers concerned with the structure and classification of Banach spaces whose duals are linearly isometric to a space of the type  $L_1(\mu)$  (see [25] and its references). General methods for the generation of such spaces have been developed in [6], [8], and [16]. The purpose of this paper is to present these theorems in their most general framework.

In §2 a general result is presented which indicates several ways in which Banach spaces whose duals are  $L$ -spaces can be constructed. Using this as a base, §3 begins with an application of the main result of section two, Theorem 2.2, to show the way in which certain affine symmetric maps generate these spaces. Finally it is shown how some of these maps from compact Hausdorff spaces to the corresponding space of regular Borel measures generate the Banach spaces whose duals are  $L$ -spaces.

All Banach spaces considered in this paper are over the real field. If  $X$  is a compact Hausdorff space then  $C(X)$  denotes the Banach space of all real-valued continuous functions on  $X$  and  $M(X)$  the Banach space dual of  $C(X)$ . If  $\mu$  is a measure,  $L_1(\mu)$  is the Banach space of all integrable real-valued functions (sometimes called an (abstract)  $L$ -space). The dual of a Banach space  $A$  is denoted by  $A^*$ . If  $A$  is a Banach space and  $S \subset A$  and  $\{a_x: x \in S\}$  is a set of numbers, then  $\sum_{x \in S} (a_x)x$  denotes the limit (provided it exists) of the net of all sums  $\sum_{x \in F} (a_x)x$  for finite sets  $F \subset S$ . The notation and terminology regarding Choquet simplexes, maximal measures, affine functions, etc. is that of [26].

2. Methods of generating Banach spaces whose duals are  $L$ -spaces. The purpose of this section is to prove a general result concerning the generation of Banach spaces whose duals are  $L$ -spaces. How to construct such spaces as subspaces of given Banach spaces having the same property is revealed in Theorem 2.2.

If  $A$  is a Banach space and  $B$  is a subset of  $A$  then  $B^\perp = \{x^* \in A^*: x^*(x) = 0 \text{ for all } x \in B\}$ . For  $M \subset A^*$ ,  ${}^\perp M = \{x \in A: x^*(x) = 0$

for all  $x^* \in M$ ). A *contractive* operator on  $A$  is a linear operator  $T: A \rightarrow A$  satisfying  $\|T\| \leq 1$ . Of course the term “projection” has its usual meaning.

A Banach space  $A$  whose isometric image in any other Banach space  $B$  is the range of a contractive projection on  $B$  is called a  $P_1$  space. The relevant facts concerning  $P_1$  spaces can be found in [9].

The following proposition serves a dual purpose. It provides motivation to the main theorem of this section as well as the key to the main result of §3.

**PROPOSITION 2.1.** *Let  $A$  and  $B$  be Banach spaces with  $B \subset A$  and  $B^*$  an  $L$ -space. Then there is a contractive projection  $P: A^* \rightarrow A^*$  whose kernel is  $B^\perp$  and whose range is linearly isometric to  $B^*$  under the restriction map  $x^* \rightarrow x^*|_B$  for  $x^* \in A^*$ .*

*Proof.* Let  $J: B^* \rightarrow B^{***}$ ,  $K: A \rightarrow A^{**}$ ,  $L: B \rightarrow B^{**}$  and  $i: B \rightarrow A$  be the natural embeddings. Since  $B^*$  is an  $L$ -space,  $B^{**}$  is a  $P_1$  space [see 9, 10, or 17]. Hence there is a contractive projection  $Q: A^{**} \rightarrow A^{**}$  whose range is  $i^{**}(B^{**})$ . Set  $T = [(i^{**})^{-1}QK]^*J$ . Then  $\|T\| \leq 1$  and  $Tx^*(ix) = x^*(x)$  for all  $x^* \in B^*$  and  $x \in B$ . For,

$$\begin{aligned} & \langle [(i^{**})^{-1}QK]^*Jx^*, ix \rangle \\ &= \langle Jx^*, (i^{**})^{-1}QKix \rangle = \langle (i^{**})^{-1}QKix, x^* \rangle \\ &= \langle (i^{**})^{-1}Qi^{**}Lx, x^* \rangle = \langle Lx, x^* \rangle = x^*(x). \end{aligned}$$

Hence  $T$  is a linear isometry of  $B^*$  into  $A^*$  and  $P = Ti^*$  is the required projection.

For motivational purposes, let  $K$  be a compact Choquet simplex and  $X$  any closed subset of  $K$  which contains the extreme points,  $EK$ , of  $K$ . Setting  $A = \{f|X: f \in A(K)\}$ , where  $A(K)$  is the space of continuous affine functions on  $K$ , it follows from the Bauer maximum principle [4] that  $f \rightarrow f|X$  is an order preserving linear isometric isomorphism of  $A(K)$  onto  $A \subset C(X)$ . Since each maximal measure on  $K$  vanishes on  $K \setminus X$ , the maximal measures have support contained in  $X$ . For each  $x \in X$  let  $\mu_x$  be the maximal measure representing  $x$  (i.e.,  $f(x) = \int f d\mu_x$  for all  $f \in A$ ). The linear span,  $N$ , of the maximal probability measures on  $X$  is a band in  $M(X)$  i.e.  $M(X) = N \oplus N^\perp$  where  $N^\perp = \{\sigma \in \mu(X): |\mu| \wedge |\sigma| = 0 \text{ for all } \mu \in N\}$  and  $(N^\perp)^\perp = N$ , and so is the closed linear span,  $M$ , of the point masses  $\xi_x$  with  $x \in Y = X \setminus EK$ . For  $(\sum_{x \in Y} a_x \xi_x) \in M$ ,  $T(\sum_{x \in Y} a_x \xi_x) = \sum_{x \in Y} a_x \mu_x$  defines a contractive linear operator from  $M$  to  $N$ . It is well-known that  $A = {}^\perp\{T\nu - \nu: \nu \in M\} = \left\{f \in C(X): f(x) = \int f d\mu_x \text{ for all } x \in X\right\}$  and  $A^\perp$  is the weak\*-closure of  $\{T\nu - \nu: \nu \in M\}$ . Note that  $A^\perp \cap N = \{0\}$ .

For, suppose  $0 \neq \nu \in A^\perp \cap N$ . Then since  $1 \in A$ ,  $\|\nu^+\| = \|\nu^-\|$  and we can assume that  $\|\nu^+\| = \|\nu^-\| = 1$ . Since  $\nu^+(h) = \nu^-(h)$  for all  $h \in A$ ,  $\nu^+$  and  $\nu^-$  represent the same point of  $K$  and since  $\nu^+$ ,  $\nu^-$  are both maximal measures and  $K$  is a simplex,  $\nu^+ = \nu^-$  (see [26]). From this it follows immediately that  $N \cap A^\perp = \{0\}$ .

The main results of this section, Theorem 2.2 below, demonstrates that by placing the preceding remarks in a more general setting, one is able to generate conjugate  $L$ -spaces in a nice way. The notation is as follows:  $A$  is a Banach space whose dual is an  $L$ -space,  $P: A^* \rightarrow A^*$  is a contractive projection,  $N$  is the range of  $P$ ,  $M$  is the kernel of  $P$ ,  $Q = I - P$ ,  $T: M \rightarrow N$  is a bounded linear operator,  $M_0$  is the weak\*-closure of  $\{Tx^* - x^*: x^* \in M\}$  and  $B = \{x \in A: Tx^*(x) = x^*(x) \text{ for } x^* \in M\}$ .

**THEOREM 2.2.** *Preserve the notation above. Then each of the conditions below ensure that  $B^*$  is an  $L$ -space linearly isometric to  $N$ .*

(1)  $M_0 \cap N = \{0\}$  and  $\|P + TQ\| \leq 1$ .

(2)  $M$  is weak\*-closed,  $T$ , is weak\*-continuous,  $\|T\| < 1$ , and  $\|P + TQ\| \leq 1$ .

(3)  $N$  is a band,  $P$  is the band projection,  $M_0 \cap N = \{0\}$  and  $T$  is contractive.

(4)  $N$  is a band,  $P$  is the band projection,  $M$  is weak\*-closed,  $T$  is weak\*-continuous, and  $\|T\| < 1$ .

*Proof.* The range of any contractive projection on an  $L$ -space is also linearly isometric to an  $L$ -space (see [17] or [30]).

Hence it is only required to show that  $B^*$  in each case is linearly isometric to  $N$ . The proofs of this fact in statements (1)–(4) are given in the correspondingly numbered paragraphs below.

(1) Let  $S: N \rightarrow B^*$  be the restriction map. The  $\|Sx^*\| \leq \|x^*\|$  for all  $x^* \in N$  and  $Sx^* = 0$  implies that  $x^* \in B^\perp$ . Since  $M_0 \cap N = \{0\}$ ,  $S$  is one-to-one. For  $x^* \in B^*$  let  $y^* \in A^*$  with  $y^*|_B = x^*$  and  $\|y^*\| = \|x^*\|$ . Then  $z^* = (P + TQ)y^*$  belongs to  $N$  and

$$\begin{aligned} z^*(x) &= (P + TQ)y^*(x) \\ &= Py^*(x) + TQy^*(x) \\ &= y^*(x) \end{aligned}$$

for  $x \in B$ . Hence  $Sz^* = x^*$ . For  $x \in A$ ,  $|z^*(x)| \leq \|P + TQ\| |y^*(x)| \leq \|y^*\| \|x\|$  and  $S$  is a linear isometry.

(2) The conditions  $T$  is weak\*-continuous and  $\|T\| < 1$  imply that  $\{Tx^* - x^*: x^* \in M\}$  is weak\*-closed when  $M$  is. For if  $\|Tx^* - x^*\| \leq 1$ , then  $\|x^*\| \leq \|Tx^* - x^*\| + \|Tx^*\|$  and so  $\|x^*\| \leq$

$(\|Tx^* - x^*\|)/(1 - \|T\|)$ . By the usual compactness argument the unit sphere of  $\{Tx^* - x^*: x^* \in M\}$  is weak\*-compact and thus  $\{Tx^* - x^*: x^* \in M\}$  is weak\*-closed. Clearly,  $N \cap \{Tx^* - x^*: x^* \in M\} = \{0\}$  and (2) now follows from (1).

(3) The only difference in the proof of (3) and (1) is to notice that because  $P$  is the band projection,  $\|x^*\| = \|Px^*\| + \|Qx^*\|$  for every  $x^* \in A^*$ . Hence for  $y^* = (P + TQ)x^*$ ,

$$\begin{aligned} \|y^*\| &\leq \|Px^*\| + \|TQx^*\| \\ &\leq \|Px^*\| + \|T\| \|Qx^*\| \\ &\leq \|Px^*\| + \|Qx^*\| \\ &= \|Px^* + Qx^*\| \\ &= \|x^*\|. \end{aligned}$$

It now follows that the restriction map in (1) is an isometry.

(4) Combine the remarks of (2) and (3) above.

Contractive projections (in fact band projections) are plentiful in  $L$ -spaces and by Theorem 2.2, many of these projections in dual  $L$ -spaces generate Banach spaces whose duals are  $L$ -spaces.

To get a less abstract idea of situations analogous to the requirements of Theorem 2.2, some important special cases of the above theorem are summarized below.

1. If  $T = 0$  and  $M$  weak\*-closed (or (weak\*-closure of  $M) \cap N = \{0\}$ ), then  $B = {}^\perp M$ .

2. If  $N$  has finite codimension, then  $M$  and  $M_0$  are automatically weak\*-closed. So,  $T$  can be any linear operator satisfying  $\|P + TQ\| \leq 1$ , or in case  $P$  is the band projection onto a band  $N$ , just  $\|T\| \leq 1$ .

3. If  $P: A \rightarrow A$  is a contractive projection, then  $P^*: A^* \rightarrow A^*$  is a contractive projection and kernel  $P^* = (\text{Range } P)^\perp$  is already weak\*-closed.

4. Let  $X$  be a compact Hausdorff space and  $F \subset X$  a nonempty closed subset. Then  $A = \{f \in C(X): f(F) = 0\}$  is a closed sub-lattice of  $C(X)$  and  $A^\perp = \{\mu \in C(X)^*: \mu(X \setminus F) = 0\}$  is a weak\*-closed band in  $C(X)^*$ . Taking  $N$  as the complementary band to  $A^\perp$  and  $P$  the band projection provides a nice example of (4).

3. Generation of general Banach spaces whose duals are  $L$ -spaces. At this point it is natural to consider a rather simple application of Theorem 2.2. which allows the generation of Banach spaces whose duals are  $L$ -spaces. This application is presented in Theorem 3.2 below. First; however, additional terminology is need.

**DEFINITION 3.1.** Let  $R$  be an  $L$ -space with unit sphere  $S$ . A convex set  $F \subset S$  is called a *biface* if its linear span is a band in  $R$  (see [14] for a discussion of bifaces).

Examples of such sets are plentiful in  $L$ -spaces. Specifically the norm closed absolutely convex hull of any face of a compact Choquet simplex is such a set.

Within the notation of the definition above if  $E$  is any subset of the extreme points of  $S$  (assuming that  $S$  has extreme points), then  $E = \{\sum_{e \in E} (a_e)e : \sum_{e \in E} |a_e| \leq 1\}$  is a biface in  $L$ .

An additional example of a biface is given by the set of all regular Borel measures on a compact Hausdorff space  $X$  whose total variation is 0 on some Borel subset  $B$ . If  $X \setminus B$  is closed this biface is weak\*-closed and hence compact.

**THEOREM 3.2.** *Let  $A$  be a Banach space whose dual is an  $L$ -space and  $F$  a weak\*-closed biface in  $S(A^*)$ , the unit sphere of  $A^*$ . Let  $M$  be the span of  $F$  and  $N$  be the complementary band to  $M$ . If  $\alpha: F \rightarrow N \cap S$  is an affine symmetric weak\*-continuous function, then for  $B = \{x \in A: \alpha(x^*)(x) = x^*(x) \text{ for all } x^* \in F\}$ ,  $B^*$  is linearly isometric to  $N$ .*

*Proof.* Since  $F = M \cap S(A^*)$  and  $F$  is weak\*-closed,  $M$  is weak\*-closed (see [11]). Because  $\alpha$  is affine and symmetric, it has a unique weak\*-continuous linear extension  $T: M \rightarrow N$  with  $\|T\| \leq 1$ . Theorem 2.2 now completes the proof.

An interesting special case of the above is the following situation. Let  $x_1^*, \dots, x_n^*$  be positive extreme points of  $A^*$  and  $F = \{\sum_{i=1}^n a_i x_i^* : \sum_{i=1}^n |a_i| \leq 1\}$ . Letting  $y_1^*, \dots, y_n^*$  be any elements of  $A^*$  such that  $|y_i^*| \wedge |x_j^*| = 0$  for  $i, j = 1, \dots, n$ , define  $\alpha(\sum_{i=1}^n a_i x_i^*) = \sum_{i=1}^n a_i y_i^*$  to obtain a weak\*-continuous affine symmetric map from  $F$  to the complementary band determined by  $F$ . Then  $B = \{x \in A: y_i^*(x) = x_i^*(x) \text{ for } i = 1, 2, \dots, n\}$  is a Banach space whose dual is an  $L$ -space.

In particular, if  $X$  is a compact Hausdorff space  $x_1, \dots, x_n$  are in  $X$ ,  $\mu_1, \dots, \mu_n$  are regular Borel measures with  $\|\mu_i\| \leq 1$  and  $|\mu_i|(\{x_1, \dots, x_n\}) = 0$  for  $i = 1, 2, \dots, n$  then  $A = \{f \in C(X): f(x_i) = \int f d\mu_i; i = 1, 2, \dots, n\}$  is a Banach space whose dual is an  $L$ -space. This result was obtained by the first author in [6], later refined in [8] and later by Gleit in a different setting in [15].

In view of the preceding theorem, one is naturally lead to ask under what conditions maps from compact Hausdorff spaces to sets of measures induce Banach spaces whose duals are  $L$ -spaces. A. Gleit [15] has obtained some partial results. It is possible to use Theorems 2.2 and 3.2 to provide a complete answer.

To motivate subsequent results, return to the setting immediately following Proposition 2.1. That is, let  $K$  be a compact Choquet simplex,  $X$  any compact set containing  $EK$  and  $\mu_x$  be the unique maximal measure with resultant  $x \in K$ . For  $x \in X$  let  $\rho(x) = \mu_x$ . From [26, p. 71],  $\rho: X \rightarrow M(X)$  is Borel measurable (i.e.  $x \rightarrow \int f d\rho(x)$  is Borel measurable for each  $f \in C(X)$ ). It is also true [27] that  $f \rightarrow f_\rho$  where  $f_\rho(x) = \int f d\rho(x)$  for all Borel measurable functions is a positive projection when restricted to the space  $H = \{f + g_\rho: f, g \in C(X)\}$ . Equivalently,  $(f_\rho)_\rho = f_\rho$  for each continuous  $f$  on  $X$ . Denoting this projection by  $P$ , one has  $\|P\| = 1$  when  $H$  is given the uniform norm.

Effros has shown that the continuous functions on  $\overline{EK}$  which satisfy  $f(x) = \int f d\mu_x$  for  $x \in \overline{EK}$  are precisely those having a unique continuous affine extension to all  $K$  [13]. Obviously the statement remains valid if  $\overline{EK}$  is replaced by  $X$ . So  $A_\rho = \{f: f_\rho = f, f \in C(X)\}$  is linearly order isometric to  $A(K)$ .

Observe that if  $\int f d\mu = \int f d\nu$  for all  $f \in A_\rho$  where  $\mu, \nu$  are probability measures on  $X$ , then  $\mu$  and  $\nu$  have the same resultant in  $K$ . Appealing to [26, p. 63] it follows that the function defined by  $f_\rho(x) = \int f d\mu_x$  where  $f \in C(X)$  is affine and  $\int f_\rho d\mu = f_\rho(x)$  for any probability measure  $\mu$  on  $K$  having  $x$  as a resultant. Thus  $\int f_\rho d\mu = \int f_\rho d\nu$  for all  $f \in C(X)$ .

Thus  $A(K)$  is generated by a natural Borel measurable map  $\rho$  defined on a compact set  $X$  with  $EK \subset X \subset K$ . This map satisfies the following three conditions.

- (i)  $f_\rho$  is Borel measurable for each  $f \in C(X)$ ,
- (ii)  $\|\rho(x)\| \leq 1$  for all  $x \in X$ ,
- (iii) if  $\mu, \nu \in M(X)$  and  $\int f d\mu = \int f d\nu$  for all  $f \in A$ , then  $\int f_\rho d\mu = \int f_\rho d\nu$  for all  $f \in C(X)$ .

As will be demonstrated any map  $\rho: X \rightarrow M(X)$  having these three properties generates a Banach space whose dual is an  $L$ -space. The essential ingredient of condition (i) is that  $f_\rho$  is universally integrable (i.e. integrable for each  $\mu \in M(X)$ ).

DEFINITION 3.2. A function  $\rho: X \rightarrow M(X)$  is said to be *affine* if

- (i)  $f_\rho$  is universally integrable for all  $f \in C(X)$ ,
- (ii)  $\|\rho(x)\| \leq 1$  for all  $x \in X$
- (iii) if  $\mu, \nu \in M(X)$  and  $\int f d\mu = \int f d\nu$  for all  $f \in C(X)$  such that

$f = f_\rho$ , then  $\int f_\rho d\mu = \int f_\rho d\nu$  for all  $f \in C(X)$ .

If  $p$  is an affine map on a compact Hausdorff space  $X$ , then for  $\mu \in M(X)$ ,  $P\mu$  denotes the unique element of  $M(X)$  such that  $\int f dP\mu = \int f_\rho d\mu$  for all  $f \in C(X)$ , which exists since the map  $f \rightarrow \int f_\rho d\mu$  is a continuous linear functional on  $C(X)$ .

LEMMA 3.4. *Preserve the notation above. Then*

(1)  *$P$  is a contractive projection,*

(2) *the range of  $P$  is the set  $R = \{\mu: \mu \in M(X), \int (f - f_\rho) d\mu = 0$  for all  $f \in C(X)\}$ .*

(3) *the kernel of  $P$  is weak\*-closed.*

*Proof.* (1) that  $P$  is linear is trivial. Because  $\|\rho(x)\| \leq 1$ ,  $\|f_\rho\| \leq \|f\|$  and hence  $\|P\| \leq 1$ . Since for  $f = f_\rho \in C(X)$  one has  $\int f dP_\mu = \int f_\rho d\mu = \int f d\mu$ , it follows from (iii) that  $\int f dP[P\mu] = \int f_\rho dP\mu = \int f dP\mu$  for all  $f \in C(X)$  and, thus,  $P[P\mu] = P\mu$ .

(2) This follows immediately from the definition of  $P$  and (1).

(3) Let  $\{\mu_i\}$  be a net element in the kernel of  $P$  which is weak\*-convergent to  $\mu$ . Then  $\int f d\mu_i \rightarrow \int f d\mu$  for  $f \in C(X)$  and hence  $\int f d\mu = 0$  when  $f = f_\rho$ . Using (iii) it follows that  $\int f_\rho d\mu = 0$  for all  $f \in C(X)$  and thus  $P\mu = 0$ .

The following theorem is an immediate consequence of the above lemma and Theorem 2.2.

THEOREM 3.5. *Let  $\rho$  be an affine map from  $X$  to  $M(X)$ . Then the Banach space*

$$A\rho = \{f: f = f_\rho \in C(X)\}$$

*is a Banach space whose dual is the  $L$ -space  $\left\{\mu: \int (f - f_\rho) d\mu = 0 \text{ for } f \in C(X)\right\}$ .*

*Proof.* It suffices to show that  $A_\rho^\perp$  is the kernel of the contractive projection  $P$  defined through  $\rho$ . But trivially, by (iii) of the definition of affine map  $\int f d\mu = 0$  when  $f \in A$  implies  $\int f_\rho d\mu = 0$  for  $f \in C(X)$ . Hence  $P\mu = 0$ . As the converse is just as trivial,  $A_\rho^\perp$  is the kernel of  $P$ . Theorem 2.2 now completes the proof.



The significance of Theorem 3.5 is enhanced by the fact that every Banach space whose dual is an  $L$ -space arises in this fashion. The proof of this is based upon recent works of Lazar [23] and Effros [14].

Let  $A$  be a Banach space whose dual is an  $L$ -space and let  $S(A^*)$  denote the unit ball in  $A^*$ . Then  $S(A^*)$  is an absolutely convex weak\*-compact set and so the general Choquet theory may be used to obtain the maximal measures on  $S(A^*)$ . Such measures are supported by the weak\*-closure,  $X$  of the set of extreme points of  $S(A^*)$ .

Define the homeomorphism  $\sigma: S(A^*) \rightarrow S(A^*)$  by  $\sigma(x) = -x$ . Then  $\sigma$  induces natural order preserving isometries  $f \rightarrow \sigma f$  and  $\mu \rightarrow \sigma\mu$  on  $C(S(A^*))$  and  $M(S(A^*))$  respectively. These are defined by the formulas

$$\sigma f(x) = f(\sigma x)$$

and

$$\sigma\mu(C) = \mu(\sigma C) .$$

For  $\mu \in M(S(A^*))$  let  $\text{odd}(\mu) = 1/2(\mu - \sigma\mu)$ . From [23] it follows that  $\text{odd}(\mu) = \text{odd}(\nu)$  for any pair of maximal measures on  $S(A^*)$  having the same resultant in  $S(A^*)$ . Since each maximal measure on  $S(A^*)$  vanishes off  $X$ , they can be considered as maximal measures on  $X$ . For  $x \in X$  let  $\mu$  be any maximal measure which represents  $x$  and let  $\rho(x) = \text{odd} \mu$ . Then  $\rho: X \rightarrow M(X)$  and  $\|\rho(x)\| \leq 1$  for all  $x \in X$ .

It is shown in the sequel that  $\rho$  is an affine map. A positive measure  $\mu$  on  $S(A^*)$  is maximal if and only if  $\int f d\mu = \int \bar{f} d\mu$  for all  $f \in C(S(A^*))$  (see [26]). For such a measure it follows that  $\int f d\sigma\mu = \int \bar{f} d\sigma\mu$  and  $\int \sigma f d\mu = \int \sigma f d\mu$  for all  $f \in C(S(A^*))$ . The main theorem of [23] then demonstrates that

$$\frac{1}{2}[\bar{f}(x) - \sigma\bar{f}(x)] = \int \frac{1}{2}[f - \sigma f] d\mu$$

for any maximal measure with resultant  $x$  and continuous convex function  $f$  on  $S(A^*)$ . Since

$$\frac{1}{2} \int [f - \sigma f] d\mu = \int f d \text{odd}(\mu) ,$$

it is true that

$$(*) \quad \frac{1}{2}[\bar{f}(x) - \sigma\bar{f}(x)] = \int f d[\text{odd}(\mu)] .$$

The preceding remarks provide a portion of the proof of the lemma below.

**LEMMA 3.6.** *Preserve the notation above. Then  $\rho: X \rightarrow M(X)$  is affine.*

*Proof.* Condition (ii) of the definition is already verified. That condition (i) is true is an easy consequence of (\*) above. Only the verification of (iii) is lacking.

Define  $P: M(X) \rightarrow M(X)$  as follows: For  $\nu \geq 0$  in  $M(X)$  let  $\beta$  be any maximal measure which dominates in the ordering of Choquet (see [26]) and set  $P\nu = \text{odd } \beta$ . For arbitrary  $\nu = \nu^+ - \nu^- \in M(X)$  let  $P\nu = P\nu^+ - P\nu^-$ . Then  $P$  is linear,  $\|P\| \leq 1$  and because  $\text{odd } (\text{odd } \beta) = \text{odd } (\beta)$  for any  $\beta \in M(X)$ ,  $P$  is a projection.

The kernel of the projection  $P$  is the set of measures  $\mu$  in  $M(X)$  which satisfy  $\int f \rho d\mu = 0$  for all  $f \in C(X)$ . If  $P = 0$  and  $f$  is continuous and convex, then

$$\int f_\rho d\mu = \frac{1}{2} \int [\bar{f} - \sigma \bar{f}] d\mu = \frac{1}{2} \int [\bar{f} - \sigma \bar{f}] d\mu^+ - \frac{1}{2} \int [\bar{f} - \sigma \bar{f}] d\mu^-.$$

Now, let  $\mu_1, \mu_2$  be maximal measures majorizing  $\mu^+, \mu^-$  respectively in Choquet's ordering. Since  $\bar{f} - \sigma \bar{f}$  is affine and satisfies the barycentric formula [26, p. 100] one has

$$\begin{aligned} \int f_\rho d\mu &= \frac{1}{2} \int [\bar{f} - \sigma \bar{f}] d\mu_1 - \frac{1}{2} \int [\bar{f} - \sigma \bar{f}] d\mu_2 \\ &= \frac{1}{2} \int [f - \sigma f] d\mu_1 - \frac{1}{2} \int [f - \sigma f] d\mu_2 \\ &= \int f dP\mu^+ - \int f dP\mu^- = \int f dP\mu = 0. \end{aligned}$$

It is now clear that  $P\mu = 0$  implies  $\int f_\rho d\mu = 0$  for all  $f \in C(X)$ . On the other hand the condition  $\int f_\rho dP\mu = 0$  for all  $f \in C(X)$  together with the fact that  $\sigma f = -f$  if  $f \in A \subset A(S(A^*))$  implies that  $\int f d\mu = 0$  when  $f \in A$ . Let  $\mu_1$  and  $\mu_2$  be maximal measures whose resultants  $x_1$  and  $x_2$  are the same as those of  $\mu^+$  and  $\mu^-$ . If  $x_1 \neq x_2$  there is some  $g \in A$  such that  $g(x_1) \neq g(x_2)$ . For  $g$ ,

$$\int g d(\mu_1 - \mu_2) = g(x_1) - g(x_2) = \int g d(\mu^+ - \mu^-) \neq 0$$

is a contradiction and so  $x_1 = x_2$ . Thus  $\text{odd } (\mu_1) = \text{odd } (\mu_2)$  and  $P\mu = 0$ .

A consequence of the above is the fact that  $\int f_\rho d(I - P)\mu = 0$  for all  $\beta \in M(X)$ . This is so because the range of  $I - P$  is precisely the kernel of  $P$ . Suppose  $\beta \geq 0$  and  $\|\beta\| = 1$ . If  $\mu$  is a maximal measure which dominates  $\beta$  in Choquet's ordering and  $f \in C(S(A^*))$ , then since

$$f_\rho = \frac{1}{2}(\bar{f} - \sigma\bar{f}),$$

$$\begin{aligned} \int f_\rho dP\beta &= \frac{1}{4} \left[ \int \bar{f} d\mu - \int \bar{f} \sigma d\mu - \int \sigma \bar{f} d\mu + \int \sigma f d\sigma\mu \right] \\ &= \frac{1}{2} \left[ \int f d\mu - \int f d\sigma\mu \right] = \int f dP\beta. \end{aligned}$$

From this it follows that  $\int g dP\beta = \int g_\rho dP\beta$  for all  $g \in C(X)$  and all  $\beta \in M(X)$ . Hence  $\int f dP\beta = \int f_\rho d\beta$  for all  $f \in C(X)$ .

Arguing as in the case which showed that  $\{\mu: P\mu = 0\} = \{\mu: \int f_\rho d\mu = 0\}$  it is possible to show that  $\int f d\beta = 0$  for  $f = f_\rho$  implies that  $P\beta = 0$ . Hence if  $\mu, \nu \in M(X)$  and  $\int f d\mu = \int f d\nu$  when  $f = f_\rho$ ,  $P(\mu - \nu) = 0$  so that  $\int g dP\mu = \int g dP\nu$  for all  $g \in C(X)$ . Hence  $\int g_\rho d\nu = \int g_\rho d\mu$  for all  $g \in C(X)$ . This completes the proof of the lemma.

Lemma 3.6 is fundamental in the proof of 3.7 below.

**THEOREM 3.7.** *Suppose  $A$  is a Banach space whose dual is an  $L$ -space. Then there is a compact Hausdorff space  $X$  and affine map  $\rho: X \rightarrow M(X)$  such that*

$$A = \left\{ f \in C(X): f(x) = \int f d\rho(x) \right\}.$$

*Proof.* Let  $\rho$  and  $X$  be as in Lemma 3.6 and the remarks preceding it. By Lemma 3.6 and Theorem 3.5,  $A_\rho = \{f: f = f_\rho \in C(X)\}$  is a Banach space whose dual is an  $L$ -space. Since  $A^\perp = \{\mu: P\mu = 0\}$ ,  $A_\rho = A$  and the proof is complete.

These affine mappings can be used to obtain some known results in an efficient manner. For example, if  $\{K_i\}_{i \in I}$  is a family of compact Choquet simplexes and  $K = \prod_{i \in I} K_i$  is the product of the  $K_i$ 's, let  $\rho: K \rightarrow M(K)$  be defined by  $\rho(\{X_i\}) = \prod_{i \in I} \mu_i$  where  $\mu_i$  is the maximal measure representing  $x_i$  for each  $i \in I$ . Then  $\rho$  is an affine mapping and  $A_\rho = A(L)$  where  $L$  is the product simplex as defined in [21].

Another example is the following. Suppose  $A$  is a Banach space whose dual is an  $L$ -space and that the extreme points  $X$  of  $S(A^*)$

form a closed set. Let  $\Sigma: X \rightarrow X$  be defined by  $\Sigma(x^*) = -x^*$ . Then  $\Sigma$  is an involutory homeomorphism. The  $\rho$  of Theorem 3.7 is given by  $\rho(x^*) = 1/2[\xi_{x^*} - \xi_{\Sigma(x^*)}]$  and  $A = \{f \in C(X): f(x^*) = -f(\Sigma x^*)\}$ , i.e.,  $A = C_\Sigma(X)$ . (This was proved in [25].)

We close with an application which seems to be new. Let  $A$  be a Banach space whose dual is an  $L$ -space and suppose that the weak\*-closure  $X$  of the set  $E$  of extreme points of the unit sphere of  $A^*$  is  $E \cup \{0\}$ . The mapping defined by  $\sigma x^* = -x^*$  is an involutory homeomorphism on  $X$ , i.e.,  $\sigma^2 = \text{identity}$ .

**THEOREM 3.8.**  $A = C_o(X) = \{f \in C(X): f(x^*) = -f(\sigma x^*) \text{ for all } x^* \in X\}$ .

*Proof.* The mapping  $\rho$  is given by  $\rho(x^*) = 1/2[\xi_{x^*} - \xi_{\sigma x^*}]$  for  $x^* \in E$  and  $\rho(0) = 0$ . By Theorem 3.7,

$$A = A_\rho = \left\{ f \in C(X): f(x^*) = \int f d\rho(x^*) \right\} = C_o(X).$$

This result is similar to the one for  $C_\Sigma(X)$  above when the extreme points are closed.

We wish to thank the referee for his many valuable comments.

## REFERENCES

1. E. Alfsen, *On the Geometry of Choquet Simplexes*, Math. Scand. **15** (1964), 97-110.
2. ———, *Boundary values for homomorphisms of compact convex sets*, Math. Scand., **19** (1966), 113-121.
3. ———, *On the Dirichlet problem of the Choquet boundary* Acta. Math., **120** (1968), 149-159.
4. H. Bauer, *Konvexität in Topologischen Vektorräumen*, Lecture Notes, University of Hamburg, 1963/1964.
5. ———, *Schilowschev Rand und Dirichletsches Problem*, Amer. Inst. Fourier, **11** (1961), 89-136.
6. J. Bee Bednar, *Compact convex Sets and Continuous Affine Functions*, Dissertation, The University of Texas at Austin, Austin, Texas (1968).
7. ———, *On the Dirichlet problem for functions on the extreme boundary of a compact convex set*, to appear in Math. Scand.
8. J. Bee Bednar and H. E. Lacey, *On spaces whose duals are abstract (L)-spaces and Choquet simplexes*, unpublished notes.
9. M. M. Day, *Normed Linear Spaces*, Academic Press, Inc., New York (1962).
10. J. Dixmier, *Sur certaines espaces consideres par M. H. Stone*, Sum Bras. Math. (1951), 151-181.
11. N. Dunford, and J. T. Schwartz, *Linear Operators, Part I*, Interscience Publishers, Inc., New York, (1964).
12. E. G. Effros, *Structure in simplexes*, Acta Math., **117** (1967), 103-121.
13. ———, *Structure in simplexes II*, J. Functional Analysis **1** (1967), 379-391.
14. ———, *On some real Banach spaces*, To appear.
15. E. G. Effros and Gleit A., *Structure in simplexes III*, Trans. Amer. Math. Soc., **142** (1969), 355-379.

16. A. Gleit, *On the existence of simplex spaces*, Israel J. Math., **9**, no. 2, (1971), 199-209.
17. A. Grothendieck, *Une caracterization Vectorielle Metrique des Espaces  $L^1$* , Canad. J. Math., **7** (1955), 552-561.
18. S. Kakutani, *Concrete representation of abstract ( $M$ ) spaces*, Annals of Math., **42**, 994-1024.
19. ———, *Concrete Representations of abstract ( $L$ ) spaces*, Annals of Math., **42**, 523-537.
20. Johannes Kohn, *Barycenters of Unique maximal Measures*, J. Fnal. Anal., **6** (1970), 78-82.
21. A. J. Lazar, *Affine products of simplexes*, Math. Scand., **22** (1968), 165-175.
22. ———, *Spaces of affine continuous functions on simplexes*, Trans. Amer. Math. Soc., **134** (1968), 503-525.
23. ———, *The unit ball in conjugate ( $L$ ) spaces*, to appear.
24. A. J. Lazar and J. Lindenstrauss, *On Banach spaces whose duals are  $L^1$ -spaces*, Israel J. Math., **4** (1966), **3**, 205-206.
25. J. Lindenstrauss and D. Wulbert, *On the classification of the Banach spaces whose duals are  $L_1$  spaces*, J. Fnal. Anal. **4**, No. 4, (1969).
26. R. R. Phelps, *Lectures on Choquet's Theorem*, D. Van Nostrand, Princeton, N. J., (1966).
27. Marc Rogalski, *Espaces de Banach ordonne's simplexes, frontieres de Silov et Probleme de Dirichlet*, C. R. Acad. Sci. Paris, **263**, Nov. (1966), Sec. A, 726-729.
28. ———, *Operateurs de Lion, projections boriliens et simplexes analytiques*, J. Fnal. Analysis **2** (1968), 458-488.
29. Z. Semadeni, *Free compact convex sets*, Bull. de l'Acad. Pol. des. Sci., **13** 2, (1965), 141-146.
30. L. Tzafriri, *Remarks on contractive Projections in  $L_p$ -spaces*, Israel J. Math., **7**, No. 1, (1969), 9-15.

Received February 11, 1971 and in revised form June 11, 1971.

UNIVERSITY OF TULSA  
AND  
UNIVERSITY OF TEXAS

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

Anatole Beck and Peter Warren, <i>Weak orthogonality</i> .....	1
Jonnie Bee Bednar and Howard E. Lacey, <i>Concerning Banach spaces whose duals are abstract L-spaces</i> .....	13
Louis Harvey Blake, <i>Canonical extensions of measures and the extension of regularity of conditional probabilities</i> .....	25
R. A. Brooks, <i>Conditional expectations associated with stochastic processes</i> .....	33
Theodore Allen Burton and Ronald Calvin Grimmer, <i>On the asymptotic behavior of solutions of <math>x'' + a(t)f(x) = e(t)</math></i> .....	43
Stephen LaVern Campbell, <i>Operator-valued inner functions analytic on the closed disc</i> .....	57
Yuen-Kwok Chan, <i>A constructive study of measure theory</i> .....	63
Alexander Munro Davie and Bernt Karsten Oksendal, <i>Peak interpolation sets for some algebras of analytic functions</i> .....	81
H. P. Dikshit, <i>Absolute total-effective <math>(N, p_n)(c, 1)</math> method</i> .....	89
Robert E. Edwards, Edwin Hewitt and Kenneth Allen Ross, <i>Lacunarity for compact groups. II</i> .....	99
James Daniel Halpern, <i>On a question of Tarski and a maximal theorem of Kurepa</i> .....	111
Gerald L. Itzkowitz, <i>A characterization of a class of uniform spaces that admit an invariant integral</i> .....	123
Mo Tak Kiang, <i>Semigroups with diminishing orbital diameters</i> .....	143
Glenn Richard Luecke, <i>A class of operators on Hilbert space</i> .....	153
R. James Milgram, <i>Group representations and the Adams spectral sequence</i> .....	157
G. S. Monk, <i>On the endomorphism ring of an abelian <math>p</math>-group, and of a large subgroup</i> .....	183
Yasutoshi Nomura, <i>Homology of a group extension</i> .....	195
R. Michael Range, <i>Approximation to bounded holomorphic functions on strictly pseudoconvex domains</i> .....	203
Norman R. Reilly, <i>Inverse semigroups of partial transformations and <math>\theta</math>-classes</i> .....	215
Chris Rorres, <i>Strong concentration of the spectra of self-adjoint operators</i> .....	237
Saharon Shelah, <i>A combinatorial problem; stability and order for models and theories in infinitary languages</i> .....	247
George Gustave Weill, <i>Vector space decompositions and the abstract imitation problem</i> .....	263
Arthur Thomas White, <i>On the genus of the composition of two graphs</i> .....	275