

Pacific Journal of Mathematics

NON-OPENNESS AND NON-EQUIDIMENSIONALITY IN ALGEBRAIC QUOTIENTS

JOHN DUSTIN DONALD

NON-OPENNESS AND NON-EQUIDIMENSIONALITY IN ALGEBRAIC QUOTIENTS

JOHN D. DONALD

Suppose given an equivalence relation R on an algebraic variety V and the associated fibering of V by a family of subvarieties. This paper treats the question of the existence of a quotient structure for this situation when the fibering is non-equidimensional. For this purpose a general definition of quotient variety for algebraic equivalence relations is used which contains no topological requirements.

The results are of two types. In §1 it is shown that certain maps into nonsingular varieties are quotient maps for the induced equivalence relation whenever the union of the excessive orbits has codimension ≥ 2 . This theorem yields many examples of non-equidimensional quotients. Section 2 contains a converse showing that no excessive orbit containing a normal hypersurface can be fitted into a quotient. This theorem depends on a stronger and less conceptual field-theoretic result which fails without the normality hypothesis. Section 3 contains a counterexample.

These results are a first attempt at answering the question: when can a good algebraic structure be assigned to a non-equidimensional fibering of V ? Analogously, one might ask whether there is a good sense in which an equidimensional family of subvarieties of V can degenerate into subvarieties of different dimension. It is a standard result that no fibers of a morphism have dimension less than that of the generic fiber, so that we must concern ourselves only with degeneration into higher-dimensional subvarieties. Thus if the equivalence relation R comes from action of an algebraic group G , in which case any exceptional orbits are *smaller* than the generic ones, there can be no non-equidimensional quotients. In this classical situation, then, existence of a quotient map $p: V \rightarrow V/G$ depends on equidimensionality of R . Moreover, such a quotient map is always open, whence topological. Since the substantial volume of recent work on the quotient problem mostly focuses on group actions, the typical definition of quotient map includes the requirement of openness, or even universal openness (cf. Mumford's book [2], p. 4, for example). Now a non-equidimensional morphism is never open (cf. Proposition 1), so we must abandon the topological point of view here.

A number of questions arise that we do not consider in this paper. Is there any relationship between exceptional fibers of a quotient and complete (or noncomplete) intersections? What local

data at an exceptional orbit relate to the existence locally of a quotient? For example, flatness is out. Finally, is there any further use for the concept of level field introduced in §2 and 3 for the purpose of proving Theorem 2?

O. Preliminaries. We use the following definitions: An *algebraic equivalence relation* or *ER* R on a variety V (assumed hereafter to be *irreducible*, for simplicity) is an algebraic subset of $V \times V$ which, as a set, is an equivalence relation. An *R -invariant function* f on V is one whose values are constant on the intersection of domain (f) with any R -equivalence class. The R -invariant functions thus constitute a subsheaf of the structure sheaf \mathcal{O}_V of V . A *quotient* for the pair (V, R) is a morphism of varieties $p: V \rightarrow W$ which as a map on sets is a quotient, and which satisfies the local condition

LC: the sheaf of functions on W is identified via p with the sheaf of R -invariant functions on V .

These general quotients have many expected properties, for which my paper [1] provides an elementary exposition. In particular we have

PROPOSITION 1. *Let R be an ER on the irreducible variety V with quotient $p: V \rightarrow W$. Then p is an open map if and only if it is equidimensional.*

Proof. If p is equidimensional, then p is open by a result of Rosenlicht ([4], Lemma 2) on algebraic group quotients. His proof goes through in our more general setting word for word.

For the converse, we apply the following general result.

PROPOSITION 2. *An open morphism of irreducible varieties is equidimensional.*

We give an elementary proof of this to conclude the proof of Proposition 1. Let $f: V \rightarrow W$ be the morphism. There is an open dense U of W such that for all $u \in U$, $f^{-1}(u)$ is equidimensional and $\dim f^{-1}(u) = \dim V - \dim W$. Let $w \in S = W - U$ be a point such that $\dim f^{-1}(w) > \dim V - \dim W$. We get a contradiction as follows.

Let C be an irreducible curve in W through w but not contained in S . Then $f^{-1}(C) \not\subset f^{-1}(S)$. Let X be the union of the components of $f^{-1}(C)$ not contained in $f^{-1}(S)$. X is the closure of $f^{-1}(C - S)$ and $\dim X = \dim V - \dim W + 1$. Now

$$\dim (X - f^{-1}(C - S)) < \dim X \leq \dim f^{-1}(w)$$

and $f^{-1}(w) \cap f^{-1}(C - S)$ is empty, so not every component of $f^{-1}(w)$ is in X . Let $v \in f^{-1}(w) - X$. Then $V - X$ is open and contains v but $f(V - X)$ can't be open—it contains w but misses a dense of C .

We see that for our initial problem to have any meaning, we must abandon the obvious topological restriction on quotients, realizing also that the quotients of the new type will probably fail to be topological.

Our notations are essentially standard. Ω stands for a universal domain. $\mathcal{O}_{X,Y}$ is the local ring of a subvariety X of the variety Y . $\text{Tr. deg. } A/\Omega$ denotes the transcendence degree of A over Ω .

1. **An existence theorem.** We can restate the local condition LC for quotients as follows: We are considering the set-wise quotient map $p: V \rightarrow W$ for the ER R on V . Let Ω be a universal domain and let $g \in \Omega(V)$ be an R -invariant function, so that within its domain, g is constant on orbits. Let $p(v) = w$ for $v \in V$.

From LC we know

LC_1 : given an R -invariant $g \in \Omega(V)$ there exists $f \in \Omega(W)$ such that $f \circ p = g$.

Furthermore, LC then implies

LC_2 : for all $f \in \Omega(W)$, $f \circ p \in \mathcal{O}_{v,V}$ if and only if $f \in \mathcal{O}_{w,W}$.

Now, conversely, LC_1 and LC_2 yield LC . In fact it is well-known that $g \in \Omega(V)$ is constant (or generically constant) on the fibers of $p: V \rightarrow W$ if and only if g is a purely inseparable element in the induced field extension $\Omega(V)/\Omega(W)$. Thus a set-wise quotient satisfying LC_2 is a quotient if and only if $\Omega(V)/\Omega(W)$ is a field extension with no pure inseparability. Since our first result is unrelated to any field-theoretic problem, we define a *quotient within inseparability*, QWI , to be a set-wise quotient satisfying LC_2 .

THEOREM 1. *Let V be an irreducible variety, let W be a nonsingular variety, and let $p: V \rightarrow W$ be a surjective morphism. Let $E = \{w \in W: \dim p^{-1}(w) > \dim V - \dim W\}$. If $\text{codimension } p^{-1}(E) > 1$, then p is a QWI for the induced ER , i.e. the one whose orbits are the fibers of p .*

Proof. We must show LC_2 . Let $f \in \Omega(W)$ be a function for which $f \circ p \in \mathcal{O}_{v,V}$. Replacing V by an open subvariety, if necessary, we may express f as g/h in lowest terms over the UFD $\mathcal{O}_{p(v),W}$. If

$f \notin \mathcal{O}_{p(v), W}$, then f has a pole at $p(v)$, so $h(p(v)) = 0$. From $f \circ p \in \mathcal{O}_{v, V}$, we see $g(p(v)) = 0$ also. Denote by $\mathcal{Z}(\cdot)$ the zeros of (\cdot) . Then $\mathcal{Z}(h \circ p) \subset \mathcal{Z}(g \circ p)$, so that in a suitable neighborhood of $p(v)$ we have $p(\mathcal{Z}(h \circ p)) \subset \mathcal{Z}(h)$. Now $\mathcal{Z}(g) \cap \mathcal{Z}(h)$ has codimension 2 near $p(v)$ by the relative primeness of g and h , whence $p^{-1}(E) \supset \mathcal{Z}(h \circ p)$. This contradicts codimension $p^{-1}(E) > 1$.

COROLLARY. *If p as above is birational, then either p is an isomorphism or codimension $p^{-1}(E) = 1$.*

Proof. Birational quotients are isomorphisms.

NOTE: The corollary is essentially *ZMT* for nonsingular varieties, and I have found precisely the same proof of it in this case in Mumford ([3], pp. 415–416).

The theorem also shows that with obvious meanings the product of two quotients is the quotient of a product ER when the quotients are of the type indicated in the theorem. I don't know if this standard categorical property is true in all non-equidimensional cases.

EXAMPLE. Using Theorem 1 we can easily construct non-equidimensional quotients.

(a) A general technique is to use subvarieties of $\Omega^n \times \Omega^m$ that project onto one factor. For example, let $V = \langle x_1x_2 + x_1^2x_3 + x_1^3x_4 = x_2 \rangle$ in $\Omega \times \Omega^3$, with $p = pr_2: V \rightarrow \Omega^3$. Then $E = \{0, 0, 0\}$, and codimension $p^{-1}(E) = 2$. Thus p is a *QWI* and, by the separability of p , even a quotient. By the next theorem this is the lowest dimensional example for a normal variety of a non-equidimensional quotient— $\dim V = \dim p(V) = 3$.

Disturbingly enough, though this quotient apparently has the quotient topology (though of course the map is not open), there are many open denses of V for which the restricted quotient fails to have the quotient topology! For example, let $V' = V - \{1, 0, 0, 0\}$. Then the line $\langle x_3 = x_4 = 0 \rangle$ in Ω^3 has an inverse image in V' consisting of two disjoint R_V -saturated closed sets, $X_1 = \langle x_1 = 1, x_3 = x_4 = 0 \rangle$ and $X_2 = \langle x_2 = x_3 = x_4 = 0 \rangle$. Now $p(V' - X_1)$ is not open, but $V' - X_1$ is open saturated.

(b) An example with messier fibers, but for which V and W are nonsingular, is the following: $p: \Omega^4 \rightarrow \Omega^3: (x, y, z, w) \rightarrow (xy - zw, y^2 - zw, x^2 - yz)$. p is surjective, with $E = \{(0, 0, 0)\}$ and $p^{-1}(E) = \text{cone of the twisted cubic} = \{(u^2t, ut^2, t^3, u^3) \mid u, t \in \Omega\}$.

(c) Examples similar to (a) but without the condition on codimension $p^{-1}(E)$ do not seem to yield quotients, even when the fibers over E are not hypersurfaces. Let $V = \langle x_1x_3 + x_1^2x_4 = x_2x_3 \rangle$ in $\Omega \times \Omega^3$

with $p = pr_2: V \rightarrow \Omega^3$. Then $E = \langle x_3 = x_4 = 0 \rangle$ and $p^{-1}(E)$ is a hyper-surface, while the exceptional fibers all have codim 2. Here p is not a quotient: for example, $x_3/x_4 = x_3^2/x_2 - x_1$ is not defined at $(0, 0, 0)$, but composing with p gives a function defined along $p^{-1}((0, 0, 0)) = \{(x, 0, 0, 0) \mid x \in \Omega\}$.

Since $pr_2: \Omega \times \Omega^3 \rightarrow \Omega^3$ is a quotient, another consequence of the example is that restricting quotients to arbitrary subvarieties needn't even yield QWI 's. (Of course, there are simpler examples of this phenomenon involving loss of normality. Consider $pr_2: \Omega \times \Omega^2 \rightarrow \Omega^2$ restricted to $\{(x, x^2, x^3) \mid x \in \Omega\}$.)

2. A non-existence theorem. The following partial converse to Theorem 1 limits unknown cases to ones like that of Example (c) above.

THEOREM 2. *If $p: V \rightarrow W$ is a quotient of irreducible varieties, then no fiber of excessive dimension contains a normal hypersurface.*

Proof. Identify $\Omega(W)$ with a subfield of $\Omega(V)$ via $\text{cohom } p$. Now if F is a fiber of p , the elements of $\Omega(W)$ are either constant on F or nowhere defined on F . Call any subfield K of $\Omega(V)$ with this property for a subvariety F of V a *level field* for F .

Exceptional orbits can exist only if $\text{tr. deg. } \Omega(W)/\Omega > 1$. Note also that we can delete all but one component of any orbit without destroying the quotient property. Our theorem can thus be restated in slightly stronger form:

THEOREM 3. *If H is a normal irreducible hypersurface and if K is a level field for H , then $\text{tr. deg. } K/\Omega \leq 1$.*

By example (a), Theorem 3 is false for codimension $H > 1$. We proceed now in two steps to prove Theorem 3 and hence Theorem 2.

Step 1. Here we do not use the level field concept. Let f be any minimal prime in the discrete valuation ring \mathcal{O}_H . Let L denote the residue field $\mathcal{O}_H/f\mathcal{O}_H$, and let $\pi: \mathcal{O}_H \rightarrow L$ be the residue map. Although we do not introduce the technique of completion for the following lemmas, one can see that from one point of view we are considering the first nonzero coefficients in the f -power series expansions of various functions.

LEMMA 1. *Let x_1, \dots, x_n in \mathcal{O}_H form a transcendence base for $\Omega(V)/\Omega$. Then the image $\pi(R)$ in L of $R = \mathcal{O}_H \cap \Omega[x_1, \dots, x_n, f^{-1}]$ contains a transcendence base of L/Ω .*

Proof of Lemma 1. Suppose the contrary and let $c \in L$ be algebraically independent of $\pi(R)$. Let $\pi(y) = c$ for $y \in \mathcal{O}_H$. There is an equation

$$(*) \quad y^m + P_{m-1}(x)y^{m-1} + \cdots + P_0(x) = 0$$

of algebraic dependence of y on $\Omega(x)$. We want to have the coefficients of $(*)$ in R and to guarantee this we adjust y .

First we need an element $u \neq 0$ in $R \cap f\mathcal{O}_H$. There exists a nonzero polynomial Q such that $Q(\pi(x_1), \dots, \pi(x_n)) = 0$ since the transcendence degree of L/Ω is less than that of $\Omega(V)/\Omega$. Now $Q(x_1, \dots, x_n) \neq 0$ by algebraic independence of the x_i , so this is our u .

For sufficiently large N , multiplying $(*)$ through by u^{mN} gives us an equation of integral dependence of $u^N y$ on R . Thus $u^N y \in \mathcal{O}_H$. For some M , $f^{-M}u^N y$ is in $\mathcal{O}_H - f\mathcal{O}_H$. Moreover its image in L is still algebraically independent of $\pi(R)$. Thus replacing y by $u^N y$, we may take $(*)$ to be an equation of integral dependence of $y \in f^M \mathcal{O}_H$ on R such that $\pi(f^{-M}y) = c$. Let n_r be the order of $P_r(x)$ in the valuation ring \mathcal{O}_H . Then $rM + n_r$ is the order of $y^r P_r(x)$. Let p be the minimal order among the $rM + n_r$. Dividing out f^p from $(*)$ and mapping down to L , we get 0 on the right and a sum of nonzero terms

$$\sum_{\text{various } r} c^r \pi(f^{-n_r} P_r(x))$$

on the left. The algebraic independence of c over the set of coefficients gives us a contradiction.

LEMMA 2. For any elements $x_1, \dots, x_r \in \mathcal{O}_H$,

$$\text{tr. deg. } \pi(\mathcal{O}_H \cap \Omega[x_1, \dots, x_r, f^{-1}])/\Omega \leq r.$$

Proof of Lemma 2. We may assume $\text{tr. deg. } \Omega[x_1, \dots, x_r, f^{-1}]/\Omega = r + 1$. Let P_1, \dots, P_s be polynomials in the x_i and f^{-1} with algebraically independent images under π . For simplicity, we can take the P_i to be monomials in f^{-1} . Since the case of no x_i 's is immediate we also assume inductively that at most $r - 1$ P_i 's do not involve x_r and that the π -images of these form a transcendence base for $\pi(\mathcal{O}_H \cap \Omega[x_1, \dots, x_{r-1}, f^{-1}])$. If $s \geq r$, then we have $P_r = (f^{-ab}x_r^b)(f^{-c}M)$ for some polynomial M in x_1, \dots, x_{r-1} and with neither factor in parentheses divisible by f . We conclude that $\pi(f^{-ab}x_r^b)$ is algebraically independent of $\{\pi(P_1), \dots, \pi(P_{r-1})\}$ and algebraically dependent on $\{\pi(P_1), \dots, \pi(P_r)\}$; i.e. we can exchange $f^{-ab}x_r^b$ for P_r . Since no two polynomials of the form $f^{-ab}x_r^b$ are algebraically independent, we obtain $s \leq r$.

NOTE. A less elementary, but quicker, proof of Lemma 2 results from considering the embedding of $\mathcal{O}_H/f\mathcal{O}_H$ in the completion of \mathcal{O}_H

with respect to $f\mathcal{O}_H$. In that case it is easy to check that f is transcendental over the embedded image of $\pi(\mathcal{O}_H \cap \Omega[x_1, \dots, x_r, f^{-1}])$.

Step 2. We now introduce the level field property of K into the situation. We have by definition $\pi(K \cap \mathcal{O}_H) = \Omega$. We use this fact and some easy manipulations to contradict the lemmas of Step 1 if $\text{tr. deg. } K/\Omega \geq 2$.

We may always choose a transcendence base for K from $f\mathcal{O}_H$ by subtracting suitable constants if necessary. Let $g \in K \cap f\mathcal{O}_H$ have minimal order r , so that $g = f^r u$ with u a unit. Let $h \in K \cap f\mathcal{O}_H$ have order m . By minimality of r , $m = qr$ for an integer q , and hg^{-q} takes a constant value, say $c_0 \neq 0$, on H . Thus $hg^{-q} - c_0 \in K \cap f^r \mathcal{O}_H$, whence $h = c_0 g^q + h_1$, $h_1 \in K \cap f^{qr} \mathcal{O}_H$. Continuing by induction, h has a finite expansion

$$(*) \quad h = \sum c_i g^{q^i} + h_s, \quad h_s \in K \cap f^{q^r s} \mathcal{O}_H.$$

Suppose now that for $\alpha \in \Omega[g, h, x_3, \dots, x_n, f^{-1}]$, $\pi(\alpha) \notin \Omega$. Let s be such that $q^r s > \text{highest power of } f^{-1} \text{ appearing in } \alpha$. Then making the substitution $(*)$ for h in α , we see that $\pi(\alpha) \in \pi(\Omega[g, x_3, \dots, x_n, f^{-1}])$. Thus (using Lemma 2 also)

$$\text{tr. deg. } \pi(\Omega[g, h, x, \dots, x_n, f^{-1}]) \leq n - 1.$$

Since we may take $x_3, \dots, x_n \in \mathcal{O}_H$ as independent transcendentals independent of $\Omega(g, h)$, we conclude from Lemma 1 that h is algebraic over $\Omega(g)$.

An immediate corollary of the above reasoning is the

COROLLARY. *With notation as in Lemma 1, there is at most one x_i such that*

$$\text{tr. deg. } \pi(\mathcal{O}_H \cap \Omega[x_i, f^{-1}])/\Omega = 0.$$

In terms of power series in a minimal prime, this implies that any function in \mathcal{O}_H whose f -power series has constant coefficients is algebraic over $\Omega(f)$.

We remark that in general for subvarieties of higher codimension one may write down level fields of excessive transcendence degree without reference to quotient maps. For example, consider $\Omega(x, y, xu + yv)$ as a level field for the plane $x = y = 0$ in 4-space.

We also note that it is not clear what can happen if the union of excessive orbits is a normal hypersurface, a situation intermediate between Theorems 1 and 2. A much more delicate problem is here

not considered at all, namely to give a decent local condition on an excessive orbit which would guarantee existence of a quotient nearby.

3. **A counterexample.** We give an example of a level field of transcendence degree 2 for a singular curve Z on a surface V in characteristic 0. This counterexample to a generalization of Theorem 3 suggests that if Theorem 2 is true without normality hypotheses, a more geometric approach will be needed to prove it.

THEOREM 4. *Let V be the product of the line Ω with the curve with a cusp $\{(x^2, x^3)\} \subset \Omega^2$, so that V embeds as $\{(x^2, x^3, y)\} \subset \Omega^3$. Let Z be the singular line $\{(0, 0, y)\}$. Then the field $K = \Omega(x^2(x + y), x^2y^2)$ is a level field for Z in characteristic 0.*

Proof. Set $u = x^2(x + y)$, $v = x^2y^2$. One checks that the map $(x^2, x^3, y) \mapsto (u, v)$ is a surjection from V onto Ω^2 whose only excessive fiber is Z . (The map is not quite a quotient, however, since for example $v/(v + u^2)$ is defined along Z but not at its image $(0, 0)$.) Thus relatively prime polynomials f and g in u, v , when viewed as polynomials in x, y , can have only powers of x as a common factor.

Let $f(u, v) = H_1(u, v) + F$, $g(u, v) = H_2(u, v) + G$, where H_1, H_2 are homogeneous in u, v , and F, G consist of higher powers. By the above remark, a typical element f/g of $\Omega(u, v)$ is in lowest terms when viewed as a quotient of polynomials in x, y unless $\deg. H_1 > 0$, $\deg. H_2 > 0$. Cancelling x^{2d} where $d = \min(\deg. H_1, \deg. H_2)$ to yield a rational function in x, y in lowest terms, we see that f/g has a pole along Y unless $\deg. H_1 \geq \deg. H_2$ and has a zero along Z if $\deg. H_1 > \deg. H_2$. Let $\deg. H_1 = \deg. H_2 = d$ and set $H_1 = \sum a_i u^i v^{d-i}$, $H_2 = \sum b_i u^i v^{d-i}$. We have $f/g = (Y_1 + A + R)/(Y_2 + B + S)$ where

$$Y_1 = \sum a_i y^{2d-i}, \quad Y_2 = \sum b_i y^{2d-i},$$

$$A = x \sum i a_i y^{2d-i-1} = x \left(2dy^{-1} Y_1 - \frac{d}{dy} (Y_1) \right),$$

$$B = x \sum i b_i y^{2d-i-1} = x \left(2dy^{-1} Y_2 - \frac{d}{dy} (Y_2) \right),$$

and both R and S are divisible by x^2 . Multiplying top and bottom by $B - (Y_2 + S)$, we see that f/g is defined on Z if and only if

$$(Y_1 + A + R)(B - Y_2 - S) \in \Omega[x^2, x^3, y]$$

$$\Leftrightarrow Y_1 B = Y_2 A$$

$$\Leftrightarrow Y_2 \frac{d}{dy} (Y_1) = Y_1 \frac{d}{dy} (Y_2)$$

$$\Leftrightarrow Y_1 = c Y_2, \quad c \in \Omega \quad (\text{char. } \Omega = 0).$$

But then $(f/g)|_Z = c$, a constant.

REFERENCES

1. John Donald, *Quotients for algebraic equivalence relations*, I: *Elementary existence theorems*, Unpublished.
2. David Mumford, *Geometric Invariant Theory*, New York, Springer-Verlag, 1965.
3. ———, *Introduction to Algebraic Geometry* (preliminary version). Harvard University Math. Dept.
4. Maxwell Rosenlicht, *On quotient varieties and the affine embedding of certain homogeneous spaces*, Trans. Amer. Math. Soc., **101**, pp. 211-223.

Received February 5, 1971 and in revised form December 27, 1971. Much of this material appeared in my Ph. D. thesis (Berkeley 1967). I would like to thank Professor Maxwell Rosenlicht again for his guidance in that effort.

UNIVERSITY OF CALIFORNIA, SAN DIEGO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 108 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Tom M. (Mike) Apostol, <i>Arithmetical properties of generalized Ramanujan sums</i>	281
David Lee Armacost and William Louis Armacost, <i>On p-thetic groups</i>	295
Janet E. Mills, <i>Regular semigroups which are extensions of groups</i>	303
Gregory Frank Bachelis, <i>Homomorphisms of Banach algebras with minimal ideals</i>	307
John Allen Beachy, <i>A generalization of injectivity</i>	313
David Geoffrey Cantor, <i>On arithmetic properties of the Taylor series of rational functions. II</i>	329
Václav Chvátal and Frank Harary, <i>Generalized Ramsey theory for graphs. III. Small off-diagonal numbers</i>	335
Frank Rimi DeMeyer, <i>Irreducible characters and solvability of finite groups</i>	347
Robert P. Dickinson, <i>On right zero unions of commutative semigroups</i>	355
John Dustin Donald, <i>Non-openness and non-equidimensionality in algebraic quotients</i>	365
John D. Donaldson and Qazi Ibadur Rahman, <i>Inequalities for polynomials with a prescribed zero</i>	375
Robert E. Hall, <i>The translational hull of an N-semigroup</i>	379
John P. Holmes, <i>Differentiable power-associative groupoids</i>	391
Steven Kenyon Ingram, <i>Continuous dependence on parameters and boundary data for nonlinear two-point boundary value problems</i>	395
Robert Clarke James, <i>Super-reflexive spaces with bases</i>	409
Gary Douglas Jones, <i>The embedding of homeomorphisms of the plane in continuous flows</i>	421
Mary Joel Jordan, <i>Period H-semigroups and t-semisimple periodic H-semigroups</i>	437
Ronald Allen Knight, <i>Dynamical systems of characteristic 0</i>	447
Kwangil Koh, <i>On a representation of a strongly harmonic ring by sheaves</i>	459
Hui-Hsiung Kuo, <i>Stochastic integrals in abstract Wiener space</i>	469
Thomas Graham McLaughlin, <i>Supersimple sets and the problem of extending a retracing function</i>	485
William Nathan, <i>Open mappings on 2-manifolds</i>	495
M. J. O'Malley, <i>Isomorphic power series rings</i>	503
Sean B. O'Reilly, <i>Completely adequate neighborhood systems and metrization</i>	513
Qazi Ibadur Rahman, <i>On the zeros of a polynomial and its derivative</i>	525
Russell Daniel Rupp, Jr., <i>The Weierstrass excess function</i>	529
Hugo Teufel, <i>A note on second order differential inequalities and functional differential equations</i>	537
M. J. Wicks, <i>A general solution of binary homogeneous equations over free groups</i>	543