

Pacific Journal of Mathematics

**ON THE ABSOLUTE MATRIX SUMMABILITY OF A FOURIER
SERIES**

ARIBINDI SATYANARAYAN RAO

ON THE ABSOLUTE MATRIX SUMMABILITY OF A FOURIER SERIES

ARIBINDI SATYANARAYAN RAO

In this paper, the author gives sufficient conditions for a Fourier series at an arbitrary but fixed point to be absolutely matrix summable.

1. Introduction. Let $\sum_{n=0}^{\infty} u_n$ be an infinite series with partial sums s_n , and let $A = (a_{nk})$ be a triangular infinite matrix of real numbers (see Hardy [2]). The series $\sum u_n$ is said to be absolutely summable A , or summable $|A|$, if

$$\sum_{n=1}^{\infty} |\tau_n - \tau_{n-1}| < \infty ,$$

where

$$\tau_n = \sum_{k=0}^n a_{nk} s_k .$$

Let $f(t)$ be a Lebesgue-integrable function of period 2π , with Fourier series

$$(1.1) \quad \frac{1}{2} a_0 + \sum_{n=1}^{\infty} (a_n \cos nt + b_n \sin nt) \equiv \sum_{n=0}^{\infty} A_n(t) .$$

With a fixed point x , we set

$$(1.2) \quad \phi(t) = \phi_x(t) = \frac{1}{2}[f(x+t) + f(x-t)] ,$$

$$(1.3) \quad \Phi(t) = \int_0^t |\phi(u)| du .$$

We establish the following theorem for the absolute matrix summability of the Fourier series (1.1) of $f(t)$ at $t = x$.

THEOREM. *Let $A = (a_{nk})$ be a triangular infinite matrix of real numbers such that $\Delta a_{nk} = a_{nk} - a_{n,k+1}$ is monotonic with respect to $n \geq k$ for each fixed $k \geq 0$.*

Let $\alpha(t)$ be a positive function such that $t^r/\alpha(t)$, for some r with $0 < r < 1$, is nondecreasing for $t \geq t_0$. Suppose that

$$(1.4) \quad \sum_{n=1}^{\infty} \frac{n |a_{nn}|}{\alpha(n)} < \infty ,$$

$$(1.5) \quad |\Delta a_{m,0}| + \sum_{n=1}^{m-1} \frac{n|\Delta a_{mn}|}{\alpha(n)} = O(1) \quad \text{as } m \rightarrow \infty .$$

Further, let

$$(1.6) \quad \Phi(t) = O\left[\frac{t}{\alpha(1/t)}\right] \quad \text{as } t \rightarrow 0+.$$

If all of the above conditions hold, then the Fourier series (1.1) of $f(t)$ at $t = x$ is summable $|A|$.

We shall require the following lemmas.

LEMMA 1. If $\alpha(t)$ is defined as in the theorem, then

$$(2.1) \quad \int_{t_0}^t \frac{du}{\alpha(u)} = O\left[\frac{t}{\alpha(t)}\right] \quad \text{for all } t \geq t_0 .$$

Proof.

$$\begin{aligned} \int_{t_0}^t \frac{du}{\alpha(u)} &= \int_{t_0}^t \frac{u^r}{\alpha(u)} \cdot \frac{du}{u^r} \\ &\leqq \frac{t^r}{\alpha(t)} \int_{t_0}^t \frac{du}{u^r} \leqq \frac{t^r}{\alpha(t)} \cdot \frac{t^{-r+1}}{1-r} = O\left[\frac{t}{\alpha(t)}\right]. \end{aligned}$$

LEMMA 2. If $A = (a_{nk})$ is defined as in the theorem and if

$$(2.2) \quad \sum_{n=0}^{\infty} |t_n| \cdot |a_{nn}| < \infty ,$$

$$(2.3) \quad \sum_{n=0}^{m-1} |t_n| \cdot |\Delta a_{mn}| = O(1) \quad \text{as } m \rightarrow \infty ,$$

where

$$t_n = \sum_{k=0}^n s_k ,$$

then $\sum u_n$ is summable $|A|$.

Proof. By Abel's transformation,

$$\begin{aligned} \tau_n - \tau_{n-1} &= \sum_{k=0}^n (a_{nk} - a_{n-1,k})s_k \\ &= \sum_{k=0}^{n-1} (\Delta a_{nk} - \Delta a_{n-1,k})t_k + a_{nn}t_n . \end{aligned}$$

Now

$$\begin{aligned} &\sum_{n=1}^m \sum_{k=0}^{n-1} |\Delta a_{nk} - \Delta a_{n-1,k}| \cdot |t_k| \\ &= \sum_{k=0}^{m-1} |t_k| \cdot \left(\sum_{n=k+1}^m |\Delta a_{nk} - \Delta a_{n-1,k}| \right) = \sum_{k=0}^{m-1} |t_k| \cdot |\Delta a_{mk} - a_{kk}| . \end{aligned}$$

Thus,

$$\sum_{n=1}^m |\tau_n - \tau_{n-1}| \leq \sum_{n=0}^{m-1} |t_n| \cdot |\Delta a_{mn}| + 2 \sum_{n=0}^m |t_n| \cdot |a_{nn}| = O(1)$$

as $m \rightarrow \infty$, by (2.2) and (2.3).

This completes the proof of the lemma.

3. Proof of the Theorem. We write

$$s_n(x) = \sum_0^n A_k(x), \quad t_n(x) = \sum_0^n s_k(x).$$

By (1.6), there exists $\delta (0 < \delta < 1)$ such that

$$(3.1) \quad \Phi(t) \leq K \frac{t}{\alpha(1/t)} \quad \text{for } 0 < t \leq \delta,$$

where K is a positive constant (not necessarily the same at each occurrence). Now, for $n > \delta^{-1}$,

$$(3.2) \quad \begin{aligned} \pi t_n(x) &= \int_0^\pi \phi(t) \left[\frac{\sin(n+1)(t/2)}{\sin(t/2)} \right]^2 dt \\ &= \int_0^{\delta^{-1}} + \int_{\delta^{-1}}^\delta + \int_\delta^\pi = I_1 + I_2 + I_3, \text{ say.} \end{aligned}$$

We observe that

$$(3.3) \quad \left[\frac{\sin(n+1) \cdot (t/2)}{\sin(t/2)} \right]^2 = \begin{cases} O(n^2) & \text{for } \sin t/2 \neq 0 \text{ and } n \geq 1, \\ O(1/t^2) & \text{for } 0 < t \leq \pi. \end{cases}$$

So, by (3.1),

$$(3.4) \quad |I_1| \leq K n^2 \int_0^{\delta^{-1}} |\phi(t)| dt \leq K \frac{n}{\alpha(n)}.$$

Further, assuming $t^r/\alpha(t)$ nondecreasing for $t \geq \delta^{-1}$,

$$(3.5) \quad \begin{aligned} |I_2| &\leq K \int_{\delta^{-1}}^\delta \frac{|\phi(t)|}{t^2} dt \\ &= K \left\{ \left[\frac{\Phi(t)}{t^2} \right]_{\delta^{-1}}^\delta + 2 \int_{\delta^{-1}}^\delta \frac{\Phi(t)}{t^3} dt \right\} \\ &\leq K \left[\frac{\Phi(\delta)}{\delta^2} + \int_{\delta^{-1}}^\delta \frac{dt}{t^2 \alpha(1/t)} \right] \\ &= K \left[\frac{\Phi(\delta)}{\delta^2} + \int_{\delta^{-1}}^n \frac{du}{\alpha(u)} \right] \\ &\leq K \frac{n}{\alpha(n)} \quad \text{as } n \rightarrow \infty, \text{ by (2.1).} \end{aligned}$$

Obviously,

$$(3.6) \quad I_3 = O(1) .$$

From (3.2), (3.4)–(3.6), it follows that

$$(3.7) \quad t_n(x) = O\left[\frac{n}{\alpha(n)}\right] \quad \text{as } n \rightarrow \infty .$$

Hence

$$(3.8) \quad \sum_n^{\infty} |t_k(x)| \cdot |a_{kk}| = O\left[\sum_n^{\infty} \frac{k}{\alpha(k)} |a_{kk}|\right] = o(1) \\ \text{as } n \rightarrow \infty , \text{ by (1.4).}$$

Moreover,

$$(3.9) \quad \sum_0^{m-1} |t_n(x)| \cdot |\Delta a_{mn}| = |t_0(x)| \cdot |\Delta a_{m0}| + O\left[\sum_1^{m-1} \frac{n}{\alpha(n)} \cdot |\Delta a_{mn}|\right] \\ = O(1) \quad \text{as } m \rightarrow \infty , \text{ by (1.5).}$$

Now the theorem follows from Lemma 2.

4. NOTE. Let $A = (a_{nk})$ be a triangular infinite matrix of real numbers such that $a_{nn} \geq 0$ for all $n \geq 0$ and Δa_{nk} is nondecreasing with respect to $n \geq k$ for each fixed $k \geq 0$. Let $\alpha(t)$ be defined as in the theorem, and let

$$(4.1) \quad \Delta a_{m0} + \sum_{n=1}^m \frac{n(\Delta a_{mn})}{\alpha(n)} = O(1) \quad \text{as } m \rightarrow \infty .$$

Then, if the condition (1.6) holds, the Fourier series (1.1) of $f(t)$ at $t = x$ is summable $|A|$.

Proof. Let

$$\tau_n(x) = \sum_{k=0}^n a_{nk} s_k(x) .$$

Then

$$(4.2) \quad \begin{aligned} & \sum_{n=1}^m |\tau_n(x) - \tau_{n-1}(x)| \\ & \leq \sum_{n=1}^m \sum_{k=0}^n |\Delta a_{nk} - \Delta a_{n-1,k}| \cdot |t_k(x)| \\ & = \sum_{k=1}^m |t_k(x)| \left(\sum_{n=k}^m |\Delta a_{nk} - \Delta a_{n-1,k}| \right) + |t_0(x)| \sum_{n=1}^m |\Delta a_{n0} - \Delta a_{n-1,0}| \\ & = \sum_{k=1}^m |t_k(x)| (\Delta a_{mk}) + |t_0(x)| (\Delta a_{m0} - a_{00}) \end{aligned}$$

$$\leq |t_0(x)|(\Delta a_{m,0}) + O\left[\sum_{k=1}^m \frac{k}{\alpha(k)} (\Delta a_{mk})\right], \quad \text{by (3.7)}$$

$$= 0(1) \quad \text{as } m \rightarrow \infty, \text{ by (4.1).}$$

So the required result follows.

I thank Professors A. Meir and A. Sharma for providing me financial support from their N.R.C. grants during the preparation of this paper.

REFERENCES

1. S. N. Bhatt, *An aspect of local property of $|N, p_n|$ summability of a Fourier series*, Indian J. Math. **5** (1963), 87-91.
2. G. H. Hardy, *Divergent Series*, (Oxford, 1949).
3. L. McFadden, *Absolute Nörlund summability*, Duke Math. J., **9** (1942), 168-207.

Received May 7, 1971.

UNIVERSITY OF ALBERTA
EDMONTON, CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 42, No. 1

January, 1972

| | |
|--|-----|
| Tage Bai Andersen, <i>On Banach space valued extensions from split faces</i> | 1 |
| David Marion Arnold, <i>A duality for quotient divisible abelian groups of finite rank</i> | 11 |
| Donald Pollard Ballou, <i>Shock sets for first order nonlinear hyperbolic equations</i> | 17 |
| Leon Brown and Lowell J. Hansen, <i>On the range sets of H^p functions</i> | 27 |
| Alexander Munro Davie and Arne Stray, <i>Interpolation sets for analytic functions</i> | 33 |
| M. G. Deshpande, <i>Structure of right subdirectly irreducible rings. II</i> | 39 |
| Barry J. Gardner, <i>Some closure properties for torsion classes of abelian groups</i> | 45 |
| Paul Daniel Hill, <i>Primary groups whose subgroups of smaller cardinality are direct sums of cyclic groups</i> | 63 |
| Richard Allan Holzsager, <i>When certain natural maps are equivalences</i> | 69 |
| Donald William Kahn, <i>A note on H-equivalences</i> | 77 |
| Joong Ho Kim, <i>R-automorphisms of $R[t][[X]]$</i> | 81 |
| Shin'ichi Kinoshita, <i>On elementary ideals of polyhedra in the 3-sphere</i> | 89 |
| Andrew T. Kitchen, <i>Watts cohomology and separability</i> | 99 |
| Vadim Komkov, <i>A technique for the detection of oscillation of second order ordinary differential equations</i> | 105 |
| Charles Philip Lanski and Susan Montgomery, <i>Lie structure of prime rings of characteristic 2</i> | 117 |
| Andrew Lenard, <i>Some remarks on large Toeplitz determinants</i> | 137 |
| Kathleen B. Levitz, <i>A characterization of general Z.P.I.-rings. II</i> | 147 |
| Donald A. Lutz, <i>On the reduction of rank of linear differential systems</i> | 153 |
| David G. Mead, <i>Determinantal ideals, identities, and the Wronskian</i> | 165 |
| Arunava Mukherjea, <i>A remark on Tonelli's theorem on integration in product spaces</i> | 177 |
| Hyo Chul Myung, <i>A generalization of the prime radical in nonassociative rings</i> | 187 |
| John Piepenbrink, <i>Rellich densities and an application to unconditionally nonoscillatory elliptic equations</i> | 195 |
| Michael J. Powers, <i>Lefschetz fixed point theorems for a new class of multi-valued maps</i> | 211 |
| Aribindi Satyanarayan Rao, <i>On the absolute matrix summability of a Fourier series</i> | 221 |
| T. S. Ravisankar, <i>On Malcev algebras</i> | 227 |
| William Henry Ruckle, <i>Topologies on sequences spaces</i> | 235 |
| Robert C. Shock, <i>Polynomial rings over finite dimensional rings</i> | 251 |
| Richard Tangeman, <i>Strong heredity in radical classes</i> | 259 |
| B. R. Wenner, <i>Finite-dimensional properties of infinite-dimensional spaces</i> | 267 |