

# Pacific Journal of Mathematics

## **ON THE ABSOLUTE MATRIX SUMMABILITY OF FOURIER SERIES**

B. KUTTNER AND B. N. SAHNEY

# ON THE ABSOLUTE MATRIX SUMMABILITY OF FOURIER SERIES

B. KUTTNER AND B. N. SAHNEY

**The paper investigates sufficient conditions under which a summability method of a certain general type absolutely sums the Fourier series of any function of bounded variation. The main theorem includes a recent theorem of M. Izumi and S. Izumi, who considered the problem for the special case of Nörlund summability.**

The summability methods considered are those given by a series-to-series transformation  $A = (\alpha_{n,k})$ . That is to say, given any series

$$(1) \quad \sum_{k=0}^{\infty} \alpha_k,$$

we describe (1) as summable  $A$  to  $s$  if

$$b_n = \sum_{k=0}^{\infty} \alpha_{n,k} \alpha_k$$

is defined for all  $n$ , and if

$$(2) \quad \sum_{n=0}^{\infty} b_n$$

converges to  $s$ . We describe (1) as absolutely summable  $|A|$  if (2) converges absolutely. Under certain quite weak restrictions on  $A$ , necessary and sufficient conditions under which the Fourier series of any function of bounded variation should be absolutely summable  $|A|$  have been given by Tripathy [10, Lemma 2]; his result will be stated later as Lemma 1. But the conditions obtained by Tripathy are of such a nature that it is not usually easy in any given example to determine whether they are satisfied or not. The object of the present paper is to obtain sufficient conditions which, while less general, are simpler than those of Tripathy. However, it does not seem possible to obtain reasonably general sufficient condition in any very simple form.

We will be concerned with the case in which  $A$  is absolutely conservative, that is to say, it is such that, whenever (1) converges absolutely, so does (2). It is known [4, 6] that in order that this should hold it is necessary and sufficient that, for  $k \geq 0$ ,

$$(3) \quad \sum_{n=0}^{\infty} |\alpha_{n,k}| = O(1).$$

We remark that, in order that  $A$  should be absolutely regular, that is to say, that in order that, whenever (1) converges absolutely then (2) converges absolutely to the same sum, it is necessary and sufficient that (3) should hold and that, further, for all  $k \geq 0$ ,

$$(4) \quad \sum_{n=0}^{\infty} \alpha_{n,k} = 1.$$

2. We now state our main result.

**THEOREM.** *Let  $A = (\alpha_{n,k})$  be an absolutely conservative series-to-series transformation, with  $\alpha_{n,k} \geq 0$  for all  $n, k$ . Suppose that either*

(a) *For each fixed  $n$ , there is a positive integer  $r(n)$  such that  $\alpha_{n,k}$  is nondecreasing for  $1 \leq k \leq r(n)$ , and nonincreasing for  $k \geq r(n)$ , or*

(b) *For each fixed  $n$ , there is a positive integer  $s(n)$ , such that  $\alpha_{n,k}/k$  is nondecreasing for  $1 \leq k \leq s(n)$ , and nonincreasing for  $k \geq s(n)$ . Suppose also in case (a) that, for  $K \geq 1$ ,*

$$(5) \quad \sum_{r(n) \geq 2K} \frac{1}{r(n)} \sum_{k=r(n)-K}^{r(n)+K} \alpha_{n,k} = O(1),$$

and in case (b) that, for  $K \geq 1$ ,

$$(6) \quad \sum_{s(n) \geq 2K} \frac{1}{s(n)} \sum_{k=s(n)-K}^{s(n)+K} \alpha_{n,k} = O(1).$$

*Then the Fourier series of any function of bounded variation is absolutely summable  $|A|$ .*

**REMARK.** It is clear that (5) is equivalent to

$$(5') \quad \sum_{r(n) \geq 2K} \sum_{k=r(n)-K}^{r(n)+K} \frac{\alpha_{n,k}}{k} = O(1)$$

and it is sometimes more convenient to express (5) in this form. Since there are  $2K + 1$  terms in the inner sum in (5), and since the middle term is the greatest, a sufficient condition for (5) is that

$$(7) \quad \sum_{r(n) \geq 2K} \frac{\alpha_{n,r(n)}}{r(n)} = O\left(\frac{1}{K}\right).$$

However (7), while much simpler than (5) is less general, and, as will be shown later, fails to be satisfied in some important cases. In a similar way, (6) is equivalent to

$$(6') \quad \sum_{s(n) \geq 2K} \sum_{k=s(n)-K}^{s(n)+K} \frac{\alpha_{n,k}}{k} = O(1);$$

also, a sufficient condition for (6) is that

$$(8) \quad \sum_{s(n) \geq 2K} \frac{\alpha_{n,s(n)}}{s(n)} = O\left(\frac{1}{K}\right).$$

It is clear that either one of (a), (b) could be satisfied without the other holding. If, however, they both hold, then (5) is a weaker assumption than (6). Thus, in this case, the first form of the theorem is preferable. To prove this assertion, we write  $\theta_n, \phi_n$  for the inner sums in (5'), (6') respectively, and shall first show that

$$(9) \quad \theta_n \leq 2\phi_n.$$

To this end, we first note that  $s(n) \leq r(n)$ . Consider first the case in which  $r(n) - s(n) \geq K$ . Since  $\alpha_{n,k}/k$  is nonincreasing for  $k \geq s(n)$ , we have<sup>1</sup>, for  $\mu = 0, 1, \dots, K-1$ ,

$$(10) \quad \begin{aligned} & \frac{\alpha(n, r(n) + K - 2\mu)}{r(n) + K - 2\mu} + \frac{\alpha(n, r(n) + K - 2\mu - 1)}{r(n) + K - 2\mu - 1} \\ & \leq \frac{2\alpha(n, s(n) + K - \mu)}{s(n) + K - \mu}. \end{aligned}$$

Also,

$$\frac{\alpha(n, r(n) - K)}{r(n) - K} \leq \frac{\alpha(n, s(n))}{s(n)},$$

whence

$$\theta_n \leq 2 \sum_{k=s(n)}^{s(n)+K} \frac{\alpha_{n,k}}{k} \leq 2\phi_n,$$

where the dash indicates the term  $k = s(n)$ , is multiplied by  $1/2$ . If  $r(n) - s(n) = t(n) < K$ , then (10) still holds for  $\mu \leq t(n) - 1$ . Hence

$$(11) \quad \theta_n \leq 2 \sum_{\mu=0}^{t(n)-1} \frac{\alpha(n, s(n) + K - \mu)}{s(n) + K - \mu} + \sum_{\nu=2t(n)}^{2K} \frac{\alpha(n, r(n) + K - \nu)}{r(n) + K - \nu}$$

where the first sum on the right is taken as 0 if  $t(n) = 0$ . Since the second sum on the right of (11) can be written

$$\sum_{\mu=t(n)}^{2K-t(n)} \frac{\alpha(n, s(n) + K - \mu)}{s(n) + K - \mu},$$

we again deduce (9).

It now follows from (9) that

$$\sum_{s(n) \geq 2K} \theta_n \leq 2 \sum_{s(n) \geq 2K} \phi_n.$$

However, since  $s(n) \leq r(n)$ , there may be values of  $n$  for which  $r(n) \geq 2K$ , but  $s(n) < 2K$ ; these values will occur in the sum (5'), but not

<sup>1</sup> To avoid complicated suffixes, we write  $\alpha(n, k)$  for  $\alpha_{n,k}$  whenever  $n, k$  are replaced by more complicated expressions

in (6'). If we show that, in any case, the contribution of these terms to the sum (5') is bounded, the conclusion will now follow. If  $r(n) \geq 2K$  but  $s(n) \leq K$ , then, since  $\alpha_{n,k}/k$  is nonincreasing for  $k \geq K$  we deduce that

$$\theta_n \leq \frac{(2K+1)\alpha_{n,K}}{K}.$$

If  $r(n) \geq 2K$  and  $K < s(n) < 2K$ , then  $\alpha_{n,k}$  is nondecreasing for  $k \leq 2K$ . Hence, for all  $k \geq 1$

$$\frac{\alpha_{n,k}}{k} \leq \frac{\alpha_{n,s(n)}}{s(n)} \leq \frac{\alpha_{n,2K}}{K}$$

so that

$$\theta_n \leq \frac{(2K+1)\alpha_{n,2K}}{K}.$$

Thus the sum of the terms in question does not exceed

$$\frac{(2K+1)}{K} \sum_{n=0}^{\infty} (\alpha_{n,K} + \alpha_{n,2K}) = O(1),$$

by (3).

3. We now state the lemma of Tripathy already mentioned.

LEMMA 1. *Let  $A = (\alpha_{n,k})$  be a series-to-series transformation such that*

$$(12) \quad \sum_{n=0}^{\infty} |\alpha_{n,0}| < \infty$$

*and such that, for every fixed  $n$ ,*

$$(13) \quad L_n(t) = \sum_{k=1}^{\infty} \alpha_{n,k} \frac{\sin kt}{k}$$

*converges boundedly in  $t$ . Then in order that the Fourier series of any function of bounded variation should be absolutely summable  $|A|$ , it is sufficient that*

$$(14) \quad \sum_{n=0}^{\infty} |L_n(t)| = O(1),$$

*and necessary that the sum (14) should be essentially bounded.*

It may be remarked that the result is not quite correctly stated in [10], where it is asserted that the essential boundedness of (14) is necessary and sufficient. But on examining the proof of sufficiency

in [10], we find that it requires the boundedness, and not just the essential boundedness, of (14). The point is not of great importance, since if we assume that, for every fixed  $n$ ,

$$\sum_{k=1}^{\infty} |\alpha_{n,k} - \alpha_{n,k+1}| < \infty ;$$

in other words, that  $b_n$  is defined whenever (1) converges, it is easy to prove that the essential boundedness of (14) is equivalent to its boundedness. This result is not, however, required for our present purposes.

In what follows, we will suppose throughout that  $0 < t \leq \pi$ . We will apply the hypothesis (5) or (6) with  $K = [\pi/t]$ ; thus, in any equations involving both  $K$  and  $t$ , it will be assumed that this relation holds.

We require two further lemmas.

**LEMMA 2.** *Let  $A = (\alpha_{n,k})$  be an absolutely conservative series-to-series transformation. If, for every fixed  $n$ ,  $\alpha_{n,k}/k$  is ultimately nonnegative nonincreasing (and thus, in particular, if the hypotheses of the theorem are satisfied) then the hypotheses of Lemma 1 are satisfied.*

Equation (12) follows at once as a special case of (3). Thus, taking  $n$  as fixed, we have only to verify that (13) converges boundedly. Suppose that  $\alpha_{n,k}/k$  is nonnegative nonincreasing for  $k \geq M$ . Then we have, uniformly in  $k_1, k_2$  for  $K, M \leq k_1 \leq k_2$ ,

$$(15) \quad \left| \sum_{k=k_1}^{k_2} \alpha_{n,k} \frac{\sin kt}{k} \right| \leq \frac{\alpha(n, k_1)}{\left(2 \sin \frac{1}{2}t\right)k_1}.$$

But (3) implies that  $\alpha(n, k)$  is bounded; hence the expression on the right of (15) is  $O(1)$  uniformly in the range considered, and, for fixed  $t$ , tends to 0 (uniformly in  $k_2$ ) as  $k_1 \rightarrow \infty$ . Since  $M$  is a constant,

$$\sum_{k=1}^{M-1} \left| \alpha_{n,k} \frac{\sin kt}{k} \right|$$

is bounded; also, if  $K \geq M$ ,

$$\sum_{k=M}^K \left| \alpha_{n,k} \frac{\sin kt}{k} \right| \leq t \sum_{k=M}^K |\alpha_{n,k}| = O(1)$$

(by the boundedness of  $\alpha_{n,k}$  and the definition of  $K$ ). Hence the result.

**LEMMA 3.** *Suppose that  $\theta_k \geq 0$ . Suppose that  $\theta_k$  is nondecreasing*

for  $1 \leq k \leq s$ , and nonincreasing for  $k \geq s$ . Then, for any positive integers  $a, b$ , and any  $t$  with  $0 < t \leq \pi$ ,

$$(16) \quad \left| \sum_{k=a}^b \theta_k e^{ikt} \right| \leq A \sum_{k=\max(1, s-K)}^{s+K} \theta_k,$$

where  $A$  is an absolute constant.

That portion (if any) of the sum on the left for which  $s - K \leq k \leq s + K$  clearly satisfies the required inequality. Also, by partial summation, that portion (if any) for which  $k > s + K$  does not, in modulus, exceed

$$\begin{aligned} \frac{2}{|1 - e^{it}|} \theta_{s+K} &\leq \frac{2}{|1 - e^{it}|(K+1)} \sum_{k=s}^{s+K} \theta_k \\ &\leq \frac{2t}{\pi |1 - e^{it}|} \sum_{k=s}^{s+K} \theta_k. \end{aligned}$$

That portion of the sum (if any) for which  $k < s - K$  may be dealt with in a similar way, and the conclusion follows.

This lemma is a slight generalisation of a lemma due to McFadden [5] which has been much used in investigations on the Nörlund summability of Fourier series.

4. We now come to the proof of the theorem. It follows from Lemmas 1 and 2 that it is enough to show that the hypotheses of the theorem imply (14). Consider first those values of  $n$  (if any) for which  $r(n) < 2K$  in case (a), and for which  $s(n) < 2K$  in case (b). In case (b), we are given that  $\alpha_{n,k}/k$  is nonincreasing for  $k \geq 2K$ ; in case (a), we are given that  $\alpha_{n,k}$  is nonincreasing for  $k \geq 2K$ ; hence, a fortiori, so is  $\alpha_{n,k}/k$ . Thus, in either case, since the partial sums of  $\sum \sin kt$  are  $O(1/t)$ , we have

$$\sum_{k=2K}^{\infty} \alpha_{n,k} \frac{\sin kt}{k} = O\left(\frac{\alpha_{n,2K}}{2Kt}\right) = O(\alpha_{n,2K})$$

by definition of  $K$ . For those terms in the sum (13) for which  $k \leq 2K$ , we use  $|\sin kt| \leq kt$ ; and it follows that

$$|L_n(t)| = O\left\{t \sum_{k=1}^{2K-1} \alpha_{n,k}\right\} + O(\alpha_{n,2K}).$$

Hence the contribution to the sum (14) of those values of  $n$  now under consideration is

$$(17) \quad O\left\{t \sum_{k=1}^{2K-1} \sum_{n=0}^{\infty} \alpha_{n,k}\right\} + O\left\{\sum_{n=0}^{\infty} \alpha_{n,2K}\right\} = O(1)$$

by (3) and the definition of  $K$ .

We now investigate the remaining values of  $n$ . Consider first

case (b). For any fixed  $n$ , we apply Lemma 3 with  $\theta_k = \alpha_{n,k}/k$ , and take the imaginary part of (16). It follows at once that

$$L_n(t) = O\left\{\sum_{k=s(n)-K}^{s(n)+K} \frac{\alpha_{n,k}}{k}\right\};$$

and (14) therefore follows from (6') and (17).

Now consider case (a). Since  $\alpha_{n,k}$  is nonincreasing for  $k \geq r(n)$  so is  $\alpha_{n,k}/k$ ; thus the part of the sum (13) for which  $k > r(n) - K$  may be dealt with as in case (b). The part for which  $k < K$  may be dealt with by using  $|\sin kt| \leq kt$ , as in the proof of (17). Thus, writing

$$R_n(t) = \sum_{k=K}^{r(n)-K} \alpha_{n,k} \frac{\sin kt}{k},$$

it remains only to show that

$$(18) \quad \sum_{r(n) \geq 2K} |R_n(t)| = O(1).$$

Now,

$$\begin{aligned} R_n(t) &= \frac{1}{2 \sin \frac{1}{2}t} \sum_{k=K}^{r(n)-K} \frac{\alpha_{n,k}}{k} \left[ \cos\left(k - \frac{1}{2}\right)t - \cos\left(k + \frac{1}{2}\right)t \right] \\ &= \frac{1}{2 \sin \frac{1}{2}t} \left\{ - \sum_{k=K}^{r(n)-K} \cos\left(k + \frac{1}{2}\right)t \Delta_k\left(\frac{\alpha_{n,k}}{k}\right) \right. \\ &\quad + \frac{\alpha_{n,K}}{K} \cos\left(K - \frac{1}{2}\right)t \\ &\quad \left. - \frac{\alpha(n, r(n) - K + 1)}{r(n) - K + 1} \cos\left(r(n) - K + \frac{1}{2}\right)t \right\}. \end{aligned}$$

Since

$$\Delta_k\left(\frac{\alpha_{n,k}}{k}\right) = \frac{\alpha_{n,k}}{k(k+1)} + \frac{\Delta_k(\alpha_{n,k})}{k+1},$$

it follows that

$$\begin{aligned} R_n(t) &= O\left\{\frac{1}{t} \left[ \sum_{k=K}^{r(n)-K} \frac{\alpha_{n,k}}{k(k+1)} + \sum_{k=K}^{r(n)-K} \frac{|\Delta_k(\alpha_{n,k})|}{k+1} \right] \right. \\ &\quad \left. + \frac{\alpha_{n,K}}{K} + \frac{\alpha(n, r(n) - K + 1)}{r(n) - K + 1} \right\} \\ &= O\{R_n^1(t) + R_n^2(t) + R_n^3(t) + R_n^4(t)\}, \end{aligned}$$

say. Now, since  $\alpha_{n,k}$  is nondecreasing in the relevant range



$$\begin{aligned}
 R_n^{n_2}(t) &= -\frac{1}{t} \sum_{k=K}^{r(n)-K} \frac{\Delta_k(\alpha_{n,k})}{k+1} \\
 &= \frac{1}{t} \sum_{k=K}^{r(n)-K} \frac{\alpha_{n,k}}{k(k+1)} - \frac{1}{t} \frac{\alpha_{n,K}}{K} + \frac{1}{t} \frac{\alpha(n, r(n) - K + 1)}{r(n) - K + 1} \\
 &= R_n^1(t) - R_n^3(t) + R_n^4(t),
 \end{aligned}$$

so that

$$R_n(t) = O\{R_n^1(t) + R_n^4(t)\}.$$

Next,

$$\sum_{r(n) \geq 2K} R_n^1(t) \leq \frac{1}{t} \sum_{k=K}^{\infty} \frac{1}{k(k+1)} \sum_{n=0}^{\infty} \alpha_{n,k} = O(1)$$

by (3) and the definition of  $K$ . Finally, if  $r(n) \geq 2K$ ,

$$\begin{aligned}
 R_n^4(t) &= O\left\{\frac{\alpha(n, r(n) - K + 1)}{tr(n)}\right\} \\
 &= O\left\{\frac{1}{tKr(n)} \sum_{k=r(n)-K+1}^{r(n)} \alpha_{n,k}\right\} \\
 &= O\left\{\frac{1}{r(n)} \sum_{k=r(n)-K+1}^{r(n)} \alpha_{n,k}\right\},
 \end{aligned}$$

so that

$$\sum_{r(n) \geq 2K} R_n^4(t) = O(1),$$

by (5). The proof of the theorem is thus completed.

5. We now consider an application of our general theorem to the special case of Nörlund summability. We recall that, given a sequence  $p = \{p_n\}$ , Nörlund summability  $(N, p)$  is defined as given by the sequence-to-sequence transformation

$$(19) \quad t_n = \frac{1}{P_n} \sum_{k=0}^n p_{n-k} s_k,$$

where we write

$$P_n = p_0 + p_1 + \cdots + p_n;$$

it is assumed that  $p$  is such that, for all  $n$ ,  $P_n \neq 0$ . If we write

$$t_n = b_0 + b_1 + \cdots + b_n; s_k = a_0 + a_1 + \cdots + a_k$$

we see that (19) can be expressed as the series-to-series transformation

$$b_0 = a_0;$$

$$b_n = \sum_{k=1}^n \left( \frac{P_{n-k}}{P_n} - \frac{P_{n-k-1}}{P_{n-1}} \right) a_n \quad (n \geq 1),$$

where we adopt the convention that  $P_{-1} = 0$ . Thus we have, with the notation of our main theorem,  $\alpha_{n,k} = 0$  for  $k > n$ , while, for  $1 \leq k \leq n$

$$\begin{aligned} \alpha_{n,k} &= \frac{P_{n-k}}{P_n} - \frac{P_{n-k-1}}{P_{n-1}} \\ (20) \quad &= \frac{P_n p_{n-k} - P_{n-k} p_n}{P_n P_{n-1}}. \end{aligned}$$

Now consider the case in which  $\{p_n\}$  is nonnegative nonincreasing. We remark that, since  $P_0 \neq 0$ , we then have  $p_0 > 0$ . Further (since  $p_n \geq 0$ )  $\{P_n\}$  is nondecreasing; thus it follows from (20) that  $\alpha_{n,k} \geq 0$ . Thus we may omit the modulus signs in (3); and it is now easy to see that (4), and hence (3), holds. Thus, in the case now considered,  $(N, p)$  is absolutely regular. Further, for fixed  $n$ ,  $p_{n-k}$  is nondecreasing and  $P_{n-k}$  nonincreasing as  $k$  increases from 1 to  $n$ . Since  $\alpha_{n,k} = 0$  for  $k > n$ , it follows that condition (a) is satisfied, with  $r(n) = n$ . Also equation (5) becomes

$$(21) \quad \sum_{n=2K}^{\infty} \frac{1}{nP_n P_{n-1}} \sum_{k=n-K}^n (P_n p_{n-k} - P_{n-k} p_n) = O(1).$$

The inner sum in (21) does not exceed

$$\sum_{k=n-K}^n P_n p_{n-K} = P_n P_K,$$

and thus a sufficient condition for (21) to hold is that

$$(22) \quad \sum_{n=2K}^{\infty} \frac{1}{nP_{n-1}} = O\left(\frac{1}{P_K}\right).$$

However, since the hypotheses on  $p$  imply that  $P_{n-1} \sim P_n$ , and that  ${}_K P \leq P_{2K} \leq 2P_K$ , it is easily seen that (22) is equivalent to the slightly simpler condition

$$(23) \quad \sum_{n=K}^{\infty} \frac{1}{nP_n} = O\left(\frac{1}{P_K}\right).$$

Thus our theorem includes the following result;

**THEOREM A.** *Suppose that  $\{p_n\}$  is nonnegative nonincreasing, and that (23) holds. Then the Fourier series of any function of bounded variation is absolutely summable  $|N, p|$ .*

The assumption that  $\{p_n\}$  is nonnegative nonincreasing is not, without some further condition, sufficient for the conclusion, for it

has been shown by Pati [8] that, when  $p_n = 1/(n+1)$ , it is not true that the Fourier series of any function of bounded variation is absolutely summable  $|N, p|$ . This example also shows that, in our main theorem, the assumptions that  $A$  is absolutely conservative and that (a) holds would not alone suffice for the conclusion.

Theorem A is included in a recent, slightly more general, theorem of M. Izumi and S. Izumi [3]. It includes earlier theorems of H. P. Dikshit [2, Theorem 2] and T. Singh [9]; the result of Singh itself generalises a theorem of Pati [7]. The theorems of Dikshit and of Singh are respectively as follows.

**THEOREM B.** *Suppose that  $p_n > 0$ , and that  $p_{n+1}/p_n$  is non decreasing, and less than or equal to 1 for all  $n$ . Suppose that (23) holds. Then the Fourier series of any function of bounded variation is absolutely summable  $|N, p|$ .*

**THEOREM C.** *Suppose that, for all  $n$ ,  $p_n \geq p_{n+1} > 0$ , and that  $p_n - p_{n+1}$  is nonincreasing. Suppose also that*

$$(24) \quad \sum_{n=0}^K \frac{P_n}{n+1} = O(P_K) .$$

*Then the fourier series of any function of bounded variation is absolutely summable  $|N, p|$ .*

It is immediately evident that Theorem A includes Theorem B. The result that Theorem A includes Theorem C follows from the following lemma, which shows that, in Theorem C, we may replace (24) by (23).

**LEMMA 4.** *Suppose that  $p_0 > 0$ ,  $p_n \geq 0$ . Then (23), (24) are equivalent. In fact, either is equivalent to the assertion*

(c) *There is a constant integer  $r > 1$ , and a constant  $\lambda > 1$  such that, for all sufficiently large  $n$ ,*

$$(25) \quad P_{rn} \geq \lambda P_n .$$

We first prove that (23) implies (c). Suppose, then, that (23) holds. Thus there is a constant  $M$  such that, for all sufficiently large  $K$ ,

$$\sum_{n=K}^{\infty} \frac{1}{nP_n} \leq \frac{M}{P_K} .$$

Since  $P_n$  is nondecreasing, this gives

$$\frac{M}{P_K} > \sum_{n=K}^{rK} \frac{1}{nP_n} \geq \frac{1}{P_{rK}} \sum_{n=K}^{rK} \frac{1}{n} .$$

But

$$\sum_{n=K}^{rK} \frac{1}{n} \longrightarrow \log r$$

as  $K \rightarrow \infty$ , and (c) therefore follows if  $r$  has been chosen so that

$$(26) \quad \log r > M.$$

If (24) holds, we have, for all sufficiently large  $K$ ,

$$\sum_{n=0}^K \frac{P_n}{n+1} \leq MP_K.$$

Thus, replacing  $K$  by  $rK$ ,

$$MP_{rK} > \sum_{n=K}^{rK} \frac{P_n}{n+1} \geq P_K \sum_{n=K}^{rK} \frac{1}{n+1},$$

and we again deduce (c) if  $r$  has been chosen so that (26) holds.

We now consider the converse implications. Suppose, then, that (25) holds for  $n \geq n_0$ . Then, for  $\nu \geq n_0$

$$\sum_{n=r\nu}^{r(\nu+1)-1} \frac{1}{nP_n} \leq \frac{r}{r\nu P_{r\nu}} \leq \frac{1}{\nu \lambda P_\nu}.$$

Hence, for  $K \geq n_0$  and  $s \geq 1$ ,

$$(27) \quad \sum_{n=r^s K}^{r^{s+1}K-1} \frac{1}{nP_n} \leq \frac{1}{\lambda} \sum_{n=r^{s-1}K}^{r^s K-1} \frac{1}{nP_n}.$$

By successive applications of (27), we deduce that for  $s \geq 0$ ,

$$\sum_{n=r^s K}^{r^{s+1}K-1} \frac{1}{nP_n} \leq \frac{1}{\lambda^s} \sum_{n=K}^{rK-1} \frac{1}{nP_n} \leq \frac{1}{\lambda^s P_K} \sum_{n=K}^{rK-1} \frac{1}{n} = O\left(\frac{1}{\lambda^s P_K}\right).$$

Hence

$$\sum_{n=K}^{\infty} \frac{1}{nP_n} = O\left(\frac{1}{P_K} \sum_{s=0}^{\infty} \frac{1}{\lambda^s}\right) = O\left(\frac{1}{P_K}\right),$$

which gives (23). To prove (24), we have, for  $\nu \geq n_0$ ,

$$\sum_{n=r\nu}^{r(\nu+1)-1} \frac{P_n}{n+1} \geq \frac{rP_{r\nu}}{r(\nu+1)} \geq \frac{\lambda P_\nu}{\nu+1}.$$

Hence, for  $s \geq 1$ ,

$$\sum_{n=r^s n_0}^{r^{s+1}n_0-1} \frac{P_n}{n+1} \geq \lambda \sum_{n=r^{s-1}n_0}^{r^s n_0-1} \frac{P_n}{n+1},$$

so that, for  $0 \leq s \leq t-1$ ,

$$(28) \quad \sum_{n=r^s n_0}^{r^{s+1} n_0 - 1} \frac{P_n}{n+1} \leq \frac{1}{\lambda^{t-1-s}} \sum_{n=r^{t-1} n_0}^{r^t n_0 - 1} \frac{P_n}{n+1}.$$

Now take any  $K \geq n_0$ . Choose  $t$  so that  $r^t n_0 \leq K < r^{t+1} n_0$ . Then, by (28),

$$(29) \quad \sum_{n=0}^K \frac{P_n}{n+1} \leq \sum_{n=0}^{n_0-1} \frac{P_n}{n+1} + \sum_{s=0}^{t-1} \frac{1}{\lambda^{t-1-s}} \sum_{n=r^{t-1} n_0}^{r^t n_0 - 1} \frac{P_n}{n+1} \\ + \sum_{n=r^t n_0}^K \frac{P_n}{n+1},$$

where the second term on the right is omitted when  $t = 0$ . The first term on the right of (29) is a constant, and is thus certainly  $O(P_K)$ , since  $P_K \geq p_0 > 0$ . Also

$$\sum_{n=r^{t-1} n_0}^{r^t n_0 - 1} \frac{P_n}{n+1} \leq P_K \sum_{n=r^{t-1} n_0}^{r^t n_0 - 1} \frac{1}{n+1} = O(P_K); \\ \sum_{n=r^t n_0}^K \frac{P_n}{n+1} \leq P_K \sum_{n=r^t n_0}^K \frac{1}{n+1} = O(P_K)$$

(since  $K < r^{t+1} n_0$ ). Thus (24) follows.

The conditions (7), (8) have been mentioned as giving simple sufficient conditions. But, while simpler than (5) or (6), they appear to be insufficiently general to be of great use. Consider, for example, the case of Cesàro summability  $(C, \delta)$ . This is a Nörlund method with

$$p_n = \binom{n + \delta - 1}{n}.$$

If  $0 < \delta \leq 1$ , then the conditions of Theorem A are satisfied. Thus that theorem includes the result that the Fourier series of any function of bounded variation is absolutely summable  $|C, \delta|$ ; this result was long ago proved by Bosanquet [1]. Now, in this case,  $\alpha_{n,k} = 0$  for  $k > n$ , while, for  $1 \leq k \leq n$ ,

$$\alpha_{n,k} = \frac{k}{n} \frac{\binom{n-k+\delta-1}{n-k}}{\binom{n+\delta}{n}}.$$

Thus (a), (b) are both satisfied, with  $r(n) = s(n) = n$ . But either (7) or (8) reduces to

$$\sum_{n=2K}^{\infty} \frac{1}{n \binom{n+\delta}{n}} = O\left(\frac{1}{K}\right),$$

and this is satisfied only if  $\delta = 1$ .

6. As another application of our main theorem, we let  $\{k(n)\}$  be an increasing sequence of nonnegative integers, with  $k(0) = 0$ , and define

$$\alpha_{n,k} = \begin{cases} 1 & (k(n) \leq k < k(n+1)); \\ 0 & \text{otherwise.} \end{cases}$$

Thus absolute summability  $|A|$  of a given series reduces to the absolute convergence of the series formed from it by bracketing together, for every  $n$ , those terms whose suffixes  $k$  satisfy  $k(n) \leq k < k(n+1)$ . It is clear that (a), (b) are both satisfied, with  $r(n) = s(n) = k(n)$  (except when  $n = 0$ ). In this case, the weaker conditions (7), (8) still give a significant result. Either of these conditions is equivalent to

$$(30) \quad \sum_{k(n) \leq K} \frac{1}{k(n)} = O\left(\frac{1}{K}\right).$$

We note that (30) is satisfied, in particular, if

$$(31) \quad \liminf_{n \rightarrow \infty} \frac{k(n+1)}{k(n)} > 1.$$

Thus our theorem includes the following result. *Suppose that (31) holds. Let us bracket together, in the way indicated, the terms of the Fourier series of any function of bounded variation. Then the resulting series is absolutely convergent.*

## REFERENCES

1. L. S. Bosanquet, *Note on the absolute summability (C) of a Fourier series*, J. London Math. Soc., **11** (1936), 11-15.
2. H. P. Dikshit, *On a class of Nörlund means and Fourier series*, Pacific J. Math., **33** (1970), 559-569.
3. M. Izumi and S. Izumi, *Absolute Nörlund summability of Fourier series of functions of bounded variation*, Bull. Austral. Math. Soc., **3** (1970) 111-123.
4. K. Knopp and G. G. Lorentz, *Beiträge zur absoluten Limitierung*, Arch. Math., **2** (1949), 10-16.
5. L. McFadden, *Absolute Nörlund summability*, Duke Math. J., **9** (1942), 168-207.
6. F. M. Mears, *Absolute regularity and the Nörlund mean*, Annals of Math., **38** (1937), 594-601.
7. T. Pati, *On the absolute Nörlund summability of a Fourier series*, J. London Math. Soc., **34** (1959), 153-160.
8. ———, *The non-absolute summability of Fourier series by Nörlund methods*, J. Indian Math. Soc., **25** (1961), 197-214.
9. T. Singh, *Absolute Nörlund summability of a Fourier series*, Indian J. Math., **6** (1964), 129-136.
10. N. Tripathy, *On the absolute Hausdorff summability of Fourier series*, J. London Math. Soc., **44** (1969), 15-25.

Received June 7, 1971.

THE UNIVERSITY BIRMINGHAM, B15 2TT, ENGLAND

AND

THE UNIVERSITY OF CALGARY, CALGARY, ALBERTA, T2N-1N4 CANADA.



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California 94305

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.



Arne P. Baartz and Gary Glenn Miller, <i>Souslin's conjecture as a problem on the real line</i> .....	277
Joseph Barback, <i>On solutions in the regressive isols</i> .....	283
Barry H. Dayton, <i>Homotopy and algebraic K-theory</i> .....	297
William Richard Derrick, <i>Weighted convergence in length</i> .....	307
M. V. Deshpande and N. E. Joshi, <i>Collectively compact and semi-compact sets of linear operators in topological vector spaces</i> .....	317
Samuel Ebenstein, <i>Some <math>H^p</math> spaces which are uncomplemented in <math>L^p</math></i> .....	327
David Fremlin, <i>On the completion of locally solid vector lattices</i> .....	341
Herbert Paul Halpern, <i>Essential central spectrum and range for elements of a von Neumann algebra</i> .....	349
G. D. Johnson, <i>Superadditivity intervals and Boas' test</i> .....	381
Norman Lloyd Johnson, <i>Derivation in infinite planes</i> .....	387
V. M. Klassen, <i>The disappearing closed set property</i> .....	403
B. Kuttner and B. N. Sahney, <i>On the absolute matrix summability of Fourier series</i> .....	407
George Maxwell, <i>Algebras of normal matrices</i> .....	421
Kelly Denis McKennon, <i>Multipliers of type <math>(p, p)</math></i> .....	429
James Miller, <i>Sequences of quasi-subordinate functions</i> .....	437
Leonhard Miller, <i>The Hasse-Witt-matrix of special projective varieties</i> .....	443
Michael Cannon Mooney, <i>A theorem on bounded analytic functions</i> .....	457
M. Ann Piech, <i>Differential equations on abstract Wiener space</i> .....	465
Robert Piziak, <i>Sesquilinear forms in infinite dimensions</i> .....	475
Muril Lynn Robertson, <i>The equation <math>y'(t) = F(t, y(g(t)))</math></i> .....	483
Leland Edward Rogers, <i>Continua in which only semi-aposyndetic subcontinua separate</i> .....	493
Linda Preiss Rothschild, <i>Bi-invariant pseudo-local operators on Lie groups</i> .....	503
Raymond Earl Smithson and L. E. Ward, <i>The fixed point property for arcwise connected spaces: a correction</i> .....	511
Linda Ruth Sons, <i>Zeros of sums of series with Hadamard gaps</i> .....	515
Arne Stray, <i>Interpolation sets for uniform algebras</i> .....	525
Alessandro Figà-Talamanca and John Frederick Price, <i>Applications of random Fourier series over compact groups to Fourier multipliers</i> ....	531