

# Pacific Journal of Mathematics

## **PROXIMITY APPROACH TO SEMI-METRIC AND DEVELOPABLE SPACES**

MANI GAGRAT AND SOMASHEKHAR AMRITH NAIMPALLY

## PROXIMITY APPROACH TO SEMI-METRIC AND DEVELOPABLE SPACES

M. GAGRAT AND S. A. NAIMPALLY

In this paper we study semi-metric and developable spaces via generalized proximities and uniformities. We find sufficient conditions for a compatible semi-metric  $d$  on a space  $X$  to induce a Lodato proximity and also study the effects on a space  $X$  when  $d$  satisfies various weaker forms of continuity. We present two new characterizations of developable spaces, one of which reads: A  $T_1$ -space is developable if and only if it has a compatible upper semi-continuous semi-metric.

We give improved versions of two known metrization theorems. Finally, we generalize the concepts:  $T_1$ -map, uniform map, completely uniform map, pseudo-open map, etc., to apply to proximity spaces and improve some of the known results; for example, an open uniform image of a developable space is developable.

Our motivation for this study was a result mentioned by Arkhangel'skii [4]: A semimetric space  $(X, d)$  is metrizable if the induced nearness relation  $\delta(d)$  is an  $EF$ -proximity (where  $A \delta(d) B$  iff  $d(A, B) = \inf \{d(a, b) : a \in A, b \in B\} = 0$ ). One naturally wonders as to what properties are satisfied by  $X$  if  $\delta(d)$  satisfies weaker proximity axioms such as those of an  $S$ -, a  $LO$ - or an  $R$ -proximity. We soon found that  $\delta(d)$  is always an  $S$ -proximity and that, if  $d$  is the natural semi-metric associated with a developable space, then  $\delta(d)$  is a  $LO$ -proximity. Further impetus was provided by a result due to Pareek [15]: A  $T_1$ -space is semi-metrizable iff there exists a countable family  $\{W_n : n \in \mathbb{N}\}$  of symmetric subsets of  $X \times X$  satisfying (a)  $\bigcap_{n=1}^{\infty} W_n = \Delta$  and (b) for each  $x \in X$ ,  $\{W_n[x] : n \in \mathbb{N}\}$  forms a neighbourhood base at  $x$ . This result is analogous to the well-known Alexandroff-Urysohn theorem: A  $T_1$ -space is metrizable iff it has a compatible uniformity with a countable base. This provided us with a motivation to find one of our new characterizations of developable spaces.

The basic definitions and results are given in this section. In the next section, we study semi-metrizable spaces having compatible semi-metric which satisfy various weaker forms of continuity, e.g., those which are lower semi-continuous or are separately continuous. We also find sufficient conditions for  $\delta(d)$  to be a  $LO$ -proximity.

In the third section, we obtain two new characterizations of developable spaces and also find a sufficient condition for a space to

be a totally bounded developable space. In §4, we prove two new metrization theorems. Finally, in the fifth section, we study the images of some specific proximity spaces under some special mappings.

We now give some basic definitions and results which are needed in this paper. Let  $X$  be a nonempty set and let  $\delta$  be a binary relation on the power set of  $X$ . Consider the following axioms:

- 1.1. (i)  $A\delta B$  implies  $B\delta A$ ,  
 (ii)  $A\delta(B \cup C)$  iff  $A\delta B$  or  $A\delta C$ ,  
 (iii)  $A\delta B$  implies  $A \neq \phi$ ,  $B \neq \phi$ ,  
 (iv)  $A \cap B \neq \phi$  implies  $A\delta B$ ,  
 (v)  $a\delta B$  and  $b\delta C$  for every  $b \in B$  implies  $a\delta C$ ,  
 (vi)  $A\delta B$  and  $b\delta C$  for every  $b \in B$  implies  $A\delta C$ ,  
 (vii)  $a\delta B$  implies the existence of an  $E \subset X$  such that  $a\delta E$  and  $(X - E)\delta B$ ,  
 (viii)  $A\delta B$  implies the existence of an  $E \subset X$  such that  $A\delta E$  and  $(X - E)\delta B$ .

DEFINITION 1.2. The relation  $\delta$  is called

- (a) an *S-proximity* if it satisfies 1.1 (i)–(iv) and (v) (see [13]),  
 (b) a *LO-proximity* if it satisfies 1.1 (i)–(iv) and (vi) (see [13]),  
 (c) an *R-proximity* if it satisfies 1.1 (i)–(iv) and (vii) (see [11]),  
 (d) an *EF-proximity* if it satisfies 1.1 (i)–(iv) and (viii) (see [13]).

By a “proximity  $\delta$ ”, we mean any one of the four proximities defined above and in this case we call the pair  $(X, \delta)$  a *proximity space*. A binary relation  $\delta$  that satisfies 1.1 (i)–(iv) induces an operator  $A \rightarrow \bar{A} = \{x \in X: x\delta A\}$  on the power set of  $X$ , and this will be a Kuratowski closure operator if and only if  $\delta$  is an *S-proximity*. Since (viii) implies (vii), (viii) implies (vi), and (vii) or (vi) implies (v), then every proximity is an *S-proximity*. The topology induced by the closure operator of a proximity  $\delta$  will be written  $\tau(\delta)$ .

We assume that all proximities  $\delta$  considered in this paper are separated, i.e.,  $x\delta y$  implies  $x = y$ . Obviously in this case  $\tau(\delta)$  is  $T_1$ .

If  $(X, \tau)$  is a topological space and  $\delta$  is a proximity on  $X$  such that  $\tau = \tau(\delta)$ , then we say that  $\tau$  and  $\delta$  are compatible. It is known that: (i) every  $T_1$ -space has a compatible *LO*- (and ipso facto *S*-) proximity [13]; (ii) A topological space is  $T_3$  iff it has a compatible *R*-proximity [11]; (iii) A topological space is Tychonoff iff it has a compatible *EF*-proximity.

DEFINITION 1.3. A *semi-metric space*  $(X, d)$  is a  $T_1$ -space  $X$  together with a real-valued function  $d$  on  $X \times X$  such that

- (a)  $d(x, y) = d(y, x) \geq 0$ ,

- (b)  $d(x, y) = 0$  iff  $x = y$ ,  
 (c)  $x \in M^-$  iff  $d(x, M) = 0$ , where  $d(A, B)$   
 $= \inf \{d(a, b): a \in A, b \in B\}$ , if  $A \neq \phi \neq B$   
 $= \infty$  if  $A = \phi$  or  $B = \phi$ .

If  $(X, d)$  is a semi-metric space, we define  $\delta = \delta(d)$  by

$$A\delta B \text{ iff } d(A, B) = 0.$$

It is easily verified that  $\delta(d)$  is an  $S$ -proximity on  $X$ . For  $\varepsilon > 0$ , we set

$$V_\varepsilon = \{(x, y) \in X \times X: d(x, y) < \varepsilon\}.$$

Clearly  $V_\varepsilon[x] = S(x, \varepsilon)$ , the sphere with centre  $x$  and radius  $\varepsilon$ . We set  $\mathcal{U}_d = \{U = U^{-1} \subset X \times X: V_{1/n} \subset U \text{ for some } n \in N\}$ . Also for  $\mathcal{U} \subset \mathcal{P}(X \times X)$  (the power set of  $X \times X$ ), we define  $\delta = \delta(\mathcal{U})$  by

$$A\delta B \text{ iff } (A \times B) \cap U \neq \phi \text{ for each } U \in \mathcal{U}.$$

Clearly, if  $d$  is a semi-metric on  $X$ , then  $\delta(d) = \delta(\mathcal{U}_d)$ .

**DEFINITION 1.4.** A *refining family*  $\Sigma$  on a topological space  $(X, \tau)$  is a family  $\{\alpha_i: i \in I\}$  of open covers of  $X$  such that for each  $x \in U \in \tau$ , there exists an  $i \in I$  such that  $\text{St}(x, \alpha_i) \subset U$ . In the special case when  $I = N$ ,  $\Sigma$  is called a *development* on  $X$  and the pair  $(X, \Sigma)$  is called a *developable space*. In this case, it is well known that  $\Sigma$  may be replaced by another development  $\Sigma' = \{\beta_j: j \in N\}$  such that if  $i < j$  then  $\beta_j \subset \beta_i$ ; we will assume that  $\Sigma$  already satisfies this condition. We will also assume that the developable spaces are  $T_1$ .

**LEMMA 1.5.** Every refining family  $\Sigma$  on a  $T_1$ -space  $(X, \tau)$  induces a compatible LO-proximity  $\delta = \delta(\Sigma)$  on  $X$ , where  $\delta$  is defined by

$$A\delta B \text{ iff } \text{St}(A, \alpha_i) \cap B \neq \phi \text{ for each } i \in I.$$

*Proof.* That  $\delta$  satisfies 1.1 (i)–(iv), is separated, and is compatible with  $\tau$  is obvious. We now show that  $\delta$  satisfies 1.1 (vi). Suppose  $A\delta B$  and  $b\delta C$  for each  $b \in B$ . Then  $A\delta B$  implies that for each  $i \in I$ , there is a  $b \in B$  such that  $b \in \text{St}(A, \alpha_i)$ . Since  $\text{St}(A, \alpha_i)$  is open, there is a  $j \in I$  such that  $b \in \text{St}(b, \alpha_j) \subset \text{St}(A, \alpha_i)$ . Since  $b\delta C$ ,  $C \cap \text{St}(b, \alpha_j) \neq \phi$  and this, in turn, implies  $C \cap \text{St}(A, \alpha_i) \neq \phi$ , i.e.,  $A\delta C$ .

For a developable space  $(X, \Sigma)$ ,  $\Sigma = \{\lambda_n: n \in N\}$ , we define  $d = d(\Sigma)$  by

$$d(x, y) = \inf \left\{ \frac{1}{n+1}: y \in \text{St}(x, \lambda_n) \right\}.$$

It is easily seen that  $d$  is a compatible semi-metric on  $X$  and that

$\delta(d) = \delta(\Sigma)$ . For each  $n \in N$ , we set

$$B_n = \cup \{G \times G : G \in \lambda_n\}.$$

LEMMA 1.6.  $B_n = V_{1/n}$ .

*Proof.*  $(x, y) \in V_{1/n}$  iff  $d(x, y) \leq 1/(n+1) < 1/n$  iff  $y \in \text{St}(x, \lambda_n)$  iff  $(x, y) \in B_n$ .

DEFINITION 1.7. An  $M$ -uniformity base  $\mathcal{B}$  on  $X$  is a family of subsets of  $X \times X$  such that,

- (i)  $\cap \{U : U \in \mathcal{B}\} = \Delta$ ,
  - (ii)  $U = U^{-1}$  for each  $U \in \mathcal{B}$ ,
  - (iii) for each  $A \subset X$  and  $U, V$  in  $\mathcal{B}$ , there exists a  $W \in \mathcal{B}$  such that  $W[A] \subset U[A] \cap V[A]$ ,
  - (iv) for each  $U \in \mathcal{U}$ ,  $U \subset V = V^{-1} \subset X \times X$  implies  $V \in \mathcal{B}$ ,
  - (v) for  $A, B$  in  $P(X)$  and  $U \in \mathcal{B}$ , if  $V[A] \cap B \neq \phi$  for each  $V \in \mathcal{B}$ ; then there exists an  $x \in B$  and a  $W \in \mathcal{B}$  such that  $W[x] \subset U[A]$ .
- $\mathcal{U} \subset P(X \times X)$  is said to be an  $M$ -uniformity on  $X$  iff there exists a family  $\mathcal{B} \subset \mathcal{U}$  satisfying conditions (i)–(v) above and for each  $U \in \mathcal{U}$  there is a  $B \in \mathcal{B}$  such that  $B \subset U$ .

REMARK 1.8. Mozzochi [12] has shown that an  $M$ -uniformity  $\mathcal{U}$  need not have an open base and that if it does have one, then every convergent filter on  $X$  is  $\mathcal{U}$ -Cauchy.

DEFINITION 1.9. An  $S$ -uniformity  $\mathcal{U}$  on  $X$  is a family of subsets of  $X \times X$  which has a base  $\mathcal{B}$  satisfying 1.7 (i)–(iv) and for each  $p \in X$ ,  $B \subset X$  and  $U \in \mathcal{B}$ , if  $V[p] \cap B \neq \phi$  for each  $V \in \mathcal{B}$ , then there exists an  $x \in B$  and a  $W \in \mathcal{B}$  such that  $W[x] \subset U[p]$ .

If  $\mathcal{U}$  is an  $M$ - or an  $S$ -uniformity, then  $\tau(\mathcal{U})$  is defined as usual; viz.  $G \in \tau(\mathcal{U})$  iff for each  $x \in G$ , there exists a  $U \in \mathcal{U}$  such that  $U[x] \subset G$ . If  $\tau = \tau(\mathcal{U})$ , we say that  $\tau$  and  $\mathcal{U}$  are compatible.

THEOREM 1.10. (See [7], [12].) If  $\mathcal{U}$  consists of symmetric subsets of  $X \times X$ , then  $\mathcal{U}$  is an  $M$ -uniformity base (resp.  $S$ -uniformity base) if and only if  $\delta(\mathcal{U})$  is a LO-proximity (resp.  $S$ -proximity).

Let  $(X, d)$  be a semi-metric space and set  $V_{1/n} = \{(x, y) : d(x, y) < 1/n\}$ . Then  $\{V_{1/n} : n \in N\}$  is a countable base for a compatible  $S$ -uniformity.

2. Semi-metric spaces. In this section we suppose that  $(X, d)$  is a semi-metric space and consider the effects of various forms of continuity properties of  $d$  on the topology of  $X$  and on the proximity  $\delta(d)$ .

LEMMA 2.1. *In the following (i) and (ii) are equivalent and each implies (iii):*

- (i)  *$d$  is separately upper semi-continuous.*
- (ii) *For each  $\varepsilon > 0$  and  $x \in X$ ,  $S(x, \varepsilon)$  is open.*
- (iii)  *$\delta(d)$  is a LO-proximity on  $X$ .*

*Proof.* (i)  $\Rightarrow$  (ii). Let  $y \in S(x, \varepsilon)$ , i.e.,  $d(x, y) < \varepsilon$ . Since  $d$  is upper semi-continuous in  $y$ , for every  $\eta > 0$  there is a nbhd.  $N_y$  of  $y$  such that  $d(x, z) < d(x, y) + \eta$  for each  $z \in N_y$ . Choose  $\eta < \varepsilon - d(x, y)$ . Then clearly  $N_y \subset S(x, \varepsilon)$ , showing thereby that  $S(x, \varepsilon)$  is open.

(ii)  $\Rightarrow$  (i). Suppose  $d(x, y) = r$  and  $\varepsilon > 0$ . Clearly  $y \in S(x, r + \varepsilon)$ , which is open, and hence there exists a nbhd.  $N_y$  of  $y$  such that  $N_y \subset S(x, r + \varepsilon)$ . But this means that for each  $z \in N_y$ ,  $d(x, z) < d(x, y) + \varepsilon$ , i.e.,  $d$  is separately upper semi-continuous.

(i)  $\Rightarrow$  (iii). Let  $\delta = \delta(d)$  and suppose  $A\delta B$  and  $b\delta C$  for each  $b \in B$ . Then for each  $\varepsilon > 0$ , there exists an  $a \in A$  and a  $b \in B$  such that  $d(a, b) < \varepsilon$ . Since  $d$  is upper semi-continuous at  $b$ , there exists a nbhd.  $N_b$  of  $b$  such that  $d(a, x) < \varepsilon$  for each  $x \in N_b$ . Also  $b\delta C$  implies the existence of a point  $c \in C \cap N_b$  and hence,  $d(a, c) < \varepsilon$ , i.e.,  $A\delta C$ .

COROLLARY 2.2. *If a semi-metric  $d$  is separately upper semi-continuous, then  $\mathcal{U}_d$  is an  $M$ -uniformity (obviously with a countable base).*

LEMMA 2.3. *If  $d$  is separately lower semi-continuous, then  $X$  is regular.*

*Proof.* Let  $A$  be closed in  $X$  and  $p \in X - A$ . Then  $d(p, A) = r > 0$  and hence for each  $a \in A$ ,  $d(p, a) \geq r$ . Since  $d$  is lower semi-continuous at  $a$ , there exists a nbhd.  $N_a$  of  $a$  such that for each  $x \in N_a$ ,  $d(p, x) > r/2$ . Set  $N_A = \bigcup \{N_a : a \in A\}$ . Then  $N_A$  is a nbhd. of  $A$  and  $N_A \cap S(p, r/2) = \emptyset$ , thereby showing that  $X$  is regular.

The following result is an improvement of a similar one stated by Cook [6]; whereas Cook require the semi-metric  $d$  to be continuous we want  $d$  to be only *separately* continuous.

THEOREM<sup>1</sup> 2.4. *If  $d$  is separately continuous, then  $X$  is Tychonoff.*

*Proof.* Suppose  $A$  is a closed subset of  $X$  and  $x \in X - A$ . We may assume that  $d(x, A) = 1$ . As in the proof of Urysohn's lemma, we now show how to construct, for each positive rational  $r \in [0, 1]$ , an open set  $V_r$  such that  $x \in V_r$  for each such  $r$ ,  $V_r^- \subset V_s$  whenever  $r < s$  and each  $V_r \subset X - A$ . We set  $V_r = \{y \in X : d(x, y) < r\}$  which is open

<sup>1</sup> This result was jointly obtained by Dr. C. M. Pareek and the second author.

by 2.1 (ii). Also the separate lower semi-continuity of  $d$  is equivalent to the set  $\{y \in X: d(x, y) \leq r\}$ , being closed for each  $r$ .

The following theorem is an analogue of the result: A  $T_1$ -space is uniformizable with a countable base iff it is metrizable (and hence has a metric  $d$  such that  $\delta(d)$  is an *EF*-proximity.)

**THEOREM 2.5.** *A  $T_1$ -space is  $M$ -uniformizable with a countable base if and only if it has a compatible semi-metric  $d$  such that  $\delta(d)$  is a  $LO$ -proximity.*

*Proof.* Sufficiency is evident from 1.10 and necessity follows from 1.10 and Theorem 4.1.7 of Pareek [15], making use of the remarks just preceding 1.4.

The following analogue of the above result is proved similarly.

**THEOREM 2.6.** *A  $T_1$ -space is  $S$ -uniformizable with a countable base if and only if it has a compatible semi-metric  $d$  (and obviously  $\delta(d)$  is an  $S$ -proximity).*

**3. Developable spaces.** In this section we suppose that  $(X, \Sigma)$  is a developable space with  $\Sigma = \{\lambda_n: n \in N\}$  where each  $\lambda_n$  is an open cover of  $X$  and  $\lambda_{n+1} \subset \lambda_n$ . Let  $d = d(\Sigma)$  be the induced semi-metric on  $X$ . Then  $\delta(d) = \delta(\Sigma)$  is a  $LO$ -proximity on  $X$ .

**LEMMA 3.1.**  *$d$  is upper semi-continuous.*

*Proof.* If  $p, q \in X$ , we have to consider two cases: (i)  $d(p, q) = 0$  and (ii)  $d(p, q) = 1/(m+1)$  for some  $m \in N$ . Let  $\varepsilon > 0$  be arbitrary. In case (i),  $\delta(d)$  is a  $LO$ -proximity by 2.1 and  $p = q$ . Now choose  $n \in N$  such that  $1/n < \varepsilon$ ; then  $p, q \in G$  for some  $G \in \lambda_n$ . For all  $(x, y) \in G \times G$ ,  $d(x, y) \leq 1/(n+1) < \varepsilon$ . In case (ii),  $p, q \in G$  for some  $G \in \lambda_m$ . Then for all  $(x, y) \in G \times G$ ,  $d(x, y) \leq 1/(m+1) < 1/(m+1) + \varepsilon$ . Thus the result is proved.

The above result (in conjunction with 2.1) provides an alternate proof of the fact that  $\delta(\Sigma)$  is a compatible  $LO$ -proximity and also shows that  $\mathcal{U}_d$  is a compatible  $M$ -uniformity with a countable open base  $\{B_n: n \in N\}$ . This provides a motivation for our next result.

**THEOREM 3.2.** *A  $T_1$ -space is developable if and only if it is  $M$ -uniformizable with a countable open base.*

*Proof.* Necessity follows from the remarks preceding 3.2 and sufficiency follows from Brown's result ([5], p. 65) that a space is developable iff it has a compatible semi-metric  $d$  for which every con-

vergent sequence is Cauchy, using 1.8.

In [6] Cook states that if a compatible semi-metric  $d$  on  $X$  is continuous, then  $X$  is developable. The following characterization of developable spaces is an improvement of this result.

**THEOREM 3.3.** *A  $T_1$ -space  $X$  is developable if and only if it has a compatible upper semi-continuous semi-metric.*

*Proof.* Since the necessity has been proved in (3.1), we need prove only sufficiency. Assuming  $d$  to be an upper semi-continuous semi-metric, we prove that  $V_{1/n}$  is open for each  $n \in \mathbb{N}$  and then the result will follow from (3.2). If  $(p, q) \in V_{1/n}$ , then  $d(p, q) < 1/n$ . Choose  $\varepsilon > 0$  such that  $\varepsilon < 1/n - d(p, q)$ . Then from the upper semi-continuity of  $d$ , there exist nbhds.  $N_p, N_q$  of  $p, q$  respectively, such that  $d(x, y) < d(p, q) + \varepsilon < 1/n$  for each  $(x, y) \in N_p \times N_q$ . This shows that  $(N_p \times N_q) \subset V_{1/n}$ , showing thereby that  $V_{1/n}$  is open.

**DEFINITION 3.4.** A proximity base  $\mathcal{B}$  for a proximity space  $(X, \delta)$  is a family of subsets of  $X$  such that if  $A \delta B$ , then there exist  $C, D \in \mathcal{B}$  such that  $A \subset C, B \subset D$  and  $C \delta D$ .

In a  $LO$ -space, we may assume that the members of  $\mathcal{B}$  are closed. The following is an improvement of 2.23 of [12].

**LEMMA 3.5.** *If  $\mathcal{B}$  is a proximity base for  $LO$ -space  $(X, \delta)$ , then  $\{U_{A,B} : A, B \in \mathcal{B} \text{ and } A \delta B\}$ , where  $U_{A,B} = X \times X - [(A \times B) \cup (B \times A)]$ , is a base for an  $M$ -uniformity  $\mathcal{U}_1$  on  $X$ .*

A developable space  $(X, \Sigma)$  is said to be *totally bounded* iff for each  $n \in \mathbb{N}$ , there exists a finite set  $F \subset X$  such that  $\text{St}(F, \lambda_n) = X$ . It is known that an  $EF$ -space  $(X, \delta)$  has a countable base iff  $X$  has a compatible totally bounded metric (Theorem (8.19) of [13]). The following is a partial generalization.

**THEOREM 3.6.** *If a  $LO$ -space  $(X, \delta)$  has a countable closed base  $\mathcal{B}$ , then  $X$  is a totally bounded developable space.*

*Proof.*  $\{U_{A,B} : A, B \in \mathcal{B}, A \delta B\}$  is a countable open base for a compatible  $M$ -uniformity, which is also totally bounded. The result then follows from (3.2).

**4. Metrizable spaces.** Arkhangel'skii [4] proved that a semi-metric space  $(X, d)$  is metrizable if  $\delta(d)$  is an  $EF$ -proximity. A glance at his proof shows that the following improved version is true.



**THEOREM 4.1.** *A  $T_1$ -space is uniformizable with a countable base if and only if it has a compatible semi-metric  $d$  such that  $\delta(d)$  is an  $R$ -proximity if and only if it is metrizable.*

Nedev [14] proved that a semi-metric space  $(X, d)$  is metrizable if  $d(A, x)$  is a continuous function of  $x$  for each  $A \subset X$ . The following is an improvement.

**THEOREM 4.2.** *A semi-metric space  $(X, d)$  is metrizable if for all closed subsets  $A$  of  $X$ ,  $d(A, x)$  is lower semi-continuous.*

*Proof.* Suppose  $A$  is closed,  $B$  is compact and  $A \cap B = \phi$ . Since the function  $d(A, x)$  is lower semi-continuous and  $B$  is compact, it follows that  $d(A, B) = d(A, x)$  for some  $x \in B$ . This implies that  $X$  is metrizable (see [4]).

We now give a table which shows the relationship of our results with the classical Alexandroff-Urysohn uniform metrization theorem. Suppose  $\{W_n: n \in N\}$  is a countable family of symmetric subsets of  $X \times X$  satisfying (a)  $\bigcap_{n=1}^\infty W_n = \Delta$  and (b) for each  $x \in X$ ,  $\{W_n[x]: n \in N\}$  forms a nbhd. base at  $x$ . Let  $d$  be the semi-metric by  $\{W_n\}$ , namely

$$d(x, y) = \text{Inf} \left\{ \frac{1}{n + 1} : y \in W_n[x] \right\} .$$

(We assume without any loss of generality that  $W_{n+1} \subset W_n$ .) Finally set  $\mathcal{U} = \{U \subset X \times X: W_n \subset U \text{ for some } n \in N\}$ .

$\mathcal{U}$ is an $S$ -uniformity	$\mathcal{U}$ is an $M$ -uniformity	$\mathcal{U}$ is an $M$ -uniformity. Each $W_n$ is open	$\mathcal{U}$ is a uniformity
$\delta(d)$ is an $S$ -proximity	$\delta(d)$ is a $LO$ -proximity	$d$ is USC $\delta(d)$ is a $LO$ -proximity	$\delta(d)$ is an $EF$ -proximity
$X$ is semi-metrizable	$X$ is semi-metrizable	$X$ is developable	$X$ is metrizable

**5. Metric spaces, developable spaces, semi-metric spaces and mappings connected with them.** In this section  $f: X \rightarrow Y$  will denote a function from a proximity space  $(X, \delta)$  onto a  $T_1$ -space  $Y$ . When  $X$  is developable, semi-metrizable or metrizable,  $\delta$  will denote the corresponding naturally induced proximity relation as defined in §1. Several kinds of functions, which have been defined for the case in which

$X$  is metrizable, can be redefined more generally when  $X$  is a proximity space; this is done by replacing the condition  $d(A, B) > 0$  by  $A\delta B$ . These mappings have been systematically discussed in the metric case by Arkhangel'skii [4]. Although we consider their generalizations, for the sake of simplicity we will keep the same terminology and attach  $\delta$ -before each term<sup>2</sup>.

Our strategy consists in defining a binary relation  $\delta^1$  on the power set of  $Y$  as follows.

$$(5.1) \quad E\delta^1 F \text{ iff } f^{-1}(E)\delta f^{-1}(F).$$

It is easily verified that  $\delta^1$  satisfies 1.1 (i)—(iv) and so  $\delta^1$  is almost a quotient proximity. In order that  $\delta^1$  be a proximity, naturally we will have to put some additional conditions on  $f$ ; also it is clear that if  $\delta^1$  is to satisfy stronger proximity conditions, so must  $\delta$ .

Our first task is to find conditions on  $f$  which will make  $\delta^1$  an  $S$ -proximity compatible with the topology of  $Y$ .

**DEFINITION 5.2.** The function  $f$  is called  $\delta$ -pseudo-open iff for each  $y \in Y$  and  $A \subset X$ , if  $f^{-1}(y)\delta(X - A)$  then  $y \in \text{Int} f(A)$ .

It is easily verified that if  $f$  is open or closed, then it is also  $\delta$ -pseudo-open.

**LEMMA 5.3.** If  $\delta$  is an  $S$ -proximity on  $X$  and  $f$  is  $\delta$ -pseudo-open, then  $y\delta^1 E$  implies  $y \notin E^-$ .

*Proof.*  $y\delta^1 E$  implies  $f^{-1}(y)\delta f^{-1}(E)$ , and since  $f$  is  $\delta$ -pseudo-open, it follows that  $y \in \text{Int} f[X - f^{-1}(E)]$ . But  $E \cap f[X - f^{-1}(E)] = \emptyset$  and hence  $y \notin E^-$ .

**COROLLARY 5.4.** If  $\delta$  is an  $S$ -proximity and  $f$  is either open or closed, then  $y\delta^1 E$  implies  $y \notin E^-$ .

**DEFINITION 5.5.** The function  $f$  is called  $\delta$ -uniform iff for each  $y \in Y$  and each nbhd.  $N_y$  of  $Y$ ,  $f^{-1}(y)\delta(X - f^{-1}(N_y))$ .

**LEMMA 5.6.** If  $\delta$  is an  $S$ -proximity, then  $f$  is  $\delta$ -uniform if and only if  $y \notin E^-$  implies  $y\delta^1 E$ .

*Proof.* If  $f$  is  $\delta$ -uniform and  $y \notin E^-$ , then  $f^{-1}(y)\delta f^{-1}(E)$  and by 5.1,  $y\delta^1 E$ . Conversely, if  $N_y$  is a nbhd. of  $y$ , then  $y \notin (Y - N_y)^-$  implies  $y\delta^1 (Y - N_y)$  and this is equivalent to  $f^{-1}(y)\delta(X - f^{-1}(N_y))$ . Thus  $f$  is  $\delta$ -uniform.

<sup>2</sup> This terminology was suggested by Dr. C. M. Pareek.

LEMMA 5.7. *If  $\delta$  is an  $R$ -proximity and  $f$  is continuous and compact, then  $y \notin E^-$  implies  $y\delta^1 E$ .*

*Proof.* If  $y \notin E^-$  then  $f^{-1}(y) \cap f^{-1}(E^-) = \phi$ . Since  $f$  is compact,  $f^{-1}(y)$  is compact and since  $f$  is continuous,  $f^{-1}(E^-)$  is closed. Finally,  $\delta$  being an  $R$ -proximity on  $X$ ,  $f^{-1}(y)\delta f^{-1}(E)$ , which in turn implies  $y\delta^1 E$ .

The following result is now obvious.

THEOREM 5.8. *The relation  $\delta^1$  is a compatible  $S$ -proximity on  $Y$  in the following cases:*

- (a)  *$\delta$  is an  $S$ -proximity,  $f$  is  $\delta$ -pseudo-open and  $\delta$ -uniform;*
- (b)  *$\delta$  is an  $R$ -proximity,  $f$  is  $\delta$ -pseudo-open, continuous, and compact.*

Next we find out when  $\delta^1$  is a  $LO$ -proximity.

THEOREM 5.9. *If  $\delta$  is a  $LO$ -proximity on  $X$  and  $f$  is  $\delta$ -uniform and open, then  $\delta^1$  is a compatible  $LO$ -proximity on  $Y$ .*

*Proof.* By 5.8 (b),  $\delta^1$  is a compatible  $S$ -proximity and since  $f$  is  $\delta$ -uniform, it follows that  $f$  is continuous. This together with the openness of  $f$  implies that for each  $B \subset Y$ ,  $f^{-1}(B^-) = f^{-1}(B)^-$ . Now  $A\delta^1 B$  iff  $f^{-1}(A)\delta f^{-1}(B)$  iff  $f^{-1}(A)\delta f^{-1}(B^-)$  iff  $f^{-1}(A^-)\delta f^{-1}(B^-)$  iff  $A^-\delta^1 B^-$ , thereby showing that  $\delta^1$  is a  $LO$ -proximity.

To investigate as to when  $\delta^1$  is an  $R$ -proximity we introduce a stronger type of map.

DEFINITION 5.10. The function  $f$  is called  $\delta$ -completely uniform iff for each nbhd.  $N_y$  of  $y$  in  $Y$ , there exists a nbhd.  $N'_y$  of  $y$  such that  $f^{-1}(N'_y)\delta(X - f^{-1}(N_y))$ .

Since the identity map on  $X$  is not  $\delta$ -completely uniform unless  $\delta$  is an  $R$ -proximity, it is clear that in order to have a meaningful discussion, we must have  $\delta$  an  $R$ -proximity.

THEOREM 5.11. *If  $\delta$  is an  $R$ -proximity on  $X$  and  $f$  is open and  $\delta$ -uniform, then  $f$  is  $\delta$ -completely uniform if and only if  $\delta^1$  is a compatible  $R$ -proximity.*

*Proof.* To prove the necessity, we first note that Theorem 5.9 shows that  $\delta^1$  is a compatible  $LO$ -proximity on  $Y$ . To see that  $\delta^1$  is an  $R$ -proximity, we note that  $y\delta^1 E$  implies  $y \in Y - E^-$  which is open. Since  $f$  is  $\delta$ -completely uniform, there is a nbhd.  $N_y$  of  $y$  such that  $f^{-1}(N_y)\delta f^{-1}(E)$  i.e.  $N_y\delta^1 E$ . Since trivially  $y\delta(Y - N_y)$ , it follows that  $\delta^1$  is an  $R$ -proximity.

Conversely, suppose  $f$  is open  $\delta$ -uniform and  $\delta^1$  is an  $R$ -proximity. If  $N_y$  is a nbhd. of  $y \in Y$ ,  $y\delta^1(Y - N_y)$  and, since  $\delta^1$  is an  $R$ -proximity, there exist a set  $A \subset Y$  such that  $y\delta^1(Y - A)$  and  $A\delta^1(Y - N_y)$ , i.e.,  $y \in \text{Int}(A)$  and  $f^{-1}(A)\delta f^{-1}(Y - N_y) = X - f^{-1}(N_y)$ . This proves that  $f$  is  $\delta$ -completely uniform.

In view of Theorem 4.1, it is not necessary to investigate as to when  $\delta^1$  is an  $EF$ -proximity and we now turn our attention to the case when  $(X, d)$  is a semi-metric space. In this case we suppose that  $\delta = \delta(d)$ .

**DEFINITION 5.12.** The function  $f$  is called a  $T_1$ -map iff for every pair of distinct points  $y_1, y_2$  of  $Y$ ,  $f^{-1}(y_1)\delta f^{-1}(y_2)$ ; equivalently,  $d(f^{-1}(y_1), f^{-1}(y_2)) > 0$ .

**LEMMA 5.13.** *If  $(X, d)$  is a semi-metric space and  $f$  is a  $T_1$ -map, then*

$$\rho(y_1, y_2) = d(f^{-1}(y_1), f^{-1}(y_2))$$

*is a semi-metric on  $Y$  (not necessarily compatible with the topology of  $Y$ ).*

**COROLLARY 5.14.** *If either  $d$  is a metric and  $f$  is compact or if  $d$  is a semi-metric and  $f$  is  $\delta$ -uniform, then  $\rho$  is a semi-metric on  $Y$ .*

**LEMMA 5.15.** *Under the conditions of Lemma 5.13, for subsets  $A, B$  of  $Y$ ,*

$$\rho(A, B) = d(f^{-1}(A), f^{-1}(B)) .$$

*Proof.*  $\rho(A, B) = \text{Inf} \{d(f^{-1}(a), f^{-1}(b)): a \in A, b \in B\}$ . Now

$$d(f^{-1}(A), f^{-1}(B)) \leq d(f^{-1}(a), f^{-1}(b))$$

for each  $a \in A, b \in B$  and this implies that  $(f^{-1}(A), f^{-1}(B)) \leq \rho(A, B)$ . Conversely, for every  $\varepsilon > 0$ , there exist  $x \in f^{-1}(A), y \in f^{-1}(B)$  such that  $d(x, y) < d(f^{-1}(A), f^{-1}(B)) + \varepsilon$ , and hence

$$\rho(A, B) \leq d(x, y) < d(f^{-1}(A), f^{-1}(B)) + \varepsilon ,$$

i.e.,  $\rho(A, B) \leq d(f^{-1}(A), f^{-1}(B))$ , thus proving the result.

**COROLLARY 5.16.** *Under the conditions of Lemma 5.15,  $A \delta(\rho)B$  iff  $f^{-1}(A)\delta f^{-1}(B)$ .*

**LEMMA 5.17.** *If  $d$  is (resp. separately) upper semi-continuous and*

$f$  is open  $\delta$ -uniform, then  $\rho$  is (resp. separately) upper semi-continuous.

*Proof.* Let  $\rho(y_1, y_2) = r$  and let  $\varepsilon > 0$ . Then there are  $x_i \in f^{-1}(y_i)$ ,  $i = 1, 2$  such that  $d(x_1, x_2) < r + \varepsilon/2$ . Since  $d$  is upper semi-continuous, there exist nbhds.  $N_{x_i}$  of  $x_i$ ,  $i = 1, 2$  such that for all  $z_i \in N_{x_i}$ ,  $i = 1, 2$ ,  $d(z_1, z_2) < r + \varepsilon$ . But this shows that  $\rho(p_1, p_2) < r + \varepsilon$  for  $p_i \in f(N_{x_i})$ ,  $i = 1, 2$  and the result follows since  $f$  is open. The case of separate upper semi-continuity is similarly handled.

It is well known that if  $X$  is metrizable and  $f$  is open  $\delta$ -uniform, then  $Y$  is a developable space. We get an improvement and a more "symmetric" result from (3.3) and (5.17); viz.,

**THEOREM 5.18.** *If  $X$  is developable and  $f$  is open  $\delta$ -uniform with respect to the proximity  $\delta$  induced by the development on  $X$ , then  $Y$  is also developable.*

**THEOREM 5.19.** *If  $d$  is a metric on  $X$  and if  $f$  is continuous, open and compact, then  $Y$  is developable.*

*Proof.* By setting  $\delta = \delta(d)$ ,  $\delta^1 = \delta(\rho)$ , the result follows from the continuity of  $d$ , (5.8), (5.17) and (3.3).

Finally, we conclude with a known result but which follows easily from our analysis.

**THEOREM 5.20.** *If  $d$  is a metric on  $X$  and if  $f$  is open and  $\delta$ -completely uniform, then  $Y$  is metrizable.*

*Proof.* Set  $\delta = \delta(d)$  and  $\delta^1 = \delta(\rho)$ . It follows from (5.11) that  $\delta^1$  is a compatible  $R$ -proximity on  $Y$ . Also from (5.14),  $\rho$  is a semi-metric on  $Y$  and the compatibility of  $\delta^1$  implies the compatibility of  $\rho$ . Hence by (4.1),  $Y$  is metrizable.

## REFERENCES

1. P. S. Alexandroff, *On some results concerning topological spaces and their continuous mappings*, Proc. Symp. Prague (1961), 41-45.
2. ———, *Some results in the theory of topological spaces, obtained within the last twenty-five years*, Russian Math. Surveys, **15** (1960), 23-83.
3. ———, *Some basic directions in general topology*, Russian Math. Surveys, **19** (1964), 1-39.
4. A. V. Arkhangel'skii, *Mappings and spaces*, Russian Math. Surveys, **21** (1966), 87-114.
5. M. Brown, *Semi-metric spaces*, Summer Inst. Set Theoretic Topology, Madison, Wisconsin (1957), 64-66.
6. H. Cook, *Cartesian products and continuous semi-metrics*, Topology Conference, Arizona State University (1967), 58-63.

7. M. Gagrath, *Proximities and extensions of continuous functions*, Ph. D. Thesis, I.I.T., Kanpur (1970), (unpublished).
8. M. S. Gagrath and S. A. Naimpally, *Wallman compactifications and Wallman real-compactifications*, J. Australian Math. Soc., (To appear).
9. ———, *Proximity approach to extension problems*, Fund. Math., **71** (1971), 63-76.
10. ———, *Proximity approach to extension problems II*, Math. Japonicae, **16** (1971), 35-43.
11. D. Harris, *Regular-closed spaces and proximities*, Pacific J. Math., **34** (1970), 675-685.
12. C. J. Mozzochi, M. S. Gagrath and S. A. Naimpally, *Symmetric generalized topological structures*, (1970), (Preprint).
13. S. A. Naimpally and B. D. Warrack, *Proximity Spaces*, Cambridge Tract In Mathematics No. 59, Cambridge University Press (1970).
14. S. I. Nedev, *Continuous and semi-continuous  $\phi$ -metrics*, Soviet Math. Dokl., **11** (1970), 975-978.
15. C. M. Pareek, *A study of some generalizations of paracompact and metrizable spaces*, Ph. D. Thesis, University of Alberta, Canada (1971) (unpublished).

Received July 15, 1971 and in revised form May 1, 1972. The authors wish to thank Dr. C. M. Pareek and the referee for valuable comments and suggestions. Revision was supported by N.R.C. (Canada) grant.

INDIAN INSTITUTE OF TECHNOLOGY, KANPUR  
 UNIVERSITY OF COLORADO  
 AND  
 SOUTHERN ILLINOIS UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan



Jimmy T. Arnold, <i>Power series rings over Prüfer domains</i> .....	1
Maynard G. Arsove, <i>On the behavior of Pincherle basis functions</i> .....	13
Jan William Auer, <i>Fiber integration in smooth bundles</i> .....	33
George Bachman, Edward Beckenstein and Lawrence Narici, <i>Function algebras over valued fields</i> .....	45
Gerald A. Beer, <i>The index of convexity and the visibility function</i> .....	59
James Robert Boone, <i>A note on mesocompact and sequentially mesocompact spaces</i> .....	69
Selwyn Ross Caradus, <i>Semiclosed operators</i> .....	75
John H. E. Cohn, <i>Two primary factor inequalities</i> .....	81
Mani Gagrath and Somashekhar Amrith Naimpally, <i>Proximity approach to semi-metric and developable spaces</i> .....	93
John Grant, <i>Automorphisms definable by formulas</i> .....	107
Walter Kurt Hayman, <i>Differential inequalities and local valency</i> .....	117
Wolfgang H. Heil, <i>Testing 3-manifolds for projective planes</i> .....	139
Melvin Hochster and Louis Jackson Ratliff, Jr., <i>Five theorems on Macaulay rings</i> .....	147
Thomas Benton Hoover, <i>Operator algebras with reducing invariant subspaces</i> .....	173
James Edgar Keesling, <i>Topological groups whose underlying spaces are separable Fréchet manifolds</i> .....	181
Frank Leroy Knowles, <i>Idempotents in the boundary of a Lie group</i> .....	191
George Edward Lang, <i>The evaluation map and EHP sequences</i> .....	201
Everette Lee May, Jr., <i>Localizing the spectrum</i> .....	211
Frank Belsley Miles, <i>Existence of special K-sets in certain locally compact abelian groups</i> .....	219
Susan Montgomery, <i>A generalization of a theorem of Jacobson. II</i> .....	233
T. S. Motzkin and J. L. Walsh, <i>Equilibrium of inverse-distance forces in three-dimensions</i> .....	241
Arunava Mukherjea and Nicolas A. Tserpes, <i>Invariant measures and the converse of Haar's theorem on semitopological semigroups</i> .....	251
James Waring Noonan, <i>On close-to-convex functions of order <math>\beta</math></i> .....	263
Donald Steven Passman, <i>The Jacobian of a growth transformation</i> .....	281
Dean Blackburn Priest, <i>A mean Stieltjes type integral</i> .....	291
Joe Bill Rhodes, <i>Decomposition of semilattices with applications to topological lattices</i> .....	299
Claus M. Ringel, <i>Socle conditions for <math>QF - 1</math> rings</i> .....	309
Richard Rochberg, <i>Linear maps of the disk algebra</i> .....	337
Roy W. Ryden, <i>Groups of arithmetic functions under Dirichlet convolution</i> .....	355
Michael J. Sharpe, <i>A class of operators on excessive functions</i> .....	361
Erling Stormer, <i>Automorphisms and equivalence in von Neumann algebras</i> .....	371
Philip C. Tonne, <i>Matrix representations for linear transformations on series analytic in the unit disc</i> .....	385