

Pacific Journal of Mathematics

**TOPOLOGICAL GROUPS WHOSE UNDERLYING SPACES ARE
SEPARABLE FRÉCHET MANIFOLDS**

JAMES EDGAR KEESLING

TOPOLOGICAL GROUPS WHOSE UNDERLYING SPACES ARE SEPARABLE FRÉCHET MANIFOLDS

JAMES KEESLING

Let G be a topological group and let G^* denote the space of all Lebesgue measurable functions from the unit interval $[0, 1]$ into G with the topology of convergence in measure. With this topology and with pointwise multiplication as the group operation, G^* is a topological group. If G is separable and has a complete metric and has more than one point, then Bessaga and Pełczyński have shown that G^* is homeomorphic to l_2 , separable infinite-dimensional Hilbert space. This fact is used in this paper to show the existence of separable Fréchet manifolds which are topological groups and which have certain algebraic and topological properties.

Introduction. Let X be a metric space and X^* be the space of all Lebesgue measurable functions from $[0, 1]$ into X with the topology of convergence in measure. The space X^* is a contractible, locally contractible metric space in general. If X has a complete metric, then X^* also has a complete metric. Bessaga and Pełczyński [3] have shown that if X is $\{0, 1\}$, $[0, 1]$, or \mathbb{R} , then X^* is homeomorphic to l_2 , separable infinite-dimensional Hilbert space. More recently they have shown [4] that X^* is homeomorphic to l_2 whenever X is a complete separable metric space with more than one point. This fact is used in this paper to show the existence of topological groups which are connected separable Fréchet manifolds and which have certain algebraic and topological properties. The construction we use is a modification of one used by Hartman and Mycielski [8]. Primarily we study two types of topological groups: periodic topological groups and those topological groups whose underlying group structure is isomorphic to the additive reals. The results are contrasted with the structure theory of locally compact topological groups [10 and 16] and with the theory of infinite-dimensional Lie groups [14].

In the last section of the paper it is shown that if π is any abelian group, then π is the fundamental group of a completely metrizable topological group G_π which is a $K(\pi, 1)$. If π is countable, then G_π is a separable Fréchet manifold. If π is any group, then we show the existence of a completely metrizable homogeneous space which is a $K(\pi, 1)$. If π is countable, then the space will be

a separable Fréchet manifold. F. E. A. Johnson [12] has shown that for (finite-dimensional) manifolds if π is a group, then there is a manifold M^n of type $K(\pi, 1)$ if and only if π is countable and has finite cohomological dimension. This contrasts with infinite-dimensional manifolds which are of type $K(\pi, 1)$. Milnor [17] has shown that if π is a countable abelian group and n is any positive integer, then there is a topological group which is a $K(\pi, n)$, but the space is not a separable Fréchet manifold. The author wishes to thank the referee for referring him to the papers of Milnor and Johnson.

NOTATION. If X and Y are topological spaces, then $X \sim Y$ will mean that X and Y are homeomorphic. We let R denote the real numbers with the usual topology. We consider R as a topological group with addition as the operation. The rational numbers are denoted by Q , the integers by Z , and the set $\{0, 1, 2, \dots, n-1\}$ by Z_n . The sets Z and Z_n have the discrete topology and are considered to be groups with the usual addition. Separable infinite-dimensional Hilbert space is denoted by l_2 . A *separable Fréchet manifold* is a separable metric space X such that for each $x \in X$, there is an open set U_x in X containing x such that $U_x \sim l_2$. This terminology comes from the fact that all separable infinite-dimensional Fréchet spaces are homeomorphic. See the Introduction in [1] for a list of references to results which, taken together, prove this. If X is a topological space, then $\pi_n(X) = \pi_n(X, x)$ denotes the n th homotopy group of X with base point x .

We assume the rudiments of measure theory as contained in [18]. We let λ denote Lebesgue measure on $[0, 1]$. If $f: [0, 1] \rightarrow R$ is Lebesgue integrable, then the integral of f is denoted $\int f d\lambda$.

1. **The space of measurable functions.** Let X be a metric space with bounded metric d . Let X^* be the set of all Lebesgue measurable functions from $[0, 1]$ into X . We identify two functions if the set on which they differ has measure zero. Define $\rho_d(f, g) = \int d(f(t), g(t)) d\lambda$. Then ρ_d is a bounded metric on X^* . If $\{f_i\}_{i=1}^\infty$ is a sequence of measurable functions $f_i: [0, 1] \rightarrow X$, then $\{f_i\}_{i=1}^\infty$ converges in measure to the measurable function $f: [0, 1] \rightarrow X$ provided that $\lambda(\{t \in [0, 1]: d(f_i(t), f(t)) \geq \varepsilon\}) = \delta_i \rightarrow 0$. It should be clear that f_i converges in measure to f if and only if $\rho_d(f_i, f) \rightarrow 0$. Thus the topology induced on X^* by ρ_d is the same as that induced by ρ_δ where δ is any other bounded metric inducing the topology of X . The space X^* with the topology induced by ρ_d is called the *space of all Lebesgue measurable functions with the topology of convergence*

in measure.

If X is a topological group or a topological vector space, then X^* is also with the operations induced by the pointwise operations. Certain spaces X^* are important in analysis. The space $\{0, 1\}^*$ is isomorphic to the measure algebra of $[0, 1]$ and R^* is the standard example of a complete separable metric topological vector space which is not locally convex. Recently Bessaga and Pełczyński [3] have shown that $\{0, 1\}^*$, $[0, 1]^*$, and R^* are all homeomorphic to l_2 . More recently they have shown the following [4].

THEOREM 1.1. (Bessaga and Pełczyński): *Let X be any complete separable metric space with more than one point. Then $X^* \sim l_2$.*

This characterizes those spaces X for which X^* is homeomorphic to l_2 . In particular, if X is a countable discrete space with more than one point, then X^* is homeomorphic to l_2 .

As pointed out in [4], Theorem 1.1 together with the remarks above and results in [19] implies that: (1) every separable metric topological group can be imbedded as a subgroup of a topological group homeomorphic to l_2 ; (2) every complete separable metric topological group can be imbedded as a closed subgroup of a topological group homeomorphic to l_2 ; (3) every separable metric topological group which contains at least two points has a free action on l_2 ; and (4) if G is a separable metric topological group with at least two points and F is any closed subset of l_2 , there is a semifree action of G on l_2 with F as its set of fixed points. In [19] (4) was shown for the case that G was compact.

In this paper Theorem 1.1 will be used to give simple examples of topological groups whose underlying spaces are separable Fréchet manifolds and which in addition have certain algebraic and topological properties as well.

2. Periodic topological groups. A group G is said to be *periodic* provided that each element of G has finite order. It is said to be of *period* n provided that each element of G has order at most n and some element has order n . By a very complicated means Markov [15] showed that there exist connected periodic topological groups of period two. Hartman and Mycielski [8] pointed out that $\{0, 1\}^*$ is an easier example of this. They also pointed out that for each positive integer n , there are arcwise connected, locally arcwise connected, metrizable topological groups G_n which are periodic of period n . Here we use Theorem 1.1 to show that if H is any countable periodic group of period n , then there is a topological

group G_H which is a separable Fréchet manifold whose fundamental group is H . It should be pointed out that if G is a locally compact group which is not totally disconnected, then G contains a non-degenerate one-parameter subgroup (see [16, Theorem 1, p. 192]). Consequently, such a group could not be periodic.

THEOREM 2.1. *If G_n is a periodic topological group of period at most n , then for each positive integer k , $\pi_k(G_n)$ is a periodic group of period at most n .*

Proof. Let $\Omega(G_n)$ denote the loop space of G_n with base point $1 \in G_n$ and let $\Omega_0(G_n)$ denote the contractible loops. Let m denote pointwise multiplication of loops in $\Omega(G_n)$. With the compact open topology on $\Omega(G_n)$, m is homotopic to the ordinary loop multiplication in $\Omega(G_n)$ and thus $\pi_1(G_n) \cong \Omega(G_n)/\Omega_0(G_n)$ is periodic of period at most n . For the higher homotopy groups the result follows by induction using the fact that $\pi_{k+1}(G_n) \cong \pi_k(\Omega_0(G_n))$ [11, p. 109] and the fact that $\Omega_0(G_n)$ is a topological group which is periodic of period at most n under the multiplication m .

EXAMPLE 2.2. Let P^3 be real projective 3-space. Then P^3 is a Lie group and $P^3 \times l_2$ is a separable Fréchet manifold which is a topological group. The fundamental group of $P^3 \times l_2$ is just Z_2 which is periodic of period two. However, $P^3 \times l_2$ does not admit a topological group structure which is periodic of period two. This is because $\pi_3(P^3 \times l_2) = \pi_3(P^3) = \pi_3(S^3) = Z$ is not periodic of period two.

THEOREM 2.3. *Let H be any abelian group which is periodic of period n . Then there is a topological group G_H which is periodic of period n which is completely metrizable and locally contractible such that $\pi_1(G_H) \cong H$. If H is finite or countably infinite, then G_H is a separable Fréchet manifold.*

Proof. Let H be the given group and give it the discrete topology. Identify H with the constant functions in H^* . Let $G_H = H^*/H$ which is clearly periodic of period n . Then the natural map $\nu: H^* \rightarrow H^*/H$ is a covering projection with H^* completely metrizable, contractible, and locally contractible. Thus H^*/H is completely metrizable and locally contractible. Since H^* is contractible,

$$\pi_1(H^*/H) \cong \ker \nu = H$$

[6, Theorem 4.5, p. 14]. If H is finite or countably infinite, then H^* is homeomorphic to l_2 by Theorem 1.1 and thus H^*/H is a separable Fréchet manifold.

3. **Topological groups isomorphic to the additive reals.** The additive group of real numbers is characterized as a vector space over the rational numbers Q of cardinality 2^{\aleph_0} . The principal examples of locally compact groups whose underlying group is isomorphic to the additive reals are the reals R and the rational solenoid Σ_ω [7]. In fact, the only locally compact topological groups having this underlying group structure are of the form $R^n \times \prod_{i \in I} \Sigma_\omega$ where n is a nonnegative integer and $2^{\text{card } I} \leq 2^{\aleph_0}$ [13]. Consequently, for any locally compact topological group G with G group isomorphic to R and any positive integer m , $\pi_m(G) = 0$. This contrasts with Theorem 3.5 where we show that a topological group G which is a separable Fréchet manifold with G group isomorphic to R may have nontrivial fundamental group.

Infinite-dimensional topological groups which are homeomorphic to l_2 and whose group structure is the additive reals are abundant. Any separable infinite dimensional Banach space is an example of one. These examples are Lie groups in the sense of Maissen [14]. The Fréchet space R^∞ is an example of one which is not a Lie group in the sense of [14]. Our next theorems show that there may be a much more complicated relationship between the algebraic structure of an infinite-dimensional topological group and its topological structure than these examples would indicate.

THEOREM 3.1. *There is a topological group G whose underlying group is isomorphic to the additive reals with $G \sim l_2$ such that G contains no nontrivial continuous one-parameter subgroups.*

Proof. Let H be any countable vector space over the rationals Q and let H have the discrete topology with metric $d(x, y) = 1$ for all $x \neq y$. Let $G = H^*$. Then $G \sim l_2$ by Theorem 1.1. Clearly G is a topological group whose underlying group structure is a vector space over Q with cardinality 2^{\aleph_0} . Now suppose that $\phi: R \rightarrow G$ is a continuous one-parameter subgroup in G . Suppose that $\phi(x) = g \neq 0$ in G . Now $x/n \rightarrow 0$. Therefore $\phi(x/n) = g/n \rightarrow 0$. Now $g: [0, 1] \rightarrow H$ has the property that $\rho_a(g, 0) = \int d(g(t), 0) d\lambda = \int d((g(t))/n, 0) d\lambda = \rho_a(g/n, 0)$ for all n . Thus $\rho_a(g/n, 0) \rightarrow 0$ and $g/n \rightarrow 0$, a contradiction. Therefore ϕ must be identically zero and G has no nontrivial continuous one-parameter subgroups.

Theorem 3.1 should be contrasted with the fact that every locally compact topological group which is not zero-dimensional has a nondegenerate one-parameter subgroup as mentioned earlier. This is also true of infinite-dimensional Lie groups [14].

Let G be a group and n a positive integer. Then G is said to

be n -divisible provided that for each element x in G , there is a y in G with $y^n = x$. A torsion free abelian group is a vector space over \mathbb{Q} if and only if it is n -divisible for all n .

The following proposition was first proved by Banach [2].

PROPOSITION (Banach) 3.2. *If G and H are separable and completely metrizable topological groups and if $\phi: G \rightarrow H$ is a surjective continuous homomorphism, then ϕ is open.*

COROLLARY 3.3. *Let G be a separable, completely metrizable topological group which is abelian, torsion free, and n -divisible. Then the mapping which takes x to x/n is a topological group isomorphism.*

EXAMPLE 4.6. The following example shows that completeness is necessary in Corollary 3.3 even though a special homomorphism is involved. Let \mathbb{Q} be the rationals and let $N_n = \{k2^n: k \in \mathbb{Z}\}$. Then $\{N_n\}_{n=1}^\infty$ forms a basis for the neighborhoods of 0 in \mathbb{Q} in a topology τ on \mathbb{Q} which makes \mathbb{Q} into a metrizable topological group [10, p.18]. Now in the topology τ , the sequence $x_i = 2^i$ has the property that $x_i \rightarrow 0$. However, $\{x_i/3\}_{i=1}^\infty$ does not converge to 0 in \mathbb{Q} in the topology τ . Thus the map taking x to $x/3$ is not continuous even though \mathbb{Q} is 3-divisible.

THEOREM 3.4. *If G is a separable completely metrizable topological group whose underlying group is a vector space over \mathbb{Q} , then $\pi_n(G)$ is a vector space over \mathbb{Q} for every positive integer n .*

Proof. Let G be written additively. Let $\Omega(G)$ and $\Omega_0(G)$ be the loop space of G with base point 0 and the contractible loops, respectively. Let m be pointwise multiplication on $\Omega(G)$ and $\Omega_0(G)$. Then, as pointed out in the proof of Theorem 2.1, $\Omega(G)$ and $\Omega_0(G)$ are topological groups. Clearly $\Omega(G)$ is a vector space over \mathbb{Q} . Let n be a positive integer. Now $\Omega(G)$ is completely metrizable and separable. Therefore the map taking g to g/n is a topological group isomorphism. Thus division by n takes $\Omega_0(G)$ onto itself. Thus $\Omega_0(G)$ is n -divisible for all n and a vector space over \mathbb{Q} . Thus the factor group $\Omega(G)/\Omega_0(G)$ is also a vector space over \mathbb{Q} . Since $\pi_1(G) \cong \Omega(G)/\Omega_0(G)$ the theorem is proved for the case $n = 1$. Now $\Omega(\Omega(G)) = \Omega(\Omega_0(G))$ since each loop in $\Omega(G)$ with base point 0 must be contained in the arc-component of 0 in $\Omega(G)$ which is just $\Omega_0(G)$. Now for $n \geq 1$, $\pi_{n+1}(G) \cong \pi_n(\Omega(\Omega_0(G))) \cong \pi_n(\Omega(\Omega(G)))$ and the theorem follows by induction.

THEOREM 3.5. *Let H be any abelian group which is a countable vector space over the rationals. Then there is a topological group G which is a connected separable Fréchet manifold such that $\pi_1(G) = H$ such that the underlying group structure of G is isomorphic to the additive reals.*

Proof. Let H have the discrete topology. Identify H with the constant functions in H^* . It is clear that H^* is a vector space over \mathbb{Q} with subspace H . Therefore H^*/H is a vector space over \mathbb{Q} . Thus $G = H^*/H$ is the required group.

Similarly we have the next result.

THEOREM 3.6. *Let H be any group which is a vector space over \mathbb{Q} . Then there is a completely metrizable topological group G which is locally contractible such that the underlying group of G is a vector space over \mathbb{Q} and such that $\pi_1(G) \cong H$.*

4. Topological groups which are of type $K(\pi, 1)$. Theorem 1.1 also gives us a simple way to construct separable Fréchet manifolds which are $K(\pi, 1)$'s for any countable group π .

THEOREM 4.1. *Let π be any abelian group. Then there is a completely metrizable, locally contractible, connected topological group G_π of type $K(\pi, 1)$. If π is countable, then G_π is a separable Fréchet manifold.*

Proof. Let π be given the discrete topology and identify π with the constant functions in π^* . Then $G_\pi = \pi^*/\pi$ is the required topological group.

A homogeneous space is a topological space X such that $X \sim G/H$ where G is a topological group, H is a closed subgroup, and G/H is the decomposition space of right (or left) cosets of H with quotient topology. If π is any group, then π^*/π is a homogeneous space which is a $K(\pi, 1)$.

THEOREM 4.2. *Let π be any group. Then there is a completely metrizable, locally contractible, homogeneous space M_π which is a $K(\pi, 1)$. If π is finite or countably infinite, then M_π is a separable Fréchet manifold.*

NOTE 4.3. If X is any separable Fréchet manifold which is a $K(\pi, 1)$, then all homotopy groups of the covering space \tilde{X} of X

vanish. Since \tilde{X} is also a separable Fréchet manifold, it has the homotopy type of a CW -complex. Thus \tilde{X} is contractible. By [9], \tilde{X} must be homeomorphic to l_2 . In the examples given above this is clearly the case.

The last results in the paper have to do with the higher homotopy groups of topological groups which are separable Fréchet manifolds.

THEOREM 4.4. *Let G be a topological group which is a separable Fréchet manifold and suppose that H is any finite or countable abelian group and that N is any nonnegative integer. Then there is a topological group G' which is a separable Fréchet manifold having the property that $\pi_1(G') = H$ and $\pi_n(G') = \pi_{n+N}(G)$ for all $n \geq 2$.*

Proof. First of all suppose that $H = \{0\}$. Define

$$\Omega_n(G) = \Omega(\Omega_{n-1}(G))$$

where $\Omega_1(G) = \Omega(G)$. Clearly $\Omega_n(G)$ is a topological group for all n . We will now show that $\Omega_n(G)$ is a separable Fréchet manifold for all n . If $G \sim l_2$, then this is true since $\Omega(G)$ can be given the structure of an infinite-dimensional Banach space. If G is an arbitrary separable Fréchet manifold, then G can be imbedded as an open subset of l_2 by [9]. Thus $\Omega(G)$ can be considered to be an open subset of $\Omega(l_2) \sim l_2$ and thus is a separable Fréchet manifold. By induction, $\Omega_n(G)$ is a separable Fréchet manifold. To complete the proof for $H = \{0\}$ simply let G' be the universal covering group of the identity component of $\Omega_N(G)$ if $N \geq 1$ and the universal covering group of the identity component of G if $N = 0$.

If $H \neq \{0\}$, then let G_1 be the topological group just constructed for the case $H = \{0\}$ and let G_2 be the topological group G_H in Theorem 4.1. Then let $G' = G_1 \times G_2$.

REMARK 4.5. In this remark we state some facts about Lie groups in order to contrast them with Theorems 4.1 and 4.4. First of all, if L is a connected Lie group, then L is homeomorphic to $R^n \times K$ where K is a compact Lie group [16, pp. 188–189]. Thus every Lie group has the homotopy type of a compact Lie group and thus of a compact manifold. Therefore its fundamental group must be finitely presented. By Theorem 4.1, the fundamental group of a topological group which is a separable Fréchet manifold need not be finitely presented.

Another fact about Lie groups is that $\pi_2(L) = 0$. Theorem 4.4 shows that this need not be true for topological groups which are

separable Fréchet manifolds. To show this let L be any compact simply connected Lie group. Then $\pi_k(L) \neq 0$ for some $k \geq 3$. Then let $H = L \times l_2$ which is a topological group and a separable Fréchet manifold. Then let G' be the group given by Theorem 4.4 for $n = k - 2$ and H arbitrary. Then $\pi_2(G') = \pi_k(L) \neq 0$.

REFERENCES

1. R. D. Anderson and R. H. Bing, *A complete elementary proof that Hilbert space is homeomorphic to the countable infinite product of lines*, Bull. Amer. Math. Soc. **74** (1968), 771-792.
2. S. Banach, *Über metrische Gruppen*, Studia Math. **3** (1931), 101-113.
3. C. Bessaga and A. Pelczyński, *The space of Lebesgue measurable functions on the interval $[0, 1]$ is homeomorphic to the countable infinite product of lines*, Math. Scand. **27** (1970), 132-140.
4. ———, (to appear).
5. J. Dugundji, *Topology*, Allyn and Bacon, Boston, 1965.
6. M. Greenberg, *Lectures on Algebraic Topology*, Benjamin, New York, 1967.
7. P. R. Halmos, *Comment on the real line*, Bull. Amer. Math. Soc., **50** (1944), 877-878.
8. S. Hartman and J. Mycielski, *On the imbedding of topological groups into connected topological groups*, Colloq. Math., **5** (1958), 167-169.
9. D. W. Henderson, *Infinite-dimensional manifolds are open subsets of Hilbert space*, Topology **9** (1970), 155-161.
10. E. Hewitt and K. Ross, *Abstract Harmonic Analysis I*, Springer-Verlag, New York, 1963.
11. S. T. Hu, *Homotopy*, Academic Press, New York, 1959.
12. F. E. A. Johnson, *Manifolds of homotopy type $K(\pi, 1)$. I*, Proc. Camb. Phil. Soc., **70** (1971), 387-393.
13. G. Mackey, *A remark on locally compact abelian groups*, Bull. Amer. Math. Soc. **52** (1946), 940-944.
14. B. Maissen, *Lie-Gruppen mit Banachräumen als Parameterräume*, Acta Math. **108** (1962), 229-270.
15. A. A. Markov, *On the existence of periodic connected topological groups*, (Russian) Izv. Akad. Nauk SSSR Ser. Math., **8** (1944), 225-232 = Amer. Math. Soc. Trans. (1) **8** (1962), 186-194.
16. D. Montgomery and L. Zippin, *Topological Transformation Groups*, John Wiley, New York, 1955.
17. J. Milnor, *The geometric realization of a semi-simplicial complex*, Ann. of Math., **65** (1957), 357-362.
18. H. L. Royden, *Real Analysis*, Macmillan, New York, 1963.
19. J. E. West, *Fixed point sets of transformation groups of infinite product spaces*, Proc. Amer. Math. Soc. **21** (1969), 575-582.

Received August 10, 1971 and in revised form March 20, 1972. This research was supported by National Science Foundation Grant GP-24616.

UNIVERSITY OF FLORIDA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

C. R. HOBBY
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Jimmy T. Arnold, <i>Power series rings over Prüfer domains</i>	1
Maynard G. Arsove, <i>On the behavior of Pincherle basis functions</i>	13
Jan William Auer, <i>Fiber integration in smooth bundles</i>	33
George Bachman, Edward Beckenstein and Lawrence Narici, <i>Function algebras over valued fields</i>	45
Gerald A. Beer, <i>The index of convexity and the visibility function</i>	59
James Robert Boone, <i>A note on mesocompact and sequentially mesocompact spaces</i>	69
Selwyn Ross Caradus, <i>Semiclosed operators</i>	75
John H. E. Cohn, <i>Two primary factor inequalities</i>	81
Mani Gagrath and Somashekhar Amrith Naimpally, <i>Proximity approach to semi-metric and developable spaces</i>	93
John Grant, <i>Automorphisms definable by formulas</i>	107
Walter Kurt Hayman, <i>Differential inequalities and local valency</i>	117
Wolfgang H. Heil, <i>Testing 3-manifolds for projective planes</i>	139
Melvin Hochster and Louis Jackson Ratliff, Jr., <i>Five theorems on Macaulay rings</i>	147
Thomas Benton Hoover, <i>Operator algebras with reducing invariant subspaces</i>	173
James Edgar Keesling, <i>Topological groups whose underlying spaces are separable Fréchet manifolds</i>	181
Frank Leroy Knowles, <i>Idempotents in the boundary of a Lie group</i>	191
George Edward Lang, <i>The evaluation map and EHP sequences</i>	201
Everette Lee May, Jr., <i>Localizing the spectrum</i>	211
Frank Belsley Miles, <i>Existence of special K-sets in certain locally compact abelian groups</i>	219
Susan Montgomery, <i>A generalization of a theorem of Jacobson. II</i>	233
T. S. Motzkin and J. L. Walsh, <i>Equilibrium of inverse-distance forces in three-dimensions</i>	241
Arunava Mukherjea and Nicolas A. Tserpes, <i>Invariant measures and the converse of Haar's theorem on semitopological semigroups</i>	251
James Waring Noonan, <i>On close-to-convex functions of order β</i>	263
Donald Steven Passman, <i>The Jacobian of a growth transformation</i>	281
Dean Blackburn Priest, <i>A mean Stieltjes type integral</i>	291
Joe Bill Rhodes, <i>Decomposition of semilattices with applications to topological lattices</i>	299
Claus M. Ringel, <i>Socle conditions for $QF - 1$ rings</i>	309
Richard Rochberg, <i>Linear maps of the disk algebra</i>	337
Roy W. Ryden, <i>Groups of arithmetic functions under Dirichlet convolution</i>	355
Michael J. Sharpe, <i>A class of operators on excessive functions</i>	361
Erling Stormer, <i>Automorphisms and equivalence in von Neumann algebras</i>	371
Philip C. Tonne, <i>Matrix representations for linear transformations on series analytic in the unit disc</i>	385