

Pacific Journal of Mathematics

ON 2-TRANSITIVE COLLINEATION GROUPS OF FINITE PROJECTIVE SPACES

WILLIAM M. KANTOR

ON 2-TRANSITIVE COLLINEATION GROUPS OF FINITE PROJECTIVE SPACES

WILLIAM M. KANTOR

In 1961, A. Wagner proposed the problem of determining all the subgroups of $P\Gamma L(n, q)$ which are 2-transitive on the points of the projective space $PG(n-1, q)$, where $n \geq 3$. The only known groups with this property are: those containing $PSL(n, q)$, and subgroups of $PSL(4, 2)$ isomorphic to A_7 . It seems unlikely that there are others. Wagner proved that this is the case when $n \leq 5$. In unpublished work, D. G. Higman handled the cases $n = 6, 7$. We will inch up to $n \leq 9$. Our result is that nothing surprising happens. The same is true if $n = r^\alpha + 1$ for a prime divisor r of $q - 1$.

One of Wagner's results is that it suffices to only consider subgroups of $PGL(n, q)$. Once this is done, it becomes simpler to view the problem as one concerning linear groups: find all those subgroups G of $GL(n, q)$ which are 2-transitive on the 1-spaces of the underlying vector space V . Our approach is based primarily on three facts. (1) Wagner showed that the global stabilizer in G of any 3-space of V induces at least $SL(3, q)$ on that 3-space. (2) Unless $G \geq SL(n, q)$ or $n = 4$, $q = 2$, and $G \approx A_7$, no nontrivial element of G can fix every 1-space of some n -2-space of V . (3) $G \leq SL(n, q)$ if $|G|$ is divisible by a prime which is a primitive divisor of $q^m - 1$ for a suitable $m \leq n - 2$.

Wagner's results are in [10]. Higman's result, and the case $n = 2^\alpha + 1$ and q odd, are mentioned by Dembowski [1], p. 39. The result mentioned above in (2) is an easy consequence of results of Wagner. The idea used in (3) is due to Perin [8] and, independently, to G. Hare and E. Shult.

I am indebted to G. Seitz for several helpful remarks.

2. Notation and preliminaries. As already mentioned, we will be dealing with linear groups. Let V be an n -dimensional vector space over $GF(q)$. We write $GL(V) = GL(n, q)$ and $SL(V) = SL(n, q)$. It will be convenient to regard everything as taking place in the relative holomorphic $V \cdot GL(V)$. For any subgroups K, L of this semi-direct product we can then consider the normalizer $N_L(K)$ and centralizer $C_L(K)$. If $L \leq GL(V)$ and W is an L -invariant subspace of V , we write $L^W = L/C_L(W)$ for the subgroup of $GL(W)$ induced by L . $C_L(V/W)$ and $L^{V/W}$ are defined similarly. For any group G , as usual G' is its commutator subgroup, $Z(G)$ its center, and $\Phi(G)$ its Frattini subgroup.

A group A is said to be *involved* in a group B if $A \approx C/D$ with $B \cong C \supseteq D$.

(2.1) If $R \leq GL(V)$ has prime power order and $(|R|, q) = 1$, then $V = C_V(R) \oplus [V, R]$, where $[V, R] = \langle v - vr \mid v \in V, r \in R \rangle$ is $N_{GL(V)}(R)$ -invariant.

Proof. [3], p. 177.

(2.2) Let $R \leq GL(V)$ have prime power order with $(|R|, q) = 1$. Let W be an R -invariant subspace. Then $\dim C_V(R) = \dim C_W(R) + \dim C_{V/W}(R)$.

Proof. [3], p. 187, or (2.1).

Both (2.1) and (2.2) will be used frequently, generally without reference.

A *primitive divisor* of $q^k - 1$ is a prime r satisfying $r \mid q^k - 1$ but $r \nmid q^i - 1$ for $1 \leq i < k$; clearly $k \mid r - 1$.

(2.3) (i) If q is a prime power and $k \geq 2$, then $q^k - 1$ has a primitive divisor unless $k = 6, q = 2$, or $k = 2$ and q is a Mersenne prime.

(ii) Let r be a primitive divisor of $q^k - 1$, and let R be an r -subgroup of $GL(V)$ for a $GF(q)$ -space V . If $C_V(R) = 0$, then k divides $\dim V$.

Proof. (i) [12].

(ii) This is clear if $|R| \leq r$. Let $|R| > r$, and let $R_i \leq Z(R)$ have order r . Then $V = W \oplus [V, R_i]$, where $W = C_V(R_i)$ is R -invariant and $C_W(R) = 0$. By induction, k divides $\dim W$ and $\dim [V, R_i]$.

(2.4) Suppose $\dim V = \alpha m$, r is a primitive divisor of $q^m - 1$, and $R \leq GL(V)$ is an r -group such that $C_V(R) = 0$. Then:

(i) Each noncyclic composition factor of $N = N_{GL(V)}(R)$ is involved in $PSL(\alpha, q^m)$; and

(ii) If R is abelian, each noncyclic composition factor of $N/C_N(R)$ is involved in the symmetric group S_α .

Proof. Write $V = W_1 \oplus \cdots \oplus W_\beta$, with each W_i a sum of R -isomorphic irreducible R -spaces and no two W_i having isomorphic irreducible R -subspaces. Set $R_i = C_R(W_i)$. Then $Z(R/R_i)$ is cyclic and nontrivial; let Z_i be its subgroup of order r . By (2.3 ii), $\dim W_i = me_i$ for some e_i . Consequently, $\beta \leq \alpha$ and $e_i \leq \alpha$.

N permutes the W_i . Let K be the kernel of this permutation representation. Then N/K is involved in $S_\beta \leq S_\alpha$, and hence in $GL(\alpha, q^m)$.

Set $K_i = N_{GL(W_i)}(Z_i)$. Then K is contained in $K_1 \times \cdots \times K_\beta$. Moreover, K_i is contained in $\Gamma L(e_i, q^m)$. This proves (i).

Now assume that R is abelian. Then R/R_i is a cyclic group normalized by K . Since $\cap R_i = 1$, it follows that $K/C_K(R)$ is abelian. Since N/K is involved in $S_{\alpha'}$ this proves (ii).

(2.5) Let q be odd, and let $H \leq GL(V)$. Suppose that $H \geq A \neq 1$, where A is an elementary abelian 2-group. Set

$$m = \min \{ |H : N_H(B)| \mid B < A, |A : B| = 2 \}.$$

Then $m \leq \dim V$.

Proof. (G. Seitz.) Let \bar{V} be an H -irreducible section of V on which A acts nontrivially. Let \bar{H} and \bar{A} be the groups induced by H and A . Then $\bar{A} \neq 1$, and the corresponding $\bar{m} \geq m$. We may thus assume that $V = \bar{V}$ is H -irreducible. By Clifford's Theorem ([3], p. 70), $V = V_1 \oplus \cdots \oplus V_t$ with the V_i direct sums of A -isomorphic irreducible A -spaces, no two V_i having a common irreducible constituent. Here A induces a group of order 2 on each V_i , while H is transitive on $\{V_1, \dots, V_t\}$. Thus, $\{C_A(V_i) \mid i = 1, \dots, t\}$ is an orbit of H of subgroups of A of index 2. Consequently, $t \geq m$, so $\dim V \geq m$.

(2.6) Let L be a finite group and $K \triangleleft L$ with L/K simple. Suppose L has no proper subgroup L_0 for which $L_0/L_0 \cap K \approx L/K$. Then:

- (i) K is nilpotent; and
- (ii) Each proper normal subgroup of L is contained in K .

Proof. (i) Let S be a Sylow subgroup of K . By the Frattini argument, $L = KN_L(S)$, so our conditions on L imply that $L = N_L(S)$.

(ii) Let $M \trianglelefteq L$ and $M \not\leq K$. Since $1 \neq MK/K \trianglelefteq L/K$, $MK = L$ and hence $M = L$.

(2.7) Let $d > e \geq 2$ and $t \geq 1$. Then $PSL(d, q)$ is not involved in $PSL(e, q^t)$.

Proof. If p is the prime dividing q , then p -Sylow subgroups of $PSL(d, q)$ and $PSL(e, q^t)$ have nilpotence class $d - 1$ and $e - 1$, respectively.

We now come to our main technical lemma.

(2.8) Let $q = p^e$, where p is a prime, and $m = \dim V$. Suppose either $m = 3, 4$, or 5 , or $m = 6$ and $p = 2$. Let $L \leq GL(V)$ and $H, K \triangleleft L$, where $H \leq K$, $L/K \approx PSL(3, q)$, and $L/H \approx PSL(3, q)$ or $SL(3, q)$. Assume that L has no proper subgroup L_0 for which $L_0/L_0 \cap K \approx PSL(3, q)$. Finally, assume: (#) If $1 \neq h \in H$ and $p \nmid |h|$, then $\dim C_V(h) \leq m - 3$.

Then there are L -invariant subspaces X, Y with $X > Y$ such that the following hold.

- (a) $K = P \times C$ with P a p -group, $|C| = (3, q - 1)$, and $H = P$ or K .
- (b) $L/P \approx SL(3, q)$.
- (c) $P^{V/X}, P^{X/Y}$ and P^Y are all 1.
- (d) $\dim X/Y = 3$ and $L^{X/Y} = SL(X/Y)$.
- (e) If $m \leq 5$ and $q \neq 2$, then $L^{V/X}$ and L^Y are 1. Moreover, some element g of order p in the center of a p -Sylow subgroup of L satisfies $\dim C_V(g) \geq m - 2$, and even $\dim C_V(g) = m - 1$ if $P = 1$.

Proof. Everything is obvious if $m = 3$, so assume $m > 3$. We will proceed by a series of steps.

(i) Clearly $L = L'$. We can apply (2.6) to L . In particular, K is nilpotent.

(ii) Suppose that there are L -invariant subspaces V_1, V_2 with $V_1 \geq V_2$ and $\dim V_1/V_2 \leq 2$. We claim that L centralizes V_1/V_2 . For, $C_L(V_1/V_2) \trianglelefteq L$, and since L^{V_1/V_2} does not have $PSL(3, q)$ as a homomorphic image, (2.6) implies that $C_L(V_1/V_2) = L$.

(iii) Next, suppose that there are L -invariant subspaces X, Y with $X > Y$, $\dim X/Y = 3$ and $L^{X/Y} \neq 1$. We claim that (a)–(e) hold.

Arguing as in (ii) we find that $L^{X/Y} = SL(X/Y)$, while $L^{V/X}$ and L^Y are both 1 or $SL(3, q)$. Write $K = P \times C$ with P a p -group and C a p' -group. C induces a group of order 1 or $(3, q - 1)$ on $V/X, X/Y$, and Y . By (2.2), (a) holds unless $|C| = 9$ and $m = 6$. However, in this case $C \leq Z(L)$, so $L/P = (L/P)'$ is a central extension of $SL(3, q)$ by a group of order 9, and this is impossible [2].

Thus, (a), (b), (c), and (d) hold.

Now let $m \leq 5$. Then $\dim V/X$ and $\dim Y$ are ≤ 2 , so $L^{V/X}$ and L^Y are 1 by (ii). If $P \neq 1$ then, by (c), each $g \neq 1$ in P satisfies $\dim C_V(g) \geq m - 2$.

Suppose $P = 1$, so $L \approx SL(3, q)$. By results of Higman [4], §5, if $q \neq 2$ then there is an L -invariant 3-space T , and each element of L inducing a transvection on T is a transvection of V . This proves (e).

(iv) From now on we assume that m and L are chosen with m minimal such that (2.8) is false. Then $m > 3$.

L is irreducible on V . For otherwise, there is an L -invariant subspace W with $V > W > 0$.

Then $L^W \neq 1$ and $L^{V/W} \neq 1$. For suppose, say, that $L^{V/W} = 1$. Consider L^W, K^W , and H^W . By (2.2), (#) is inherited by L^W . Also, if $L_0 \leq L$ and $L_0^W/L_0^W \cap K^W \approx PSL(3, q)$ then $L_0 K/K \approx L_0/L_0 \cap K$ has $PSL(3, q)$ as a homomorphic image, so that $L_0 K = L$ and hence $L_0 = L$.

Consequently, L^w satisfies the hypotheses of (2.8). Then we can find subspaces X and Y of W such that (iii) applies, whereas (2.8) is assumed false. Thus, $L^w \neq 1$ and $L^{v/w} \neq 1$.

By (ii) we must have $m = 6$ and $\dim W = 3$. Then (iii) again applies, and this is again impossible.

(v) By (iv) and the nilpotence of K , $(|K|, q) = 1$.

K is not central in L . For suppose $K \leq Z(L)$. Since $L = L'$, L is a homomorphic image of the covering group of $PSL(3, q)$. Then L is $PSL(3, q)$ or $SL(3, q)$ (see, e.g., [2]).

On the other hand, L has an irreducible $GF(q)$ -representation of degree m , where $4 \leq m \leq 6$ and q is even if $m = 6$. No such representation exists by [7] and [9].

(vi) Let r be a prime and R_1 an r -Sylow subgroup of K such that $R_1 \not\leq Z(L)$. Set $R = R_1 \cap H$. Then $R \not\leq Z(L)$ and $R \triangleleft L$.

Let A be a characteristic elementary abelian subgroup of R . By (#), $|A| \leq r^{m-3}$.

We claim that $A \leq Z(L)$. For otherwise, L has a nontrivial $GF(r)$ -representation of degree $\leq m - 3 \leq 3$. By (2.6 ii), $PSL(3, q)$ is involved in $GL(3, r)$. Thus, $q = 2$ and $r \neq 3$. Since A is a non-cyclic elementary abelian subgroup of $GL(6, 2)$, $|A| = 7^2$. Then L acts transitively on $A - \{1\}$. However, not all elements of $A - \{1\}$ are conjugate in $GL(6, 2)$.

Thus, $A \leq Z(L)$. In (iv), $|A| = r$. In particular, $Z(R)$ is cyclic.

(vii) Suppose $r \nmid q - 1$. By (vi), $R \leq GL(6, q)$ is nonabelian, so $r = 3|q + 1$ and $m = 6$. Moreover, $R \triangleright B$ with $|R:B| = 3$ and B abelian. By (vi) we can find $B_1 \neq B$ with $R \triangleright B_1$, $|R:B_1| = 3$, and B_1 abelian. Then $B \cap B_1 \leq Z(R)$ and $|R/Z(R)| \leq 9$. Consequently, L centralizes $Z(R)$, $R/Z(R)$, and hence also R , which is not the case.

Thus, $r|q - 1$. In (iv), $A \leq L \cap Z(GL(V)) \leq Z(SL(V))$, so $r|(q - 1, m)$.

There are now just three possibilities: $m = 4, r = 2$; $m = 5, r = 5$; and $m = 6, r = 3$.

(viii) Let $m = 4, r = 2$. By (vii), $-1 \in R$. There is an involution $t \neq -1$ in R . Either $\dim C_v(t) \geq 2$ or $\dim C_v(-t) \geq 2$. This contradicts (#).

(ix) Let $m = 5, r = 5$. A 5-Sylow subgroup of $GL(5, q)$ has a normal abelian subgroup of index 5 (the "diagonal subgroup"). Thus, we can find $B \leq R$ with B abelian and $|R:B| = 1$ or 5 . By (vi), $|R:B|$ is 5 and B is not characteristic in R . Let $B_1 < R$, $B_1 \neq B$, satisfy the same conditions as B . Then $B_1 \cap B \leq Z(R)$ and $|R:Z(R)| \leq 5^2$. By (vi), $Z(R)$ is cyclic, so L centralizes $Z(R)$, $R/Z(R)$, and hence also R , which is not the case.

(x) Finally, let $m = 6, r = 3$, and $q = 2^i$. Here $3|q - 1$. On the one hand, $L/C_L(R/\Phi(R))$ can be regarded as a subgroup of $GL(e, 3)$ for some e ; on the other hand, using (2.6) and $(|K|, q) = 1$, we

find that this group has an elementary abelian 2-subgroup of order q^2 whose normalizer is transitive on the nontrivial elements. By (2.5), $e \geq q^2 - 1$. However, a 3-Sylow subgroup of $SL(6, q)$ has order $\leq 3(q - 1)^6$. Thus, $3^{q^2-1} \leq 3^e \leq |R| < 3q^6$, and since $q \geq 4$ this is ridiculous.

This contradiction completes the proof of (2.8).

3. Wagner's results and some corollaries. Let V be n -dimensional over $GF(q)$, $n \geq 3$, and let $G \leq GL(V)$ be 2-transitive on 1-spaces.

(3.1) For each 3-space T , $N_G(T)^T \geq SL(T)$.

Proof. Wagner [10], p. 417.

(3.2) If $n \leq 5$ then $G \geq SL(V)$, unless $n = 4$, $q = 2$, and $G \approx A_7$.

Proof. Wagner [10], p.422.

(3.3) For each n -1-space W , $N_G(W)$ is 2-transitive on the 1-spaces of V not in W .

Proof. [6], p. 6.

(3.4) If G has an element $g \neq 1$ such that $\dim C_V(g) \geq n - 2$, then $G \geq SL(V)$ or $n = 4$, $q = 2$, and $G \approx A_7$.

Proof. We may assume that $|g|$ is prime and $n > 5$. Since $\dim [V, g] \leq 2$ and g centralizes $V/[V, g]$, there is a 3-space $T > [V, g]$ such that $g^T \neq 1$. Then $1 \neq C_G(V/T)^T \leq N_G(T)^T$. By (3.1), $C_G(V/T)^T \geq SL(T)$. Choose $g' \in C_G(V/T)$ with $|g'| \mid q + 1$ and $\dim C_T(g') = 1$. Then $\dim C_V(g') = n - 2$.

We may thus assume that $(|g|, q) = 1$. Since $g^{[V, g]} \neq 1$, as before $C_G(V/T)^T \geq SL(T)$ for each 3-space $T > [V, g]$. By the 2-transitivity of G , this holds for every 3-space of V .

Choose $m \leq n$ maximal with respect to $C_G(V/U)^U \geq SL(U)$ for all m -spaces U . Suppose $m < n$. By Wagner [10], p. 420, $m \leq n - 2$. Take any subspace W of dimension $m + 1$ or $m + 2$. For each m -space $U < W$, $C_G(V/U)$ fixes W and centralizes V/W , while $C_G(V/U)^U \geq SL(U)$. By Wagner [10], p. 420, and (3.2), $C_G(V/W)^W \geq SL(W)$ for each $m + 1$ -space W . This contradicts the maximality of m .

(3.5) Let s be a prime and S an s -group maximal with respect to $\dim C_V(S) \geq 3$. Then $N_G(S)$ is 2-transitive on the 1-spaces of $C_V(S)$.

Proof. Take any 3-space $T \leq C_V(S)$. Then S is Sylow in $C_G(T)$. By the Frattini argument and (3.1), $(N_G(S) \cap N_G(T))^T = N_G(T)^T \geq SL(T)$. Our assertion follows immediately.

4. The case $n = r^\alpha + 1$. There is one very easy case of our problem.

(4.1) **THEOREM.** *Let r be a prime divisor of $q - 1$, and let $\alpha \geq 1$. Then every collineation group of $PG(r^\alpha, q)$ which is 2-transitive on points contains $PSL(r^\alpha + 1, q)$.*

We first prove:

(4.2) Let r be a prime divisor of $q - 1$, and let $\alpha \geq 1$. Let V be an r^α -dimensional vector space over $GF(q)$. If $G \leq GL(V)$ is transitive on $V - \{0\}$, then $r \parallel G \cap Z(GL(V))$.

Proof. Let r^β be the largest power of r dividing $q^d - 1$, where $d = r^\alpha$. Then q is not an r^β th power, so $r \parallel G \cap GL(V)$.

Let R be an r -Sylow subgroup of G . By [11], p. 6, each orbit of R on $V - \{0\}$ has length divisible by r^β .

R fixes no nontrivial proper subspace of V . For, if it did we would have $r^\beta \mid q^m - 1$ with $1 \leq m < d$. Set $e = (d, m)$. Then $r^\beta \mid q^e - 1$. However, as d/e is a power of r , $(q^d - 1)/(q^e - 1)$ is divisible by r , and this contradicts the definition of r^β .

Let $x \in Z(R) \cap GL(V)$ have order r . Since $r \mid q - 1$, x can be diagonalized. By the preceding paragraph, x is a scalar transformation, that is, $x \in Z(GL(V))$.

(4.3) Let r be a prime divisor of $q - 1$, and let $\alpha \geq 1$. Then a collineation group of the affine space $AG(r^\alpha, q)$ which is 2-transitive on points contains the translation group.

Proof. (4.2).

Now (4.1) follows immediately from (3.3) and (4.3).

5. Primes dividing $|G|$. We will consider the following situation in the remainder of this paper.

Let V be an n -dimensional $GF(q)$ -space, $n \geq 6$, and G be a subgroup of $GL(V)$, 2-transitive on 1-spaces, such that $G \not\cong SL(V)$. We may clearly assume that $G > Z = Z(GL(V))$.

In this section let s be a prime dividing $(|G|, q^m - 1)$, $1 < m \leq n - 2$, such that s is a primitive divisor of $q^m - 1$. (5.1) is essentially due to Perin [8] and, independently, to E. Shult and G. Hare.

(5.1) If $m = n - 2$ then $q = 2$ and n is even.

(5.2) Suppose that $n = \alpha m + \beta$, $\alpha < \beta \leq m + 2$, and an element of order s centralizes some 3-space X . Then, for some n' satisfying $5 < n' < n$ and $n' \equiv n \pmod{m}$, there is a subgroup of $GL(n', q)$, not containing $SL(n', q)$, which is 2-transitive on the points of $PG(n' - 1, q)$.

Clearly (5.2) has an inductive flavor. Since the proofs are similar, we will only prove the second of the above results.

Proof of (5.2). Choose $S \leq C_G(X)$ as in (3.5). Set $W = C_V(S)$, $W^* = [V, S]$, and $N = N_G(S)$. Then $V = W \oplus W^*$, $C_{W^*}(S) = 0$, and N^W is 2-transitive on 1-spaces.

Set $n' = \dim W$, so $n' \geq 3$. By (2.3 ii), since $\beta \leq m + 2$ we have $\dim W^* = \gamma m$ with $\gamma \leq \alpha$. Then $n' = n - \gamma m \geq n - \alpha m = \beta > \alpha \geq \gamma$.

We must show that $n' > 5$ and $N^W \not\cong SL(W)$. Deny this. Then either $N^W \geq SL(W)$ or $n' = 4$, $q = 2$, and $N^W \approx A_7$. In particular, the commutator subgroup N'^W contains a nontrivial element centralizing an n' -2-space.

In this situation, $C_{N'}(W^*)^W \leq Z(GL(W))$. For otherwise, $C_{N'}(W^*)^W \leq N'^W$ implies that $C_{N'}(W^*)^W = N'^W$. Then $C_{N'}(W^*)$ has a nontrivial element g centralizing an n' -2-space of W . Hence, $\dim C_V(g) \geq n - 2$, which contradicts (3.4).

It follows that N'^{W^*} has $PSL(n', q)$ as a homomorphic image, unless $n' = 4$ and $q = 2$, in which case A_7 may be a homomorphic image.

Since $C_{W^*}(S) = 0$, we can apply (2.4): each noncyclic composition factor of N'^{W^*} is involved in $PSL(\gamma, q^m)$. Since $n' > \gamma$, by (2.7) $PSL(n', q)$ cannot be such a composition factor. Thus, $n' = 4$, $q = 2$, $\gamma \leq 3$, and A_7 is a composition factor of N'^{W^*} . However, A_7 is not involved in $PSL(3, 2^m)$. This is a contradiction.

REMARK. It is useful to note that the above proof holds under slightly weaker hypotheses: s is a primitive divisor of $q^m - 1$, $S \neq 1$ is an s -subgroup of G with $W = C_V(S)$ of dimension $n' \geq 3$, $(n - n')/m < n'$, and $N_G(S)^W$ is 2-transitive on 1-spaces.

We conclude this section with two miscellaneous results.

(5.3) Assume that G has a cyclic subgroup H of order $q^n - 1$ containing an r -Sylow subgroup of G for some prime r dividing $q^2 + q + 1$. Then $q = 2$ and n is even.

Proof. Suppose $q \neq 2$ or $q = 2$ and n is odd. By (2.3), H is transitive on $V - \{0\}$. Thus, H is transitive on the 3-spaces fixed by its subgroup R of order r .

On the other hand, by (3.1) each 3-space is fixed by a conjugate of R . Thus, G is transitive on 3-spaces, and this contradicts Perin [8] or (5.1) since $n \geq 6$.

(5.4) Assume that G has a cyclic subgroup of order $q^{n-1} - 1$ fixing some $n - 1$ -space W and transitive on $W - \{0\}$. Then $N_G(W)$ is 2-transitive on the 1-spaces of W , $q = 2$, and n is even.

Proof. We may assume that $G - Z$ has no element fixing all 1-spaces in W . By [6], Lemma 7.3, $N_G(W)$ is 2-transitive on the 1-spaces of W . The result now follows from (2.3) and (5.1).

6. The case $n \leq 9$. Let n, V, G , and Z be as in §5, so $G \not\cong SL(V)$. Let p be the prime dividing q .

Assume that $6 \leq n \leq 9$.

(6.1) $n \neq 6$.

Proof. Suppose $n = 6$. If $q = 2$ then $q^5 - 1$ is a prime. By (5.4), the stabilizer of a 5-space W is 2-transitive on $W - \{0\}$. By (3.2) and (3.4), $G \geq SL(V)$, which is not the case.

Thus, $q > 2$. Let r be a prime dividing $q - 1$.

Suppose that there is 3-space T for which $N_G(T) - Z$ contains an element inducing a scalar transformation of order r on T . Using Z , we find that $r \mid |C_G(T)|$. Let R be an r -Sylow subgroup of $C_G(T)$. By (3.4), $T = C_V(R)$. By (3.5), $N_G(R)^r \geq SL(T)$. Also, $N_G(R)$ normalizes the 3-space $[V, R]$. An element of order p in the center of a p -Sylow subgroup of $N_G(R)$ centralizes 2-spaces of both $C_V(R)$ and $[V, R]$, and hence centralizes a 4-space of $V = C_V(R) \oplus [V, R]$. This contradicts (3.4). Thus, no element of $G - Z$ of order r has an eigenspace of dimension > 2 .

Now take any 3-space T , and write $T = X \oplus Y$ with $\dim X = 2$ and $\dim Y = 1$. Set $F = N_G(X) \cap N_G(Y)$, so $F^X = GL(X)$. Take $R \leq F$ of order r with $R \not\leq Z$ and $R^T \leq Z(F^T)$. By the Frattini argument, $N_F(R)^X = GL(X)$. Let $E \leq N_F(R)$ be minimal with respect to $E^X = SL(X)$.

Since R is diagonalizable and each of its eigenspaces has dimension 1 or 2, we can write $V = X \oplus W_1 \oplus W_2$ with $W_1 > Y$, $\dim W_i = 2$, and W_i invariant under $N_G(R)$. If $q \neq 3$, $E = E'$ centralizes W_1 , so an element of E of order p centralizes a 4-space, which contradicts (3.4). If $q = 3$, R cannot have more than two eigenspaces as $|R| = 2$, which is again a contradiction.

(6.2) q is even.

Proof. Assume that q is odd. There is an involution $t \in G - Z$. Since $n \geq 6$, $\dim C_V(t)$ or $\dim C_V(-t)$ is ≥ 3 . Let S be a 2-group in G maximal with respect to $\dim C_V(S) \geq 3$. Set $W = C_V(S)$ and $W^* = [V, S]$, so $V = W \oplus W^*$. Set $M = N_G(S)$. By (3.5), M^W is 2-transitive on 1-spaces. Since $M > Z$ and all involutions in M^W centralize at most a 2-space (by the maximality of S), $\dim W \leq 4$. Consequently, by (3.2), $M^W \geq SL(W)$.

By (4.1) and (6.1), $n = 7$ or 8 , so $\dim W^* \leq 5$.

We claim that $C_M(W^*)^w \leq Z(GL(W))$. For otherwise, $C_M(W^*)^w \trianglelefteq M^w$ yields $C_M(W^*)^w \geq SL(W)$. Then $C_M(W^*)$ contains a nontrivial transvection of V , which contradicts (3.4).

Thus, $C_M(W^*)$ is cyclic and M'^{w*} has $PSL(W)$ as a homomorphic image.

Suppose that $\dim W = 4$. Then $\dim W^* = 3$ or 4 . Use of M'^{w*} yields $\dim W^* = 4$ and $M'^{w*} \geq SL(W^*)$. If $g \neq 1$ is in the center of a p -Sylow subgroup of M' then g^w and g^{w*} are transvections, and this contradicts (3.4).

Thus, $\dim W = 3$. Let $L \leq M$ be minimal with respect to having $PSL(3, q)$ as a homomorphic image. Let $H = C_L(W) \leq K \triangleleft L$ with $L/K \approx PSL(3, q)$. Then (2.8) applies to W^* , L^{w*} , K^{w*} , and H^{w*} .

Choose $g \in L$ so that g^{w*} is as in (2.8 e). If $g \in H = C_L(W)$, then $\dim C_V(g) \geq n - 2$. If $H^{w*} = 1$ then $H = 1$, and both g^w and g^{w*} are transvections, so once again $\dim C_V(g) \geq n - 2$. In either case we have contradicted (3.4).

$$(6.3) \quad n \neq 7, 8.$$

Proof. Let $n = 7$ or 8 . Fix a prime $r | q + 1$.

Take any 3-space T . By (3.1), $N_G(T)^r \geq SL(T)$. Also, $N_G(T)$ acts on V/T . By (3.4), $C_G(V/T)^r \leq Z(GL(T))$ (since otherwise, $C_G(V/T)$ would have an element of order r), so $C_G(V/T)$ is solvable. Thus, $N_G(T)^{r/r}$ has $PSL(3, q)$ as a composition factor. By (2.8), there is an r -group $R \neq 1$ in $N_G(T)$ such that $\dim C_{V/T}(R) \geq 2$, and then $\dim C_V(R) \geq 3$.

This contradicts (5.2) with $n = 2 \cdot 2 + 3$ or $2 \cdot 2 + 4$.

$$(6.4) \quad \text{If } n = 9 \text{ then } q = 2 \text{ or } 4.$$

Proof. Suppose $n = 9$ and $q > 4$ is even.

(i) By (5.2) with $n = 2 \cdot 3 + 3$, no nontrivial element of order dividing $(q^2 + q + 1)/(q + 1, 3)$ can centralize a 1-space.

(ii) Let T be any 3-space. Let $L \leq N_G(T)$ be minimal with respect to having $PSL(3, q)$ as a homomorphic image. By (3.4), $C_G(V/T)^r \leq Z(GL(T))$, so (2.8) applies to $L^{r/r}$. Consequently, by (i) there is a 6-space $Y > T$ such that $L^{r/r} = SL(Y/T)$ and $L^{r/r} = SL(V/Y)$.

(iii) Let s be a prime dividing $q + 1$. By (ii), there is an element of order s centralizing a 3-space.

Let S be an s -group maximal with respect to $\dim C_V(S) \geq 3$. By (3.5), $N_G(S)$ is 2-transitive on the 1-spaces of $C_V(S)$. In view of (i), it follows from (3.2), (6.1), and (6.3) that $\dim C_V(S) = 3$.

Let $T = C_V(S)$ in (ii), and choose $L \leq N_G(S)$ there. By (i) and the proof of (2.4), $(LS)^{[r, s]}$ acts as a subgroup of $\Gamma L(3, q^2)$, with S inducing scalar transformations.

(iv) Since $q > 4$, by (2.3 i) there is a prime $r \neq 3$ dividing $q - 1$. Moreover, if $q \neq 16$ we can choose $r \neq 5$.

We claim that some element of order r centralizes a 4-space. For, since $r \neq 3$, in (iii) we can find $g \in L - Z$ of order r such that $g^{[V, S]}$ has an eigenspace of dimension ≥ 4 . Consequently, some element of $\langle g, Z \rangle$ of order r centralizes a 4-space.

(v) Let R be an r -group maximal with respect to $\dim C_v(R) \geq 3$; by (iv), $R \neq 1$. Set $T = C_v(R)$ and $T^* = [V, R]$. By (3.5), $N_G(R)^T$ is 2-transitive on 1-spaces, so $\dim T = 3$ by (i). We can thus choose $L \leq N_G(R)$ in (ii).

We claim that LR centralizes R and that R is diagonalizable. Certainly $(LR)^{T^*} \leq GL(T^*)$. Suppose $r > 5$. Then an r -Sylow subgroup of $GL(6, q)$ is diagonalizable, and hence abelian. By (2.4 ii) (with $m = 1, \alpha = 6$), each composition factor of $L/C_L(R)$ is involved in S_6 . By (2.6 ii), $L = C_L(R)$, so $R \leq Z(LR)$.

Consider the case $r = 5, q = 16$. Suppose $L > C_L(R)$. Then L acts nontrivially on $R/\Phi(R)$, where $|R/\Phi(R)| \leq 5^7$. By (2.6 ii), $16 + 1$ divides $|GL(7, 5)|$, which is not the case.

Thus, L centralizes R . There is an s -group $S_0 < L$ such that $\dim C_{T^*}(S_0) = 2$. Since R normalizes $C_{T^*}(S_0)$ and $[T^*, S_0]$, it follows that R is again diagonalizable. Thus, $R \leq Z(LR)$.

(vi) T^* is the direct sum of R -invariant subspaces, each invariant under LR . By (ii) and (v), there are 3-spaces X and X' such that $T^* = X \oplus X'$, R^X and $R^{X'}$ consist of scalar transformations, $L^X = SL(X')$, and $L^{X'} = SL(X)$.

Consequently, for each $h \in R$, $\dim C_v(h) = 3, 6$, or 9 .

(vii) By (iv), there is an r -group $R_1 \neq 1$ maximal with respect to $\dim C_v(R_1) \geq 4$. By (vi), $W = C_v(R_1)$ has dimension 6. Set $M = N_G(R_1)$.

Take any 3-space $T < W$. Let $R \geq R_1$ be an r -Sylow subgroup of $C_G(T)$. If $R = R_1$ then $N_M(T)^T \geq SL(T)$ by the Frattini argument. If $R > R_1$ then the choice of R_1 implies that $C_v(R) = T$, and hence that R is an r -group maximal with respect to $\dim C_v(R) \geq 3$; by (v), $C_G(R)^T \geq SL(T)$, so again $N_M(T)^T \geq SL(T)$.

Consequently, M^W is 2-transitive on 1-spaces. Then $(q^6 - 1)/(q - 1)$ divides $|G|$, and this contradicts (5.2).

(6.5) If $n = 9$ then $q \neq 4$.

Proof. Suppose $n = 9$ and $q = 4$. We will try to imitate the proof of (6.4) using $r = 3$. Steps (i) and (ii) of that proof still hold.

We begin by showing the existence of $x \in G$ of order 3 such that $x^y = x^{-1}$ for some 2-element y . Take T and L as in (ii). Then we can find $x, y \in L$ with $|x| = 3, y$ a 2-element, and $x^y = x^{-1}a, a \in C_L(T)$.

By (2.8), $C_L(T) = P \times C$ with P a 2-group and $|C| = 1$ or 3. Then $\langle x \rangle$ is Sylow in $\langle x, y \rangle P$. By the Frattini argument, some element of $\langle y \rangle P$ inverts $\langle x \rangle$, and we may assume this is y .

We next claim that some element of order 3 centralizes a 4-space. For, assume that this is false, and choose x, y as above. Since $q = 4$, x is diagonalizable and has at most 3 eigenspaces. However, no element of $\langle x, Z \rangle - \{1\}$ centralizes a 4-space, so $C_V(x) = T$ is a 3-space and x has two other 3-dimensional eigenspaces T_1, T_2 . Moreover, by our assumption, $C_G(T)$ has a cyclic 3-Sylow subgroup. Thus, by the Frattini argument, $N_G(\langle x \rangle)^T \geq SL(T)$, so $C_G(x)^T \geq SL(T)$. Since $|GL(T):SL(T)| = 3$, $y^T \in SL(T)$, so we can find $c \in C_G(X)$ such that $c^{-1}y \in C_G(T)$. Clearly $c^{-1}y$ inverts x , so there is an involution $t \in \langle c^{-1}y \rangle$. Here, t centralizes T and centralizes 2-spaces of each T_i , so $\dim C_V(t) \geq 7$. This contradicts (3.4), and proves our claim.

Now define R, T, T^* , and L as in (v). We will be able to obtain a contradiction precisely as in (vi) and (vii) if we can show that $R \leq Z(LR)$ and R is diagonalizable.

By (2.6), $L \triangleright K$ with $L/K \approx PSL(3, 4)$ and K nilpotent. By (2.2) and (2.8), $K = P \times C$ with $|C| = 3$ or 9 and P a 2-group; moreover, there is an L -invariant 3-space $X < T^*$ such that $L^X = SL(X)$, $L^{T^*/X} = SL(T^*/X)$, and P centralizes T, X , and T^*/X . By (3.4), no nontrivial element of P centralizes a 4-space of T^* . Consequently, P is elementary abelian of order $\leq 4^3$. Thus, if $P \not\leq Z(L)$ then $PSL(3, 4)$ is isomorphic to a subgroup of $GL(6, 2)$, which is not the case ([7], [9]). Thus, $K \leq Z(L)$.

Now suppose that L acts nontrivially on R , and hence on $R/\Phi(R)$. Since $R \leq GL(6, 4)$, $|R/\Phi(R)| \leq 3^6 \cdot 3^2$. Thus, $PSL(3, 4)$ or $SL(3, 4)$ is isomorphic to a subgroup of $GL(8, 3)$. Then $GL(8, 3)$ has an elementary abelian subgroup of order 4^2 whose normalizer is transitive on the nontrivial elements. By (2.5), this is impossible.

Consequently, $L \leq C_G(R)$. An element of L of order 5 centralizes 1-spaces of X and T^*/X . It follows that T^* is the sum of R -invariant 2-spaces. Thus, R is diagonalizable and $R \leq Z(LR)$. This completes the proof of (6.5).

Last, and least:

(6.6) If $n = 9$ then $q \neq 2$.

Proof. Suppose $n = 9$ and $q = 2$. Using (5.1) and (5.2) we find that $|G| = 2^\alpha \cdot 3^\beta \cdot 5 \cdot 7 \cdot 17 \cdot 73$ for some α, β .

Let S be a 73-Sylow subgroup of G . By (5.3), $|C_G(S)| = 73$. Thus, $|N_G(S)| = 3^\gamma \cdot 73$ with $\gamma \leq 2$.

By Sylow's theorem, $2^\alpha \cdot 3^{\beta-\gamma} \cdot 5 \cdot 7 \cdot 17 \equiv 1 \pmod{73}$. A little arithmetic shows that this is impossible.

In view of (3.2) and the results of this section, we can now state:

THEOREM 6.7. *Let H be a subgroup of $P\Gamma L(n, q)$ which is 2-transitive on the points of $PG(n-1, q)$. If $3 \leq n \leq 9$, then $H \geq PSL(n, q)$ or $n = 4, q = 2$, and $H \approx A_7$.*

REFERENCES

1. P. Dembowski, *Finite Geometries*, Springer, Berlin-Heidelberg-New York, 1968.
2. W. Feit, *The current situation in the theory of finite simple groups*, Proc. Int. Cong. Math. Nice, 1970, Vol. 1, 55-93.
3. D. Gorenstein, *Finite Groups*, Harper and Row, New York, 1968.
4. G. Hare, *Thesis*, Southern Illinois University, 1970.
5. D. G. Higman, *Flag-transitive collineation groups of finite projective spaces*, Illinois J. Math., **6** (1962), 434-446.
6. W. M. Kantor, *2-Transitive symmetric designs*, Trans. Amer. Math. Soc., **146** (1969), 1-28.
7. C. Mark, *Thesis*, University of Toronto, 1939.
8. D. Perin, *On collineation groups of finite projective spaces*, Math. Z., **126** (1972), 135-142.
9. R. Steinberg, *Representations of algebraic groups*, Nagoya Math. J., **22** (1963), 33-56.
10. A. Wagner, *On collineation groups of finite projective spaces*, I. Math. Z., **76** (1961), 411-426.
11. H. Wielandt, *Finite Permutation Groups*, Academic Press, New York, 1964.
12. K. Zsigmondy, *Zur Theorie der Potenzreste*, Monatsh. f. Math. u. Phys., **3** (1892), 265-284.

Received November 6, 1971 and in revised form October 13, 1972. This research was supported in part by NSF Grant GP28420.

UNIVERSITY OF OREGON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Jan Aarts and David John Lutzer, <i>Pseudo-completeness and the product of Baire spaces</i>	1
Gordon Owen Berg, <i>Metric characterizations of Euclidean spaces</i>	11
Ajit Kaur Chilana, <i>The space of bounded sequences with the mixed topology</i>	29
Philip Throop Church and James Timourian, <i>Differentiable open maps of $(p+1)$-manifold to p-manifold</i>	35
P. D. T. A. Elliott, <i>On additive functions whose limiting distributions possess a finite mean and variance</i>	47
M. Solveig Espelie, <i>Multiplicative and extreme positive operators</i>	57
Jacques A. Ferland, <i>Domains of negativity and application to generalized convexity on a real topological vector space</i>	67
Michael Benton Freeman and Reese Harvey, <i>A compact set that is locally holomorphically convex but not holomorphically convex</i>	77
Roe William Goodman, <i>Positive-definite distributions and intertwining operators</i>	83
Elliot Charles Gootman, <i>The type of some C^* and W^*-algebras associated with transformation groups</i>	93
David Charles Haddad, <i>Angular limits of locally finitely valent holomorphic functions</i>	107
William Buhmann Johnson, <i>On quasi-complements</i>	113
William M. Kantor, <i>On 2-transitive collineation groups of finite projective spaces</i>	119
Joachim Lambek and Gerhard O. Michler, <i>Completions and classical localizations of right Noetherian rings</i>	133
Kenneth Lamar Lange, <i>Borel sets of probability measures</i>	141
David Lowell Lovelady, <i>Product integrals for an ordinary differential equation in a Banach space</i>	163
Jorge Martinez, <i>A hom-functor for lattice-ordered groups</i>	169
W. K. Mason, <i>Weakly almost periodic homeomorphisms of the two sphere</i>	185
Anthony G. Mucci, <i>Limits for martingale-like sequences</i>	197
Eugene Michael Norris, <i>Relationally induced semigroups</i>	203
Arthur E. Olson, <i>A comparison of c-density and k-density</i>	209
Donald Steven Passman, <i>On the semisimplicity of group rings of linear groups. II</i>	215
Charles Radin, <i>Ergodicity in von Neumann algebras</i>	235
P. Rosenthal, <i>On the singularities of the function generated by the Bergman operator of the second kind</i>	241
Arthur Argyle Sagle and J. R. Schumi, <i>Multiplications on homogeneous spaces, nonassociative algebras and connections</i>	247
Leo Sario and Cecilia Wang, <i>Existence of Dirichlet finite biharmonic functions on the Poincaré 3-ball</i>	267
Ramachandran Subramanian, <i>On a generalization of martingales due to Blake</i>	275
Bui An Ton, <i>On strongly nonlinear elliptic variational inequalities</i>	279
Seth Warner, <i>A topological characterization of complete, discretely valued fields</i>	293
Chi Song Wong, <i>Common fixed points of two mappings</i>	299