

Pacific Journal of Mathematics

**FINITELY GENERATED F -ALGEBRAS WITH APPLICATIONS
TO STEIN MANIFOLDS**

E. R. HEAL AND MICHAEL WINDHAM

FINITELY GENERATED F -ALGEBRAS WITH APPLICATIONS TO STEIN MANIFOLDS

E. R. HEAL AND M. P. WINDHAM

Let A be a commutative n -generated F -algebra with identity, M_A the spectrum of A . Conditions are given which insure that M_A can be mapped homeomorphically into C^n and that A is topologically isomorphic to the algebra of all holomorphic functions on a polynomially convex open subset of C^n . A characterization is given of those Stein manifolds X for which $\mathcal{O}(X)$ is finitely generated.

1. Introduction. We seek conditions on a commutative F -algebra A with identity that force A to be topologically isomorphic to an algebra of all functions analytic on a polynomially convex open subset of C^n . Several conditions for the case A is singly-generated have been given by Arens in [1], Brooks in [4], and Carpenter [6]. Birtel in [3] characterized the algebra of all entire functions on C .

Our results are for the case A is n -generated. We first show that, under various conditions, the spectrum of A can be mapped homeomorphically into C^n (see Theorems 3.4 and 3.6). Theorem 3.7 shows that if the spectrum of A is a topological $2n$ -manifold without boundary, then A is an algebra of all analytic functions on an open, polynomially convex subset of C^n . In Theorem 3.8 we obtain an F -algebra characterization of the algebra of all entire functions on C^n .

In § 4, we apply the results of § 3 to Stein manifolds. We obtain a characterization of those Stein manifolds whose algebras of analytic functions are finitely generated.

2. Preliminaries. An F -algebra A is a complete topological algebra over the complex numbers in which the topology is given by a countable family $\{\|\cdot\|_k : k = 1, 2, \dots\}$ of algebraic seminorms. It can be assumed that the seminorms are increasing; i.e., $\|f\|_k \leq \|f\|_{k+1}$ for each positive integer k and each f in A . All algebras in this paper are assumed to be commutative and contain units.

For each seminorm $\|\cdot\|_k$, we can obtain a Banach algebra A_k by letting A_k be the completion of the quotient algebra $A/(\ker \|\cdot\|_k)$ with respect to the norm induced on $A/(\ker \|\cdot\|_k)$ by $\|\cdot\|_k$. It follows that A is the inverse limit of the Banach algebras A_k , denoted $A = \lim_{\text{inv}} A_k$. For details of this construction the reader is referred to [10].

The spectrum of A , denoted by M_A , is the space of all continuous homomorphisms of A onto the complex numbers with the Gelfand

topology (the relative weak * topology). For each positive integer k , the spectrum M_k of A_k is a compact Hausdorff space and is imbedded homeomorphically in M_A by the dual map π_k^* of π_k , where π_k is the natural projection of A into A_k . It follows that $M_A = \bigcup_{k=1}^{\infty} M_k$ and any compact set in M_A is a subset of M_k for some k . Hence, M_A is a hemicompact Hausdorff space.

For each f in A , we define $\hat{f}: M_A \rightarrow C$ by $\hat{f}(\phi) = \phi(f)$ for each $\phi \in M_A$. This mapping is called the Gelfand transform of f , and the mapping $f \rightarrow \hat{f}$ is a homomorphism of A onto a separating subalgebra \hat{A} of $C(M_A)$. We define seminorms on the algebra \hat{A} by letting $|\hat{f}|_k = \sup \{|\hat{f}(x)|; x \in M_k\}$ for each f in A . The topology on \hat{A} defined by these seminorms is the compact-open topology.

If each of the seminorms for an F -algebra A satisfies $\|f^2\|_k = \|f\|_k^2$ for each f in A , then A is called a uniform algebra. If A is a uniform algebra, then $\|f\|_k = |\hat{f}|_k$ for each f and each k . Hence, for uniform algebras, the mapping $f \rightarrow \hat{f}$ is a topological isomorphism of A onto a complete subalgebra of $C(M_A)$. Therefore, a uniform F -algebra is a complete, hence closed, subalgebra of $C(X)$ where X is a hemicompact Hausdorff space.

The spectrum M_A also satisfies "convexity" conditions which have been particularly prominent in the study of algebras of analytic functions on complex manifolds. In particular if E is compact in M_A then the \hat{A} -hull of E is

$$\hat{E} = \{\phi \in M_A: |\hat{f}(\phi)| \leq \|\hat{f}\|_E \text{ for all } \hat{f} \in \hat{A}\}.$$

If $\hat{E} = E$ then E is said to be \hat{A} -convex. Since each M_k is the spectrum of the Banach algebra A_k it follows easily that M_k is \hat{A} -convex. Furthermore, this implies that \hat{E} is compact in M_A for any compact set E ; in this case M_A is said to be \hat{A} -convex. For a discussion of convexity in function algebras see Rickart [12].

For a hemicompact complex analytic manifold X , the algebra $\mathcal{O}(X)$ of analytic functions is a uniform F -algebra (compact-open topology). It is possible that the spectrum of $\mathcal{O}(X)$ is also a complex manifold which is $\mathcal{O}(X)$ convex (more commonly called holomorphically convex); such a manifold is called a Stein manifold. A particular example is C^n itself. In this case, since analytic polynomials are dense in $\mathcal{O}(C^n)$, the phrase polynomial convexity is used rather than holomorphic convexity. We will say that an arbitrary subset of C^n is polynomially convex if it is the hemicompact union of polynomially convex compact subsets of C^n .

3. Finitely generated F -algebras. An F -algebra is n -generated provided there exist elements f_1, \dots, f_n in A such that A is the closure of the polynomials in f_1, \dots, f_n . In order that a family $\{f_1, \dots, f_n\}$

generate A it is necessary and sufficient that for each k the family $\{\pi_k f_1, \dots, \pi_k f_n\}$ generate A_k .

In this section we will give conditions on the spectrum of A which will insure that an n -generated algebra A is topologically isomorphic to the algebra of all functions holomorphic on some polynomially convex open subset U of C^n (denoted $\mathcal{O}(U)$). We will first consider the problem of mapping the spectrum of A homeomorphically into C^n .

DEFINITION 3.1. Suppose A is an n -generated F -algebra with spectrum M_A . A map $F: M_A \rightarrow C^n$ is called a *spectrum map* provided there exist generators f_1, \dots, f_n for A such that $F(\phi) = (\hat{f}_1(\phi), \dots, \hat{f}_n(\phi))$ for each ϕ in M_A .

It is easy to show that any spectrum map is one-to-one and continuous. In [5] examples are given where (i) one spectrum map is a homeomorphism (onto its image), while another is not, and (ii) no spectrum map is a homeomorphism. Since M_k is the spectrum of a finitely generated Banach algebra, $F|_{M_k}$ is a homeomorphism and $F(M_k)$ is a polynomially convex compact set in C^n (see [9]). The following theorem, which is Theorem 1.3 of [5] for the case A is an F -algebra, gives a necessary and sufficient condition in order that a spectrum map be a homeomorphism.

THEOREM 3.2. Suppose A is an n -generated F -algebra with spectrum M_A and $M_A = \bigcup_{k=1}^{\infty} M_k$. A spectrum map $F: M_A \rightarrow C^n$ is a homeomorphism if and only if for each compact $E \subseteq F(M_A)$ there exists a positive integer k such that $E \subseteq F(M_k)$.

COROLLARY 3.3. If F is a homeomorphism then $F(M_A)$ is polynomially convex.

Proof. Since $F(M_A) = \bigcup_{k=1}^{\infty} F(M_k)$, F is a homeomorphism, and $F(M_k)$ is polynomially convex, it follows that $F(M_A)$ is the hemi-compact union of polynomially convex compact subsets of C^n ; hence, $F(M_A)$ is polynomially convex.

We will now give a sufficient condition in order that a spectrum map be a homeomorphism and have a closed image in C^n .

THEOREM 3.4. Suppose A is an n -generated F -algebra with spectrum M_A and $F: M_A \rightarrow C^n$ is a spectrum map defined by generators f_1, \dots, f_n . If for each $c > 0$ the set $K_c = \{\phi \in M_A: |\phi(f_i)| \leq c \text{ for } i = 1, \dots, n\}$ is compact in M_A , then F is a homeomorphism and $F(M_A)$ is closed in C^n .

Proof. Let E be a compact subset of $F(M_A)$. Then there exists

$c > 0$ such that $F^{-1}(E) \subseteq K_c$. Since F is continuous and K_c is compact, $F^{-1}(E)$ is compact; hence, $E \subseteq F(M_k)$ for some k . Theorem 3.2 now implies that F is a homeomorphism.

Suppose λ is a limit point of $F(M_A)$. Then there exists a sequence $\{F(\phi_n)\}$ converging to λ . There exists $c > 0$ such that $\{\phi_1, \phi_2, \dots\}$ is a subset of K_c . Hence, some subsequence $\{\phi_{k_n}\}$ converges to $\phi \in K_c$. Since F is continuous, $F(\phi) = \lambda$ and $\lambda \in F(M_A)$. Thus $F(M_A)$ is closed in C^n .

Suppose A is an n -generated uniform F -algebra with spectrum $M_A = \bigcup_{k=1}^{\infty} M_k$ and $F: M_A \rightarrow C^n$ is a spectrum map defined by the generators f_1, \dots, f_n . Let $\text{Hol } F(M_A)$ denote the algebra of all functions h such that h is holomorphic in some open set U_h containing $F(M_A)$. Let the topology on $\text{Hol } F(M_A)$ be generated by the family of seminorms $\{\|\cdot\|_{F(M_k)}: k = 1, 2, \dots\}$. The map $F^*: \text{Hol } F(M_A) \rightarrow \hat{A}$ is defined by $F^*(h) = h \circ F$ for each h in $\text{Hol } F(M_A)$. The function calculus for F -algebras (see [2]) implies that $h \circ F$ is in \hat{A} . The following lemma is immediate.

LEMMA 3.5. *F^* is an isometric isomorphism of $\text{Hol } F(M_A)$ onto a dense subalgebra of \hat{A} .*

THEOREM 3.6. *F is a homeomorphism if and only if the topology on $\text{Hol } F(M_A)$ is equivalent to the compact open topology.*

Proof. If F is a homeomorphism, then any compact subset of $F(M_A)$ is a subset of $F(M_k)$ for some k ; hence the topology on $\text{Hol } F(M_A)$ is the compact open topology.

Suppose that the topology on $\text{Hol } F(M_A)$ is the compact open topology. Let E be a compact subset of $F(M_A)$. If for each k , $E \not\subseteq F(M_k)$ then there exists a sequence $\{\lambda_k\}$ such that $\lambda_k \in E \setminus F(M_k)$. Since $F(M_k)$ is polynomially convex there exists a polynomial P_k of n -variables such that $P_k(\lambda_k) \geq 1$ and $\|P_k\|_{F(M_k)} < 1/k$. Then the sequence $\{P_k\}$ is a sequence in $\text{Hol } F(M_A)$ which converges to 0. Since the topology on $\text{Hol } F(M_A)$ is equivalent to the compact open topology the sequence $\{P_k\}$ converges uniformly to 0 on the set E . But $P_k(\lambda_k) \geq 1$ for each k , hence $\{P_k\}$ cannot converge uniformly to 0 on E . This is a contradiction, therefore, it must be that $E \subseteq F(M_k)$ for some k . Consequently, by Theorem 3.2, F is a homeomorphism.

We will now give an F -algebra characterization of the algebra of all functions holomorphic on a polynomially convex open subset of C^n .

THEOREM 3.7. *An n -generated, uniform F -algebra is topologically*

isomorphic to the algebra of all functions holomorphic on a polynomially convex open subset of C^n if and only if the spectrum of A is a topological $2n$ -dimensional manifold (without boundary).

Proof. If A is topologically isomorphic to $\mathcal{O}(U)$ where U is a polynomially convex open subset of C^n , then the spectrum of A is homeomorphic to the spectrum of $\mathcal{O}(U)$ which is U . Hence M_A is a topological $2n$ -manifold.

Suppose A is generated by f_1, \dots, f_n and F is the associated spectrum map. Since M_A is a $2n$ -manifold, for each point ϕ of M_A there exists an open set V_ϕ in M_A and a map $h_\phi: V_\phi \rightarrow R^{2n}$ such that $\phi \in V_\phi$, \bar{V}_ϕ is compact, and h_ϕ is a homeomorphism of V_ϕ onto an open set in R^{2n} . Since \bar{V}_ϕ is compact, $F|_{\bar{V}_\phi}$ is a homeomorphism, hence $F|_{V_\phi}$ is a homeomorphism of V_ϕ into C^n . The Brouwer domain invariance theorem (see [7], page 358) implies that $F(V_\phi)$ is an open subset of C^n . It now follows that F is a homeomorphism and that $U = F(M_A)$ is an open subset of C^n . U is polynomially convex since $F(M_k)$ is polynomially convex for each k and U is the hemicompact union of these sets. Since A is a uniform algebra, A is topologically isomorphic to \hat{A} . Lemma 3.5 implies that $\text{Hol } F(M_A)$ is topologically isomorphic to a dense subalgebra of \hat{A} . Since F is a homeomorphism, the topology on $\text{Hol } F(M_A)$ is equivalent to the compact open topology (Theorem 3.6). But $F(M_A)$ is open, so $\text{Hol } F(M_A)$ is topologically isomorphic to $\mathcal{O}(U)$. Since $\mathcal{O}(U)$ is complete, it follows that \hat{A} is topologically isomorphic to $\mathcal{O}(U)$; hence A is topologically isomorphic to $\mathcal{O}(U)$.

THEOREM 3.8. *Suppose A is a uniform F -algebra generated by f_1, \dots, f_n , M_A is a topological $2n$ -manifold, and for each $c > 0$ the set $K_c = \{\phi \in M_A: |\phi(f_i)| \leq c \text{ for } i = 1, \dots, n\}$ is compact in M_A . Then A is topologically isomorphic to $\mathcal{O}(C^n)$.*

Proof. From the proof of Theorem 3.7 we know that $U = F(M_A)$ is open in C^n and A is topologically isomorphic to $\mathcal{O}(U)$. Theorem 3.4 implies that U is closed in C^n . Since C^n is connected, $U = C^n$ and A is therefore topologically isomorphic to $\mathcal{O}(C^n)$.

4. Stein manifolds. If X is a Stein manifold, then $\mathcal{O}(X)$ is a uniform F -algebra with the compact open topology. It follows from the proper imbedding theorem (Hörmander [9], Theorem 5.3.9) that Stein manifolds are just closed submanifolds of C^n . Hence from Rickart [12] Theorem 1.3, it follows that $X = M_{\mathcal{O}(X)}$. So, if $\mathcal{O}(X)$ is n -generated then a spectrum map F can be viewed as an analytic map of X into C^n . In this section we give a more complete description of the image $F(X)$ when F is a homeomorphism and finally give a charac-

terization of those Stein manifolds for which $\mathcal{O}(X)$ is finitely-generated. For properties of analytic maps, see Narasimhan [11] and Herve [8].

THEOREM 4.1. *If X is a Stein manifold and $\mathcal{O}(X)$ is generated by f_1, \dots, f_n then*

- (i) $n \geq \dim X$;
- (ii) *if F , the associated spectrum map, is a homeomorphism then $F(X)$ is a closed polynomially convex submanifold of an open set in C^n and F is bianalytic;*
- (iii) *if $n = \dim X$ then $F(X)$ is a polynomially convex open set in C^n .*

Proof. Part (i) follows from the fact that an injective analytic map can not decrease dimension.

To prove (ii), first note that since F is injective, each $x \in X$ has a fundamental system of neighborhoods whose images have germs at $F(x)$ which are the germs of analytic varieties. But since F is a homeomorphism, the germ of the images of the neighborhood at $F(x)$ are the same as the germ of $F(X)$ at $F(x)$ so $F(X)$ is an analytic subvariety of some open set in C^n . To show that $F(X)$ is a submanifold it suffices to show that $F^{-1}: F(X) \rightarrow X$ is analytic.

$\mathcal{O}(X)$ provides local coordinate system for X , so to show F^{-1} is analytic it suffices to show $g \circ F^{-1} \in \mathcal{O}(F(X))$ for $g \in \mathcal{O}(X)$. But since F is a homeomorphism $\text{Hol } F(X)$ has the compact open topology and $g \circ F^{-1}$ is in the closure of $\text{Hol } F(X)$ in $C(F(X))$. Furthermore, $\text{Hol } F(X) \subset \mathcal{O}(F(X))$ and $\mathcal{O}(F(X))$ is closed in $C(F(X))$ so $g \circ F^{-1} \in \mathcal{O}(X)$.

If X is given as a hemicompact union of $\{M_k\}$ where $M_k = \hat{M}_k$ then $F(M_k)$ is polynomially convex; since F is a homeomorphism $F(X)$ is the hemicompact union of $\{F(M_k)\}$ hence is also polynomially convex.

Part (iii) follows immediately from Theorem 3.7.

THEOREM 4.2. *If X is a Stein manifold then $\mathcal{O}(X)$ is finitely generated if and only if X can be imbedded as a closed polynomially convex submanifold of C^N for some N .*

Proof. The proper imbedding theorem states that there is an analytic, regular, proper map $G: X \rightarrow C^n$. If $\mathcal{O}(X)$ is finitely generated and $F: X \rightarrow C^m$ is any spectrum map then $G \times F$ is a spectrum map. Furthermore, it is regular and proper, hence is an imbedding. That the image is polynomially convex follows from 4.1. (ii).

The converse follows immediately from the following lemma.

LEMMA 4.3. *If Ω is a domain of holomorphy in C^n and X is a polynomially convex submanifold of Ω then $\mathcal{O}(X)$ is generated by $\{z_1|_X, \dots, z_n|_X\}$.*

Proof. Since Ω is a domain of holomorphy $\mathcal{O}(X)$ is precisely the restriction of $\mathcal{O}(\Omega)$ to X . Furthermore, each function in $\mathcal{O}(\Omega)$ can be approximated on polynomially convex compact subsets of X by analytic polynomials (Hörmander [9], Theorem 2.7.7).

REFERENCES

1. R. Arens, *Dense inverse limit rings*, Mich. Math. J., **5** (1958), 169-182.
2. ———, *The analytic-functional calculus in commutative topological algebras*, Pacific J. Math., **11** (1961), 405-429.
3. F. T. Birtel, *Singly-generated Liouville F -algebras*, Mich. Math. J., **11** (1964), 89-94.
4. R. M. Brooks, *On singly-generated locally m -convex algebras*, Duke, Math J., **38** (1970), 529-536.
5. ———, *On the spectrum of finitely-generated locally m -convex algebras*, Studia Math., **29** (1968), 143-150.
6. R. L. Carpenter, *Singly-generated homogeneous F -algebras*, Trans. Amer. Math. Soc., **150** (1970), 457-468.
7. J. Dugundji, *Topology*, Allyn and Bacon, Inc., 1967.
8. M. Herve, *Several Complex Variables, Local Theory*, Oxford University Press, 1963.
9. L. Hörmander, *An Introduction to Complex Analysis in Several Variables*, Van Nostrand, 1966.
10. E. A. Michael, *Locally multiplicatively-convex topological algebras*, Mem. Amer. Math. Soc., **11** (1952).
11. R. Narasimhan, *Introduction to the Theory of Analytic Spaces*, Springer-Verlag, 1966.
12. C. E. Rickart, *Holomorphic convexity for general function algebras*, Canad. J. Math., **20** (1968), 272-290.

Received February 15, 1973.

UTAH STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Robert F. V. Anderson, <i>Laplace transform methods in multivariate spectral theory</i>	339
William George Bade, <i>Two properties of the Sorgenfrey plane</i>	349
John Robert Baxter and Rafael Van Severen Chacon, <i>Functionals on continuous functions</i>	355
Phillip Wayne Bean, <i>Helly and Radon-type theorems in interval convexity spaces</i>	363
James Robert Boone, <i>On k-quotient mappings</i>	369
Ronald P. Brown, <i>Extended prime spots and quadratic forms</i>	379
William Hugh Cornish, <i>Crawley's completion of a conditionally upper continuous lattice</i>	397
Robert S. Cunningham, <i>On finite left localizations</i>	407
Robert Jay Daverman, <i>Approximating polyhedra in codimension one spheres embedded in S^n by tame polyhedra</i>	417
Burton I. Fein, <i>Minimal splitting fields for group representations</i>	427
Peter Fletcher and Robert Allen McCoy, <i>Conditions under which a connected representable space is locally connected</i>	433
Jonathan Samuel Golan, <i>Topologies on the torsion-theoretic spectrum of a noncommutative ring</i>	439
Manfred Gordon and Edward Martin Wilkinson, <i>Determinants of Petrie matrices</i>	451
Alfred Peter Hallstrom, <i>A counterexample to a conjecture on an integral condition for determining peak points (counterexample concerning peak points)</i>	455
E. R. Heal and Michael Windham, <i>Finitely generated F-algebras with applications to Stein manifolds</i>	459
Denton Elwood Hewgill, <i>On the eigenvalues of a second order elliptic operator in an unbounded domain</i>	467
Charles Royal Johnson, <i>The Hadamard product of A and A^*</i>	477
Darrell Conley Kent and Gary Douglas Richardson, <i>Regular completions of Cauchy spaces</i>	483
Alan Greenwell Law and Ann L. McKerracher, <i>Sharpened polynomial approximation</i>	491
Bruce Stephen Lund, <i>Subalgebras of finite codimension in the algebra of analytic functions on a Riemann surface</i>	495
Robert Wilmer Miller, <i>TTF classes and quasi-generators</i>	499
Roberta Mura and Akbar H. Rhemtulla, <i>Solvable groups in which every maximal partial order is isolated</i>	509
Isaac Namioka, <i>Separate continuity and joint continuity</i>	515
Edgar Andrews Rutter, <i>A characterization of $QF - 3$ rings</i>	533
Alan Saleski, <i>Entropy of self-homeomorphisms of statistical pseudo-metric spaces</i>	537
Ryōtarō Satō, <i>An Abel-maximal ergodic theorem for semi-groups</i>	543
H. A. Seid, <i>Cyclic multiplication operators on L_p-spaces</i>	549
H. B. Skerry, <i>On matrix maps of entire sequences</i>	563
John Brendan Sullivan, <i>A proof of the finite generation of invariants of a normal subgroup</i>	571
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, VI</i>	573
Ronson Joseph Warne, <i>Generalized $\omega - \mathcal{L}$-unipotent bisimple semigroups</i>	631
Toshihiko Yamada, <i>On a splitting field of representations of a finite group</i>	649