

Pacific Journal of Mathematics

GENERALIZED ω – \mathcal{L} -UNIPOTENT BISIMPLE SEMIGROUPS

RONSON JOSEPH WARNE

GENERALIZED ω - \mathcal{L} -UNIPOTENT BISIMPLE SEMIGROUPS

R. J. WARNE

Let S be a bisimple semigroup and let $E(S)$ be the set of idempotents of S . If $E(S)$ is an ω -chain of rectangular bands (E_n : $n \in N$, the nonnegative integers) and \mathcal{L} , Green's equivalence relation, is a left congruence on $E(S)$, we term S a generalized ω - \mathcal{L} -unipotent bisimple semigroup. We characterize S in terms of (I, o) , an ω -chain of left zero semigroups (I_k : $k \in N$); $(J, *)$ an ω -chain of right groups (J_k : $k \in N$); a homomorphism $(n, r) \rightarrow \alpha_{(n,r)}$ of C , the bicyclic semigroup, into $\text{End}(I, o)$, the semigroup of endomorphisms of (I, o) (iteration); a homomorphism $(n, r) \rightarrow \beta_{(n,r)}$ of C into $\text{End}(J, *)$; and an (upper) anti-homomorphism $j \rightarrow A_j$ of $(J, *)$ into T_I , the full transformation semigroup on I (A_j is "almost" an endomorphism). In fact, $S \cong ((i, (n, k), j): i \in I_n, j \in J_k, n, k \in N)$ under the multiplication $(i, (n, k), j)(u, (r, s), v) = (i \circ (u A_j \alpha_{(k,n)}), (n + r - \min(k, r), k + s - \min(k, r)), j \beta_{(r,s)}^* v)$ (Theorem 4.1). We then characterize $(J, *)$ as a semi-direct product of an ω -chain of right zero semigroups by an ω -chain of groups. Finally, we specialize Theorem 4.1 to obtain our previous characterization of ω - \mathcal{L} -unipotent bisimple semigroups $S(E(S))$ is an ω -chain of right zero semigroups).

We will use the definitions of Clifford and Preston [1] unless otherwise specified. In particular, \mathcal{R} , \mathcal{L} , \mathcal{H} , and \mathcal{D} will denote Green's equivalence relations on a semigroup S , i.e., $((a, b) \in \mathcal{R}$ if $a \cup aS = b \cup bS$; $(a, b) \in \mathcal{L}$ if $a \cup Sa = b \cup Sb$; $\mathcal{H} = \mathcal{R} \cap \mathcal{L}$; $\mathcal{D} = \mathcal{R} \circ \mathcal{L}$ ($(a, b) \in \mathcal{D}$ if there exists $x \in S$ such that $(a, x) \in \mathcal{R}$ and $(x, b) \in \mathcal{L}$). R_a will denote the \mathcal{R} -class containing $a \in S$. A semigroup consisting of a single \mathcal{D} -class is termed a bisimple semigroup. This bicyclic semigroup is $C = N \times N$ under the multiplication $(n, m)(p, q) = (n + p - \min(m, p), m + q - \min(m, p))$. A semigroup S which is a union of a collection of pairwise disjoint subsemigroups $(S_y: y \in Y)$ where Y is a semilattice and $S_y S_t \subseteq S_{y \wedge t}$ for all $y, t \in Y$ is termed a semilattice Y of the semigroups $(S_y: y \in Y)$.

If $Y = N$ with $n \wedge m = \max(n, m)$, S is termed an ω -chain of the semigroups $(S_n: n \in N)$. A semigroup is termed regular if $a \in aSa$ for every $a \in S$. A rectangular band is the algebraic direct product of a left zero semigroup $U(x, y \in U \text{ implies } xy = x)$ and a right zero semigroup. A right group is a semigroup X such that $a, b \in X$ implies there exists a unique $x \in S$ such that $ax = b$. If V is a subset of a semigroup S , $E(V)$ will always denote the set of idempotents of V .

In [4], we defined a generalized \mathcal{L} -unipotent semigroup to be a

regular semigroup S such that $E(S)$ satisfy the condition: $e, f \in E(S)$ and $ef = e$ imply that $gegfe = ge$ for all $g \in E(S)$. Combining [4, Lemma 1] and a result of Clifford and McLean [2, 1, p. 129, Exercise 1], a regular semigroup S is generalized \mathcal{L} -unipotent if and only if $E(S)$ is a semilattice Y of rectangular bands $(E_y: y \in Y)$ and \mathcal{L} is a left congruence on $E(S)$. Since any bisimple semigroup containing an idempotent is regular by a result of Clifford and Miller [1, Theorem 2.11], the reason for the terminology “generalized ω - \mathcal{L} -unipotent bisimple semigroup” is clear. We introduced the term \mathcal{L} -unipotent in [3] to denote a semigroup in which each \mathcal{L} -class contains precisely one idempotent. By [3, Proposition 5], a semigroup S is \mathcal{L} -unipotent if and only if S is regular and $E(S)$ is a semilattice Y of right zero semigroups $(E_y: y \in Y)$. Hence, the terminology “ ω - \mathcal{L} -unipotent bisimple semigroup” is also clear.

Let S be a generalized ω - \mathcal{L} -unipotent bisimple semigroup. In §1, we define a congruence t on S such that $S/t = C$, the bicyclic semigroup, and give an explicit multiplication for $(E(C))t^{-1}$, the kernel of $t(\ker t)$. In §2, we describe S as an “extension” of $\ker t$ by S/t (the converse of Theorem 4.1). In §3, we prove the direct part of Theorem 4.1. In §4, we state Theorem 4.1 and characterize an ω -chain of right groups as a semi-direct product of an ω -chain of right zero semigroups by an ω -chain of groups (Theorem 4.3). Combining Theorem 4.1, Theorem 4.3, and Clifford’s characterization of semilattices of groups [1; theorem 4.11], we have characterized generalized ω - \mathcal{L} -unipotent bisimple semigroups in terms of groups, ω -chains of left zero semigroups, ω -chains of right zero semigroups, and ‘homomorphisms’. In §5, we obtain our characterization of ω - \mathcal{L} -unipotent bisimple semigroups [5, Theorem 7.11] as a corollary of Theorem 4.1.

1. The congruence t . In this section, S will denote a generalized ω - \mathcal{L} -unipotent bisimple semigroup, i.e., S is a bisimple semigroup such that $E(S)$ is an ω -chain of rectangular bands $(E_n: n \in N)$ and \mathcal{L} is a left congruence on $E(S)$. Recall S is a regular semigroup. Thus, for each $a \in S$, there exists $y \in S$ such that $aya = a$ and $yay = y$ (for example, if $a = axa$, let $y = xax$ [1, Lemma 1.14]). The element y is termed an inverse of a . We will denote the set of all inverses of a by $\mathcal{I}(a)$.

Let $t = ((x, y) \in S^2: xx', yy' \in E_n \text{ and } x'x, y'y \in E_m \text{ for some } m, n \in N, x' \in \mathcal{I}(x), \text{ and } y' \in \mathcal{I}(y))$. We first show that t is a congruence on S and that $S/t \cong C$, the bicyclic semigroup. We also note that C may be taken as a set of representative elements for the t -classes of S and that $T = \ker t$ (the union of the collection of t -classes of S containing idempotents) is an ω -chain of rectangular groups.

Finally, we describe T in terms of (I, o) , an ω -chain of left zero

semigroups $(I_n: n \in N)$; $(J, *)$, an ω -chain of right groups $(J_n: n \in N)$; and an anti-homomorphism $j \rightarrow A_j$ of J into T_I , the full transformation semigroups on I . In fact, $T \cong U(I_n \times J_n: n \in N)$ under the multiplication $(i, j)(p, q) = (i \circ p A_j, j^* q)$.

LEMMA 1.1. *If $e_0 \in E_0$, R_{e_0} is a semigroup.*

Proof. Lemma 1.1 is a special case of [5, Lemma 3.1].

REMARK. Immediately below, we write Theorem 1.2 [5, Theorem 3.3]). This means Theorem 1.2 is obtained by taking “ Y ” to be a one element set in [5, Theorem 3.3]. $(D_\delta: \delta \in Y)$ is the collection of \mathcal{D} -classes in the semigroup of [5, Theorem 3.3]. We do the same thing in Note 1.3, Propositions 1.4 and 1.5, and Lemmas 1.9–1.12.

THEOREM 1.2 [5, Theorem 3.3]. *t is a congruence on S and $S/t \cong C$.*

Note 1.3 [5, Note 3.4]. If we let $t_{(n,k)} = (n, k)t^{-1}$, the t -classes of S are $(t_{(n,k)}: n, k \in N)$ with $t_{(n,k)}t_{(r,s)} \subseteq t_{(n+r-\min(k,r), k+s-\min(k,r))}$. We may write $E(S) = \bigcup (E_{(k,k)}: k \in N)$ where $E_{(k,k)}$ is a rectangular band and $E(t_{(k,k)}) = E_{(k,k)}$. Actually, $E_{(k,k)} = E_k$.

PROPOSITION 1.4 [5, Proposition 3.5]. *$t_{(n,k)} = (a \in S: aa' \in E_{(n,n)} \text{ and } a'a \in E_{(k,k)} \text{ for some } a' \in \mathcal{J}(a)) = \bigcup (R_e \cap L_f: e \in E_{(n,n)} \text{ and } f \in E_{(k,k)})$.*

A rectangular group is the algebraic direct product of a group and a rectangular band.

PROPOSITION 1.5 [5, Proposition 3.6]. *For each $k \in N$, $t_{(k,k)}$ is a rectangular group. In fact, $t_{(k,k)} \cong G \times E_{(k,k)}$ where G is a fixed maximal subgroup of S . Furthermore, $t_{(k,k)}t_{(s,s)} \subseteq t_{(\max(k,s), \max(k,s))}$.*

REMARK 1.6. If $b \in R_e \cap L_f (e, f \in E(S))$, there exists $x \in S$ such that $bx = e$. It is shown in the proof of [1, Theorem 2.18] that $b^{-1} = fxe$ is the unique inverse of b contained in $R_f \cap L_e$ and that $bb^{-1} = e$ and $b^{-1}b = f$.

Note 1.7. Let e_0 be a fixed element of E_0 and fix an element $e_1 \in E_{(1,1)}$ such that $e_1 < e_0$. For example, select any $f \in E_{(1,1)}$ and let $e_1 = e_0 f e_0$. Hence, $e_1 \in E_{(1,1)}$ by Note 1.3 and $e_1 < e_0$.

Note 1.8. Select and fix $a \in R_{e_0} \cap L_{e_1}$. By Remark 1.6, there exists a unique $a^{-1} \in \mathcal{J}(a) \cap R_{e_1} \cap L_{e_0}$ with $aa^{-1} = e_0$ and $a^{-1}a = e_1$. Define

$a^{-n} = (a^{-1})^n$ for all positive integers n and define $a^0 = e_0$. Utilizing Proposition 1.4 and Note 1.3, $a^{-n}a^k \in t_{(n,k)}$ for all $n, k \in N$.

LEMMA 1.9 [5, Lemma 3.9]. $a^k a^{-k} = e_0$ for all $k \in N$.

LEMMA 1.10 [5, Lemma 3.10].

$$a^k a^{-r} = \begin{cases} a^{k-r} & \text{if } k > r \\ a^{-(r-k)} & \text{if } r > k \\ e_0 & \text{if } r = k. \end{cases}$$

LEMMA 1.11 [5, Lemmas 3.11, 3.12].

$$(1) a^{-k} a^p a^{-r} a^s = a^{-(k+r-\min(r,p))} a^{p+s-\min(r,p)} \quad (2) a^{-r} a^r \in E_{(r,r)} \text{ for all } r \in N.$$

For brevity, let $T_k = t_{(k,k)}$ and let $T = \bigcup (T_k: k \in N)$. Hence, T is an ω -chain of the rectangular groups $(T_k: k \in N)$ by Proposition 1.5. Since $E(S) = E(T)$ by Note 1.3, T is generalized \mathcal{L} -unipotent. Utilizing Proposition 1.5, $T_k = G \times M_k \times N_k$ where G is a group, M_k is a left zero semigroup, and N_k is a right zero semigroup. By Lemma 1.11, $a^{-k}a^k \in E(T_k)$. Let I_k denote the set of idempotents of the \mathcal{L} -class of T_k containing $a^{-k}a^k$ and let J_k denote the \mathcal{R} -class of T_k containing $a^{-k}a^k$. We may suppose that $l_k \in M_k \cap N_k$, $a^{-k}a^k = (e, l_k, l_k)$ where e is the identity of G , $I_k = (e) \times M_k \times (l_k)$, and $J_k = G \times (l_k) \times N_k$. For brevity, let $e_k = (e, l_k, l_k)$. Hence, using Lemma 1.11, $e_m e_n = e_{\max(n,m)}$.

Let $I = \bigcup (I_n: n \in N)$ and let $J = \bigcup (J_n: n \in N)$.

LEMMA 1.12. I is an ω -chain of left zero semigroups $(I_n: n \in N)$.

Proof. By a direct calculation, I_n is a left zero semigroup for each $n \in N$. Let $x \in I_k$ and let $y \in I_n$. Hence, $x \mathcal{L} e_k$ and $y \mathcal{L} e_n$. Since T is generalized \mathcal{L} -unipotent, $xy \mathcal{L} x e_n$. Thus, since

$$x e_n \mathcal{L} e_k e_n, xy \mathcal{L} e_{\max(k,n)}.$$

Hence, $xy \in I_{\max(k,n)}$.

LEMMA 1.13. For each $n \in N$, J_n is a right group. If $x \in J_n$, $y \in J_m$, and $n \geq m$, $xy \in J_n$.

Proof. By [1, Theorem 1.27], J_n is a right group for each $n \in N$. Let $x \in J_n$, $y \in J_m$, and $n \geq m$. Hence, $y \mathcal{R} e_m$ implies $xy \mathcal{R} x e_m$. Since $e_n(x e_m) = (e_n x) e_m = x e_m$ and $x e_m \in T_n$, $x e_m \in J_n$ by a simple calculation.

Thus, $xy \in J_n$.

LEMMA 1.14. *Every element of T may be uniquely expressed in the form $x = ij$ with $i \in I_n$ and $j \in J_n$ for some $n \in N$.*

Proof. If $x = (g, i, j) \in T_n$, $x = (e, i, l_n)(g, l_n, j)$.

If X is a set, T_X will denote the semigroup (iteration) of mappings of X into X .

LEMMA 1.15. *There exists a mapping $j \mapsto A_j$ of J into T_I and a mapping $p \mapsto B_p$ of I into T_J such that $I_n A_j \subseteq I_{\max(n, m)}$ for $j \in J_m$ and $J_n B_p \subseteq J_{\max(n, m)}$ for $p \in I_m$. If $j \in J$ and $p \in I$, $jp = pA_j jB_p$. Furthermore, $jp \not\subseteq pA_j (\in T)$ and $jp \not\subseteq jB_p (\in T)$.*

Proof. Let $j \in J_m$ and $p \in I_n$. Thus, $jp \in T_{\max(m, n)}$. Hence, by Lemma 1.14, there exists a unique $u \in I_{\max(m, n)}$ and $v \in J_{\max(m, n)}$ such that $jp = uv$. Let $u = pA_j$ and $v = jB_p$. The last statement is valid by a simple calculation.

LEMMA 1.16. *If $j \in J$, $jB_{e_v} = e_v j e_v$. If $j \in J_r$ and $r \geq v$, $jB_{e_v} = j e_v$.*

Proof. Let $j \in J_r$ and suppose that $v > r$. Thus, $j e_v \in T_v$ and $(j e_v) e_v = j e_v$. Hence, $j e_v = (g, i, l_v)$ for some $g \in G$ and $i \in M_v$. By Lemma 1.15, $j e_v = e_v A_j j B_{e_v}$ with $j B_{e_v} \not\subseteq j e_v (\in T)$. Hence, $j B_{e_v} = (g, l_v, l_v)$. Thus, $j B_{e_v} = (e, l_v, l_v)(g, i, l_v) = e_v j e_v$. Next, suppose that $r \geq v$. Hence, $j e_v \in J_r$ by Lemma 1.13. Thus, utilizing Lemma 1.15, $e_r(j e_v) = j e_v = e_v A_j j B_{e_v}$ where $e_v A_j \in I_r$ and $j B_{e_v} \in J_r$. Hence, $j B_{e_v} = j e_v$ by Lemma 1.14. This establishes the second sentence of the lemma. Since, for $r \geq v$, $e_v j = e_v e_r j = e_r j = j$, $j B_{e_v} = e_v j e_v$ for $r \geq v$.

LEMMA 1.17. *$(e, f) \in \mathcal{L} \cap (E(T))^2$ and $p \in T$ imply $(pe, pf) \in \mathcal{L}(\in T)$.*

Proof. Suppose $(e, f) \in \mathcal{L} \cap (E(T))^2$. Hence, for

$$p \in T, (p^{-1}pe, p^{-1}pf) \in \mathcal{L}$$

(p^{-1} is the group inverse of p in the group containing p). Thus, $p^{-1}pe p^{-1}pf = p^{-1}pe$ and $p^{-1}pf p^{-1}pe = p^{-1}pf$. Hence, $(pe p^{-1})pf = pe$ and $(pf p^{-1})pe = pf$. Thus, $(pe, pf) \in \mathcal{L}(\in T)$.

LEMMA 1.18. *If $p \in I_r$, $B_p = B_{e_r}$.*

Proof. If $n, m \in N$, let $nm = \max(n, m)$ in this proof. Let $j \in J_s$

and $p \in I_r$. Hence $e_{rs}j = (g, l_{rs}, j')$ for some $g \in G$ and $j' \in N_{rs}$ by Lemma 1.13. By Lemma 1.12, $pe_{rs} = (e, n, l_{rs})$ for some $n \in M_{rs}$. Thus,

$$e_{rs}jpe_{rs} = (g, l_{rs}, l_{rs}) = e_{rs}je_{rs}.$$

Hence, if $jp = (w, m, n)$ and $je_r = (u, c, d)$, then $w = u$. Since $(p, e_r) \in \mathcal{L}$, $(jp, je_r) \in \mathcal{L} (\in T)$ by Lemma 1.17. Hence, $n = d$. Thus, $jp = (w, m, n) = (e, m, l_{rs})(w, l_{rs}, n)$ while $je_r = (w, c, n) = (e, c, l_{rs})(w, l_{rs}, n)$. Hence, utilizing Lemmas 1.14 and 1.15, $jB_p = jB_{e_r}$.

LEMMA 1.19. *Let $r \in J_u, s \in J_v, v \leq u$, and $z \in N$. Then, (a) $(rs)B_{e_z} = rB_{e_{\max(z, v)}}sB_{e_z}$ (b) if $x \in I_z, xA_{rs} = xA_sA_r$.*

Proof. Let $r \in J_u, s \in J_v, u \geq v$, and $x \in I_z$. Hence, utilizing Lemmas 1.13 and 1.15, $(rs)x = xA_{rs}(rs)B_x$ while

$$r(sx) = r(xA_sB_x) = (r(xA_s))(sB_x) = xA_sA_r(rB_{xA_s}B_x).$$

Thus, utilizing Lemma 1.14, $xA_{rs} = xA_sA_r$ and $(rs)B_x = rB_{xA_s}B_x$. Utilizing Lemmas 1.15 and 1.18, $(rs)B_{e_z} = rB_{e_{\max(z, v)}}sB_{e_z}$.

If $x \in J_u$ and $y \in J_v$, define $x^*y = xB_{e_v}y$.

LEMMA 1.20. *If $x \in J_u$ and $y \in J_v, x^*y = e_vxy$. If $u \geq v, x^*y = xy$.*

Proof. Let $x \in J_u$ and $y \in J_v$. Hence, utilizing Lemma 1.16,

$$x^*y = xB_{e_v}y = (e_vxe_v)y = e_vx(e_vy) = e_vxy.$$

If $u \geq v$, again utilizing Lemma 1.16, $x^*y = xB_{e_v}y = (xe_v)y = x(e_vy) = xy$.

LEMMA 1.21. *$(J, *)$ is an ω -chain of right groups $(J_n; n \in N)$.*

Proof. Utilizing Lemmas 1.13 and 1.20, $(J_n, *)$ is a right group for each $n \in N$ and $J_n^*J_m \subseteq J_{\max(n, m)}$. We must just establish associativity. Let $i \in J_s, p \in J_y$, and $w \in J_z$. Hence, utilizing Lemmas 1.15 and 1.13, $i^*(p^*w) = i^*(pB_{e_z}w) = iB_{e_{\max(y, z)}}pB_{e_z}w$ while

$$(i^*p)^*w = (iB_{e_y}p)^*w = (iB_{e_y}p)B_{e_z}w.$$

Utilizing Lemma 1.19 (a) $(iB_{e_y}p)B_{e_z} = iB_{e_y}B_{e_{\max(y, z)}}pB_{e_z}$. However, utilizing Lemma 1.16,

$$iB_{e_y}B_{e_{\max(y, z)}} = e_{\max(y, z)}e_yie_ye_{\max(y, z)} = e_{\max(y, z)}ie_{\max(y, z)} = iB_{e_{\max(y, z)}}.$$

Hence, $(i^*p)^*w = iB_{e_{\max(y, z)}}pB_{e_z}w = i^*(p^*w)$.

DEFINITION 1.22. Let the semigroup X be an ω -chain of semi-

groups $(X_n: n \in N)$ and let ϕ be a mapping of X into a semi-group Y . If $r \in X_n, s \in X_m$, and $n \geq m$ imply $(rs)\phi = s\phi r\phi$, ϕ is termed an upper anti-homomorphism of X into Y .

LEMMA 1.23. $r \mapsto A_r$ is an upper anti-homomorphism of $(J, *)$ into T_I .

Proof. Combine Lemmas 1.20 and 1.19 (b).

LEMMA 1.24. If $j \in J_v$ and $i \in I_z$, $ji = iA_jje_z = iA_jj^*e_z$.

Proof. Let $j \in J_v$ and $i \in I_z$. Hence, $ji = iA_jjB_i$ by Lemma 1.15. However, utilizing Lemmas 1.18 and 1.16, $jB_i = jB_{e_z} = e_zje_z$. Since $iA_j \in I_{\max(v,z)}$, $iA_j = iA_je_{\max(v,z)}$. Hence, $ji = iA_je_{\max(v,z)}e_zje_z = iA_jje_z$. However, $e_zje_z = j^*e_z$ by Lemma 1.20. Hence, $ji = iA_jj^*e_z$.

LEMMA 1.25. If $r, s \in I$ with $r \in I_u$, $(rs)A_x = rA_xsA_{x**e_u}$ for all $x \in J$.

Proof. Let $r, s \in I$ with $r \in I_u$ and let $x \in J$. Hence, utilizing Lemmas 1.15 and 1.12, $x(rs) = (rs)A_xxB_{r_s}$ while

$$(xr)s = (rA_xxB_r)s = rA_x(xB_r s) = rA_x(sA_{xB_r}xB_rB_s) = rA_xsA_{xB_r}xB_rB_s.$$

Hence, utilizing Lemmas 1.15, 1.12, and 1.14, $(rs)A_x = rA_xsA_{xB_r}$. Utilizing Lemmas 1.18, 1.16, and 1.20, $xB_r = xB_{e_u} = e_uxe_u = x^*e_u$.

REMARK 1.26. Results of [6] could have been applied to characterize T .

2. Structure theorem for generalized ω - \mathcal{L} -unipotent bisimple semigroups. (Proof of converse.) In this section, we complete the proof of the converse of our structure theorem for generalized ω - \mathcal{L} -unipotent bisimple semigroups (Theorem 2.21).

We will use a sequence of twenty entries to establish Theorem 2.21. S will denote a generalized ω - \mathcal{L} -unipotent bisimple semigroup.

LEMMA 2.1. Every element of S may be uniquely expressed in the form $x = ia^{-n}a^k j$ where $i \in I_n$ and $j \in J_k$.

Proof. Let $x \in t_{(n,k)}$. Hence, $(x, e) \in \mathcal{R}$ for some $e \in E_n$ by Proposition 1.4. Thus, $(x, i) \in \mathcal{R}$ for some $i \in I_n$. Thus, since $a^{-n}a^n \in I_n$, a left zero semigroup, $x = ix = (ia^{-n}a^n)x = ia^{-n}a^n x$. Since $a^n \in R_{e_0}$ by Note 1.8 and Lemma 1.1 and $a^k a^{-k} = e_0$ by Lemma 1.9, $a^k a^{-k} a^n =$

a^n . Hence $x = ia^{-n}(a^k a^{-k} a^n)x = (ia^{-n} a^k)(a^{-k} a^n x)$. However, $a^{-k} a^n x \in t_{(k,k)}$ by Notes 1.8 and 1.3. Thus, since $a^{-k} a^k (a^{-k} a^n x) = a^{-k} a^n x$ and $a^{-k} a^k \in J_k$, $a^{-k} a^n x \in J_k$ by Proposition 1.5. Hence, $x = ia^{-n} a^k j$ where $i \in I_n$ and $j \in J_k$. We next establish uniqueness. Suppose that $x = ia^{-n} a^k j = ua^{-r} a^s v$ ($i \in I_n$, $j \in J_k$, $u \in I_r$, and $v \in J_s$). Thus, using Note 1.3, $x \in t_{(n,k)} \cap t_{(r,s)}$ and, hence, $n = r$ and $k = s$. Thus, $ia^{-n} a^k j = ua^{-n} a^k v$. Hence, $a^{-n} a^n ia^{-n} a^k j = a^{-n} a^n ua^{-n} a^k v$. Thus, since $a^{-n} a^n$, i , $u \in I_n$, a left zero semigroup, $a^{-n} a^n ia^{-n} a^k j = a^{-n} a^n ua^{-n} a^k v$. Hence, $a^{-n} a^k j = a^{-n} a^k v$. Thus, $a^{-k} a^n a^{-n} a^k j = a^{-k} a^n a^{-n} a^k v$. Hence, $a^{-k} a^k j = a^{-k} a^k v$. Since $a^{-k} a^k \in E(J_k)$ and $j, v \in J_k$, a right group, $j = v$. Thus, $ia^{-n} a^k j = ua^{-n} a^k j$. Since J_k is a right group, there exists $z \in J_k$ such that $jz = a^{-k} a^k$. Hence $ia^{-n} a^k jz = ua^{-n} a^k jz$ implies $ia^{-n} a^k a^{-k} a^k = ua^{-n} a^k a^{-k} a^k$. Thus, $ia^{-n} a^k = ua^{-n} a^k$. Hence $ia^{-n} a^k a^{-k} a^n = ua^{-n} a^k a^{-k} a^n$. Thus,

$$ia^{-n} a^n = ua^{-n} a^n.$$

Since $i, u, a^{-n} a^n \in I_n$, a left zero semigroup, $i = u$.

DEFINITION 2.2. If $u \in T$ and $n, k \in N$, define $u\nu_{(k,n)} = a^{-n} a^k u a^{-k} a^n$.

LEMMA 2.3. $T_r \nu_{(k,n)} \subseteq T_{n+r-\min(k,r)}$.

Proof. Let $g \in T_r$. Hence, utilizing Note 1.3, $g\nu_{(k,n)} = a^{-n} a^k g a^{-k} a^n \in t_{(n,k)(r,r)(k,n)} = T_{n+r-\min(k,r)}$.

LEMMA 2.4. Let $g_r \in T_r$ and $h_s \in T_s$. If $k \geq r, s$ or $r = s \geq k$, $(g_r h_s)\nu_{(k,n)} = g_r \nu_{(k,n)} h_s \nu_{(k,n)}$. In particular, $\nu_{(k,n)}$ is a homomorphism of T_r into $T_{n+r-\min(k,r)}$.

Proof. Let $g_r \in T_r$ and $h_s \in T_s$ with $k \geq r, s$. Hence,

$$(g_r h_s)\nu_{(k,n)} = a^{-n} a^k g_r h_s a^{-k} a^n = a^{-n} a^k (a^{-k} a^k g_r) u_k a^{-k} a^k f_k (h_s a^{-k} a^k) a^{-k} a^n$$

where $(u_k, a^{-k} a^k g_r) \in \mathcal{L}$ with $u_k \in E(J_k)$ and $(f_k, h_s a^{-k} a^k) \in \mathcal{R}$ with $f_k \in I_k$. Hence, $(g_r h_s)\nu_{(k,n)} = a^{-n} a^k g_r a^{-k} a^k h_s a^{-k} a^n = (a^{-n} a^k g_r a^{-k} a^n)(a^{-n} a^k h_s a^{-k} a^n) = g_r \nu_{(k,n)} h_s \nu_{(k,n)}$. Next suppose that $r = s \geq k$. Then,

$$(g_r h_r)\nu_{(k,n)} = a^{-n} a^k g_r v_r a^{-k} a^k f_r h_r a^{-k} a^n$$

where $(v_r, g_r) \in \mathcal{L}$ with $v_r \in E(J_r)$ and $(f_r, h_r) \in \mathcal{R}$ with $f_r \in I_r$. Hence, $(g_r h_r)\nu_{(k,n)} = (a^{-n} a^k g_r a^{-k} a^n)(a^{-n} a^k h_r a^{-k} a^n) = g_r \nu_{(k,n)} h_r \nu_{(k,n)}$.

DEFINITION 2.5. Let $\nu_{(k,n)}|I = \alpha_{(k,n)}$ and $\nu_{(k,n)}|J = \beta_{(k,n)}$.

LEMMA 2.6. (a) $I_r \alpha_{(k,n)} \subseteq I_{n+r-\min(k,r)}$ (b) $J_r \beta_{(k,n)} \subseteq J_{n+r-\min(k,r)}$.

Proof. (a) By Lemma 2.3, $I_r \nu_{(k,n)} \subseteq T_n$ if $k \geq r$ and $I_r \nu_{(k,n)} \subseteq T_{n+r-k}$

if $r \geq k$. If $k \geq r$, $\nu_{(k,n)}$ is a homomorphism of T_r into T_n by Lemma 2.4. Hence, $I_r \nu_{(k,n)} \subseteq E(T_n)$. Let $g_r \in I_r$. Hence, $g_r \mathcal{L} a^{-r} a^r (\in T_r)$. Thus, $g_r \nu_{(k,n)} \mathcal{L} a^{-n} a^k a^{-r} a^r a^{-k} a^n (\in T_n)$. However,

$$a^{-n} a^k a^{-r} a^r a^{-k} a^n = a^{-n} a^n$$

by Lemma 1.11. Hence, $g_r \nu_{(k,n)} \in I_n$ if $k \geq r$. The case $r \geq k$ is treated similarly. To prove (b), just replace “ I ” by “ J ” and “ \mathcal{L} ” by “ \mathcal{R} ” in the proof of (a).

DEFINITION 2.7. If X is a semigroup $\text{End } X$ will denote the semigroup of endomorphisms of X (iteration).

LEMMA 2.8. $\alpha_{(k,n)} \in \text{End } I$ for each $n, k \in N$.

Proof. Let $i_r \in I_r$ and $i_s \in I_s$. If $r \geq k$, $i_r a^{-k} a^k i_s = i_r i_s$. Hence,

$$\begin{aligned} (i_r i_s) \alpha_{(k,n)} &= a^{-n} a^k i_r i_s a^{-k} a^n = a^{-n} a^k i_r a^{-k} a^k i_s a^{-k} a^n \\ &= a^{-n} a^k i_r a^{-k} a^n a^{-n} a^k i_s a^{-k} a^n = i_r \alpha_{(k,n)} i_s \alpha_{(k,n)}. \end{aligned}$$

Next, suppose that $k > r$. Since S is generalized \mathcal{L} -unipotent, $i_r \mathcal{L} a^{-r} a^r$ implies $a^{-k} a^k i_r \mathcal{L} a^{-k} a^k a^{-r} a^r$. Thus, $a^{-k} a^k i_r \mathcal{L} a^{-k} a^k$ by Lemma 1.11. Hence, $a^{-k} a^k i_r \in I_k$. Thus,

$$\begin{aligned} (i_r i_s) \alpha_{(k,n)} &= a^{-n} a^k i_r i_s a^{-k} a^n = a^{-n} a^k (a^{-k} a^k i_r) a^{-k} a^k i_s a^{-k} a^n \\ &= (a^{-n} a^k i_r a^{-k} a^n) (a^{-n} a^k i_s a^{-k} a^n) = i_r \alpha_{(k,n)} i_s \alpha_{(k,n)}. \end{aligned}$$

LEMMA 2.9. $(n, k) \rightarrow \alpha_{(n,k)}$ is a homomorphism of C into $\text{End } I$.

Proof. Let $g \in I$. We will employ Lemma 1.11. Thus,

$$\begin{aligned} g \alpha_{(r,s)} \alpha_{(n,p)} &= a^{-p} a^n a^{-s} a^r g a^{-r} a^s a^{-n} a^p \\ &= a^{-(p+s-\min(n,s))} a^{n+r-\min(n,s)} g a^{-(r+n-\min(n,s))} a^{s+p-\min(n,s)} \\ &= g \alpha_{(r,s)(n,p)}. \end{aligned}$$

We next establish that $\beta_{(n,k)} \in \text{End } (J, *)$. This will be accomplished by Lemmas 2.10-2.15.

LEMMA 2.10. $\beta_{(1,0)} \in \text{End } (J, *)$.

Proof. Let $w \in J_p$ and $u_s \in J_s$. If $p = s$, $\beta_{(1,0)} \in \text{End } (J, *)$ by Lemmas 2.4, 1.20, and 2.6(b), and Definition 2.5. Let us first suppose $s = 0$. Utilizing Lemmas 1.13, 1.11, Note 1.8, and Definition 2.5,

$$\begin{aligned} (w u_0) a^{-1} &= a^{-p} a^p a^{-1} a (w u_0) a^{-1} = a^{-p} a^p a^{-1} (w u_0) \beta_{(1,0)} \\ &= a^{-p} a^p a^{-1} a^0 (w u_0) \beta_{(1,0)} = e_p a^{-p} a^{p-1} (w u_0) \beta_{(1,0)}. \end{aligned}$$

We note that $(wu_0)\beta_{(1,0)} \in J_{p-1}$ by Lemma 2.6(b). Utilizing Note 1.8, Lemmas 1.15, 1.16, and Definition 2.5, $u_0a^{-1} = u_0e_1a^{-1} = e_1A_{u_0}e_1u_0e_1a^{-1} = e_1A_{u_0}a^{-1}au_0a^{-1} = e_1A_{u_0}a^{-1}(u_0\beta_{(1,0)})$. Hence, utilizing Lemmas 1.15, 1.23, 1.18, 1.16, and 1.11, and Definition 2.5,

$$\begin{aligned} wu_0a^{-1} &= w(e_1A_{u_0})a^{-1}(u_0\beta_{(1,0)}) = e_1A_{wu_0}e_1we_1a^{-1}(u_0\beta_{(1,0)}) \\ &= e_1A_{wu_0}a^{-1}(awa^{-1}(u_0\beta_{(1,0)})) = e_1A_{wu_0}a^{-1}(w\beta_{(1,0)})u_0\beta_{(1,0)} \\ &= e_1A_{wu_0}a^{-p}a^pa^{-1}a^0(w\beta_{(1,0)})u_0\beta_{(1,0)} \\ &= e_1A_{wu_0}a^{-p}a^{p-1}(w\beta_{(1,0)})u_0\beta_{(1,0)}. \end{aligned}$$

Utilizing Lemmas 1.15, 2.6(b), and 1.13, $e_1A_{wu_0} \in I_p$ and $w\beta_{(1,0)}u_0\beta_{(1,0)} \in J_{p-1}$. Hence, $(wu_0)\beta_{(1,0)} = w\beta_{(1,0)}u_0\beta_{(1,0)}$ by Lemma 2.1. Thus, utilizing Lemma 1.20 and 2.6(b), $(w^*u_0)\beta_{(1,0)} = w\beta_{(1,0)}^*u_0\beta_{(1,0)}$. Next, we assume that $p \geq s \geq 1$. Hence, utilizing Lemmas 1.11, 2.6(b), and 1.20,

$$\begin{aligned} (w^*u_s)\beta_{(1,0)} &= (wu_s)\beta_{(1,0)} = awu_s a^{-1} = awa^{-s}a^s u_s a^{-1} \\ &= awa^{-1}aa^{-s}a^s u_s a^{-1} = (awa^{-1})(au_s a^{-1}) \\ &= w\beta_{(1,0)}u_s\beta_{(1,0)} = w\beta_{(1,0)}^*u_s\beta_{(1,0)}. \end{aligned}$$

Finally, we assume $s > p$. Utilizing Lemmas 1.20, 1.13, 1.10 and the case $(p \geq s)$ just established,

$$\begin{aligned} (w^*u_s)\beta_{(1,0)} &= ((e_s w)u_s)\beta_{(1,0)} = (e_s w)\beta_{(1,0)}u_s\beta_{(1,0)} = e_s\beta_{(1,0)}w\beta_{(1,0)}u_s\beta_{(1,0)} \\ &= aa^{-s}a^s a^{-1}w\beta_{(1,0)}u_s\beta_{(1,0)} = e_{s-1}w\beta_{(1,0)}u_s\beta_{(1,0)}. \end{aligned}$$

Since $u_s\beta_{(1,0)} \in J_{s-1}$ by Lemma 2.6(b), $e_{s-1}w\beta_{(1,0)}u_s\beta_{(1,0)} = w\beta_{(1,0)}^*u_s\beta_{(1,0)}$ by Lemma 1.20. Hence, $(w^*u_s)\beta_{(1,0)} = w\beta_{(1,0)}^*u_s\beta_{(1,0)}$.

LEMMA 2.11. $\beta_{(0,0)} \in \text{End}(J, *)$.

Proof. Let $u_r \in J_r$ and $v_s \in J_s$. Utilizing Lemmas 1.20 and 2.6(b),

$$\begin{aligned} (u_r^*v_s)\beta_{(0,0)} &= (e_s u_r v_s)\beta_{(0,0)} = e_0 e_s u_r v_s e_0 = e_s e_0 u_r e_0 e_s v_s e_0 = e_s e_0 u_r e_0 e_0 v_s e_0 \\ &= e_s (u_r \beta_{(0,0)}) v_s \beta_{(0,0)} = u_r \beta_{(0,0)}^* v_s \beta_{(0,0)}. \end{aligned}$$

LEMMA 2.12. $(n, k) \rightarrow \beta_{(n,k)}$ is a homomorphism of C into T_J .

Proof. Replace “ I ” by “ J ” and “ α ” by “ β ” in the proof of Lemma 2.9.

LEMMA 2.13. $\beta_{(k,0)} \in \text{End}(J, *)$ for all $k \in N$.

Proof. We have shown that $\beta_{(0,0)} \in \text{End}(J, *)$ (Lemma 2.11) and that $\beta_{(1,0)} \in \text{End}(J, *)$ (Lemma 2.10). Suppose that $\beta_{(n,0)} \in \text{End}(J, *)$.

We show that $\beta_{(n+1,0)} \in \text{End}(J, *)$. Let $g, h \in J$. Hence, utilizing Lemma 2.12,

$$\begin{aligned} (g^*h)\beta_{(n+1,0)} &= (g^*h)\beta_{(n,0)}\beta_{(1,0)} = (g\beta_{(n,0)}^*h\beta_{(n,0)})\beta_{(1,0)} \\ &= g\beta_{(n,0)}\beta_{(1,0)}^*h\beta_{(n,0)}\beta_{(1,0)} = g\beta_{(n+1,0)}^*h\beta_{(n+1,0)}. \end{aligned}$$

LEMMA 2.14. $\beta_{(0,k)} \in \text{End}(J, *)$ for all $k \in N$.

Proof. Let $u_r \in J_r$ and $v_s \in J_s$. First, assume $s \geq r$. Utilizing Lemma 1.20, $(u_r^*v_s)\beta_{(0,k)} = (e_s u_r v_s)\beta_{(0,k)}$. Since $u_r \mathcal{R} e_r$, $e_s u_r \mathcal{R} e_s e_r = e_s$. Hence, utilizing Lemma 2.4, $((e_s u_r) v_s)\beta_{(0,k)} = (e_s u_r)\beta_{(0,k)} v_s \beta_{(0,k)}$. Utilizing Definition 2.5, Note 1.8, Lemma 1.1, and Lemma 1.9, $(e_s u_r)\beta_{(0,k)} = a^{-k} e_0 (e_s u_r) e_0 a^k = a^{-k} e_s u_r a^k = a^{-k} e_s a^k a^{-k} u_r a^k = (a^{-k} a^{-s} a^s a^k) (a^{-k} a^0 u_r a^{-0} a^k) = e_{s+k} (u_r \beta_{(0,k)})$. Since $v_s \beta_{(0,k)} \in J_{s+k}$ by Lemma 2.6(b), $e_{s+k} u_r \beta_{(0,k)} v_s \beta_{(0,k)} = u_r \beta_{(0,k)}^* v_s \beta_{(0,k)}$ by Lemma 1.20. Thus, $(u_r^* v_s)\beta_{(0,k)} = u_r \beta_{(0,k)}^* v_s \beta_{(0,k)}$. We utilize Lemmas 1.20 and 1.9, and Definition 2.5 for the case $r > s$.

LEMMA 2.15. $\beta_{(n,k)} \in \text{End}(J, *)$ for all $n, k \in N$.

Proof. Let $g, h \in J$. Hence, utilizing Lemmas 2.12, 2.13, and 2.14,

$$\begin{aligned} (g^*h)\beta_{(n,k)} &= (g^*h)\beta_{(n,0)(0,k)} = (g^*h)\beta_{(n,0)}\beta_{(0,k)} = (g\beta_{(n,0)}^*h\beta_{(n,0)})\beta_{(0,k)} \\ &= g\beta_{(n,0)}\beta_{(0,k)}^*h\beta_{(n,0)}\beta_{(0,k)} = g\beta_{(n,k)}^*h\beta_{(n,k)}. \end{aligned}$$

LEMMA 2.16. $(n, k) \rightarrow \beta_{(n,k)}$ is a homomorphism of C into $\text{End}(J, *)$.

Proof. Combine Lemmas 2.12 and 2.15.

If $a, b \in I$, define $a \circ b = ab$.

LEMMA 2.17. $S \cong ((i, (n, k), j): i \in I_n, j \in J_k, n, k \in N)$ under the multiplication $((i, (n, k), j)(u, (r, s), v) = (i \circ (u A_j \alpha_{(k,n)}), (n + r - \min(k, r), k + s - \min(k, r)), j \beta_{(r,s)}^* v)$.

Proof. Let $i \in I_n, j \in J_k, u \in I_r$, and $v \in J_s$. Hence, utilizing Lemmas 1.24, 1.15, 1.11, 1.9, 2.6(b), 1.20, and Definition 2.5,

$$\begin{aligned} (i a^{-n} a^k j)(u a^{-r} a^s v) &= i a^{-n} a^k (j u) a^{-r} a^s v = i a^{-n} a^k u A_j j a^{-r} a^{-r} a^s v \\ &= i a^{-n} a^k u A_j a^{-k} a^k a^{-r} a^r j a^{-r} a^s v \\ &= i (a^{-n} a^k (u A_j) a^{-k} a^n) (a^{-n} a^k a^{-r} a^s) a^{-s} a^r j a^{-r} a^s v \\ &= i ((u A_j) \alpha_{(k,n)}) a^{-(n+r-\min(k,r))} a^{k+s-\min(k,r)} j \beta_{(r,s)} v \\ &= i \circ ((u A_j) \alpha_{(k,n)}) a^{-(n+r-\min(k,r))} a^{k+s-\min(k,r)} (j \beta_{(r,s)}^* v). \end{aligned}$$

Utilizing Lemma 2.6, $i \circ ((u A_j) \alpha_{(k,n)}) \in I_{n+r-\min(k,r)}$ and

$$j \beta_{(r,s)}^* v \in J_{k+s-\min(k,r)}.$$

Hence, $((i, (n, k), j): i \in I_n, j \in J_k, n, k \in N)$ under the multiplication given in the statement of the lemma is a groupoid. The required isomorphism is given by the mapping $(ia^{-n}a^k j)\varphi = (i, (n, k), j)$ by virtue of the above and Lemma 2.1.

LEMMA 2.18. $\alpha_{(r,s)}A_j\alpha_{(s,r)} = A_{j\beta(s,r)}$ for all $j \in J$ and $r, s \in N$.

Proof. Let $j \in J_p$ and $w \in I_q$. Utilizing Definitions 2.2 and 2.5, and Lemmas 1.9, 1.12, 1.15, and 2.6,

$$\begin{aligned} (j(w\alpha_{(r,s)}))\nu_{(s,r)} &= a^{-r}a^s j(a^{-s}a^r w a^{-r}a^s) a^{-s}a^r \\ &= a^{-r}a^s j a^{-s}a^r (a^{-r}a^r w) a^{-r}a^r = a^{-r}a^s j a^{-s}a^r a^{-r}a^r w \\ &= a^{-r}a^s j a^{-s}a^r w = j\beta_{(s,r)} w = w A_{j\beta(s,r)} j\beta_{(s,r)} B_w. \end{aligned}$$

Utilizing Lemmas 1.15 and 2.6, $w A_{j\beta(s,r)} \in I_{\max(q, r+p-\min(s,p))}$ and

$$j\beta_{(s,r)} B_w \in J_{\max(q, r+p-\min(s,p))}.$$

Utilizing Definitions 2.2 and 2.5, and Lemmas 1.15 and 2.6,

$$\begin{aligned} (j(w\alpha_{(r,s)}))\nu_{(s,r)} &= a^{-r}a^s j(w\alpha_{(r,s)}) a^{-s}a^r \\ &= a^{-r}a^s (w\alpha_{(r,s)} A_j) (j B_{w\alpha_{(r,s)}}) a^{-s}a^r \\ &= a^{-r}a^s (w\alpha_{(r,s)} A_j) a^{-s}a^s (j B_{w\alpha_{(r,s)}}) a^{-s}a^r \\ &= a^{-r}a^s (w\alpha_{(r,s)} A_j) a^{-s}a^r (a^{-r}a^s j B_{w\alpha_{(r,s)}} a^{-s}a^r) \\ &= (w\alpha_{(r,s)} A_j) \alpha_{(s,r)} (j B_{w\alpha_{(r,s)}}) \beta_{(s,r)}. \end{aligned}$$

Utilizing Lemmas 1.15 and 2.6, $w\alpha_{(r,s)} A_j \alpha_{(s,r)} \in I_{\max(p, s+q-\min(r,q))+r-s}$ and $(j B_{w\alpha_{(r,s)}}) \beta_{(s,r)} \in J_{\max(p, s+q-\min(r,q))+r-s}$. Hence, $w\alpha_{(r,s)} A_j \alpha_{(s,r)} = w A_{j\beta(s,r)}$ by Lemma 1.14.

LEMMA 2.19. (a) $g\alpha_{(s,s)} = e_s \circ g$ for all $g \in I$. (b) $g\beta_{(s,s)} = g^* e_s$ for all $g \in J$.

Proof. (a) Let $g \in I$. Utilizing Lemma 1.12 $g\alpha_{(s,s)} = (e_s g) e_s = e_s g = e_s \circ g$. (b) Let $g \in J$. Utilizing Lemma 1.20, $g\beta_{(s,s)} = e_s g e_s = g^* e_s$.

In the following definition, we will describe the objects we will use to represent generalised ω - \mathcal{L} -unipotent bisimple semigroups.

DEFINITION 2.20. Let (I, o) be an ω -chain of left zero semigroups $(I_k: k \in N)$; let $(n, r) \rightarrow \alpha_{(n,r)}$ be a homomorphism of C into $\text{End}(I, o)$; let $(J, *)$ be an ω -chain of right groups $(J_k: k \in N)$; let $(n, r) \rightarrow \beta_{(n,r)}$ be a homomorphism of C into $\text{End}(J, *)$; let $j \rightarrow A_j$ be an upper anti-homomorphism of $(J, *)$ into T_I ; and let $I_k \cap J_k = (e_k)$, a single idempotent, for each $k \in N$ such that

- (1) $g\beta_{(s,s)} = g^*e_s$ for all $g \in J$.
 (2) $I_r\alpha_{(n,k)} \subseteq I_{k+r-\min(n,r)}$ and $J_r\beta_{(n,k)} \subseteq J_{k+r-\min(n,r)}$.
 (3) $I_rA_j \subseteq I_{\max(r,k)}$ if $j \in J_k$.
 (4) $(r \circ s)A_x = rA_x \circ sA_{x * e_u}$ for $r, s \in I$ with $r \in I_u$ and $x \in J$.
 (5) $\alpha_{(r,s)}A_j\alpha_{(s,r)} = A_{j\beta_{(s,r)}}$ for all $j \in J$ and $r, s \in N$.
 We denote $((i, (n, k), j): i \in I_n, j \in J_k)$ under the multiplication

$$(6) \quad (i, (n, k), j)(u, (r, s), v) \\ = (i \circ (uA_j\alpha_{(k,n)}), (n + r - \min(k, r), k + s - \min(k, r)), j\beta_{(r,s)}^*v)$$

by (I, J, α, β, A) .

THEOREM 2.21. *Let S be a generalized ω - \mathcal{L} -unipotent bisimple semigroup. Then, S is isomorphic to some (I, J, α, β, A) .*

Proof. The theorem is a consequence of the definition of “ \circ ”, Lemmas 1.12, 2.9, 1.21, 2.16, 1.23, the choice of “ e_k ”, Lemmas 2.19, 2.6, 1.15, 1.25, 2.18, and 2.17.

We thank the referee for the following remark.

REMARK 2.22. In Definition 2.20, the middle component (m, n) of $(i, (m, n), j)$ serves only as a marker. Hence, S is actually represented by the cartesian product $I \times J$ under the multiplication

$$(i, j)(u, v) = (i \circ (uA_j\alpha_{(k,n)}), j\beta_{(r,s)}^*v)$$

where $i \in I_n, j \in J_k, u \in I_r$, and $v \in J_s$.

3. Structure theorem for generalized ω - \mathcal{L} -unipotent bisimple semigroups (proof of direct half). In this section, we show that (I, J, α, β, A) is a generalized ω - \mathcal{L} -unipotent bisimple semigroup.

LEMMA 3.1. *(I, J, α, β, A) is a semigroup.*

Proof. We use (2) and (3) of Definition 2.20 to establish closure. We next establish associativity. Let $(i, (n, k), j)_1 = i$ and $(i, (n, k), j)_{23} = ((n, k), j)$. Let $a = (i, (n, k), j)$, $b = (u, (r, s), v)$, and $c = (z, (p, q), w) \in (I, J, \alpha, \beta, A)$. Utilizing the fact that $(n, r) \rightarrow \beta_{(n,r)}$ is a homomorphism,

$$\begin{aligned} ((ab)c)_{23} &= ((i \circ (uA_j\alpha_{(k,n)}), (n, k)(r, s), j\beta_{(r,s)}^*v)(z, (p, q), w))_{23} \\ &= ((n, k)(r, s)(p, q), (j\beta_{(r,s)}^*v)\beta_{(p,q)}^*w) \\ &= ((n, k)(r, s)(p, q), j\beta_{(r,s)(p,q)}^*v\beta_{(p,q)}^*w) \end{aligned}$$

while

$$\begin{aligned} (a(bc))_{23} &= ((i, (n, k), j)(u \circ (zA_v \alpha_{(s,r)}), (r, s)(p, q), v\beta_{(p,q)}^* w))_{23} \\ &= ((n, k)(r, s)(p, q), j\beta_{(r,s)(p,q)}^* v\beta_{(p,q)}^* w). \end{aligned}$$

Hence, $((ab)c)_{23} = (a(bc))_{23}$. Utilizing the fact $(k, n) \rightarrow \alpha_{(k,n)}$ is a homomorphism of C into $\text{End}(I, o)$, the fact $j \rightarrow A_j$ is an upper anti-homomorphism of $(J, *)$ into T_I , (5), (1), and (4),

$$\begin{aligned} ((ab)c)_1 &= i \circ uA_j \alpha_{(k,n)} \circ zA_{j\beta_{(r,s)} * v} \alpha_{(s,r)(k,n)} \\ &= i \circ ((uA_j \circ zA_{j\beta_{(r,s)} * v} \alpha_{(s,r)}) \alpha_{(k,n)}) \\ &= i \circ ((uA_j \circ zA_v A_{j\beta_{(r,s)}} \alpha_{(s,r)}) \alpha_{(k,n)}) \\ &= i \circ ((uA_j \circ zA_v \alpha_{(s,r)} A_j \alpha_{(r,s)} \alpha_{(s,r)}) \alpha_{(k,n)}) \\ &= i \circ ((uA_j \circ zA_v \alpha_{(s,r)} \alpha_{(r,r)} A_j \alpha_{(r,r)}) \alpha_{(k,n)}) \\ &= i \circ ((uA_j \circ zA_v \alpha_{(s,r)} A_{j\beta_{(r,r)}}) \alpha_{(k,n)}) \\ &= i \circ ((uA_j \circ zA_v \alpha_{(s,r)} A_{j * e_r}) \alpha_{(k,n)}) \\ &= i \circ ((u \circ zA_v \alpha_{(s,r)}) A_j \alpha_{(k,n)}) \\ &= (a(bc))_1. \end{aligned}$$

Hence, $(ab)c = a(bc)$.

LEMMA 3.2. Let $(i, (n, k), j), (w, (p, q), z) \in (I, J, \alpha, \beta, A)$.

(a) $(i, (n, k), j) \mathcal{R}(w, (p, q), z)$ if and only if $i = w$ and $n = p$.

(b) $(i, (n, k), j) \mathcal{L}(w, (p, q), z)$ if and only if $k = q$ and $(j, z) \in \mathcal{H}(\in J_q)$.

Proof. (a) Let us show that $(i, (n, k), j) \mathcal{R}(i, (n, q), z)$. Let $u \in I_k$. Hence, $uA_j \alpha_{(k,n)} \in I_n$ by (2) and (3). Thus, since (I_n, o) is a left zero semigroup, $i \circ uA_j \alpha_{(k,n)} = i$. By (2), $j\beta_{(k,q)} \in J_q$. Hence, since $(J_q, *)$ is a right group, there exists $v \in J_q$ such that $j\beta_{(k,q)}^* v = z$. Hence, utilizing (6), $(i, (n, k), j)(u, (k, q), v) = (i, (n, q), z)$. Similarly, there exists $a \in I_q$ and $b \in J_k$ such that $(i, (n, q), z)(a, (q, k), b) = (i, (n, k), j)$. Utilizing (6), the converse follows from the fact that \mathcal{R} is the identity on $(I, 0)$ and $(n, k) \mathcal{R}(p, q)$ in C implies $n = p$. Let us show that $(i, (n, k), j) \mathcal{L}(w, (p, k), z)$ if $(j, z) \in \mathcal{H}(\in J_k)$. Since $(j, z) \in \mathcal{H}(\in J_k)$, there exists $u \in J_k$ such that $u^* j = z$. By (2), $u\beta_{(k,n)} \in J_n$. Utilizing (1) and the fact $(n, k) \rightarrow \beta_{(n,k)}$ is a homomorphism of C into $\text{End}(J, *)$, $u\beta_{(k,n)} \beta_{(n,k)} = u\beta_{(k,k)} = u^* e_k$. Hence, $(u\beta_{(k,n)} \beta_{(n,k)})^* j = u^* e_k^* j = u^* j = z$. Thus, utilizing (2), (3), and (6), $(w, (p, n), u\beta_{(k,n)})(i, (n, k), j) = (w, (p, k), z)$. Similarly, there exists $v \in J_k$ such that $(i, (n, p), v\beta_{(k,p)})(w, (p, k), z) = (i, (n, k), j)$. Utilizing (6), the converse follows from the fact that $\mathcal{H} = \mathcal{L}$ in $(J, *)$ and $(n, k) \mathcal{L}(p, q)$ in C implies $k = q$.

LEMMA 3.3. (I, J, α, β, A) is a bisimple semigroup.

Proof. Let $(i, (n, k), j), (u, (r, s), v) \in (I, J, \alpha, \beta, A)$. Hence, utilizing Lemma 3.2, $(i, (n, k), j) \mathcal{R} (i, (n, s), v) \mathcal{L} (u, (r, s), v)$. (I, J, α, β, A) is a semigroup by Lemma 3.1.

LEMMA 3.4. $E(I, J, \alpha, \beta, A) = ((i, (n, n), j) : j \in E(J_n), n \in N)$.

Proof. Let $(i, (n, k), j) \in E(I, J, \alpha, \beta, A)$. Hence, $(i, (n, k), j)(i, (n, k), j) = (i, (n, k), j)$. Using (6), $n = k$ since $(n, k)^2 = (n, k)$ in C . Hence, using (6) and (1), $j = j\beta_{(n, n)}^* j = j^* e_n^* j = j^2$. Utilizing (6), (2), (3), and (1), $j \in E(J_n)$ implies $(i, (n, n), j) \in E(I, J, \alpha, \beta, A)$ for $n \in N$ and $i \in I_n$.

LEMMA 3.5. (I, J, α, β, A) is a regular bisimple semigroup.

Proof. It follows from a result of Clifford and Miller [1, Theorem 2.11] that any bisimple semigroup containing an idempotent is regular. Hence, we just apply Lemmas 3.3 and 3.4.

LEMMA 3.6. $E(I, J, \alpha, \beta, A)$ is a semigroup.

Proof. We will utilize Lemma 3.4. Let $a = (i, (n, n), j)$, $b = (u, (s, s), v) \in E(I, J, \alpha, \beta, A)$. Hence, $j \in E(J_n)$ and $v \in E(J_s)$. Thus, using (1), $j\beta_{(s, s)}^* v = j^* e_s^* v = j^* v$. However, $E(T)$ is a semigroup for any chain of right groups T . Thus, it follows that $j^* v \in E(J_{\max(n, s)})$. Hence, $ab \in E(I, J, \alpha, \beta, A)$ by Lemma 3.4.

LEMMA 3.7. \mathcal{L} is a congruence on the semigroup $E(I, J, \alpha, \beta, A)$.

Proof. Let X be any semigroup such that $E(X)$ is a semigroup. Then, it is easily seen that if $e, f \in E(X)$, $(e, f) \in \mathcal{L}(\in X)$ if and only if $(e, f) \in \mathcal{L}(\in E(X))$. Let $j \in E(J_n)$ and $v \in E(J_s)$. Hence, utilizing Lemmas 3.4 and 3.2(b), $(i, (n, n), j) \mathcal{L} (u, (s, s), v) (\in E(I, J, \alpha, \beta, A))$ if and only if $n = s$ and $j = v$. Thus, using (6), \mathcal{L} is a left congruence on $E(I, J, \alpha, \beta, A)$ by a routine calculation.

LEMMA 3.8. $E(I, J, \alpha, \beta, A)$ is an ω -chain of rectangular bands $(E_n : n \in N)$ where $E_n = ((i, (n, n), j) : i \in I_n, j \in E(J_n))$.

Proof. Let $(i, (n, n), j), (u, (n, n), v) \in E_n$. Utilizing (6), (2), (3), and a routine calculation, $(i, (n, n), j)(u, (n, n), v) = (i, (n, n), v)$. Hence, E_n is a rectangular band. Again, utilizing (6), (2), (3), and a routine calculation, $E_n E_k \subseteq E_{\max(n, k)}$.

THEOREM 3.9. (I, J, α, β, A) is a generalized ω - \mathcal{L} -unipotent bisimple semigroup.

Proof. Combine Lemmas 3.5–3.8.

4. Structure of generalized ω - \mathcal{L} -unipotent bisimple semigroups. Combining Theorems 4.1 and 4.3 (below) will give a description of generalized ω - \mathcal{L} -unipotent bisimple semigroups in terms of groups, ω -chains of left zero semigroups, and ω -chains of right zero semigroups.

THEOREM 4.1. (I, J, α, β, A) is a generalized ω - \mathcal{L} -unipotent bisimple semigroup, and conversely every such semigroup is isomorphic to some (I, J, α, β, A) .

Proof. Combine Theorems 3.9 and 2.21.

REMARK. In contrast to the structure theorem for generalized \mathcal{L} -unipotent semigroups given in [4], no factor systems are required in Theorem 4.1.

We will next characterize an ω -chain J of right groups $(J_n: n \in N)$ as a semi-direct product of an ω -chain X of right zero semigroups $(X_n: n \in N)$ by an ω -chain G of groups $(G_n: n \in N)$.

We first need a definition.

DEFINITION 4.2. Let the semigroup U be an ω -chain of semigroups $(U_n: n \in N)$ and let θ be a mapping of U into a semigroup V such that $r \in U_n, s \in U_m$, and $m \geq n$ imply $(rs)\theta = r\theta s\theta$. We term θ a lower homomorphism of U into V .

Let (G, o) be an ω -chain of groups $(G_n: n \in N)$ and let $(X, *)$ be an ω -chain of right zero semigroups $(X_n: n \in N)$ such that $G_n \cap X_n = (e_n)$, a single idempotent element, for each $n \in N$. Let $g \mapsto B_g$ be a lower homomorphism of G into T_X subject to the conditions (1) $X_n B_g \subseteq X_{\max(n, m)}$ if $g \in G_m$ (2) if $r \in X_m, s \in X_n$ and $m \geq n$, $(r^*s)B_g = rB_{e_n \circ g^*} B_g$. Let (G, X, B) denote $\bigcup (G_n \times X_n: n \in N)$ under the multiplication $(i, j)(p, q) = (i \circ p, jB_p^*q)$.

THEOREM 4.3. J is an ω -chain of right groups if and only if $J \cong (G, X, B)$ for some collection G, X, B .

Proof. We just specialize [6, Theorem 7.2].

Note 4.4. The structure of G is known mod groups and homomorphisms by a well known result of Clifford [1, Theorem 4.11].

5. ω - \mathcal{L} -unipotent bisimple semigroups. In this section, we specialize Theorem 4.1 to obtain [5, Theorem 7.11] (our previous structure theorem for ω - \mathcal{L} -unipotent bisimple semigroups).

A bisimple semigroup S is termed ω - \mathcal{L} -unipotent if $E(S)$ is an ω -chain of right zero semigroups.

THEOREM 5.1. *Let S be an ω - \mathcal{L} -unipotent bisimple semigroup. Then, there exists an ω -chain $(J, *)$ of right groups $(J_n; n \in N)$ and a homomorphism $(n, r) \rightarrow \beta_{(n,r)}$ of C into $\text{End}(J, *)$ such that for each $k \in N$ there exists $e_k \in E(J_k)$ and*

$$(1) \quad g\beta_{(k,k)} = g^*e_k \text{ for all } g \in J.$$

(2) $J_r\beta_{(n,k)} \subseteq J_{k+r-\min(n,r)}$. Furthermore, $S \cong (((n, k), j): j \in J_k, n, k \in N)$ under the multiplication.

(3) $((n, k), j)((r, s), v) = ((n, k)(r, s), j\beta_{(r,s)}^*v)$ where juxtaposition denotes multiplication in C .

Conversely, let $(J, *)$ be an ω -chain of right groups and let $(n, r) \rightarrow \beta_{(n,r)}$ be a homomorphism of C into $\text{End}(J, *)$ such that (1) and (2) are valid. Then, $S = (((n, k), j): j \in J_k, n, k \in N)$ under (3) is an ω - \mathcal{L} -unipotent bisimple semigroup.

Proof. We first establish the converse. We employ Theorem 4.1 and its notation. Let $I_v = (e_v)$ for each $v \in N$ and define $e_u \circ e_v = e_{\max(u,v)}$. Let $I = \bigcup (I_v: v \in N)$. Then, (I, \circ) is an ω -chain of left zero semigroups $(I_n; n \in N)$. Define $e_n\alpha_{(r,s)} = e_{s+n-\min(n,r)}$ and $e_nA_v = e_{\max(n,m)}$ if $v \in J_m$. By a routine calculation, $(n, r) \rightarrow \alpha_{(n,r)}$ is a homomorphism of C into $\text{End}(I, \circ)$ and $p \rightarrow A_p$ is an upper anti-homomorphism of $(J, *)$ into T_I such that (2)–(5) of Theorem 4.1 is valid. The multiplication (6) of Theorem 4.1 becomes (6') $(e_n, (n, k), j)(e_r, (r, s), v) = (e_{n+r-\min(k,r)}, (n, k)(r, s), j\beta_{(r,s)}^*v)$ where juxtaposition is multiplication in C . Hence, $U = (I, J, \alpha, \beta, A)$ (notation of §3) is a generalized ω - \mathcal{L} -unipotent bisimple semigroup by Theorem 4.1. Utilizing Lemma 3.4, $E(U) = ((e_n, (n, n), j): j \in E(J_n), n \in N)$. Utilizing Lemma 3.2, $(e_n, (n, n), j)\mathcal{L}(e_k, (k, k), u)(j \in E(J_n)$ and $u \in E(J_k))$ implies $n = k$ and $j = u$. Hence, $E(U)$ is an ω -chain of right zero semigroups and, thus, U is an ω - \mathcal{L} -unipotent bisimple semigroup. Since $(e_n, (n, k), j)\varphi = ((n, k), j)$ define an isomorphism of $(U, (6'))$ onto $(S, (3))$. S is an ω - \mathcal{L} -unipotent bisimple semigroup.

Next, let T be an ω - \mathcal{L} -unipotent bisimple semigroup. Hence, T is a generalized ω - \mathcal{L} -unipotent bisimple semigroup and the structure of T is given by Theorem 4.1. Thus, utilizing Lemmas 3.8 and 3.2, $I_n = (e_n)$ for each $n \in N$. Hence, utilizing (2) and (3) of Theorem 4.1, $e_r\alpha_{(n,k)} = e_{k+r-\min(n,r)}$ and $e_rA_j = e_{\max(r,k)}$ if $j \in J_k$. Thus, (6) of Theorem 4.1 becomes (6') and $(U, (6')) \cong (S, (3))$. The conditions of Theorem 5.1 are given by Theorem 4.1 ((1) and (2)).

REFERENCES

1. A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, Math. Surveys of the Amer. Math. Soc., 7, Vol. 1, Providence, R. I., 1961; Vol. 2, Providence, R. I., 1967.
2. David McLean, *Idempotent semigroups*, Amer. Math. Monthly, **61** (1954), 110-113.
3. R. J. Warne, *\mathcal{L} -unipotent semigroups*, Nigerian J. Science, **5** (1972), 245-248.
4. ———, *Generalized \mathcal{L} -unipotent semigroups*, Bulletino della Unione Matematica Italiana, **5** (1972), 43-47.
5. ———, *ωY - \mathcal{L} -unipotent semigroups*, to appear in JÑÑABHA.
6. ———, *On the structure of semigroups which are unions of groups*, Trans. Ameer. Math. Soc., **186** (1973), 385-401.

Received February 1, 1973.

UNIVERSITY OF ALABAMA IN BIRMINGHAM

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Robert F. V. Anderson, <i>Laplace transform methods in multivariate spectral theory</i>	339
William George Bade, <i>Two properties of the Sorgenfrey plane</i>	349
John Robert Baxter and Rafael Van Severen Chacon, <i>Functionals on continuous functions</i>	355
Phillip Wayne Bean, <i>Helly and Radon-type theorems in interval convexity spaces</i>	363
James Robert Boone, <i>On k-quotient mappings</i>	369
Ronald P. Brown, <i>Extended prime spots and quadratic forms</i>	379
William Hugh Cornish, <i>Crawley's completion of a conditionally upper continuous lattice</i>	397
Robert S. Cunningham, <i>On finite left localizations</i>	407
Robert Jay Daverman, <i>Approximating polyhedra in codimension one spheres embedded in S^n by tame polyhedra</i>	417
Burton I. Fein, <i>Minimal splitting fields for group representations</i>	427
Peter Fletcher and Robert Allen McCoy, <i>Conditions under which a connected representable space is locally connected</i>	433
Jonathan Samuel Golan, <i>Topologies on the torsion-theoretic spectrum of a noncommutative ring</i>	439
Manfred Gordon and Edward Martin Wilkinson, <i>Determinants of Petrie matrices</i>	451
Alfred Peter Hallstrom, <i>A counterexample to a conjecture on an integral condition for determining peak points (counterexample concerning peak points)</i>	455
E. R. Heal and Michael Windham, <i>Finitely generated F-algebras with applications to Stein manifolds</i>	459
Denton Elwood Hewgill, <i>On the eigenvalues of a second order elliptic operator in an unbounded domain</i>	467
Charles Royal Johnson, <i>The Hadamard product of A and A^*</i>	477
Darrell Conley Kent and Gary Douglas Richardson, <i>Regular completions of Cauchy spaces</i>	483
Alan Greenwell Law and Ann L. McKerracher, <i>Sharpened polynomial approximation</i>	491
Bruce Stephen Lund, <i>Subalgebras of finite codimension in the algebra of analytic functions on a Riemann surface</i>	495
Robert Wilmer Miller, <i>TTF classes and quasi-generators</i>	499
Roberta Mura and Akbar H. Rhemtulla, <i>Solvable groups in which every maximal partial order is isolated</i>	509
Isaac Namioka, <i>Separate continuity and joint continuity</i>	515
Edgar Andrews Rutter, <i>A characterization of $QF - 3$ rings</i>	533
Alan Saleski, <i>Entropy of self-homeomorphisms of statistical pseudo-metric spaces</i>	537
Ryōtarō Satō, <i>An Abel-maximal ergodic theorem for semi-groups</i>	543
H. A. Seid, <i>Cyclic multiplication operators on L_p-spaces</i>	549
H. B. Skerry, <i>On matrix maps of entire sequences</i>	563
John Brendan Sullivan, <i>A proof of the finite generation of invariants of a normal subgroup</i>	571
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, VI</i>	573
Ronson Joseph Warne, <i>Generalized $\omega - \mathcal{L}$-unipotent bisimple semigroups</i>	631
Toshihiko Yamada, <i>On a splitting field of representations of a finite group</i>	649