

Pacific Journal of Mathematics

CATEGORY THEORY APPLIED TO PONTRYAGIN DUALITY

DAVID W. ROEDER

CATEGORY THEORY APPLIED TO PONTRYAGIN DUALITY

DAVID W. ROEDER


A proof of the Pontryagin duality theorem for locally compact abelian (LCA) groups is given, using category-theoretical ideas and homological methods. The proof is guided by the structure within the category of LCA groups and does not use any deep results except for the Peter-Weyl theorem. The duality is first established for the subcategory of elementary LCA groups (those isomorphic with $T^i \oplus Z^j \oplus R^k \oplus F$, where T is the circle group, Z the integers, R the real numbers, and F a finite abelian group), and through the study of exact sequences, direct limits and projective limits the duality is expanded to larger subcategories until the full duality theorem is reached.

Introduction. In this note we present a fairly economical proof of the Pontryagin duality theorem for locally compact abelian (LCA) groups, using category-theoretic ideas and homological methods. This theorem was first proved in a series of papers by Pontryagin and van Kampen, culminating in van Kampen's paper [5], with methods due primarily to Pontryagin. In [10, pp. 102-109], Weil introduced the simplifying notion of compactly generated group and explored the functorial nature of the situation by examining adjoint homomorphisms and projective limits. Proofs along the lines of Pontryagin-van Kampen-Weil appear in the books by Pontryagin [7, pp. 235-279] and Hewitt and Ross [2, pp. 376-380]. A different proof based on abstract Fourier analysis was given by Cartan and Godement [1]; similar methods are also used by Rudin [9, pp. 27-29] and Heyer [3, pp. 148-161]. Negrepontis [6, pp. 239-252] presented the theorem in light of category theory, but for the most part used different methods and more structure theory than is used here.

The proof we present is based on ideas used previously by Hofmann [4, pp. 109-117] and the author [8]. It rests neither on the structure theorem for compactly generated LCA groups (see [2], [6], [7]) nor on the structure of the L^1 -algebra of G (see [6] and [9]). The only deep result we require is the classical Peter-Weyl theorem, applied in our case to compact abelian groups, of course.

Definitions and preliminaries. L is the category of LCA groups, with continuous homomorphisms as morphisms. Groups will be written additively. Pointwise addition of functions makes each $\text{Hom}(G, H)$ an abelian group and L an additive category. R denotes the additive real numbers with the usual topology, Z is the subgroup of R con-

sisting of the integers with the induced (discrete) topology, T is the compact quotient group R/Z , and Z_n is Z/nZ . If $G \in L$, then the character group of G is $\chi(G) = G^* = \text{Hom}(G, T)$, taken with the compact-open topology, which makes G^* an LCA group. For $f \in \text{Hom}(G, H)$, the adjoint homomorphism $\chi(f) = f^* \in \text{Hom}(H^*, G^*)$ is defined by $f^*(\alpha) = \alpha \circ f$ for $\alpha \in H^*$. Thus χ is a contravariant functor from L to L . There is a natural transformation ρ from the identity functor η in L to the covariant functor $\chi^2 = \chi \circ \chi$; we specify $\rho_G: G \rightarrow G^{**}$ by $[\rho_G(x)](\alpha) = \alpha(x)$ for $x \in G, \alpha \in G^*$. The Pontryagin duality theorem is now the statement that ρ is a natural isomorphism; i.e., that each ρ_G is an isomorphism between G and its second character group G^{**} . The method of proof will be to show that ρ restricted to a suitable subcategory is an isomorphism, and then enlarge the subcategory in successive steps until we have the theorem proved. We indicate this schematically in the following diagram, where each name stands for the full subcategory of L with the named property, and the labels on connecting lines indicate the methods used to extend the duality to the larger category.



First, $T \simeq Z^*$ with $x \in T$ corresponding to the character $n \mapsto nx$ of Z . Conversely, $Z \simeq T^*$ with $n \in Z$ corresponding to the character $x \mapsto nx$ of T . Also $R \simeq R^*$ with $x \in R$ corresponding to the character $y \mapsto xy + Z$ of R . And $Z_n \simeq Z_n^*$ by corresponding $x \in Z_n$ to the character $y \mapsto xy$ of Z_n . Thus the relation $G \simeq G^{**}$ holds in a natural way for $G = T, Z, R$, or Z_n , respectively, and in each case the isomorphism obtained is precisely the map ρ_G defined earlier.

It is easy to show that χ is additive (that is, for $f, g \in \text{Hom}(G, H)$ we have $\chi(f + g) = \chi(f) + \chi(g)$); as a consequence χ preserves direct sums. (See Lemma 2, p. 162, in [8].) From this we see that Pontryagin duality holds at least for the elementary groups in L ;

that is, groups isomorphic to $T^i \oplus Z^j \oplus R^k \oplus F$, where F is a finite abelian group. The full subcategory of elementary groups in L will be denoted by E . Then we have the following:

THEOREM 1. $\rho \mid E: \eta \mid E \rightarrow \chi^2 \mid E$ is an isomorphism.

Duality for compact groups in L . We define a morphism $f \in \text{Hom}(G, H)$ to be *proper* if f is open considered as a function from G to its image in H . (Thus a "proper morphism" as the term is used here may not be a "proper map" as the term is used in general topology.)

PROPOSITION 1. *If $f: G \rightarrow H$ is proper and $f(G)$ is open in H , then f^* is proper.*

Proof. Let M be a compact neighborhood of 0 in H and W a closed neighborhood of 0 in T which contains no proper subgroups of T . (M, W) denotes the set of all characters α in H^* for which $\alpha(M) \subset W$. The set of (M, W) obtained from all possible such choices of M and W forms a base of compact neighborhoods of 0 in H^* . Suppose now that M is chosen to be a compact neighborhood of 0 in $f(G)$. Since f is proper and G is locally compact, we may find a compact neighborhood N of 0 in G such that $f(N) = M$. Then (N, W) is a neighborhood of 0 in G^* and $f^*(M, W) = (N, W) \cap f^*(H^*)$ is a compact neighborhood of 0 in $f^*(H^*)$. The properness of f^* follows from the open mapping theorem [2, Theorem 5.29, p. 42] applied to the open subgroup of H^* generated by (M, W) .

PROPOSITION 2. χ takes a short proper exact sequence $0 \rightarrow K \xrightarrow{i} G \xrightarrow{j} H \rightarrow 0$ to a sequence $0 \leftarrow K^* \xleftarrow{i^*} G^* \xleftarrow{j^*} H^* \leftarrow 0$ in which j^* is proper and exactness holds at G^* and H^* . If in addition K is an open subgroup of G , then the sequence induced by χ is also proper exact.

Proof. It is now easy to see that H^* is isomorphic to the closed subgroup of G^* consisting of those characters of G which are trivial on K , via the proper injection j^* . The kernel of i^* is also the set of characters on G which are trivial on K . This proves exactness at G^* and H^* . Now, because T is a divisible group, any character on H extends to a homomorphism (not necessarily continuous) from G to T . If H is an open subgroup, then any such extension will be continuous on G , so the proper map i^* will be a surjection.

Using the (M, W) notation of the proof of Proposition 1, we see that if G is discrete and $M = \{0\}$ in G , then $(M, W) = G^*$, so G^* is

compact. On the other hand, if G is compact and $M = G$, then $(M, W) = \{0\}$, so G^* is discrete. We now let A be the subcategory of discrete groups in L and A_0 the subcategory of discrete elementary groups (isomorphic to $Z^i \oplus F$). C is the subcategory of compact groups and C_0 the subcategory of compact elementary groups (isomorphic to $T^i \oplus F$) in L .

We can make a directed set I into a category by declaring, for $i, j \in I$, that $\text{Hom}(i, j)$ consist of exactly one element if $i \leq j$ and be empty otherwise. Then, for our purposes, a direct system in A_0 is a covariant functor U from a directed set to A_0 . We shall often write U_i for $U(i)$ and u_{ij} for $U(\text{Hom}(i, j))$. We let DA_0 be the collection of direct systems in A_0 whose morphisms are all injective. DA_0 becomes a category when we define a morphism from $U: I \rightarrow A_0$ to $V: J \rightarrow A_0$ to be a pair (m, φ) , where $m: I \rightarrow J$ is a functor (order-preserving map) and φ is a natural transformation from U to $V \circ m$.

Well-known properties of abelian groups include the fact that each element U of DA_0 has a direct limit $\varinjlim U$ (or $\varinjlim U_i$) which will be an object in D . In fact, a necessary and sufficient condition for a discrete group G to be isomorphic to $\varinjlim U$ is the existence of injective morphisms $g_i: U_i \rightarrow G$, one for each i , such that $g_j \circ u_{ij} = g_i$ whenever $i \leq j$, and the union of the images $g_i(U_i)$ is all of G . The result is that $\varinjlim: DA_0 \rightarrow A$ can be regarded as a covariant functor, since if (m, φ) is a functor from U to V in DA_0 , the universal property of $\varinjlim U$ guarantees existence of a unique morphism $\varinjlim (m, \varphi): \varinjlim U \rightarrow \varinjlim V$ making the following diagram commutative for every $i \in I$:

$$\begin{array}{ccc} U_i & \longrightarrow & \varinjlim U \\ \downarrow \varphi_i & & \downarrow \varinjlim (m, \varphi) \\ V_{m(i)} & \longrightarrow & \varinjlim V. \end{array}$$

Similarly, an inverse system in C_0 is a contravariant functor U from a directed set to C_0 . The category of all inverse systems in C_0 all of whose morphisms are surjective is denoted by IC_0 . A morphism from $U: I \rightarrow C_0$ to $V: J \rightarrow C_0$ in IC_0 is a pair (m, φ) where $m: J \rightarrow I$ is a functor and $\varphi: U \circ m \rightarrow V$ is a natural transformation. Also, any inverse system in IC_0 has a projective limit $\varprojlim U$ which will be an object of C . Further, any object G of C is isomorphic to $\varprojlim U$ if and only if there exists a surjective morphism $g_i: G \rightarrow U_i$ for each i such that $u_{ij} \circ g_j = g_i$ whenever $i \leq j$ and the intersection of the kernels of the g_i is $\{0\}$ in G . (The usual condition [10, p. 25] is that every neighborhood of 0 in G contain $\text{Ker } g_i$ for some i , but if

the intersection of the $\text{Ker } g_i$ is $\{0\}$ and N is any open neighborhood of 0 in G , then by compactness we can find a finite number of g_i the intersection of whose kernels is contained in N . Picking an index j greater than these i gives us $\text{Ker } g_j \subset N$.) Again in this situation $\varprojlim(m, \varphi)$ for a morphism (m, φ) in IC_0 is defined by the universal property, and \varprojlim becomes a covariant functor. We call DA_0 and IC_0 convergence structures on A and C , respectively.

By Proposition 2, if $U \in DA_0$ then $\chi \circ U \in IC_0$. This correspondence $U \mapsto \chi \circ U$ gives us a functor which we denote by $D\chi: DA_0 \rightarrow IC_0$. Similarly, χ induces the functor $I\chi: IC_0 \rightarrow DA_0$.

PROPOSITION 3. *χ is continuous on A with respect to DA_0 ; that is, the two functors $\chi \circ \varinjlim$ and $\varprojlim \circ D\chi: DA_0 \rightarrow C$ are naturally isomorphic.*

Proof. Let $U \in DA_0$ and $G = \varinjlim U_i$ with $g_i: U_i \rightarrow G$ the associated injections. We must show that $G^* \simeq \varinjlim (U_i^*)$. It is clear that each $g_i^*: G^* \rightarrow U_i^*$ is surjective. Let $0 \neq \alpha \in G^*$. We shall show the existence of an index i with $g_i^*(\alpha) \neq 0$. We know $\alpha(x) \neq 0$ for some $x \in G$, and then $x = g_i(y)$ for some i and $y \in U_i$. Then for this i , $g_i^*(\alpha)(y) = \alpha(g_i(y)) \neq 0$. So $(\varinjlim U_i)^* \simeq \varinjlim (U_i^*)$. The fact that we have a natural isomorphism follows from the universal property.

PROPOSITION 4. *χ is continuous on C with respect to IC_0 ; that is, the functors $\chi \circ \varprojlim$ and $\varinjlim \circ I\chi: IC_0 \rightarrow A$ are naturally isomorphic.*

Proof. Let $U \in IC_0$ and $G = \varprojlim U_i$ with $g_i: G \rightarrow U_i$ the associated surjections. Clearly each g_i^* is injective. To show that G^* is isomorphic with $\varinjlim (U_i^*)$, we must show that every $\alpha \in G^*$ is equal to $g_i^*(\beta)$ for some i and some $\beta \in U_i^*$. Let W be a neighborhood of 0 in T containing no proper subgroups of T . Let M be a neighborhood of 0 in G with $\alpha(M) \subset W$. Then we may find U_i with $\text{Ker } g_i \subset M$. Then $\alpha(\text{Ker } g_i) = 0$, so α factors; $\alpha = \beta \circ g_i$ for some $\beta \in U_i^*$. But $\beta \circ g_i = g_i^*(\beta)$, so we are done.

We remark that Proposition 3 above could be proved as in [6] by showing that χ has a left adjoint functor, which implies that χ takes direct limits to limits. However, Proposition 4 does not admit an analogous proof.

PROPOSITION 5. *The category C_0 is dense in C . That is, there is a functor $S: C \rightarrow IC_0$ such that the functor $\varprojlim \circ S$ and the identity functor on C are naturally isomorphic.*

Proof. This follows from the Peter-Weyl theorem [7, Theorem 33, p. 229], which says for our abelian case that the characters of any G in C separate the points of G . Let $G \in C$ and define $S(G) \in IC_0$ to be the collection of quotient groups G/K of G which are in C_0 . We order them by $G/K \leq G/N$ if $N \subset K$. Note that $G/(K \cap N)$ is isomorphic to a subgroup of $(G/K) \oplus (G/N)$, so we have a directed set. We can define $g_{KN}: G/N \rightarrow G/K$ to be the natural projection when $N \subset K$. For a morphism $f: G \rightarrow H$ in C , $S(f)$ is defined as follows: H/K in $S(H)$ corresponds to $G/f^{-1}(K)$ in $S(G)$, and the map $G/f^{-1}(K) \rightarrow H/K$ is the natural one induced by f . Then if $x \in G$, let $\alpha \in G^*$ with $\alpha(x) \neq 0$. Then G/K is in $S(G)$, where $K = \text{Ker } \alpha$, and $g_K(x) \neq 0$, where $g_K: G \rightarrow G/K$ is the natural map. The collection of g_K exhibits G as $\lim S(G)$.

←

We are now ready to consider our two covariant functors η and χ^2 again.

THEOREM 2. *Pontryagin duality holds in C ; that is, $\rho | C: \eta | C \rightarrow \chi^2 | C$ is an isomorphism.*

Proof. $\rho | C_0$ is already an isomorphism. By Proposition 1.18 in [4, p. 115], $\rho | C_0$ extends uniquely to a natural transformation between $\eta | C$ and $\chi^2 | C$. This extension must also be an isomorphism. But $\rho | C$ already extends $\rho | C_0$ so $\rho | C$ must be an isomorphism.

Duality of compactly generated groups in L . Let CG be the full subcategory of the compactly generated groups in L .

LEMMA. *Suppose $G \in CG$ is generated by the compact neighborhood M of 0 in G . Then there is a subgroup K of G , $K \simeq Z^n$ for some n , such that $K \cap M = \{0\}$ and G/K is compact.*

Proof. This is Lemma 2.42 in [9, p. 41].

PROPOSITION 6. *If $G \in CG$, then ρ_G is injective.*

Proof. This is an easy consequence of the above lemma. Let $x \in G$, $x \neq 0$. Apply the lemma to $M \cup \{x\}$, which is also a compact neighborhood of 0 which generates G . The coset $x + K$ is not the identity element in the compact group G/K . Therefore, there is a character α of G/K such that $\alpha(x + K) \neq 0$. Composing α with the natural projection $G \rightarrow G/K$ gives us a character on G which is not trivial on x . Therefore $\rho_G(x) \neq 0$.

THEOREM 3. *Pontryagin duality holds in CG .*

Proof. Let $G \in CG$. Let M be a compact neighborhood of 0 in G and let $S \simeq Z^n$ be a subgroup of G such that $S \cap M = \{0\}$ and G/S is compact, as guaranteed by the lemma above. Let $Q = G/S$ and $p: G \rightarrow Q$ the natural map. Let N be a compact symmetric neighborhood of 0 in G such that $N + N + N \subset M$. Then p maps N homeomorphically onto $p(N)$. Since Q is compact and p is proper, there is a compact subgroup Q_1 of Q such that $Q_1 \subset p(N)$ and Q/Q_1 is compact elementary, by Proposition 5. Then $p^{-1}(Q_1)$ is a closed subgroup of G contained in $S + N$. Letting $K = p^{-1}(Q_1) \cap N$, we have by the choice of N that K is a compact subgroup of G satisfying $p(K) = Q_1$.

Now let $H = G/K$; we shall show that H is an elementary group. First, p gives rise to proper surjection $H \rightarrow Q/Q_1$ with kernel $S + K$ which is discrete in H by the construction of K . Therefore H is locally isomorphic with Q/Q_1 , which in turn is locally isomorphic with R^n for some n . This means we have an isomorphism $f: B \rightarrow V$, where B is an open ball about 0 in R^n and V is a neighborhood of 0 in H . Then we can extend f to a proper surjective homomorphism $g: R^n \rightarrow H_1$, where H_1 is the open subgroup of H generated by V , by defining $g(x) = n \cdot f(x/n)$, for $x \in R^n$ and n large enough so that $x/n \in B$. Thus $H_1 \simeq R^a \oplus T^b$ for some integers a and b (a quotient group of R^n). Since H_1 is a divisible open subgroup of H , we can obtain a morphism $H \rightarrow H_1$ which is the identity on H_1 , so $H \simeq H_1 \oplus H/H_1$. But H/H_1 is an elementary group since it is discrete and compactly generated. Therefore H is also elementary. Now we examine the following commutative diagram:

$$(1) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & K & \xrightarrow{i} & G & \xrightarrow{j} & H & \longrightarrow & 0 \\ & & \downarrow & & \downarrow \rho_K & & \downarrow \rho_G & & \downarrow \rho_H \\ & & 0 & \longrightarrow & K^{**} & \xrightarrow{i^{**}} & G^{**} & \xrightarrow{j^{**}} & H^{**} \longrightarrow 0 \end{array}$$

Now $K \in C$ and $H \in E$, so ρ_K and ρ_H are isomorphisms, while ρ_G is injective by Proposition 6. Since i is injective, we conclude that i^{**} is injective. Now briefly consider $i^*: G^* \rightarrow K^*$. If i^* were not surjective, then there would be a nontrivial character on $K^*/i^*(G^*)$. Composing with the natural projection $K^* \rightarrow K^*/i^*(G^*)$, we get a character on K^* which is trivial on $i^*(G^*)$. This character would then be in the kernel of i^{**} , contradicting the injectivity of i^{**} . Therefore, i^* is surjective and Proposition 2 now tells us that the induced sequence

$$0 \longleftarrow K^* \xleftarrow{i^*} G^* \xleftarrow{j^*} H^* \longleftarrow 0$$

is a proper exact sequence, since K^* is discrete. Thus H^* can be regarded as an open subgroup of G^* , and so the lower sequence in

diagram (1) is also proper exact. The 5-lemma and the open mapping theorem show that ρ_G is an isomorphism, and we are done.

Duality for arbitrary locally compact groups. We begin with discrete groups.

PROPOSITION 7. *The category A_0 is dense in A . That is, there is a functor $T: A \rightarrow DA_0$ such that $\lim_{\rightarrow} T$ and the identity functor on A are naturally isomorphic.*

Proof. Every abelian group is the direct limit of its finitely generated subgroups. The functor T assigns to each G in A the direct system (U_i) of finitely generated subgroups of G ordered by $i \leq j$ if $U_i \subset U_j$. A morphism $f: G \rightarrow H$ in A is carried by T to $T(f)$, in which each finitely generated subgroup of G is mapped via the restriction of f to its image in H .

THEOREM 4. *Pontryagin duality holds in A .*

Proof. This is completely analogous to the proof of Theorem 2.

THEOREM 5. *The Pontryagin duality theorem. ρ is a natural equivalence.*

Proof. Let $G \in L$. Let M be a compact neighborhood of 0 in G and let K be the subgroup of G generated by M . Let $H = G/K$. Since K is open, the induced sequence

$$0 \longleftarrow K^* \longleftarrow G^* \longleftarrow H^* \longleftarrow 0$$

is proper exact by Proposition 2. Consider diagram (1) for this K , G , and H . We have exactness at K^{**} and G^{**} in the bottom row. H is discrete since K is open, so ρ_H is an isomorphism, and so j^{**} is surjective. Both rows of the diagram are proper exact and ρ_K is also an isomorphism. Again the 5-lemma applies and ρ_G is algebraically an isomorphism. But ρ_G restricted to the open subgroup K is the isomorphism ρ_K , so ρ_G is also an isomorphism.

Had we extended the concepts of convergence to include direct and projective limits of nondiscrete and noncompact groups, respectively, and had we modified Propositions 3 and 4 and their proofs accordingly, we could have proved Theorem 5 differently by showing that CG is dense in L and χ is continuous on L with respect to the modified convergence. Theorem 3 may be treated similarly; in fact, the proof of Theorem 3 already shows that E is dense in CG .

REFERENCES

1. H. Cartan and R. Godement, *Théorie de la dualité et analyse harmonique dans les groupes abéliens localement compacts*, Ann. Sci École Norm. Sup., (3) **64** (1947), 79-99.
2. E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis*, Volume I, Academic Press, New York and Springer-Verlag, Berlin, 1963.
3. H. Heyer, *Dualität Lokalkompakter Gruppen*, Lecture Notes in Mathematics, Volume 150, Springer-Verlag, New York, 1970.
4. K. H. Hofmann, *Categories with convergence, exponential functors, and the cohomology of compact abelian groups*, Math Z., **104** (1968), 106-140.
5. E. R. van Kampen, *Locally bicomact abelian groups and their character groups*, Ann. of Math., (2) **36** (1935), 448-463.
6. J. W. Negreponitis, *Duality in analysis from the point of view of triples*, J. Algebra, **19** (1971), 228-253.
7. L. S. Pontryagin, *Topological Groups*, Second Edition, Gordon and Breach, New York, 1966.
8. D. W. Roeder, *Functorial characterizations of Pontryagin duality*, Trans. Amer. Math. Soc., **154** (1971), 151-175.
9. W. Rudin, *Fourier Analysis on Groups*, Interscience, New York, 1962.
10. A. Weil, *L'intégration dans les groupes topologiques et ses applications*, Hermann, Paris, 1941 and 1951.

Received August 22, 1972 and in revised form January 26, 1973. This research was supported in part by the College Science Improvement Program of the National Science Foundation (NSF grant GY-5700).

COLORADO COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Harm Bart, <i>Spectral properties of locally holomorphic vector-valued functions</i>	321
J. Adrian (John) Bondy and Robert Louis Hemminger, <i>Reconstructing infinite graphs</i>	331
Bryan Edmund Cain and Richard J. Tondra, <i>Biholomorphic approximation of planar domains</i>	341
Richard Carey and Joel David Pincus, <i>Eigenvalues of seminormal operators, examples</i>	347
Tyrone Duncan, <i>Absolute continuity for abstract Wiener spaces</i>	359
Joe Wayne Fisher and Louis Halle Rowen, <i>An embedding of semiprime P.I.-rings</i>	369
Andrew S. Geue, <i>Precompact and collectively semi-precompact sets of semi-precompact continuous linear operators</i>	377
Charles Lemuel Hagopian, <i>Locally homeomorphic λ connected plane continua</i>	403
Darald Joe Hartfiel, <i>A study of convex sets of stochastic matrices induced by probability vectors</i>	405
Yasunori Ishibashi, <i>Some remarks on high order derivations</i>	419
Donald Gordon James, <i>Orthogonal groups of dyadic unimodular quadratic forms. II</i>	425
Geoffrey Thomas Jones, <i>Projective pseudo-complemented semilattices</i>	443
Darrell Conley Kent, Kelly Denis McKennon, G. Richardson and M. Schroder, <i>Continuous convergence in $C(X)$</i>	457
J. J. Koliha, <i>Some convergence theorems in Banach algebras</i>	467
Tsang Hai Kuo, <i>Projections in the spaces of bounded linear operations</i>	475
George Berry Leeman, Jr., <i>A local estimate for typically real functions</i>	481
Andrew Guy Markoe, <i>A characterization of normal analytic spaces by the homological codimension of the structure sheaf</i>	485
Kunio Murasugi, <i>On the divisibility of knot groups</i>	491
John Phillips, <i>Perturbations of type I von Neumann algebras</i>	505
Billy E. Rhoades, <i>Commutants of some quasi-Hausdorff matrices</i>	513
David W. Roeder, <i>Category theory applied to Pontryagin duality</i>	519
Maxwell Alexander Rosenlicht, <i>The nonminimality of the differential closure</i>	529
Peter Michael Rosenthal, <i>On an inversion theorem for the general Mehler-Fock transform pair</i>	539
Alan Saleski, <i>Stopping times for Bernoulli automorphisms</i>	547
John Herman Scheuneman, <i>Fundamental groups of compact complete locally affine complex surfaces. II</i>	553
Vashishtha Narayan Singh, <i>Reproducing kernels and operators with a cyclic vector. I</i>	567
Peggy Strait, <i>On the maximum and minimum of partial sums of random variables</i>	585
J. L. Brenner, <i>Maximal ideals in the near ring of polynomials modulo 2</i>	595
Ernst Gabor Straus, <i>Remark on the preceding paper: "Ideals in near rings of polynomials over a field"</i>	601
Masamichi Takesaki, <i>Faithful states on a C^*-algebra</i>	605
R. Michael Tanner, <i>Some content maximizing properties of the regular simplex</i>	611
Andrew Bao-hwa Wang, <i>An analogue of the Paley-Wiener theorem for certain function spaces on $SL(2, \mathbb{C})$</i>	617
James Juei-Chin Yeh, <i>Inversion of conditional expectations</i>	631