

Pacific Journal of Mathematics

**ON CHARACTERIZING CERTAIN CLASSES OF FIRST
COUNTABLE SPACES BY OPEN MAPPINGS**

KENNETH ABERNETHY

ON CHARACTERIZING CERTAIN CLASSES OF FIRST COUNTABLE SPACES BY OPEN MAPPINGS

KENNETH ABERNETHY

This paper has three main results. These are characterizations of Nagata spaces, γ -spaces, and semi-metric spaces, respectively, as images of metrizable spaces under certain kinds of continuous open mappings.

1. Introduction. A basic area of research in general topology is the study of how various classes of spaces are related through mappings (see [3] and [5]). More specifically, many important classes of spaces have been characterized as the image of a metrizable space under an open continuous mapping of some sort. For example, Heath [10] has characterized developable spaces in this way and Hanai and Ponomarev independently have given an elegant characterization of first countable spaces (see Theorem 2.1). In recent years considerable attention has been given to the problem of characterizing generalized metrizable spaces in this way. We mention some of these results in § 2. In this paper we characterize Nagata, semi-metric, and γ -spaces as the image of a metrizable space under certain types of open continuous mappings. Definitions and some known results are given in § 2, Nagata spaces are characterized with Theorem 3.3, γ -spaces with Theorem 4.3, and semi-metric spaces with Theorem 5.3. Throughout the paper the set of natural numbers will be denoted by N .

2. Definitions and background results. The spaces which interest us in this paper can be described in terms of sequences of open covers. It should be pointed out that many of the definitions which follow are not the original definitions, but are actually characterizations which were proved, after the particular concept had been introduced, in efforts to unify the various concepts. Consequently, the definitions we give, in terms of a *COC*-function, display some degree of this unification.

Let (X, T) be a topological space and let g be a function from $N \times X$ into T . Then g is called a *COC-function* for X (*COC* = countably many open covers) if it satisfies these two conditions: (1) $x \in \bigcap_{n=1}^{\infty} g(n, x)$ for all $x \in X$; (2) $g(n+1, x) \subseteq g(n, x)$ for all $n \in N$ and $x \in X$. Note that if g is a *COC-function* for X , we obtain countably many open covers of X , $\langle G_n \rangle$, by taking $G_n = \{g(n, x) : x \in X\}$ for each n .

Now let X be a space with *COC-function* g , and consider the

following conditions on g :

(A) $y_n \in g(n, x)$ for each $n \in N$ implies that the sequence $\langle y_n \rangle$ has x as a cluster point.

(B) $g(n, x) \cap g(n, y_n) \neq \emptyset$ for each $n \in N$ implies that $\langle y_n \rangle$ has x as a cluster point.

(C) $y_n \in g(n, x)$ and $p_n \in g(n, y_n)$ for each $n \in N$ implies that $\langle p_n \rangle$ has x as a cluster point.

(D) $\{g(n, x): n = 1, 2, \dots\}$ is a fundamental system of neighborhoods for x , for each x , and $x \in g(n, y_n)$ for each $n \in N$ implies that $\langle y_n \rangle$ has x as a cluster point.

(E) If H is closed and $p \in \overline{\bigcup \{g(n, x): x \in H\}}$ for each $n \in N$, then $p \in H$.

(F) $x \in g(n, y_n)$ for each $n \in N$ implies that $\langle y_n \rangle$ has x as a cluster point.

If X is a space with a COC-function g satisfying (A), X is called a *first countable space* and g a *first countable function* for X ; X is called a *Nagata space* and g is called a *Nagata function* for X if g satisfies (B); if g satisfies (C), X is called a γ -*space* and g a γ -*function* for X ; if g satisfies (D), X is called a *semi-metric space* and g a *semi-metric function* for X ; X is called a *stratifiable space* and g a *stratifiable function* for X if g satisfies (E); and finally if g satisfies (F), X is called a *semi-stratifiable space* and g a *semi-stratifiable function* for X .

Ceder [6] first studied stratifiable spaces under the name " M_3 -spaces". Borges [4] renamed them "stratifiable" and investigated them in more detail. Creede [7] introduced semi-stratifiable spaces. Our definition of semi-metric spaces is a characterization given by Heath in [9] where he studies semi-metric spaces. Hodel [12] introduced γ -spaces. Ceder [6] also introduced Nagata spaces, but our definition is a characterization due to Heath [9].

It is clear from our definitions that a space is a semi-metric space if and only if it is a first countable semi-stratifiable space. Also, it is true (c.f. [4]) that a space is a Nagata space if and only if it is a first countable stratifiable space.

Each class of spaces which we characterize below is included in the class of first countable spaces. Consequently we are able to make use of the following theorem proved independently by Hanai [8] and Ponomarev [16].

THEOREM 2.1. *A T_1 -space Y is first countable if and only if there is a metrizable space X and a continuous open mapping from X onto Y .*

Let us now mention some characterizations of Nagata spaces

and semi-metric spaces. Heath [10] has characterized each of these as follows.

THEOREM 2.2. *A T_1 -space is a semi-metric space if and only if there exists a continuous open mapping ψ from some metric space (X, d) onto Y , and a subset X' of X such that (1) $\psi(X') = Y$ and (2) if $y \in Y$, W an open set containing y , then there exists an $\varepsilon > 0$ such that $\psi(\text{Bd}(\psi^{-1}(y), \varepsilon) \cap X') \subseteq W$.*

THEOREM 2.3. *A T_2 -space Y is a Nagata space if and only if there exists an open continuous mapping ψ from some metric space (X, d) onto Y and a subset X' of X such that (1) $\psi(X') = Y$ and (2) if K is compact in Y , W an open set with $K \subseteq W$, then there exists an $\varepsilon > 0$ such that $\psi(\text{Bd}(\psi^{-1}(K), \varepsilon) \cap X') \subseteq W$.*

Nagata [15] has recently given quite similar characterizations of Nagata spaces and semi-metric spaces using the concept of a q -closed mapping. In comparing these results (of both Heath and Nagata) with results such as Theorem 2.1 and Heath's characterization of developable spaces, we can see that one natural way to try to improve them is to avoid having to consider a subset X' of the metric space X .

3. Nagata spaces.

DEFINITION 3.1. Let X and Y be topological spaces, let $\psi: X \rightarrow Y$ be a surjection, and let g be a COC-function for X . Then ψ is an N -mapping relative to g (N =Nagata) if given any $y \in Y$ and neighborhood W of y , there is a neighborhood V of y and a positive integer n such that if $g(n, x) \cap \psi^{-1}(V) \neq \emptyset$, then $\psi(x) \in W$. A surjection $\psi: X \rightarrow Y$ is an N -mapping if there is a COC-function g for X such that ψ is an N -mapping relative to g .

We note that our N -mapping is quite similar to Arhangel'skii's [3] regular mapping. Indeed our definition was suggested by his definition. In [3] he proved a theorem showing that conditions on the range space of a mapping can force the mapping to be regular. This theorem motivated the following proposition on N -mappings.

PROPOSITION 3.2. *Let (X, T) and Y be topological spaces with Y a stratifiable space, and let $\psi: X \rightarrow Y$ be a continuous surjection. Then ψ is an N -mapping.*

Proof. Let h be a stratifiable function for Y , and define $g: N \times X \rightarrow T$ by $g(n, x) = \psi^{-1}[h(n, \psi(x))]$. Then g is a COC-function for X .

Now let $y \in Y$, and let W be an open set containing y . Then $Y - W$ is closed and $y \notin Y - W$; hence there exists an $n_0 \in N$ such that $y \notin \bigcup \{h(n_0, p) : p \in Y - W\}$. Let $V = Y - \bigcup \{h(n_0, p) : p \in Y - W\}$. Now if $g(n_0, x) \cap \psi^{-1}(V) \neq \emptyset$, then $h(n_0, \psi(x)) \cap V \neq \emptyset$. But this means that $\psi(x) \notin Y - W$, i.e., $\psi(x) \in W$.

Using Theorem 2.1 and Proposition 3.2, we are able to characterize Nagata spaces.

THEOREM 3.3. *Let Y be a T_1 -space. Then Y is a Nagata space if and only if there is a metrizable space X and an open continuous N -mapping from X onto Y .*

Proof. First assume that Y is a Nagata space. Then Y is first countable; so by Theorem 2.1, there is a metrizable space X and an open continuous surjection from X onto Y . By Proposition 3.2, ψ is an N -mapping.

Now assume that X is metrizable and ψ is an open continuous N -mapping from X onto Y . Clearly Y is first countable, so it suffices to show that Y is stratifiable. Let g be a COC -function for X relative to which ψ is an N -mapping. Let $y \in Y$, $n \in N$. Then choose any $s \in \psi^{-1}(y)$ and define $h(n, y) = \psi[g(n, s)]$, for every n . We claim that h is a stratifiable function for Y . Let H be closed in Y , and suppose that $p \in \bigcup \{h(n, z) : z \in H\}$, for each $n \in N$. Suppose $p \in H$; then $p \in Y - H = W$, which is open. Thus there exist a neighborhood V of p and an $n_0 \in N$ such that if $g(n_0, x) \cap \psi^{-1}(V) \neq \emptyset$ then $\psi(x) \in W$.

Now since V is a neighborhood of p , $V \cap (\bigcup \{h(n, z) : z \in H\}) \neq \emptyset$ for each $n \in N$. Thus there is a $z \in H$ such that $h(n_0, z) \cap V \neq \emptyset$. Therefore, if t is such that $h(n_0, z) = \psi[g(n_0, t)]$, we have $g(n_0, t) \cap \psi^{-1}(V) \neq \emptyset$. But this implies that $\psi(t) = z \in W$, an obvious contradiction.

In reference to the remark at the end of § 2, we note that we are able to do away with having to look at a subset X' of the metric space X in our characterization of Nagata spaces. (A similar remark applies to our characterization of semi-metric spaces in § 5.)

4. γ -spaces. This section proceeds almost exactly as § 3. We begin by giving the definition of the kind of mapping we need in order to characterize γ -spaces.

DEFINITION 4.1. Let X and Y be topological spaces, let $\psi: X \rightarrow Y$ be a surjection, and let g be a COC -function for X . Then ψ is a G -mapping relative to g ($G = \text{gamma}$) if given any $y \in Y$ and neighbor-

hood W of y , there is a neighborhood V of y and an $n \in N$ such that $\psi[\bigcup \{g(n, x): x \in \psi^{-1}(V)\}] \subseteq W$. A surjection $\psi: X \rightarrow Y$ is a G -mapping if there is a COC-function g for X such that ψ is a G -mapping relative to g .

PROPOSITION 4.2. *Let (X, T) and Y be topological spaces with Y a γ -space, and let $\psi: X \rightarrow Y$ be a continuous surjection. Then ψ is a G -mapping.*

Proof. Let h be a γ -function for Y , and define g as in the proof of Proposition 3.2. Let $y \in Y$ and let W be a neighborhood of y . We claim that there exists an $n_0 \in N$ such that $\bigcup \{h(n_0, z): z \in h(n_0, y)\} \subseteq W$. For suppose not; then we can choose sequence $\langle z_n \rangle$ and $\langle u_n \rangle$ such that $z_n \in h(n, y)$ and $u_n \in h(n, z_n) - W$ for each $n \in N$. But since h is a γ -function, this means that y is a cluster point of $\langle u_n \rangle$; obviously a contradiction since $u_n \notin W$ for any $n \in N$.

Now let $V = h(n_0, y)$. Then we have $\psi[\bigcup \{g(n_0, x): x \in \psi^{-1}(V)\}] = \bigcup \{h(n_0, z): z \in h(n_0, y)\} \subseteq W$.

THEOREM 4.3. *Let Y be a T_1 -space. Then Y is a γ -space if and only if there is a metrizable space X and an open continuous G -mapping from X onto Y .*

Proof. Suppose that Y is a γ -space. Then by Theorem 2.1 there is a metrizable space X and an open continuous surjection $\psi: X \rightarrow Y$. By Proposition 4.2 ψ is a G -mapping.

On the other hand, suppose that X is metrizable and $\psi: X \rightarrow Y$ is an open continuous surjection, which is a G -mapping. Then Y is first countable. Let f be a first countable function for Y , and let $y \in Y$. Choose an $s \in \psi^{-1}(y)$ and define $h(n, y) = \psi[g(n, s)] \cap f(n, y)$. Now suppose $y_n \in h(n, p)$ and $x_n \in h(n, y_n)$ for each $n \in N$. Then we must show that p is a cluster point of $\langle x_n \rangle$. Assume not, and choose a neighborhood U of p and an integer k so that for $n \geq k$, $x_n \notin U$. Now there exists a neighborhood V of p and an n_0 (which we may choose to be $\geq k$ since $g(n+1, x) \subseteq g(n, x)$ for all x and n) such that $\psi[\bigcup \{g(n_0, x): x \in \psi^{-1}(V)\}] \subseteq U$. But we can choose an $m_0 \geq n_0$ such that $h(m_0, p) \subseteq V$. Then we have $\psi[\bigcup \{g(m_0, x): x \in \psi^{-1}(V)\}] \subseteq U$. Now since $y_{m_0} \in h(m_0, p)$, we get $\psi[g(m_0, s_{m_0})] \subseteq U$, where $h(m_0, y_{m_0}) = \psi[g(m_0, s_{m_0})] \cap f(m_0, y_{m_0})$ and $s_{m_0} \in \psi^{-1}(y_{m_0})$. But there is a $t_{m_0} \in \psi^{-1}(x_{m_0}) \cap g(m_0, s_{m_0})$ and so $\psi(t_{m_0}) = x_{m_0} \in U$, a contradiction.

5. Semi-metric spaces. Since the proofs of Proposition 5.2 and Theorem 5.3 are very similar to the proofs of Proposition 3.2 and Theorem 3.3 respectively, we omit them here.

DEFINITION 5.1. Let X and Y be topological spaces, let $\psi: X \rightarrow Y$ be a surjection, and let g be a COC-function for X . Then ψ is an *SM-mapping relative to g* (*SM*=semi-metric) if given any $y \in Y$ and neighborhood W of y , there is an $n \in N$ such that $g(n, x) \cap \psi^{-1}(y) \neq \emptyset$ implies that $\psi(x) \in W$. A surjection $\psi: X \rightarrow Y$ is an *SM-mapping* if there is a COC-function g for X such that ψ is an *SM-mapping relative to g* .

We remark that our *SM-mapping* is a generalization of Ponomarev's [16] π -mapping. In fact if (X, d) is a metric space and $g(n, x) = B_d(x, 1/n)$ then a π -mapping and an *SM-mapping relative to g* are identical concepts. Notice, however, that in Theorem 5.3 we get an *SM-mapping relative to a COC-function g* which is not directly related to the metric on X . Consequently, that *SM-mapping* is not necessarily a π -mapping. In connection with this remark, the question arises as to whether we can find characterizations similar to the ones we give, but with mappings which are directly related to the metric on X . Such characterizations, if they exist, would in a sense be improvements of our theorems.

PROPOSITION 5.2. Let (X, T) and Y be topological spaces with Y a semi-stratifiable space, and let $\psi: X \rightarrow Y$ be a continuous surjection. Then ψ is an *SM-mapping*.

THEOREM 5.3. Let Y be a T_1 -space. Then Y is a semi-metric space if and only if there is a metrizable space X and an open continuous *SM-mapping* from X onto Y .

In addition to comparing this characterization with those of Heath and Nagata mentioned above, the reader should compare it with results by Alexander [1] and Burke [5].

6. Some properties of the mappings. We note that each of the kinds of mappings we have defined in this paper (*N-mapping*, *G-mapping*, and *SM-mapping*) is countably productive. We can use this property to get relatively simple proofs that Nagata spaces, γ -spaces, and semi-metric spaces are countably productive as follows (the results are known, with the possible exception of the γ -space case).

THEOREM 6.1. Let $\{Y_n: n = 1, 2, \dots\}$ be a sequence of T_1 -spaces.

- (1) If each Y_n is a Nagata space, so is $\prod_{n=1}^{\infty} Y_n$.
- (2) If each Y_n is a γ -space, so is $\prod_{n=1}^{\infty} Y_n$.
- (3) If each Y_n is a semi-metric space, so is $\prod_{n=1}^{\infty} Y_n$.

Proof. (1) By Theorem 3.3 for each n , there is a metric space X_n and an open continuous N -mapping $\psi_n: X_n \rightarrow Y_n$. Then $\prod_{n=1}^{\infty} X_n$ is a metric space and $\prod_{n=1}^{\infty} \psi_n$ is an open continuous mapping from $\prod_{n=1}^{\infty} X_n$ onto $\prod_{n=1}^{\infty} Y_n$. It is not difficult to show that this mapping is in fact an N -mapping. Consequently, again by Theorem 3.3, $\prod_{n=1}^{\infty} Y_n$ is a Nagata space.

The proofs of (2) and (3) are similar.

Now we ask when finite-to-one and compact mappings are SM -mappings, and derive several corollaries concerning images under these mappings.

THEOREM 6.2. *Let $\psi: X \rightarrow Y$ be a continuous finite-to-one surjection, and let g be a semi-stratifiable function for X . Then ψ is a SM -mapping relative to g .*

Proof. Let $y \in Y$, W an open set in Y containing y . Suppose that $\psi^{-1}(y) = \{x_1, \dots, x_k\}$. For each i , $1 \leq i \leq k$, there exists an n_i such that $x_i \notin \cup \{g(n_i, p): p \in X - \psi^{-1}(W)\}$. Let $n_0 = \max\{n_i: i=1, \dots, k\}$. Then (recall that g is decreasing) if $g(n_0, x) \cap \psi^{-1}(y) \neq \emptyset$, we know $x \notin X - \psi^{-1}(W)$, i.e., $\psi(x) \in W$.

THEOREM 6.3. *Let $\psi: X \rightarrow Y$ be a continuous compact surjection, and let g be a K -semi-stratifiable function for X . Then ψ is a SM -mapping relative to g .*

The proof of Theorem 6.3 is very similar to the proof of Theorem 6.2. For the definition and a discussion of K -semi-stratifiable spaces, the reader should see Lutzer [13].

Next we note that we can weaken one implication of Theorem 5.3 to get the following.

THEOREM 6.4. *Let X be a space with a COC-function g , let $\psi: X \rightarrow Y$ be an almost-open SM -mapping relative to g . Then Y is semi-stratifiable.*

Proof. (Sketch). Let $y \in Y$, $n \in N$. Then there is an $s \in \psi^{-1}(y)$ with a system of neighborhoods $\{N_\alpha: \alpha \in A\}$ such that each $\psi(N_\alpha)$ is open. Now choose $N_{\alpha_n} \subseteq g(n, s)$ and define $h(n, y) = \psi(N_{\alpha_n})$. Then h can be shown to be a semi-stratifiable function for Y .

COROLLARY 6.5. *An almost-open continuous finite-to-one image of a semi-stratifiable space is semi-stratifiable.*

Proof. Combine Theorems 6.2. and 6.4.

It should be noted that Henry [11] has proved a slightly stronger result than Corollary 6.5 (pseudo-open rather than almost-open).

COROLLARY 6.6. *An almost-open continuous compact image of a K -semi-stratifiable space is semi-stratifiable.*

Proof. Combine Theorems 6.3 and 6.4.

REFERENCES

1. Charles C. Alexander, *Semi-developable spaces and quotient images of metric spaces*, Pacific J. Math., **37** (1971), 277-293.
2. A. V. Arhangel'skii, *On mappings of metric spaces*, Soviet Math. Doklady, **3** (1962), 953-956.
3. ———, *Mappings and spaces*, Russian Math. Surveys, **21** (1966), 115-162.
4. Carlos J. Borges, *On stratifiable spaces*, Pacific J. Math., **17** (1966), 1-16.
5. Dennis K. Burke, *Cauchy sequences in semi-metric spaces*, Proc. Amer. Math. Soc., **33** (1972), 161-165.
6. Jack Ceder, *Some generalizations of metric spaces*, Pacific J. Math., **11** (1961), 105-125.
7. Geoffrey C. Creede, *Concerning semi-stratifiable spaces*, Pacific J. Math., **32** (1970), 47-54.
8. S. Hanai, *On open mappings, II*, Proc. Japan Acad., **37** (1961), 233-238.
9. R. W. Heath, *Arc-wise connectedness in semi-metric spaces*, Pacific J. Math., **12** (1962), 1301-1319.
10. ———, *On open mappings and certain spaces satisfying the first countability axiom*, Fund. Math., **57** (1965), 91-96.
11. Michael Henry, *Stratifiable spaces, semi-stratifiable spaces and their relation through mappings*, Pacific J. Math., **37** (1971), 697-700.
12. R. E. Hodel, *Spaces defined by sequences of open covers which guarantee that certain sequences have cluster points*, Duke Math. J., **39** (1972), 253-265.
13. David Lutzer, *Semi-metrizable and stratifiable spaces*, General Topology and Appl., **1** (1971), 43-48.
14. E. A. Michael, *On representing spaces as images of metrizable and related spaces*, General Topology and Appl., **1** (1971), 329-344.
15. Jun-iti Nagata, *On generalized metric spaces and q -closed mappings*, to appear.
16. V. I. Ponomarev, *Axioms of countability and continuous mappings*, Bull. Pol. Akad. Nauk., **8** (1960), 127-134.

Received September 24, 1973 and in revised form December 1, 1973. The results in this paper are taken from the author's Ph.D. dissertation, done at Duke University under the supervision of R. E. Hodel. The author wishes to express his gratitude for Professor Hodel's help and encouragement.

WAYNESBURG COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Kenneth Abernethy, <i>On characterizing certain classes of first countable spaces by open mappings</i>	319
Ross A. Beaumont and Donald Lawver, <i>Strongly semisimple abelian groups</i>	327
Gerald A. Beer, <i>The index of convexity and parallel bodies</i>	337
Victor P. Camillo and Kent Ralph Fuller, <i>On Loewy length of rings</i>	347
Stephen LaVern Campbell, <i>Linear operators for which T^*T and TT^* commute.</i> <i>II</i>	355
Charles Kam-Tai Chui and Philip Wesley Smith, <i>Characterization of a function by certain infinite series it generates</i>	363
Allan L. Edelson, <i>Conjugations on stably almost complex manifolds</i>	373
Patrick John Fleury, <i>Hollow modules and local endomorphism rings</i>	379
Jack Tilden Goodykoontz, Jr., <i>Connectedness in kleinen and local connectedness in 2^X and $C(X)$</i>	387
Robert Edward Jamison, II, <i>Functional representation of algebraic intervals</i>	399
Athanassios G. Kartsatos, <i>Nonzero solutions to boundary value problems for nonlinear systems</i>	425
Soon-Kyu Kim, Dennis McGavran and Jingyal Pak, <i>Torus group actions on simply connected manifolds</i>	435
David Anthony Klarner and R. Rado, <i>Arithmetic properties of certain recursively defined sets</i>	445
Ray Alden Kunze, <i>On the Frobenius reciprocity theorem for square-integrable representations</i>	465
John Lagnese, <i>Existence, uniqueness and limiting behavior of solutions of a class of differential equations in Banach space</i>	473
Teck Cheong Lim, <i>A fixed point theorem for families on nonexpansive mappings</i> ...	487
Lewis Lum, <i>A quasi order characterization of smooth continua</i>	495
Andy R. Magid, <i>Principal homogeneous spaces and Galois extensions</i>	501
Charles Alan McCarthy, <i>The norm of a certain derivation</i>	515
Louise Elizabeth Moser, <i>On the impossibility of obtaining $S^2 \times S^1$ by elementary surgery along a knot</i>	519
Gordon L. Nipp, <i>Quaternion orders associated with ternary lattices</i>	525
Anthony G. O'Farrell, <i>Equiconvergence of derivations</i>	539
Dorte Olesen, <i>Derivations of AW^*-algebras are inner</i>	555
Dorte Olesen and Gert Kjærgaard Pedersen, <i>Derivations of C^*-algebras have semi-continuous generators</i>	563
Duane O'Neill, <i>On conjugation cobordism</i>	573
Chull Park and S. R. Paranjape, <i>Probabilities of Wiener paths crossing differentiable curves</i>	579
Edward Ralph Rozema, <i>Almost Chebyshev subspaces of $L^1(\mu; E)$</i>	585
Lesley Millman Sibner and Robert Jules Sibner, <i>A note on the Atiyah-Bott fixed point formula</i>	605
Betty Salzberg Stark, <i>Irreducible subgroups of orthogonal groups generated by groups of root type 1</i>	611
N. Stavrakas, <i>A note on starshaped sets, (k)-extreme points and the half ray property</i>	627
Carl E. Swenson, <i>Direct sum subset decompositions of Z</i>	629
Stephen Tefteller, <i>A two-point boundary problem for nonhomogeneous second order differential equations</i>	635
Robert S. Wilson, <i>Representations of finite rings</i>	643