

# Pacific Journal of Mathematics

## **ON SOME GROUP ALGEBRA MODULES RELATED TO WIENER'S ALGEBRA $M_1$**

TENG-SUN LIU, ARNOUD C. M. VAN ROOIJ AND JU-KWEI WANG

# ON SOME GROUP ALGEBRA MODULES RELATED TO WIENER'S ALGEBRA $M_1$

TENG-SUN LIU, ARNOUD VAN ROOIJ AND JU-KWEI WANG

Along with his study of the general Tauberian theorem in  $L_1, N$ . Wiener introduced the algebra  $M_1$  which consists of all those continuous functions  $f$  on the real line  $R$  for which

$$\sum_{n=-\infty}^{\infty} \max_{x \in [n, n+1]} |f(x)| < \infty.$$

He proved that many features of  $L_1$ , including the general Tauberian theorem, are shared by  $M_1$ . In this paper to generalize  $M_1$  to an arbitrary locally compact group  $G$ . While doing this, a host of  $L_1(G)$ -modules mutually related by conjugation and the operation of forming multiplier modules.  $\mathcal{M}_1(G)$  is among them. In case  $G$  is abelian,  $\mathcal{M}_1(G)$  is a Segal algebra, so that it has the same ideal-theoretical structure as  $L_1(G)$ . If further  $G = R$ ,  $\mathcal{M}_1(G)$  reduces to the Wiener algebra  $M_1$  with an equivalent norm.

1. Our notations are basically the same as those used in [3]. We use, however,  $C$  to denote the complex number field. Throughout the paper,  $G$  is a locally compact group with a left Haar measure  $\lambda$ . Instead of  $C_{00}(G)$ ,  $L_p(G)$  etc. we write  $C_{00}$ ,  $L_p$  etc. We view  $L_1$  as a subspace of  $M$ . We identify two functions that are equal almost everywhere.

For a function  $f$  on  $G$  define  $f'$  by

$$f'(x) = f(x^{-1})\Delta(x^{-1}),$$

where  $\Delta$  denotes the modular function of  $G$ . Then  $f'' = f$  and  $(f * g)' = g' * f'$  for  $f, g \in L_1$ .

If  $B$  is a left Banach module over  $L_1$  (see [3; 32.14]), then  $B^*$  becomes a left Banach module by

$$(j, f * \phi) = (f' * j, \phi) \quad (j \in B; \phi \in B^*; f \in L_1).$$

If  $B = L_p (1 \leq p < \infty)$  or  $B = C_0$  the module operation on  $B^*$  coincides with the convolution operation on  $L_q (q = p/(p-1))$  or  $M$ .

Let  $B$  be a left Banach module over  $L_1$ . By [3; 32.22],  $\{f * j; f \in L_1; j \in B\}$  is a closed submodule of  $B$ . We denote this submodule by  $L_1 * B$  or  $B_{abs}$ . We call  $B$  *absolutely continuous* if  $B_{abs} = B$ .

Suppose  $B$  is a Banach space, and there is a map  $(j, x) \mapsto j_x$  of  $B \times G$  into  $B$  such that

$$(1) \quad j_1 = j(j \in B),$$

- (2)  $(j_x)_y = j_{xy}(j \in B; x, y \in G)$ ,
- (3) for every  $x \in G$ ,  $j \mapsto j_x$  is a linear isometry,
- (4) for every  $j \in B$ ,  $x \mapsto j_x$  is a continuous map  $G \rightarrow B$ .

Then  $(j, x) \mapsto j_x$  is called a *continuous shift* in  $B$ . Such a continuous shift makes  $B$  into an absolutely continuous left Banach module over  $L_1$  by

$$(f * j, \phi) = \int f(x)(j_{x^{-1}}, \phi) dx, (f \in L_1; j \in B; \phi \in B^*).$$

For details, see [2], [4].

We can define continuous shifts in  $L_p (1 \leq p < \infty)$  and  $C_0$  by

$$j_x(y) = j(xy) \quad (j \in L_p \text{ or } C_0; x, y \in G).$$

The resulting module operation is ordinary convolution.

Let  $B$  be a left Banach module over  $L_1$ . The continuous module-homomorphism  $L_1 \rightarrow B$  (the multipliers of  $B$ ) form a Banach space  $\text{Mult } B$  that can be turned into a left Banach module by

$$(f * T)(g) = T(g * f) \quad (f, g \in L_1; T \in \text{Mult } B).$$

Every  $j \in B$  induces the multiplier  $f \mapsto f * j$ . The following theorem is essentially due to Rieffel [6] and is proved in [2] as Theorem 5.2:

**THEOREM 1.1.** *Let  $B$  be an absolutely continuous module. For  $\phi \in B^*$  let  $T_\phi$  be the multiplier  $f \mapsto f * \phi$  of  $B^*$ . Then  $T$  is a module isomorphism and a linear homeomorphism of  $B^*$  onto  $\text{Mult } B^*$ .*

A *Radon measure* on  $G$  is a linear functional  $\mu: C_0 \rightarrow \mathbb{C}$  such that for every compact set  $C \subset G$  there exists a number  $c$  such that

$$|(j, \mu)| \leq c \|j\|_\infty \quad (j \in C_0, \text{Supp } j \subset C).$$

The Radon measures form a vector space which we denote by  $R$ .

For  $\mu \in R$  and for an  $f \in L_1$  with compact support we define  $f * \mu \in R$  and  $\mu * f \in R$  by

$$\left. \begin{aligned} (j, f * \mu) &= (f' * j, \mu) \\ (j, \mu * f) &= (j * \Delta f', \mu) \end{aligned} \right\} (j \in C_0).$$

These formulas reduce to the familiar convolution formulas in case  $\mu \in L_p$ .

Every Radon measure is a linear combination of positive Radon measures [1]. Thus, if  $\mu \in R$  and if  $X$  is a relatively compact Borel subset of  $G$ , we can in a natural way define  $\mu(X)$ . Further, if  $\mu \in R$  and if  $A$  is any Borel set there is a unique  $\xi_A \mu \in R$  such that  $\xi_A \mu(X) = \mu(X \cap A)$  for all relatively compact Borel sets  $X$ . There

exists a unique  $|\mu| \in R$  such that  $\xi_A |\mu| = |\xi_A \mu|$  for every compact set  $A$  (see [1; Ch. 13]).

By the Radon-Nikodým Theorem [3; 12.17] we may identify  $L_{1,loc}$  with  $\{\mu \in R: \mu \ll \lambda\}$ .

**THEOREM 1.2.** *If  $f \in L_1$  has compact support and if  $\mu \in R$ , then  $f * \mu$  and  $\mu * f$  lie in  $L_{1,loc}$ . If, in addition,  $f$  is bounded, and  $\mu \in L_{1,loc}$ , then  $f * \mu$  and  $\mu * f$  are continuous.*

*Proof.* Let  $C = \text{Supp } f$ . If  $D \subset G$  is compact, then

$$(f * \xi_{G^{-1}D} \mu)(X) = (f * \mu)(X)$$

for all Borel set  $X \subset D$ . It follows that  $f * \mu \ll \lambda$ . Further by [3; 20.16], if  $f$  is bounded and  $\mu \in L_{1,loc}$ , then  $f * \mu$  is continuous on every compact  $D \subset G$ .

The proof for  $\mu * f$  is similar.

*From now on,  $K$  will be a nonempty compact subset of  $G$  which is the closure of its interior.*

For  $\mu \in R$  we define  $\mu^\sharp: G \rightarrow [0, \infty)$  by

$$\mu^\sharp(x) = \|\xi_{xK} \mu\| = |\mu|(xK) \quad (x \in G).$$

It is easy to check that

$$\mu^\sharp = |\mu| * \xi_{K^{-1}}.$$

Thus,  $\mu^\sharp \in L_{1,loc}$ , and  $\mu^\sharp$  is continuous if  $\mu \in L_{1,loc}$ .

Further, if  $f$  is a measurable function on  $G$ , define  $f^\sharp: G \rightarrow [0, \infty]$  by

$$f^\sharp(x) = \|f \xi_{xK}\|_\infty.$$

$f^\sharp$  is lower semicontinuous, hence measurable. (*Proof:* For  $r \in R$  put  $f_r(x) = \min(r, |f(x)|)$ ; then  $f^\sharp = \sup_r f_r^\sharp$ . Thus, we may assume that  $f$  is bounded and  $\geq 0$ . For  $j \in L_1$ ,  $j \geq 0$ ,  $\int j = 1$ , the function  $fj * \xi_{K^{-1}}$

is continuous [3; 20.16], and for every  $x \in G$ ,

$$f^\sharp(x) = \sup_j \int \xi_{xK} f j = \sup_j f j * \xi_{K^{-1}}.$$

Thus,  $f^\sharp$  is a supremum of continuous functions.)

**LEMMA 1.3.** *Let  $\mu \in R$ , let  $f$  be a measurable function. Assume*

that either  $\mu \ll \lambda$  or  $f$  is continuous. Then

$$(f^*, \mu^\sharp) \geq \lambda(K^{-1})(|f|, |\mu|).$$

*Proof.* By the assumption, for every  $x \in G$  we have  $|f| \leq f^*(x)|\mu|$  a.e. on  $xK$ , so that

$$f^*(x)\mu^\sharp(x) \geq \int_{xK} |f(y)| d|\mu|(y).$$

We may assume  $f^*\mu^\sharp \in L_1$ . Then there exists a  $\sigma$ -compact set  $X$  such that  $f^*\mu^\sharp = 0$  a.e. outside  $X$ . Since the  $X$  is  $\sigma$ -compact, it follows from [3; 5.7] that there exists a closed and open  $\sigma$ -compact subgroup  $H$  of  $G$  containing  $X$  and  $K$ . Every relatively compact Borel subset  $C$  of  $G \setminus H$  can be covered by finitely many cosets  $a_1K, \dots, a_mK$  where  $f^*(a_i)\mu^\sharp(a_i) = 0$  for each  $i$ . Then

$$\int_C |f| d|\mu| \leq \sum f^*(a_i)\mu^\sharp(a_i) = 0.$$

Put  $f_1 = \xi_H |f|$ ,  $\mu_1 = \xi_H |\mu|$ . Then  $\int |f| d|\mu| = \int f_1 d\mu_1$ . Further,  $f_1^*\mu_1^\sharp = f^*\mu^\sharp$  on  $H$  and  $f_1^* \leq f^*$ ,  $\mu_1^\sharp \leq \mu^\sharp$  everywhere; therefore  $(f^*, \mu^\sharp) = (f_1^*, \mu_1^\sharp)$ . It follows that we may replace  $f$  by  $f_1$  and  $\mu$  by  $\mu_1$ ; i.e., we may assume that  $f \geq 0$ ,  $\mu \geq 0$  and that  $G$  is  $\sigma$ -compact. This enables us to apply the Fubini Theorem:

$$\begin{aligned} (f^*, \mu^\sharp) &\geq \iint \xi_{xK}(y) |f(y)| d|\mu|(y) dx \\ &= \iint \xi_{yK^{-1}}(x) |f(y)| dx d|\mu|(y) \\ &= \int \lambda(yK^{-1}) |f(y)| d|\mu|(y) = \lambda(K^{-1}) \int |f(y)| d|\mu|(y). \end{aligned}$$

*Note.* This lemma, and also its applications, Theorems 3.1 and 6.1, should be read with some caution. In the case where  $f$  is continuous the “ $f$ ” in the right hand member of the formula denotes a single function, but the “ $f$ ” in the left hand member stands for the class of all functions that are l.a.e. equal to  $f$ .

**COROLLARY 1.4.** For all  $\mu \in R$ ,  $\int \mu^\sharp = \lambda(K^{-1})|\mu|(G)$ . Thus, if  $\mu^\sharp = 0$  a.e., then  $\mu = 0$ .

*Proof.* It is clear that  $\int \mu^\sharp \leq \|\xi_{K^{-1}}\|_1 |\mu|(G) = \lambda(K^{-1})|\mu|(G)$ . On taking  $f = 1$ , we get  $\int \mu^\sharp \geq \lambda(K^{-1})|\mu|(G)$ . Hence the equality. If  $C$

is compact, then

$$\int \mu^{\sharp} \geq \int (\xi_C \mu)^{\sharp} = \int |\xi_C \mu| * \xi_{K^{-1}} = \lambda(K^{-1}) |\xi_C \mu|(G) = \lambda(K^{-1}) |\mu|(C) .$$

Hence we have the second statement.

2. For  $1 \leq p < \infty$  let

$$\mathcal{V}_p^{\wedge} = \mathcal{V}_{p0}^{\wedge} = \{j \in L_{1,loc}; j^{\sharp} \in L_p\} .$$

Further, put

$$\mathcal{V}_{\infty}^{\wedge} = \{j \in L_{1,loc}; j^{\sharp} \in L_{\infty}\} ,$$

$$\mathcal{V}_{\infty 0}^{\wedge} = \{j \in L_{1,loc}; j^{\sharp} \in C_0\} .$$

Clearly  $\mathcal{V}_{\infty 0}^{\wedge} \subset \mathcal{V}_{\infty}^{\wedge}$ . As  $\int j^{\sharp} = \lambda(K^{-1}) \int |j|$ , we have  $\mathcal{V}_1^{\wedge} = L_1$ . For  $j \in \mathcal{V}_p^{\wedge}$ , we set

$$\|j\|_p^{\sharp} = \|j^{\sharp}\|_p .$$

**THEOREM 2.1.** *Let  $1 \leq p \leq \infty$ .  $\mathcal{V}_p^{\wedge}$  and  $\mathcal{V}_{p0}^{\wedge}$  are Banach spaces.  $L_1$  is a dense subset of  $\mathcal{V}_{p0}^{\wedge}$ ; the natural injection  $L_1 \rightarrow \mathcal{V}_{p0}^{\wedge}$  is continuous. The formula*

$$j_x(y) = j(xy) \quad (j \in \mathcal{V}_{p0}^{\wedge}; x, y \in G)$$

*defines a continuous shift in  $\mathcal{V}_{p0}^{\wedge}$ .*

*Proof.* Clearly  $\mathcal{V}_p^{\wedge}$ ,  $\mathcal{V}_{p0}^{\wedge}$  are vector spaces and  $\|\cdot\|_p^{\sharp}$  is a norm. To prove completeness of  $\mathcal{V}_p^{\wedge}$ , let  $\{j_n\}$  be a sequence in  $\mathcal{V}_p^{\wedge}$  such that  $\sum \|j_n\|_p^{\sharp} < \infty$ ; it suffices to prove that  $\sum j_n$  converges in  $\mathcal{V}_p^{\wedge}$ . We know that  $\sum j_n^{\sharp}(x)$  converges for all  $x \in G$  outside a certain locally null set  $X$ . Take a compact set  $C \subset G$ .  $C$  is covered by finitely many translates  $a_1 K, \dots, a_m K$  of  $K$  and we can choose all  $a_i$  in  $G \setminus X$ . Then

$$\sum_n \int_C |j_n| \leq \sum_n \sum_i j_n^{\sharp}(a_i) < \infty ,$$

so that  $\sum |j_n| < \infty$  a.e. on  $C$ . By [3; 11.39 and 11.42] there exists a measurable function  $j$  such that  $\sum j_n = j$  l.a.e. Then  $\int_C |j| \leq \sum \int_C |j_n|$  for every compact  $C$ , so  $j \in L_{1,loc}$  and  $j^{\sharp} \leq \sum j_n^{\sharp}$ . Hence,  $j \in \mathcal{V}_p^{\wedge}$ . We also obtain  $(j - \sum_1^N j_n)^{\sharp} \leq \sum_{N+1}^{\infty} j_n^{\sharp}$ , so

$$\left\| j - \sum_1^N j_n \right\|_p^{\sharp} \leq \left\| \sum_{N+1}^{\infty} j_n^{\sharp} \right\|_p \leq \sum_{N+1}^{\infty} \|j_n^{\sharp}\|_p = \sum_{N+1}^{\infty} \|j_n\|_p^{\sharp} .$$

Therefore,  $j = \Sigma j_n$  in the sense of  $\mathcal{N}_p$ .

For  $\mathcal{V}_{\infty}$  we simply observe that  $j \mapsto j^\sharp$  is a continuous map  $\mathcal{V}_{\infty} \rightarrow L_{\infty}$ , so that  $\mathcal{V}_{\infty}$  is closed in  $\mathcal{V}_{\infty}$ .

Next, take  $j \in L_1$ . Since  $\xi_{K^{-1}} \in L_p$ ,  $j^\sharp = |j| * \xi_{K^{-1}} \in L_p$  and  $\|j\|_p^\sharp \leq \|j\|_1 \|\xi_{K^{-1}}\|_p$ . It is easy to see that  $j^\sharp \in C_0$ . Hence,  $L_1 \subset \mathcal{V}_{p_0}$  and the injection is continuous. To prove that  $L_1$  is dense, take  $j \in \mathcal{V}_{p_0}$ ,  $\varepsilon > 0$ . There is a compact set  $C$  such that  $\|j^\sharp(1 - \xi_C)\|_p \leq \varepsilon$ . Then  $(1 - j_{CK}^\sharp)^\sharp = 0$  on  $C$  and  $(j - j_{CK}^\sharp)^\sharp \leq j^\sharp$  everywhere. For  $p < \infty$  it follows that  $\|j - j_{CK}^\sharp\|_p^\sharp \leq \|j^\sharp(1 - \xi_C)\|_p \leq \varepsilon$ . But  $j_{CK}^\sharp \in L_1$ . Thus, for  $p < \infty$ ,  $L_1$  is dense in  $\mathcal{V}_{p_0}$ . A similar proof works for  $p = \infty$ .

Trivially, if  $j \in \mathcal{V}_{p_0}$ , then  $j_x \in \mathcal{V}_{p_0}$  for every  $x$ . We only have to prove that  $x \mapsto j_x$  is continuous. As  $L_1$  is dense, we may assume  $j \in L_1$ . Now for such  $j$  we know that the formula  $x \mapsto j_x$  defines a continuous map  $G \rightarrow L_1$ . Now observe that the injection  $L_1 \rightarrow \mathcal{V}_{p_0}$  is continuous.

**THEOREM 2.2.** *For  $1 \leq p \leq \infty$ ,  $L_p$  is a dense subset of  $\mathcal{V}_p$ , and the injection  $L_p \rightarrow \mathcal{V}_p$  is continuous. Further,  $C_0$  is a dense subset of  $\mathcal{V}_{\infty}$  and  $C_0 \rightarrow \mathcal{V}_{\infty}$  is continuous.*

*Proof.* The first statement follows from the formula

$$j^\sharp = |j| * \xi_{K^{-1}}$$

and [3; 20.13]. Further, for  $j \in C_0$  we have  $|j| * \xi_{K^{-1}} \in C_0$  as is easy to prove.

According to the remarks made in §1,  $\mathcal{V}_{p_0}$  can be made into an absolutely continuous module. The module operation  $*$  is given by the familiar convolution formula:

**THEOREM 2.3.** *If  $f \in L_1$  and  $j \in \mathcal{N}_{p_0}$  ( $1 \leq p \leq \infty$ ), one has for locally almost every  $x \in G$ ,*

$$f * j(x) = \int f(y)j(y^{-1}x)dy.$$

*Proof.* We may assume  $f, j \geq 0$ . If  $h \in C_{00}$  and  $h \geq 0$ , then

$$\begin{aligned} (h, f * j) &= \int f(y)(h, j_{y^{-1}})dy = \iint f(y)h(x)j(y^{-1}x)dxdy \\ &= \int h(x) \int f(y)j(y^{-1}x)dydx. \end{aligned}$$

3. For  $1 \leq p \leq \infty$  let  $\mathcal{N}_p$  be the set of all measurable functions  $f$  on  $G$  for which  $f^\sharp \in L_p$ . Then  $\mathcal{N}_p$  is a vector space, and the formula

$$\|f\|_p^* = \|f^*\|_p \quad (f \in \mathcal{N}_p)$$

defines a norm on  $\mathcal{N}_p$ .

**THEOREM 3.1.** *Let  $1 \leq p \leq \infty$ ,  $q = p/(p-1)$ . Every  $f \in \mathcal{N}_q$  determines a  $\Phi f \in \mathcal{V}_{p_0}^*$  by*

$$(j, \Phi f) = \int j(x)f(x)dx \quad (j \in \mathcal{V}_{p_0}).$$

$\Phi$  is a linear homeomorphism on  $\mathcal{N}_q$  onto  $\mathcal{V}_{p_0}^*$ . We have

$$\|\Phi\| \leq 1/\lambda(K^{-1}) \quad \text{and} \quad \|\Phi^{-1}\| \leq \int_{KK^{-1}} \Delta^{-1/q}.$$

*Proof.* It follows directly from Lemma 1.3 that  $\Phi$  maps  $\mathcal{N}_q$  into  $\mathcal{V}_{p_0}^*$  and that  $\|\Phi\| \leq \lambda(K^{-1})^{-1}$ . For the converse, take  $\phi \in \mathcal{V}_{p_0}^*$ . As  $L_1 \subset \mathcal{V}_{p_0}$  and as the injection  $L_1 \rightarrow \mathcal{V}_{p_0}$  is continuous, there is an  $f \in L_\infty$  such that

$$(j, \phi) = \int j(x)f(x)dx \quad (j \in L_1).$$

If  $f \in \mathcal{N}_q$ , then  $\phi = \Phi f$  on a dense subspace of  $\mathcal{V}_{p_0}$ , hence on all of  $\mathcal{V}_{p_0}$ . We proceed to prove that  $f \in \mathcal{N}_q$  and  $\|f\|_q^* \leq \|\phi\| \int_{KK^{-1}} \Delta^{-1/q}$ . Assume for the time being that  $f \geq 0$ . Take  $\varepsilon \in (0, 1)$ . The set

$$S = \{(x, y) \in G^2: y \in xK; f(y) \geq (1 - \varepsilon)f^*(x)\}$$

is measurable. Then for l.a.e.  $x \in G$  the set

$$S_x = \{y \in G: (x, y) \in S\} = \{y \in xK: f(y) \geq (1 - \varepsilon)f^*(x)\}$$

is measurable. Moreover, the function  $x \rightarrow \lambda(S_x)$  is measurable, and  $\lambda(S_x) > 0$  l.a.e.

Let  $h \in L_1 \cap L_p$ ,  $h \geq 0$ ,  $\text{Supp } h$  compact. Then

$$(1 - \varepsilon) \int f^*(x)h(x)dx \leq \int \left[ \int \frac{f(y)}{\lambda(S_x)} \xi_{S_x}(y)dy \right] h(x) dx = \int f(y)j(y)dy,$$

where

$$j(y) = \int \frac{h(x)}{\lambda(S_x)} \xi_{S_x}(y)dx.$$

One easily sees that  $j \in L_1$  so that

$$(1 - \varepsilon) \int f^*(x)h(x)dx \leq \int f(y)j(y)dy = (j, \phi) \leq \|\phi\| \|j\|_p^*.$$

To find an estimate for  $\|j\|_p^*$  we observe that for every  $a \in G$ ,



$$j^{\sharp}(a) = \iint \tilde{\xi}_{aK}(y) \frac{h(x)}{\lambda(S_x)} \tilde{\xi}_{S_x}(y) dy dx = \int h(x) \frac{\lambda(aK \cap S_x)}{\lambda(S_x)} dx .$$

As  $S_x \subset xK$  we have  $\lambda(aK \cap S_x) = 0$  unless  $x \in aKK^{-1}$ , so that

$$j^{\sharp}(a) \leq \int_{aKK^{-1}} h(x) dx = h_1^* \tilde{\xi}_{KK^{-1}}(a) .$$

By again applying [3; 20.13] we find

$$\|j\|_p^{\sharp} \leq \|h * \tilde{\xi}_{KK^{-1}}\|_p \leq \|h\|_p \int_{KK^{-1}} \Delta(y)^{-1/q} dy .$$

Thus, for all  $\varepsilon > 0$  and all  $h \in L_1 \cap L_p$ ,  $h \geq 0$  we get

$$(1 - \varepsilon) \int f^{\sharp} h \leq \|\phi\| \|h\|_p \int_{KK^{-1}} \Delta^{-1/q} .$$

Then  $f^{\sharp} \in L_q$ , i.e.,  $f \in \mathcal{N}_q$ , and  $\|f\|_q^{\sharp} \leq \|\phi\| \int_{KK^{-1}} \Delta^{-1/q}$ .

We have proved our point for the case  $f \geq 0$ . For the general case we notice that there exists a measurable function  $\tau: G \rightarrow \mathbb{C}$  such that  $|\tau(x)| = 1$  for all  $x$  and  $\tau f \geq 0$ . Define  $\psi: \mathcal{Y}_{p_0} \rightarrow \mathbb{C}$  by

$$(j, \psi) = (j\tau^{-1}, \phi) .$$

Then  $\psi \in \mathcal{Y}_{p_0}^*$ ,  $\|\psi\| = \|\phi\|$ , and  $(j, \psi) = \int j(x)(\tau f)(x) dx$  for  $j \in L_1$ . It follows that  $\tau f \in \mathcal{N}_q$  and  $\|\tau f\|_q^{\sharp} \leq \|\psi\| \int_{KK^{-1}} \Delta^{-1/q}$ ; so  $f \in \mathcal{N}_q$  and  $\|f\|_q^{\sharp} \leq \|\phi\| \int_{KK^{-1}} \Delta^{-1/q}$ .

**COROLLARY 3.2.**  $\mathcal{N}_q \subset L_{\infty} \cap L_q$ . The injections  $\mathcal{N}_q \rightarrow L_{\infty}$  and  $\mathcal{N}_q \rightarrow L_q$  are continuous.

*Proof.* By Theorem 2.1, there is a constant  $c$  such that for every  $j \in L_1$ ,  $\|j\|_p^{\sharp} \leq c\|j\|_1$ . Then if  $f \in \mathcal{N}_q$ , we have

$$\|jf = (j, \Phi f) \leq c\|\Phi\| \|f\|_q^{\sharp} \|j\|_1 .$$

Hence  $f \in L_{\infty}$  and  $\|f\|_{\infty} \leq c\|\Phi\| \|f\|_q^{\sharp}$ . The proof for the statements concerning  $L_q$  is simimilar, using Theorem 2.2 instead of Theorem 2.1.

**COROLLARY 3.3.**  $\mathcal{N}_q$  is a Banach space which is also an  $L_1$ -module under convolution.

*Proof.* We know that  $\mathcal{Y}_{p_0}^*$  can be made into a left Banach module over  $L_1$ . After Theorem 3.1 we only have to show that the induced module operation in  $\mathcal{N}_q$  is convolution. Now for  $f \in L_1$ ,  $g \in \mathcal{N}_q$  and

$j \in \mathcal{V}_{p_0}$  we have

$$\begin{aligned} (j, f_* g) &= (f' * j, g) = \iint f'(x) j(x^{-1}y) g(y) dx dy \\ &= \iint f(x^{-1}) j(x^{-1}y) g(y) \Delta(x^{-1}) dx dy \\ &= \iint f(x) j(xy) g(y) dy dx = \iint f(x) j(y) g(x^{-1}y) dy dx \\ &= \int j(y) \int f(x) g(x^{-1}y) dx dy . \end{aligned}$$

4. For  $1 \leq p < \infty$  let

$$\mathcal{M}_p = \{f \in \mathcal{N}_p : f \text{ is continuous}\} .$$

THEOREM 4.1.  $C_{00}$  is a dense subspace of  $\mathcal{M}_p$ .

*Proof.* Take  $f \in \mathcal{M}_p$ ,  $\varepsilon > 0$ . There exists a compact  $C \subset G$  such that  $\|f^*(1 - \xi_C)\|_p \leq \varepsilon$ , and there exists a  $g \in C_{00}$ ,  $g = f$  on  $CK$ , such that  $|g| \leq |f|$ . Then  $(f - g)^* = 0$  on  $C$ , and

$$\|f - g\|_p^* \leq \|(f - g)^* \xi_C\|_p + \|f^*(1 - \xi_C)\|_p + \|g^*(1 - \xi_C)\|_p \leq 2\varepsilon .$$

THEOREM 4.2. For  $1 \leq p < \infty$ ,  $\mathcal{M}_p = (\mathcal{N}_p)_{abs}$ . The formula

$$T_f g = g * f \quad (f \in \mathcal{N}_p; g \in L_1)$$

establishes a one-to-one correspondence between  $\mathcal{N}_p$  and  $\text{Mult } \mathcal{M}_p$ .

*Proof.* First, take  $f \in C_{00}$ . Let  $S = \text{Supp } f$ . Let  $U$  be a compact neighborhood of  $S$ . For every  $\varepsilon > 0$  there exists a neighborhood  $V_\varepsilon$  of 1 such that  $V_\varepsilon^{-1}S \subset U$  and  $\|f - f_x\|_\infty \leq \varepsilon$  for all  $x \in V_\varepsilon$ . Then for  $x \in V_\varepsilon$ , we have  $\|(f - f_x)^*\|_\infty \leq \varepsilon$  and  $(f - f_x)^* = 0$  off  $UK^{-1}$ . Thus, for  $x \in V_\varepsilon$ ,  $\|f - f_x\|_p^* \leq \varepsilon[\lambda(UK^{-1})]^{1/p}$ .

It follows that  $x \mapsto f_x$  is a continuous map  $G \rightarrow \mathcal{M}_p$  for every  $f \in C_{00}$ , hence for every  $f \in \mathcal{M}_p$  (see Theorem 4.1). Thus,  $\mathcal{M}_p$  can be made into an absolutely continuous left Banach module over  $L_1$ . It is easy to see that the induced module operation is convolution, which is the same as the module operation in  $\mathcal{N}_p$ . Thus,  $\mathcal{M}_p = L_1 * \mathcal{M}_p = (\mathcal{N}_p)_{abs}$ .

Now  $(\mathcal{N}_p)_{abs} = L_1 * \mathcal{N}_p \subset L_1 * L_\infty$  and every element of  $L_1 * L_\infty$  is continuous [3; 20.16]. Hence,  $(\mathcal{N}_p)_{abs} \subset \mathcal{M}_p$ , so that  $(\mathcal{N}_p)_{abs} = \mathcal{M}_p$ .

We also see that every  $T_f$  maps  $L_1$  into  $\mathcal{M}_p$ , hence is an element of  $\text{Mult } \mathcal{M}_p$ . Let  $q = p/(p - 1)$ , ( $q = \infty$  if  $p = 1$ ). As  $\mathcal{V}_{p_0}$  is an absolutely continuous module, it follows from Theorem 1.1 that for every  $T \in \text{Mult } \mathcal{V}_{p_0}^*$  there exists a  $\phi \in \mathcal{V}_{p_0}^*$  such that  $Tf = f * \phi$  ( $f \in L_1$ ).

The rest of the theorem follows from Theorem 3.1 and the trivial observation that  $\text{Mult } \mathcal{M}_p \subset \text{Mult } \mathcal{N}_p$ .

COROLLARY 4.3.  $\mathcal{M}_p$  is a Banach space.

Let us consider the case  $p = \infty$ . Obviously,  $\mathcal{N}_\infty = L_\infty$ . Thus,  $\{f \in N_\infty: f \text{ is continuous}\}$  is the space of all bounded continuous functions. It is known, however, that  $(L_\infty)_{abs} = C_{ru}$  (see [3; 32.45]). We could save the situation by defining, for  $1 \leq p \leq \infty$ ,

$$\mathcal{M}_p = C_{ru} \cap \mathcal{N}_p.$$

Then Theorem 4.2 remains valid if we change “ $1 \leq p < \infty$ ” into “ $1 \leq p \leq \infty$ ”.

5.  $\mathcal{M}_1$  deserves some special attention.

THEOREM 5.1.  $\mathcal{N}_1$  and  $\mathcal{M}_1$  are Banach algebras under convolution. If  $G$  is abelian,  $\mathcal{M}_1$  is a Segal algebra (as defined in [5; Ch. 6, §2]).

*Proof.* The injection  $I: \mathcal{N}_1 \rightarrow L_1$  is continuous (Corollary 3.2). For  $f, g \in \mathcal{N}_1$  we have  $\|f * g\|_1^\# \leq \|f\|_1 \|g\|_1^\# \leq \|I\| \|f\|_1^\# \|g\|_1^\#$ . Further,  $\mathcal{M}_1$  is a left ideal in  $L_1$  by Theorem 4.2. The second statement follows from Theorem 4.1 and the continuity of the shift (see proof of Theorem 4.2).

Consider in particular the case  $G = \mathbf{R}$ , the additive reals. Wiener defines his Banach algebra  $M_1$  as the set of all continuous functions  $f$  on  $\mathbf{R}$  for which  $\|f\|_{M_1} < \infty$ , where

$$\|f\|_{M_1} = \sum_{n=-\infty}^{\infty} \max_{[n, n+1]} |f(x)|.$$

$M_1$  was discussed in [7; Ch. 2] and [5; pp. 12, 127], [3; II pp. 506, 600]. To show that  $M_1 = \mathcal{M}_1$ , for a continuous function  $f$  on  $\mathbf{R}$  define  $f^\flat$  on  $\mathbf{R}$  by

$$f^\flat(x) = \sup_{[n, n+1]} |f| \quad \text{if } n \in \mathbf{Z}, n \leq x < n+1.$$

Then  $\|f\|_{M_1} = \|f^\flat\|_1$ . By taking  $K = [0, 1]$  we find  $f^\sharp(x) \leq f^\flat(x) + f^\flat(x+1)$  and  $f^\flat(x) \leq f^\sharp(x-1) + f^\sharp(x)$  for all  $x$ . Hence  $M_1 = \mathcal{M}_1$  and

$$\frac{1}{2} \| \cdot \|_1^\# \leq \| \cdot \|_{M_1} \leq 2 \| \cdot \|_1^\#.$$

6. Finally, for  $1 \leq p \leq \infty$  we set

$$\mathcal{W}_p = \{\mu \in \mathbf{R}: \mu^\sharp \in L_p\}, \quad \|\mu\|_p^\sharp = \|\mu^\sharp\|_p.$$

For all  $\mu \in R$  we have  $\int \mu^\sharp(x) dx = \lambda(K^{-1}) |\mu| (G)$ . Hence  $\mathcal{W}_1 = M$ .

**THEOREM 6.1.** *Let  $1 \leq p < \infty$ ,  $q = p/(p-1)$ . The formula*

$$(f, \Psi\mu) = \int f d\mu \quad (f \in \mathcal{M}_p; \mu \in \mathcal{W}_q)$$

*establishes a linear homeomorphism of  $\mathcal{W}_q$  onto  $\mathcal{M}_p^*$ . Further,*

$$\|\Psi\| \leq 1/\lambda(K^{-1}), \quad \|\Psi^{-1}\| \leq \int_{KK^{-1}} \Delta^{-1/q}.$$

*Proof.* It follows from Lemma 1.3 that  $\Psi: \mathcal{W}_q \rightarrow \mathcal{M}_p^*$  and that  $\|\Psi\| \leq \lambda(K^{-1})^{-1}$ . Conversely, take  $\phi \in \mathcal{M}_q^*$ . If  $C \subset G$  is compact, then for every  $f \in C_{00}$  that has its support in  $C$  we have

$$|(f, \phi)| \leq \|\phi\| \|f\|_p^\sharp \leq \|\phi\| \|f\|_\infty [\lambda(CK^{-1})]^{-1/p}.$$

Thus, there is a  $\mu \in R$  such that

$$(f, \phi) = \int f d\mu \quad (f \in C_{00}).$$

If  $\mu^\sharp \in L_q$  and  $\|\mu^\sharp\|_q \leq \|\phi\| \int_{KK^{-1}} \Delta^{-1/q}$ , then  $\Psi\mu = \phi$  on a dense subset of  $\mathcal{M}_p$ , and we are done.

Take  $h \in C_{00}$ ,  $h \geq 0$ . Then

$$\begin{aligned} \int h(x) \mu^\sharp(x) dx &= \iint h(x) \xi_{xK}(y) d|\mu|(y) dx \\ &= \iint h(x) \xi_K(x^{-1}y) dx d|\mu|(y) = (h * \xi_K, |\mu|). \end{aligned}$$

By [3; 14.5] and the continuity of  $h * \xi_K$  we obtain  $(h * \xi_K, |\mu|) = \sup_{j \in \mathcal{J}} |(j, \mu)|$  where  $\mathcal{J} = \{f \in C_{00} : |f| \leq h * \xi_K\}$ . Observe that for every  $f \in \mathcal{J}$  and  $a \in G$ ,

$$f^\sharp(a) \leq \sup_{x \in aK} h * \xi_K(x) = \sup_{x \in aK} \int_{xK^{-1}} h \leq \int_{aKK^{-1}} h = h * \xi_{KK^{-1}}(a).$$

By another application of [3; 20.13] we find

$$\begin{aligned} \int h(x) \mu^\sharp(x) dx &= (h * \xi_K, |\mu|) = \sup_{f \in \mathcal{J}} |(f, \mu)| \leq \sup_{f \in \mathcal{J}} |(f, \phi)| \\ &\leq \sup_{f \in \mathcal{J}} \|\phi\| \|f\|_p^\sharp \leq \|\phi\| \|h * \xi_{KK^{-1}}\|_p \leq \|\phi\| \|h\|_p \int_{KK^{-1}} \Delta^{-1/q}. \end{aligned}$$

Thus,  $\mu^\sharp \in L_q$  and  $\|\mu^\sharp\|_q \leq \|\phi\| \int_{KK^{-1}} \Delta^{-1/q}$ .

COROLLARY 6.2.  $\mathscr{W}_p$  is a Banach space.

By the general theory of Banach modules,  $\mathscr{W}_p$  can be made into a left Banach module over  $L_1$  (see §1, and §3 where we introduced a module operation on  $\mathscr{N}_p$ ). An application of Theorem 4.1 yields the fact that for  $f \in L_{1,\text{loc}}$  and  $\mu \in \mathscr{W}_p$  the module product and the convolution product  $f * \mu$  coincide. It follows that  $\mathscr{V}_p$  is a submodule of  $\mathscr{W}_p$ .

We close the circle by:

COROLLARY 6.3. If  $p < \infty$ , then  $\mathscr{V}_p = (\mathscr{W}_p)_{\text{abs}}$ . For all  $p$  the formula

$$S_\mu f = f * \mu \quad (\mu \in \mathscr{W}_p; f \in L_1)$$

yields a linear homeomorphism  $S$  of  $\mathscr{W}_p$  onto  $\text{Mult } \mathscr{V}_p$ .

*Proof.* For  $p = 1$  we can apply Wendel's Theorem [3; 35.5], since  $\mathscr{W}_1 = M$  and  $\mathscr{V}_1 = L_1$ . By the last part of Theorem 2.2,  $\mathscr{V}_p = (\mathscr{V}_p)_{\text{abs}} \subset (\mathscr{W}_p)_{\text{abs}}$  if  $p < \infty$ . For any  $p$  and for an  $f \in L_1$  that has compact support, we have

$$f * \mathscr{W}_p \subset (f * R) \cap \mathscr{W}_p \subset L_{1,\text{loc}} \cap \mathscr{W}_p = \mathscr{V}_p.$$

As  $\mathscr{V}_p$  is closed,  $(\mathscr{W}_p)_{\text{abs}} = L_1 * \mathscr{W}_p \subset \mathscr{V}_p$ . Thus  $\mathscr{V}_p = (\mathscr{W}_p)_{\text{abs}}$  if  $p < \infty$ , and  $S_\mu \in \text{Mult } \mathscr{V}_p$  for all  $\mu \in \mathscr{W}_p$ . The proof of the facts that for  $p \neq 1$  every element of  $\text{Mult } \mathscr{V}_p$  is of the form  $S_\mu$ , and that  $S$  is a homeomorphism is entirely analogous to the final part of the proof of Theorem 4.2 (using 6.1 instead of 3.1).

7. In order to see how the operations  $^\natural$  and  $^\sharp$  depend on  $K$ , take another nonempty compact set  $K_1$  that is the closure of its interior and define

$$\begin{aligned} \mu_1^\natural(x) &= |\mu|(xK_1) \quad (x \in G; \mu \in R), \\ f_1^\sharp(x) &= \|f\|_{\xi_{xK_1}}^\infty \quad (x \in G; f \in L_{1,\text{loc}}). \end{aligned}$$

As  $K$  is compact and  $K_1$  has nonempty interior, there exists  $a_1, \dots, a_n$  such that  $K \subset a_1 K_1 \cup \dots \cup a_n K_1$ . Then we see that

$$\begin{aligned} \mu^\natural &\leq (\mu_1^\natural)_{a_1} + \dots + (\mu_1^\natural)_{a_n} \quad (\mu \in R), \\ f^\natural &\leq (f_1^\natural)_{a_1} + \dots + (f_1^\natural)_{a_n} \quad (f \in L_{1,\text{loc}}). \end{aligned}$$

From these formulas it will follow directly that the Banach spaces  $\mathscr{V}_p, \mathscr{V}_{\infty,0}, \mathscr{N}_p, \mathscr{M}_p, \mathscr{W}_p$  are essentially independent of  $K$ : A different choice of  $K$  will only lead to a different, but equivalent norm in the same space.

In fact the proof shows that we can relax the conditions on  $K$ . We only need to require that  $K$  is a relatively compact set with nonempty interior. Any such  $K$  will lead to the same spaces  $\mathcal{V}_p$ ,  $\mathcal{V}_{\infty 0}$ ,  $\mathcal{N}_p$ ,  $\mathcal{M}_p$ , and  $\mathcal{W}_p$  with equivalent norms. The inequalities in Theorem 6.1 will no longer be true for such a general  $K$ . (The analogous inequalities in Theorem 3.1 remain valid.)

The results obtained so far can be summarized in the following table where we use the equality sign to indicate linear homeomorphism. In each formula,  $1/p + 1/q = 1$ .

$$\begin{aligned} \mathcal{V}_p^* &= \mathcal{N}_q, & \mathcal{V}_{\infty 0}^* &= \mathcal{N}_1 & (1 \leq p < \infty) \\ (\mathcal{N}_p)_{abs} &= \mathcal{M}_p, & \text{Mult } \mathcal{M}_p &= \mathcal{N}_p & (1 \leq p \leq \infty) \\ \mathcal{M}_p^* &= \mathcal{W}_q & & & (1 \leq p < \infty) \\ (\mathcal{W}_p)_{abs} &= \mathcal{V}_p & & & (1 \leq p < \infty) \\ \text{Mult } \mathcal{V}_p &= \mathcal{W}_p & & & (1 \leq p \leq \infty) \end{aligned}$$

The equalities in the first and third line clearly do not hold in general if we put  $p = \infty$ . For the fourth line this is less easy to see. Take  $G = \mathbf{R}$ ,  $K = [0, 1]$ . Let  $j$  be the function that vanishes on  $(-\infty, 0]$  and coincides with  $n$ th Rademacher function  $\phi_n$  on  $(n-1, n]$  for every positive integer  $n$  (see [8; Ch. I, §3]). Then  $j \in \mathcal{V}_{\infty}$ , but  $j \notin (\mathcal{V}_{\infty})_{abs}$ . (It is not hard to prove that, if  $j \in (\mathcal{V}_{\infty})_{abs}$ , then  $\lim_{x \rightarrow 0} \|j - j_x\|_{\infty}^n = 0$ , and that the latter formula is false.)

We do not have an adequate description of  $\text{Mult } \mathcal{V}_{\infty 0}$ .

Let us conclude with a table listing  $\mathcal{V}_p$ ,  $\mathcal{W}_p$ ,  $\mathcal{M}_p$ ,  $\mathcal{N}_p$  for compact  $G$  and for discrete  $G$ .

$G$	compact	discrete
$\mathcal{V}_p (1 \leq p \leq \infty)$	$L_1$	$L_p$
$\mathcal{V}_{\infty 0}$	$L_1$	$C_0$
$\mathcal{W}_p (1 \leq p \leq \infty)$	$M$	$L_p$
$\mathcal{W}_{\infty}$	$M$	$L_{\infty}$
$\mathcal{M}_p (1 \leq p \leq \infty)$	$C$	$L_p$
$\mathcal{N}_p (1 \leq p \leq \infty)$	$L_{\infty}$	$L_p$

## REFERENCES

1. J. Dieudonné, *Eléments d'analyse, II*, Gauthier-Villars, Paris, 1968.
2. S. L. Gulick, T.-S. Liu and A. C. M. van Rooij, *Group algebra modules, III*, Trans. Amer. Math. Soc., **152** (1970), 581-596.
3. E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis*, Springer-Verlag, Berlin-Heidelberg, I, 1963; II, 1970.

4. T.-S. Liu, A. C. M. van Rooij and J.-K. Wang, *Group representations in Banach spaces; orbits and almost-periodicity*, Studies and Essays Presented to Yu-Why Chen on His Sixtieth Birthday, Mathematics Research Center, Taipei, 1970, pp. 243-254.
5. H. Reiter, *Classical Fourier Analysis and Locally Compact Groups*, Oxford University Press, Oxford, 1968.
6. M. A. Rieffel, *Induced Banach representations of Banach algebras and locally compact groups*, J. Functional Anal., **1** (1967), 443-491.
7. N. Wiener, *The Fourier Integral and Certain of its Applications*, Cambridge University Press, Cambridge, England, 1933.
8. A. Zygmund, *Trigonometric Series, I*, Cambridge University Press, Cambridge, England, 1959.

Received November 29, 1971 and in revised form January 31, 1974. The first authors' research was supported in part by the U. S. National Science Foundation.

UNIVERSITY OF MASSACHUSETTS

AND

CATHOLIC UNIVERSITY, NIJMEGEN, THE NETHERLANDS

*Caution:* This was not proof read by the authors.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)

University of California  
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan



Walter Allegretto, <i>On the equivalence of two types of oscillation for elliptic operators</i> .....	319
Edward Arthur Bertram, <i>A density theorem on the number of conjugacy classes in finite groups</i> .....	329
Arne Brøndsted, <i>On a lemma of Bishop and Phelps</i> .....	335
Jacob Burbea, <i>Total positivity and reproducing kernels</i> .....	343
Ed Dubinsky, <i>Linear Pincherle sequences</i> .....	361
Benny Dan Evans, <i>Cyclic amalgamations of residually finite groups</i> .....	371
Barry J. Gardner and Patrick Noble Stewart, <i>A "going down" theorem for certain reflected radicals</i> .....	381
Jonathan Light Gross and Thomas William Tucker, <i>Quotients of complete graphs: revisiting the Heawood map-coloring problem</i> .....	391
Sav Roman Harasymiv, <i>Groups of matrices acting on distribution spaces</i> .....	403
Robert Winship Heath and David John Lutzer, <i>Dugundji extension theorems for linearly ordered spaces</i> .....	419
Chung-Wu Ho, <i>Deforming p. l. homeomorphisms on a convex polygonal 2-disk</i> .....	427
Richard Earl Hodel, <i>Metrizability of topological spaces</i> .....	441
Wilfried Imrich and Mark E. Watkins, <i>On graphical regular representations of cyclic extensions of groups</i> .....	461
Jozef Krasinkiewicz, <i>Remark on mappings not raising dimension of curves</i> .....	479
Melven Robert Krom, <i>Infinite games and special Baire space extensions</i> .....	483
S. Leela, <i>Stability of measure differential equations</i> .....	489
M. H. Lim, <i>Linear transformations on symmetric spaces</i> .....	499
Teng-Sun Liu, Arnoud C. M. van Rooij and Ju-Kwei Wang, <i>On some group algebra modules related to Wiener's algebra <math>M_1</math></i> .....	507
Dale Wayne Myers, <i>The back-and-forth isomorphism construction</i> .....	521
Donovan Harold Van Osdol, <i>Extensions of sheaves of commutative algebras by nontrivial kernels</i> .....	531
Alan Rahilly, <i>Generalized Hall planes of even order</i> .....	543
Joylyn Newberry Reed, <i>On completeness and semicompleteness of first countable spaces</i> .....	553
Alan Schwartz, <i>Generalized convolutions and positive definite functions associated with general orthogonal series</i> .....	565
Thomas Jerome Scott, <i>Monotonic permutations of chains</i> .....	583
Eivind Stensholt, <i>An application of Steinberg's construction of twisted groups</i> .....	595
Yasuji Takeuchi, <i>On strongly radical extensions</i> .....	619
William P. Ziemer, <i>Some remarks on harmonic measure in space</i> .....	629
John Grant, <i>Corrections to: "Automorphisms definable by formulas"</i> .....	639
Peter Michael Rosenthal, <i>Corrections to: "On an inversion for the general Mehler-Fock transform pair"</i> .....	640
Carl Clifton Faith, <i>Corrections to: "When are proper cyclics injective"</i> .....	640