

Pacific Journal of Mathematics

THE ORTHOCENTRIC SIMPLEX AS AN EXTREME SIMPLEX

LEON GERBER

THE ORTHOCENTRIC SIMPLEX AS AN EXTREME SIMPLEX

LEON GERBER

Let \mathcal{A} be a variable n -simplex containing a fixed point Q and having vertices A_i and corresponding opposite faces \mathcal{A}_i , $i = 0, 1, \dots, n$. We use the properties of orthocentric simplexes to present brief solutions to the following problems and obtain several Erdős-Mordell type inequalities as a by-product, some of which are stronger than known inequalities.

(i) Maximize the volume of \mathcal{A} given the distances $QA_i = d_i \geq 0$, $i = 0, \dots, n$.

(ii) Minimize the volume of \mathcal{A} given the distances $e_i \geq 0$ from Q to \mathcal{A}_i , $i = 0, \dots, n$.

(iii) Find the extrema of (i) and (ii) when only the power means of the distances are given.

(iv) Construct an orthocentric simplex given the lengths of the altitudes.

(v) Maximize the volume of \mathcal{A} given the $(n-1)$ -dimensional volumes of the faces.

(vi) Find the maximum in (i) given that Q must be the centroid of \mathcal{A} .

(vii) Maximize the volume of the convex hull of a skew $(n+1)$ -gon given the power means of its edges.

1. Introduction. Problems (i) and (ii) have been solved recently [1, 15]. The present solution is much shorter and shows the relation between the two problems. Problem (iii) is a generalization of [14].

The relation between the three-dimensional version of problems (iv) and (v) was first noticed in 1773 by Lagrange [11] and the n -dimensional problems completely solved in 1866 by C. W. Borchardt [4]. Unaware of this latter paper, in 1953 M. A. Marmion [12] solved the three-dimensional problems anew, obtaining similar results. The present paper obtains Borchardt's results for (iv) and (v) and shows that problem (vi) and problem (vii), which generalizes the result of [13], are also related to these two.

All these solutions depend on the properties of an orthocentric simplex, that is, a simplex whose altitudes A_iH_i , $i = 0, \dots, n$, concur at its orthocenter. (In general, the altitudes of a simplex are not concurrent but are associated [8, 9] a weaker property.) The following facts about an orthocentric simplex \mathcal{A} with orthocenter H are known [7] and easily proved. Let $\mathcal{J} = \{0, 1, \dots, n\}$, $\mathcal{J}' = \{1, \dots, n\}$ and let the summa-

tions and products Σ, Π have range \mathcal{J} and let Σ', Π' have range \mathcal{J}' .

(1.1) Each edge of \mathcal{A} is perpendicular to the opposite $(n-2)$ -face and conversely, any simplex with this property is orthocentric.

(1.2) There exist numbers $a_i, i \in \mathcal{J}$, such that $A_i A_j^2 = a_i + a_j, i \neq j$; and conversely, any simplex which has such a representation is orthocentric. Define $a^{-1} = \Sigma a_i^{-1}$. The numbers $a_i, i \in \mathcal{J}$, and a are called the parameters of \mathcal{A} .

(1.3) If H is interior to \mathcal{A} , then $a_i > 0, i \in \mathcal{J}$, and an isometric copy of \mathcal{A} can be constructed in $(n+1)$ -space by locating A_i at a distance $a_i^{1/2}$ along the i th coordinate axis, $i \in \mathcal{J}$. The coordinates of H are $(aa_0^{-1/2}, \dots, aa_n^{-1/2})$ and the distance of H from the origin is $a^{1/2}$. If H is exterior to \mathcal{A} , then some vertex, say A_0 , is the interior orthocenter of $HA_1 \cdots A_n$. In this case $a < a_0 < 0$ and $a_i > -a, i \in \mathcal{J}'$. An isometric copy of \mathcal{A} can be constructed in $(n+1)$ -space by locating H at a distance $(-a)^{1/2}$ along the zero coordinate axis and A_i at a distance $a_i^{1/2}$ along the i th coordinate axis, $i \in \mathcal{J}'$. The coordinates of A_0 are $(-a_0(-a)^{-1/2}, -a_{01}^{1/2}, \dots, -a_{0n}^{1/2})$ and the distance of A_0 from the origin is $(-a_0)^{1/2}$. If H lies on \mathcal{A} , then H is in fact one of the vertices, say A_0 , so that the edges $A_0 A_i, i \in \mathcal{J}'$, are mutually perpendicular, i.e. \mathcal{A} is a right simplex. In this case $a = a_0 = 0$ and $a_i > 0, i \in \mathcal{J}'$. An isometric copy of \mathcal{A} can be constructed in n -space by locating A_0 at the origin and A_i at a distance $a_i^{1/2}$ along the i th coordinate axis, $i \in \mathcal{J}'$.

Thus in particular, the parameters determine \mathcal{A} to within an isometry.

$$(1.4) \quad HA_i^2 = a_i - a, i \in \mathcal{J}.$$

$$(1.5) \quad \vec{HA}_i \cdot \vec{HA}_j = a, i \neq j.$$

$$(1.6) \quad \text{The volume of } \mathcal{A} \text{ is}$$

$$(n!)^{-1}(a^{-1}\Pi a_i)^{\frac{1}{2}}, \quad \text{if } a \neq 0, \text{ and}$$

$$(n!)^{-1}(\Pi' a_i)^{\frac{1}{2}}, \quad \text{if } a = a_0 = 0.$$

(1.7) $A_i H \cdot H H_i = a$, where the segments are considered to be directed.

(1.8) A simplex is regular if and only if it is orthocentric and the a_i are equal, $i \in \mathcal{J}$.

2. Solution to problem (i).

THEOREM 2.1. *The maximum volume of a simplex \mathcal{A} containing the point Q given the distances $QA_i = d_i \geq 0, i \in \mathcal{J}$, is attained by an orthocentric simplex. Assume $d_0 \leq d_1 \leq \dots \leq d_n$. If $d_0 > 0$, the param-*

ter a and the volume are determined by (2.1) and (2.2) below. If $d_0 = 0$, \mathcal{A} is a right simplex and the volume is given by (2.3).

Proof. A standard compactness argument shows that the maximum is attained.

Suppose $d_0 > 0$. Let \mathcal{A} be a maximizing n -simplex. If for some i , line QA_i is not perpendicular to \mathcal{A}_i , then the i th altitude and thus the volume can be increased by moving A_i (on a sphere of radius d_i with center at Q) to make it perpendicular, contradicting the maximality of \mathcal{A} . Thus the maximum is attained when Q is the orthocenter of \mathcal{A} . By (1.4),

$$a_i = a + QA_i^2 = a + d_i^2, \quad i \in \mathcal{I}.$$

To find the parameter a , we need only solve the equation

$$(2.1) \quad 1 = \sum a/a_i = \sum a/(a + d_i^2).$$

As a increases from 0 to infinity, the right side increases monotonely from 0 to $n + 1$ so there is a unique positive root. Thus the parameters are uniquely determined and the maximizing simplex is unique up to an isometry via (1.3). The maximum volume is

$$(2.2) \quad v_{\max} = (n!)^{-1} a^{-\frac{1}{2}} \prod (a + d_i^2)^{\frac{1}{2}}$$

and is attained when $\cos(A_iQA_i) = a/d_id_i$ via (1.5).

If $d_0 = 0$ and $d_1 > 0$, the simplex has Q as vertex A_0 and is clearly maximized when the lines QA_i , $i \in \mathcal{I}'$, are mutually orthogonal (Hadamard inequality). The maximum is then

$$(2.3) \quad v_{\max} = (n!)^{-1} \prod' d_i.$$

(Note that the function v_{\max} defined by (2.2) and (2.3) is continuous at $d_0 = 0$.) If $d_1 = 0$, the simplex is degenerate and the volume is zero.

In particular, by setting $d_i = R$, $i \in \mathcal{I}$, we have

THEOREM 2.2. *The volume of an n -simplex inscribed in a sphere of radius R satisfies*

$$(2.4) \quad v \leq (n!)^{-1} (n + 1)^{\frac{1}{2}(n+1)} n^{-\frac{1}{2}n} R^n$$

with equality only for the regular simplex.

3. Solution to problem (ii).

THEOREM 3.1. *The minimum volume of the simplex \mathcal{A} given the distances $e_i \geq 0$ from Q to \mathcal{A}_i , $i \in \mathcal{J}$, is attained by an orthocentric simplex. Assume $e_0 \leq e_1 \leq \dots \leq e_n$. The parameter and volume are determined by (3.1) and (3.2) below.*

Proof. First we prove the existence of a minimizing simplex. Let B_i be the foot of the perpendicular from Q to \mathcal{A}_i so that $QB_i = e_i$, and \mathcal{A}_i is tangent at B_i to the sphere \mathcal{S}_i having center Q and radius e_i , $i \in \mathcal{J}$. If $e_0 = e_1 = 0$, the minimum volume is zero. If $e_0 = 0$ and $e_1 > 0$, then the volume exceeds one half of the volume of a sphere with radius e_1 . The reciprocal of the volume is a continuous function of the B_i and hence attains its maximum since the B_i lie in a compact set.

Let \mathcal{A} be a minimizing simplex. We shall use a device of M. M. Day [6, Lemma 4.2] to show that B_i coincides with the centroid G_i of \mathcal{A}_i , $i \in \mathcal{J}$. Suppose, for some i , that B_i and G_i are distinct. Let \mathcal{L} be the $(n-2)$ -flat of points of \mathcal{A}_i equidistant from B_i and G_i . Then \mathcal{A}_i can be rotated about \mathcal{L} through a small angle θ to a position \mathcal{A}'_i which does not intersect \mathcal{S}_i . Since the moment m of \mathcal{A}_i about \mathcal{L} is not zero, for small θ the decrease in volume due to replacing \mathcal{A}_i by \mathcal{A}'_i (B_i is moved away from Q so G_i is moved to the interior of \mathcal{A}) is approximately $m\theta$. Translating \mathcal{A}'_i to touch \mathcal{S}_i decreases the volume further. This contradicts the hypothesis that \mathcal{A} was minimizing. Thus the minimum occurs when the perpendiculars from Q to the faces meet them in their centroids.

The dilatation with constant $-n$ and with center at the centroid of \mathcal{A} carries the centroid of each face into the corresponding opposite vertex and hence carries the perpendiculars QB_i into the altitudes of \mathcal{A} . Since these altitudes concur at the image of Q , it follows that \mathcal{A} is orthocentric and the orientation of the perpendiculars is the same as in the solution to (i). The minimum volume is n^n times the values of v_{\max} , obtained by substituting the e_i for the d_i in equations (2.2) and (2.3), namely,

$$(3.1) \quad \begin{aligned} v_{\min} &= (n!)^{-1} n^n a^{-\frac{1}{2}} \prod (a + e_i^2)^{\frac{1}{2}}, & \text{if } e_0 > 0, \\ &= (n!)^{-1} n^n \prod e_i, & \text{if } e_0 = 0, \end{aligned}$$

where a is the unique positive root of

$$(3.2) \quad 1 = \sum a/(a + e_i^2).$$

In particular, by setting $e_i = r$, $i \in \mathcal{J}$, we have

THEOREM 3.2. *The volume of an n -simplex with inscribed sphere of radius r satisfies*

$$(3.3) \quad v = \geq (n!)^{-1}(n+1)^{\frac{1}{2}(n+1)} n^{\frac{1}{2}n} r^n$$

with equality only for the regular simplex.

Since we do not solve (2.1) or (3.2) explicitly, we obtain some bounds for the positive root of (3.2) which will prove useful later.

THEOREM 3.3. *The unique positive root of (3.2) satisfies*

$$(3.4) \quad \frac{1}{n} e_0 e_1 \leq a \leq \frac{1}{k} \left(\prod_{i=0}^k e_i^2 \right)^{1/(k+1)}, \quad k \in \mathcal{F}'.$$

Proof. Let $f(a)$ denote the right side of (3.2); we have

$$f\left(\frac{1}{n} e_0 e_1\right) \leq \frac{\frac{1}{n} e_0 e_1}{\frac{1}{n} e_0 e_1 + e_0^2} + \frac{n\left(\frac{1}{n} e_0 e_1\right)}{\frac{1}{n} e_0 e_1 + e_1^2} \leq 1$$

and

$$f(e_0 e_1) \geq \frac{e_0 e_1}{e_0 e_1 + e_0^2} + \frac{e_0 e_1}{e_0 e_1 + e_1^2} = 1.$$

Thus $\frac{1}{n} e_0 e_1 \leq a \leq e_0 e_1$. For $k = 2, \dots, n$, let $t_i = e_i^{-2/(k+1)}$, $i = 0, 1, \dots, k$ and $t = \prod_{i=0}^k t_i$. Then by omitting some positive terms and using the inequality of the arithmetic and geometric means with weights t_i^{k-1} we obtain

$$\begin{aligned} f\left(\frac{1}{k} \left(\prod_{i=0}^k e_i^2\right)^{1/(k+1)}\right) &\geq \sum_{i=0}^k (1/kt) / [(1/kt) + t_i^{-(k+1)}] \\ &= \sum_{i=0}^k t_i^{k-1} / (t_i^{k-1} + k t t_i^{-2}) \\ &\geq \left[\sum_{i=0}^k t_i^{k-1} \right]^2 / \left[\sum_{i=0}^k t_i^{k-1} (t_i^{k-1} + k t t_i^{-2}) \right] \\ &= \left[\sum_{i=0}^k t_i^{k-1} \right]^2 / \left[\sum_{i=0}^k t_i^{2k-2} + k t t_i^{k-3} \right] \\ &\geq 1. \end{aligned}$$

The last inequality follows from

$$\begin{aligned} & \left[\sum_{i=0}^k t_i^{k-1} \right]^2 - \left[\sum_{i=0}^k t_i^{2k-2} + k t t_i^{k-3} \right] \\ &= 2 \sum_{0 \leq i < j \leq k} t_i^{k-1} t_j^{k-1} - k \sum_{i=0}^k \left(\prod_{j=0}^k t_j \right) t_i^{k-3} \end{aligned}$$

which is positive since the exponents $(k-1, k-1, 0, \dots, 0)$ in the first term on the right majorize the exponents $(k-2, 1, \dots, 1)$ of the second term in the sense of Muirhead [10, p. 45].

We also note that Newton iteration applied to $f(a) - 1$, with an initial value of zero, seems to converge rapidly.

4. Solution to problem (iii). First we maximize the volume of \mathcal{A} given that the distances $d_i = QA_i$, $i \in \mathcal{J}$, satisfy

$$(4.1) \quad \frac{1}{n+1} \sum d_i^p = s^p$$

for some fixed s and $p > 0$.

We begin by treating two special cases. If some distance, say d_0 , equals zero, the maximum of (2.3) is clearly attained when $d_i = [(n+1)/n]^{1/p} s$, $i \in \mathcal{J}'$, so \mathcal{A} is an isosceles right simplex, and the volume is given by

$$(4.2) \quad v = (n!)^{-1} [(n+1)/n]^{n/p} s^n.$$

In the second case, if $d_i = s$, $i \in \mathcal{J}$, so that \mathcal{A} is a regular simplex, then $a + s^2 = (n+1)a$ from (2.1), so $a = s^2/n$ and the volume is found from (2.2) to be

$$(4.3) \quad v = (n!)^{-1} a^{-\frac{1}{2}} [(n+1)a]^{\frac{1}{2}(n+1)} = (n!)^{-1} n^{-\frac{1}{2}} (n+1)^{\frac{1}{2}(n+1)} s^n.$$

Observe that the value in (4.3) exceeds that in (4.2) provided

$$(4.4) \quad p \geq p_n = 2 / \left[1 + \frac{\log(n+1)}{n \log(n+1) - n \log n} \right].$$

(We have $p_1 = 1$ and $p_n < 2/[1 + \log(n+1)] < 1$ for $n \geq 2$ since $[(n+1)/n]^n < e$.) We may now state

THEOREM 4.1. *The maximum volume of \mathcal{A} subject to (4.1) is attained only by a regular simplex and is given by (4.3) if p satisfies*

(4.4). Otherwise it is attained only by an isosceles right simplex and is given by (4.2).

Proof. Again it is clear that the maximum is attained. Now the set of points (d_0, \dots, d_n) under consideration consists of that part of the surface in $(n+1)$ -dimensional space defined by (4.1) and bounded by the coordinate primes $d_i = 0$, $i \in \mathcal{J}$. On the interior of this set, i.e. when no $d_i = 0$, the maximum volume, v_{\max} , of a simplex with these d_i is given by (2.2). An easy application of the method of Lagrange to $\log(v_{\max})$ with (2.1) and (4.1) as constraints shows that the only relative extremum is when $d_i = s$, $i \in \mathcal{J}$. Thus if $p \geq p_n$ the relative extremum (4.3) is the maximum and if $p < p_n$ the boundary value (4.2) is the maximum.

If (4.1) is modified so $p \leq 0$, the volume is unbounded. In fact, for $d_i = d$, $i \in \mathcal{J}'$, arbitrarily large, we can pick $d_0 > 0$ to satisfy (4.1). Select A_i , $i \in \mathcal{J}'$, so that $QA_1 \cdots A_n$ is an isosceles right simplex of volume $(n!)^{-1} d^n$. We can then choose A_0 so that \mathcal{A} contains this simplex.

To solve the second part of problem (iii) we minimize the volume of \mathcal{A} given that the distances e_i from Q to A_i , $i \in \mathcal{J}$, satisfy for some fixed $t > 0$,

$$(4.5) \quad \frac{1}{n+1} \sum e_i^p = t^p, \text{ for some fixed } p < 0, \text{ or}$$

$$(4.6) \quad \prod e_i = t^{n+1}.$$

(For positive p we can attain a minimum of zero by setting $e_0 = e_1 = 0$.) It follows from (4.5) that $e_0 > (n+1)^{1/p} t$, and as e_0 approaches this value, the others approach infinity as does the corresponding v_{\min} .

Since $e_0 > 0$, it follows from (3.4) and (3.2) that

$$a^n \leq e_0^n e_1^n \leq e_0^n \prod' e_i = e_0^{n-1} t^{n+1}.$$

Then

$$n! n^{-n} v_{\min} = a^{-\frac{1}{2}} \prod (a + e_i^2)^{\frac{1}{2}} > (e_0^{n-1} t^{n+1})^{-1/2n} t^{n+1}$$

so here too the minimum is not attained for small values of e_0 . Thus, via Lagrange we have

THEOREM 4.2. *The minimum volume of \mathcal{A} subject to (4.5) or (4.6) is attained when $e_i = t$, $i \in \mathcal{J}$, and is given by*

$$(4.7) \quad v = (n!)^{-1} n^{\frac{1}{2}n} (n+1)^{\frac{1}{2}(n+1)} t^n.$$

This completes the solution of problem (iii). Using (4.3) and (4.2) with (4.1), and (4.7) with (4.5) and (4.6) to give bounds on the volume in terms of the power means we obtain

THEOREM 4.3. *In an arbitrary n -simplex of volume v , let Q be a point whose distances from the vertices and faces are d_i and e_i respectively, $i \in \mathcal{J}$. Then*

$$(4.8) \quad \left(\frac{1}{n+1} \sum d_i^p \right)^{1/p} \geq (n!)^{1/n} n^{\frac{1}{2}} (n+1)^{-(n+1)/2n} v^{1/n} \quad \text{for } p \geq p_n,$$

$$(4.9) \quad \left(\frac{1}{n+1} \sum d_i^p \right)^{1/p} \geq (n!)^{1/n} [n/(n+1)]^{1/p} v^{1/n} \quad \text{for } p_n \geq p > 0,$$

$$(4.10) \quad \left(\frac{1}{n+1} \sum e_i^p \right)^{1/p} \leq (n!)^{1/n} n^{-\frac{1}{2}} (n+1)^{-(n+1)/2n} v^{1/n} \quad \text{for } p < 0,$$

$$(4.11) \quad \left(\prod e_i \right)^{1/(n+1)} \leq (n!)^{1/n} n^{-\frac{1}{2}} (n+1)^{-(n+1)/2n} v^{1/n}.$$

Equality holds in (4.8), (4.10) and (4.11) only for the centroid of a regular simplex and in (4.9) only for the vertex of an isosceles right simplex.

The result of [14] is obtained by setting $p = 1$ in (4.8). Combining (4.8) with (2.4) and (4.11) with (3.3) we get the following theorem, the first part of which improves the result of [3].

THEOREM 4.4. *For the interior point Q of a simplex of inradius r and circumradius R ,*

$$\left(\frac{1}{n+1} \sum d_i^p \right)^{1/p} \geq nr \quad \text{for } p \geq p_n,$$

and

$$\left(\prod e_i \right)^{1/(n+1)} \leq R/n,$$

with equality only for the centroid of a regular simplex.

5. On the existence of simplexes with specified faces or altitudes. We shall need the following extensions of the triangle inequality to n dimensions; v_k denotes the k -dimensional volume function.

THEOREM 5.1. *Let $f_0 \geq f_1 \geq \dots \geq f_n > 0$ be $n+1$ numbers. There*

exists an n -simplex \mathcal{A} such that $v_{n-1}(\mathcal{A}_i) = f_i$, $i \in \mathcal{J}$, if and only if $f_0 < \sum' f_i$.

Proof. If \mathcal{A} is a simplex with $v_{n-1}(\mathcal{A}_i) = f_i$, $i \in \mathcal{J}$, and f_i is a vector normal to \mathcal{A}_i in the outbound direction with magnitude f_i , $i \in \mathcal{J}$, then [8]

$$(5.1) \quad -f_0 = \sum' f_i.$$

(Applying the Gauss divergence theorem to a constant function on \mathcal{A} and its boundary gives this result immediately.) The f_i are called the vector hyperareas of \mathcal{A} . Since the f_i , $i \in \mathcal{J}'$, are linearly independent, the triangle inequality for vectors yields $f_0 < \sum' f_i$.

Conversely, if $f_0 < \sum' f_i$, we can construct a skew $(n+1)$ -gon $P_0P_1 \cdots P_nP_{n+1}$ where $P_{n+1} = P_0$ with $P_iP_{i+1} = f_i$, $i \in \mathcal{J}$. For an arbitrary point Q , define Q_i by $\vec{QQ}_i = \vec{P_iP_{i+1}}$ and construct a simplex \mathcal{B} by taking a prime perpendicular to line QQ_i at Q_i , $i \in \mathcal{J}$. For some k_i , $i \in \mathcal{J}$, the vectors $k_i\vec{QQ}_i$ are normal to the faces \mathcal{B}_i of \mathcal{B} in the outbound direction with magnitude equal to $v_{n-1}(\mathcal{B}_i)$. In view of (5.1) we have

$$(5.2) \quad -k_0\vec{QQ}_0 = \sum' k_i\vec{QQ}_i.$$

However, $\sum \vec{QQ}_i = \sum \vec{P_iP_{i+1}} = \mathbf{0}$ so

$$(5.3) \quad -\vec{QQ}_0 = \sum \vec{QQ}_i.$$

Since, by skewness, the vectors \vec{QQ}_i , $i \in \mathcal{J}'$, are independent, comparison of (5.2) and (5.3) shows that all the k_i must be equal. Thus the $v_{n-1}(\mathcal{B}_i)$ are proportional to the QQ_i , i.e. to the given f_i . A dilatation completes the construction and the proof.

It is clear from the construction that if the f_i are equal there is still a great deal of freedom and the resulting simplex need not be regular.

COROLLARY 5.2. *Let f_i , $i \in \mathcal{J}$, be given vectors. There exists an n -simplex \mathcal{A} whose vector hyperareas are the f_i if and only if $\sum f_i = \mathbf{0}$.*

COROLLARY 5.3 [2]. *For any n -simplex \mathcal{A} , with $v_{n-1}(\mathcal{A}_i) = f_i$, $i \in \mathcal{J}$, $f_0^2 = \sum' f_i^2 + 2 \sum_{1 \leq i < j \leq n} f_i f_j \cos(a_i, a_j)$.*

Proof. Take the dot product of each side of (5.1) with itself.

COROLLARY 5.4. *Theorem of Pythagoras.* If \mathcal{A} is a right simplex with orthocenter at A_0 , then $f_0^2 = \sum' f_i^2$.

Since the altitudes are inversely proportional to the $(n - 1)$ volumes of the faces, we have the following.

COROLLARY 5.5. Let $0 \leq h_0 \leq h_1 \leq \cdots \leq h_n$ be numbers. There exists an n -simplex \mathcal{A} such that the altitude from A_i has length h_i , $i \in \mathcal{I}$, if and only if $h_0^{-1} < \sum' h_i^{-1}$.

COROLLARY 5.6. If \mathcal{A} is a right simplex with orthocenter $H = A_0$, then $h_0^{-2} = \sum' h_i^{-2} = \sum' A_0 A_i^{-2}$.

In conclusion we remark that Corollary 5.2 has the following physical interpretation. If a joint in a three dimensional structure is to be in equilibrium under the action of four forces, it is possible to construct a tetrahedral bearing plate with faces normal to the forces and such that the bearing load is uniform.

6. Solution to problem (iv). Essentially we are seeking the parameters of an orthocentric simplex given its altitudes.

LEMMA 6.1. Let \mathcal{A} be an orthocentric simplex with parameters, altitudes and hyperareas $a_i, h_i, f_i, i \in \mathcal{I}$. Then the following conditions are equivalent.

$$(6.1) \quad a_0 \leq a_1 \leq \cdots \leq a_n,$$

$$(6.2) \quad h_0 \leq h_1 \leq \cdots \leq h_n,$$

$$(6.3) \quad f_0 \geq f_1 \geq \cdots \geq f_n$$

and strict inequality corresponds to strict inequality.

Proof. Assume (6.1). Since the f_i are inversely proportional to the h_i , we need only show the equivalence of (6.1) and (6.2). If $a = 0$, then \mathcal{A} is a right simplex, $a_0 = 0$, $h_i^2 = A_0 A_i^2 = a_i$, $i \in \mathcal{I}'$, and $h_0^{-2} = \sum' h_i^{-2}$ so the equivalence is immediate. If $a \neq 0$, then by (1.7) and (1.4)

$$\begin{aligned} (6.4) \quad h_i^2 &= (A_i H + H H_i)^2 = A_i H^2 + 2 A_i H \cdot H H_i + H H_i^2 \\ &= A_i H^2 + 2a + a^2 / A_i H^2 = (a_i - a) + 2a + a^2 / (a_i - a) \\ &= a_i^2 / (a_i - a), \quad i \in \mathcal{I}. \end{aligned}$$

That is, $ah_i^{-2} = aa_i^{-1}(1 - aa_i^{-1})$, $i \in \mathcal{J}$, subject to $\Sigma aa_i^{-1} = 1$. With $y_i = ah_i^{-2}$ and $x_i = aa_i^{-1}$, $i \in \mathcal{J}$, we have

$$y_i = x_i(1 - x_i) \text{ subject to } \Sigma x_i = 1.$$

Suppose $a > 0$ so $x_i > 0$, $i \in \mathcal{J}$. If $x_i \leq \frac{1}{2}$, $i \in \mathcal{J}$, then clearly $\text{sign}(y_i - y_j) = \text{sign}(x_i - x_j)$. If $x_0 > \frac{1}{2}$, then $x_0 + x_i < 1$ implies that $x_0 - \frac{1}{2} < \frac{1}{2} - x_i$, $i \in \mathcal{J}'$. That is, x_0 is closest to $\frac{1}{2}$ and y_0 is the largest. For $i \in \mathcal{J}'$, $x_i < \frac{1}{2}$ and the properties follow.

Suppose $a < 0$ so $x_i < 0$, $i \in \mathcal{J}'$, and $x_0 > 1$. Since $x_0 > \frac{1}{2}$, the order properties of the x_i and y_i again follow and the equivalence of (6.1) and (6.2) is immediate.

COROLLARY 6.2. *An orthocentric simplex with equal altitudes or hyperareas is regular.*

COROLLARY 6.3. *In an orthocentric simplex \mathcal{A} , the orthocenter H lies closer to a face than to the corresponding vertex on all except possibly the shortest altitude.*

Proof. If A_0 is interior to $HA_1 \cdots A_n$, then for $i \in \mathcal{J}'$, H and A_i are on opposite sides of \mathcal{A}_i . If \mathcal{A} is a right simplex, the result is trivial. If H is interior to \mathcal{A} , then from (1.3) it follows that the barycentric coordinates of H with respect to \mathcal{A} are $(aa_0^{-1}, \dots, aa_n^{-1})$ and only the largest of these can exceed $\frac{1}{2}$.

We now state the solution to problem (iv).

THEOREM 6.4. *Let $0 < h_0 \leq h_1 \leq \dots \leq h_n$ be given numbers with $h_0^{-1} < \Sigma' h_i^{-1}$. Then there is an orthocentric simplex \mathcal{A} whose i th altitude A_iH_i has length h_i , $i \in \mathcal{J}$. \mathcal{A} is unique to within an isometry.*

Further, let

$$(6.5) \quad f(a) = \epsilon(1 - 4h_0^{-2}a)^{\frac{1}{2}} + \Sigma' (1 - 4h_i^{-2}a)^{\frac{1}{2}} - (n - 1).$$

If $h_0^{-2} < \Sigma' h_i^{-2}$, then the orthocenter H will be interior to \mathcal{A} and the parameter a is the unique positive root of $f(a) = 0$ where $\epsilon = \pm 1$ according as $\Sigma' (1 - h_0^2 h_i^{-2})^{\frac{1}{2}} - (n - 1)$ is negative or positive. If $h_0^{-2} > \Sigma' h_i^{-2}$, then A_0 will be interior to $HA_1 \cdots A_n$ and a is the unique negative root of $f(a) = 0$ with $\epsilon = -1$. In either case,

$$a_0 = \frac{1}{2} h_0^2 [1 + \epsilon(1 - 4ah_0^{-2})^{\frac{1}{2}}]$$

and

$$a_i = \frac{1}{2} h_i^2 [1 + (1 - 4ah_i^{-2})^{\frac{1}{2}}], \quad i \in \mathcal{J}'.$$

If $h_0^{-2} = \Sigma' h_i^{-2}$, then \mathcal{A} is a right simplex with $H = A_0$, $a = a_0 = 0$ and $a_i = h_i^2$, $i \in \mathcal{J}'$.

Proof. First we consider a right simplex with orthocenter $H = A_0$. From facts (1.3) and (1.2) we see that $a = a_0 = 0$ and for $i \in \mathcal{J}'$, the altitude $A_i H_i$ coincides with the edges $A_0 A_i$ so $a_i = A_0 A_i^2 - a_0 = h_i^2$. From Corollary 5.6 we have $h_0^{-2} = \Sigma' h_i^{-2}$.

Suppose now \mathcal{A} is an orthocentric simplex which is not right. Solving (6.4) for a_i we find

$$a_i = \frac{1}{2} h_i^2 [1 + \epsilon_i (1 - 4ah_i^{-2})^{\frac{1}{2}}], \quad \epsilon_i = \pm 1,$$

and

$$(6.6) \quad 1 - 2aa_i^{-1} = \epsilon_i (1 - 4ah_i^{-2})^{\frac{1}{2}}.$$

It follows from Lemma 6.1 that $a_0 \leq a_1 \leq \dots \leq a_n$ and that $aa_i^{-1} < \frac{1}{2}$ for $i \in \mathcal{J}'$, so $\epsilon_i = +1$, $i \in \mathcal{J}'$. Summing (6.6) over i , using $\Sigma aa_i^{-1} = 1$, and replacing ϵ_0 by ϵ we obtain $f(a) = 0$ where $f(a)$ is given by (6.5). We assert that if $h_0^{-2} < \Sigma' h_i^{-2}$, then only one choice of ϵ will yield a positive root and for that choice the solution is unique; and if $h_0^{-2} > \Sigma' h_i^{-2}$, then neither choice of ϵ will yield a positive root.

Suppose $\epsilon = -1$. Rearranging,

$$\begin{aligned} f(a) &= -[(1 - 4h_0^{-2}a)^{\frac{1}{2}} - 1] + \sum' [(1 - 4h_i^{-2}a)^{\frac{1}{2}} - 1] \\ &= 4a \left\{ \frac{h_0^{-2}}{(1 - 4h_0^{-2}a)^{\frac{1}{2}} + 1} - \sum' \frac{h_i^{-2}}{(1 - 4h_i^{-2}a)^{\frac{1}{2}} + 1} \right\} \end{aligned}$$

and $f(0) = 0$. Further, the derivative

$$f'(a) = 2 \left\{ \frac{h_0^{-2}}{(1 - 4h_0^{-2}a)^{\frac{1}{2}}} - \sum' \frac{h_i^{-2}}{(1 - 4h_i^{-2}a)^{\frac{1}{2}}} \right\}$$

and $f'(0) = 2(h_0^{-2} - \Sigma' h_i^{-2})$. For $i \in \mathcal{J}'$, $h_0^{-2} \geq h_i^{-2}$ so for $a > 0$

$$\frac{(1 - 4h_0^{-2}a)^{\frac{1}{2}}}{(1 - 4h_i^{-2}a)^{\frac{1}{2}}} \leq \frac{(1 - 4h_0^{-2}a)^{\frac{1}{2}} + 1}{(1 - 4h_i^{-2}a)^{\frac{1}{2}} + 1}$$

A term by term comparison shows that

$$2af'(a)(1 - 4h_0^{-2}a)^{\frac{1}{2}} \geq f(a)[(1 - 4h_0^{-2}a)^{\frac{1}{2}} + 1].$$

Thus $f'(a) > 0$ whenever $f(a) \geq 0$ (unless all the h_i are equal in which case it is clear the only root is zero) so there is at most one positive root. If $h_0^{-2} > \sum' h_i^{-2}$, then $f'(0) > 0$ and $f(a)$ and $f'(a)$ are positive $a > 0$, so there are no positive roots. Suppose $h_0^{-2} < \sum' h_i^{-2}$ so $f'(a) < 0$. Thus if the right endpoint value $f(\frac{1}{4}h_0^2) \geq 0$ there is precisely one positive root, and if $f(\frac{1}{4}h_0^2) < 0$ there is no positive root.

Suppose $\epsilon = 1$. Then $f(0) = 2$ and $f(a)$ is monotone decreasing so if $f(\frac{1}{4}h_0^2) > 0$ there is no positive root, and if $f(\frac{1}{4}h_0^2) \leq 0$ there is a unique positive root. Further, if $h_0^{-2} > \sum' h_i^{-2}$, then $h_0 < h_i$, $i \in \mathcal{J}'$, and

$$f(\frac{1}{4}h_0^2) = 1 - h_0^2 \sum' \frac{h_i^{-2}}{1 + (1 - h_0^2 h_i^2)^{\frac{1}{2}}} > 1 - h_0^2 \sum' h_i^{-2} \geq 0$$

so there are no positive roots here either.

The case $\epsilon = 1$ corresponds to $aa_0^{-1} < \frac{1}{2}$, i.e. H nearer to \mathcal{A}_0 and the case $\epsilon = -1$ corresponds to H nearer A_0 . If $\frac{1}{4}h_0^2$ is a root, H is the midpoint of A_0H_0 , and the sign of ϵ is immaterial, a petty infringement on uniqueness.

The search for negative roots is considerably simplified by the fact that $a < a_0 < 0$ and $a_i > 0$, $i \in \mathcal{J}'$, together with (6.6) imply $\epsilon_0 = -1$ and $\epsilon_i = +1$, $i \in \mathcal{J}'$. In this case we observe that at $-\infty$,

$$f(a) \sim 2(-a)^{\frac{1}{2}} \left[\sum' h_i^{-1} - h_0^{-1} \right] > 0$$

and it follows as above that there will be a negative root if and only if

$$f'(0) = 2 \left(h_0^{-2} - \sum' h_i^{-2} \right) \geq 0.$$

7. Solution to the remaining problems.

THEOREM 7.1. *Let f_i , $i \in \mathcal{J}$, be numbers. The simplex \mathcal{A} of maximum volume for which $v_{n-1}(a_i) = f_i$, $i \in \mathcal{J}$, is orthocentric with altitudes proportional to f_i^{-1} , $i \in \mathcal{J}$. If Q is a given point, the simplex \mathcal{B} of maximal volume such that $QB_i = f_i$, $i \in \mathcal{J}$, and that Q is the centroid of \mathcal{B} is constructed by setting the vectors $\vec{QB_i}$ equal to the vector hyperareas of \mathcal{A} .*

THEOREM 7.2. *Let $s > 0$ and $p \geq p_n$ be fixed, where $p_n \leq 1$ is given by (4.4). The n -simplex \mathcal{C} of maximal volume such that*

$$\frac{1}{n+1} \sum C_i C_{i+1}^p = s^p.$$

is a regular simplex.

Proofs. If the given hyperareas satisfy the extended triangle inequality there are simplexes whose faces have the prescribed volume and it is clear that a maximizing simplex exists. (The isoperimetric inequality states that the n -volume of a simplex with these hyperareas is less than that of a sphere with surface Σf_i , but the following crude estimate will also suffice: the incircled sphere of any such simplex obviously has surface less than Σf_i and the n -volume of the simplex is $1/n \Sigma f_i$ times its radius.) Let \mathcal{A} be any simplex with $v_{n-1}(\mathcal{A}_i) = f_i$, $i \in \mathcal{J}$. Let $h_i = A_i H_i$, $i \in \mathcal{J}$, and let $v = v_n(\mathcal{A})$. Let Q be any point and construct QB_i perpendicular to \mathcal{A}_i with $QB_i = f_i$, $i \in \mathcal{J}$. Since $\Sigma \overrightarrow{QB_i} = \mathbf{0}$ it follows that Q is the centroid of the simplex $\mathcal{B} = B_0 B_1 \cdots B_n$. Let \mathcal{B}_0 be the simplex $QB_1 \cdots B_n$. Then (cf. [5, p. 4])

$$(n!v)(n!v_n(\mathcal{B}_0)) = \det(\overrightarrow{A_0 A_i} \cdot \overrightarrow{QB_j}) = \prod_i' \overrightarrow{A_0 A_i} \cdot \overrightarrow{QB_i} = n^n v^n$$

since the nondiagonal elements vanish and $\overrightarrow{A_0 A_i} \cdot \overrightarrow{QB_i} = h_i QB_i = h_i f_i = nv$. Thus

$$v_n(\mathcal{B}) = (n+1)v_n(\mathcal{B}_0) = (n!)^{-2}(n+1)n^n v^{n-1}.$$

Now construct $\mathcal{C} = C_0 C_1 \cdots C_n$ with C_0 arbitrary and $\overrightarrow{C_{i-1} C_i} = \overrightarrow{QB_i}$, $i \in \mathcal{J}'$. Thus

$$\sum_{i=1}^k \overrightarrow{QB_i} = \sum_{i=1}^k \overrightarrow{C_{i-1} C_i} = \overrightarrow{C_0 C_k}, \quad k \in \mathcal{J}',$$

and in particular $\overrightarrow{C_n C_0} = -\Sigma' \overrightarrow{QB_i} = \overrightarrow{QB_0}$. If the vectors $\overrightarrow{C_0 C_k}$, $k \in \mathcal{J}'$, are chosen as a basis, the volume of \mathcal{B}_0 relative to the volume of \mathcal{C} is a lower triangular determinant with 1's on and below the main diagonal. It follows that $v_n(\mathcal{C}) = v_n(\mathcal{B}_0)$ so if we maximize $v_n(\mathcal{C})$, we maximize $v_n(\mathcal{B})$ and hence $v_n(\mathcal{A})$. Allowing C_0 to vary on the intersection of the spheres determined by $C_n C_0 = QB_0 = f_0$ and $C_0 C_1 = QB_1 = f_1$, it is clear that the altitude from C_0 to the opposite face $\mathcal{C}_0 = C_1 C_2 \cdots C_n$, and hence $v_n(\mathcal{C})$, is maximized when the plane $C_n C_0 C_1$ is perpendicular to \mathcal{C}_0 , i.e. $QB_0 B_1$ is perpendicular to $QB_2 \cdots B_n$ so $A_2 \cdots A_n$ is perpendicular to the lines $A_0 A_1$. Since the labelling is arbitrary, each edge of the maximal \mathcal{A} is perpendicular to the opposite face, so \mathcal{A} is orthocentric (fact 1.1). The orthocentric simplex with altitudes f_i^{-1} , $i \in \mathcal{J}$, constructed as in the solution to (iv), is similar to \mathcal{A} and \mathcal{A} is obtained by a dilatation. This completes the solution to (v). \mathcal{B} can be constructed immediately, which gives the solution to (vi).

Turning to problem (vii), we suppose we want to maximize $v_n(\mathcal{C})$ given the sum of the p th powers, $p \geq p_n$ as in the solution of (iii), of the edges of the skew polygon $C_0C_1 \cdots C_nC_0$. This is equivalent to maximizing $v_n(\mathcal{B})$ given the sum $\sum QB_i^p$ with the constraint that Q is the centroid of \mathcal{B} . But we have already shown in solving (iii) that the unconstrained maximum is attained when \mathcal{B} is a regular simplex and Q is its center. So \mathcal{C} is also regular; this completes the solution of (vii). When $p = 1$, the sum of the p th powers of the edges of \mathcal{C} is just the perimeter of the polygon, so we have the result of [13].

REFERENCES

1. M. M. Ali, *On some extreme simplexes*, Pacific J. Math., **33** (1970), 1–14.
2. P. Bartos, *Cosine theorems on simplexes in E^n* (Czech), Casopis Pest. Mat., **95** (1970), 150–154.
3. J. Berkes, *Einfacher Beweis und Verallgemeinerung einer Dreiecksgleichung*, Elem. Math., **22** (1967), 135–136.
4. C. W. Borchardt, *Über die Aufgabe des Maximum, welche der Bestimmung des Tetraeders von grossten Volumen bei gegebenen Flächeninhalt der Seitenflächen für mehr als drei Dimensionen entspricht*, Math. Abh. Akad. Wis. Berlin, 1866, pp. 121–155 = Gesammelte Werke, Berlin, 1888, pp. 201–232.
5. L. Brand, *Vector Analysis*, Wiley, New York, 1957.
6. M. M. Day, *Polygons circumscribed about closed convex curves*, Trans. Amer. Math. Soc., **62** (1947), 315–319.
7. E. Egervary, *On orthocentric simplexes*, Acta Lith. Sci. Szeged, **9** (1940), 218–266.
8. L. Gerber, *The altitudes of a simplex are associated*, Math. Mag., **46** (1973), 155–157.
9. ———, *Associated and perspective simplexes*, Trans. Amer. Math. Soc., **201** (1975), 43–55.
10. G. H. Hardy, J. E. Littlewood, G. Polya, *Inequalities*, Cambridge U. Press, London, 1934.
11. J. L. Lagrange, *Solutions analytiques de quelques problèmes sur les pyramides triangulaires*, Nouv. Mém. Acad. Sci. Berlin, 1773, pp. 149–176 = Ouvres completes III, pp. 659–692.
12. M. A. Marmion, *Solution to Question 3253*, Mathesis, **62** (1953), 69–72.
13. B. Misek, *The simplex polygon with the greatest volume of its convex hull* (Czech), Casopis Pest. Mat., **84** (1959), 99–104.
14. C. M. Petty and D. Waterman, *An extreme theorem for n -simplexes*, Monatsh. Math., **59** (1955), 320–322.
15. D. Slepian, *The content of some extreme simplexes*, Pacific J. Math., **31** (1969), 795–808.

Received August 2, 1973.

ST. JOHN'S UNIVERSITY

Shimshon A. Amitsur, <i>Central embeddings in semi-simple rings</i>	1
David Marion Arnold and Charles Estep Murley, <i>Abelian groups, A, such that $\text{Hom}(A, - - -)$ preserves direct sums of copies of A</i>	7
Martin Bartelt, <i>An integral representation for strictly continuous linear operators</i>	21
Richard G. Burton, <i>Fractional elements in multiplicative lattices</i>	35
James Alan Cochran, <i>Growth estimates for the singular values of square-integrable kernels</i>	51
C. Martin Edwards and Peter John Stacey, <i>On group algebras of central group extensions</i>	59
Peter Fletcher and Pei Liu, <i>Topologies compatible with homeomorphism groups</i>	77
George Gasper, Jr., <i>Products of terminating ${}_3F_2(1)$ series</i>	87
Leon Gerber, <i>The orthocentric simplex as an extreme simplex</i>	97
Burrell Washington Helton, <i>A product integral solution of a Riccati equation</i>	113
Melvyn W. Jeter, <i>On the extremal elements of the convex cone of superadditive n-homogeneous functions</i>	131
R. H. Johnson, <i>Simple separable graphs</i>	143
Margaret Humm Kleinfeld, <i>More on a generalization of commutative and alternative rings</i>	159
A. Y. W. Lau, <i>The boundary of a semilattice on an n-cell</i>	171
Robert F. Lax, <i>The local rigidity of the moduli scheme for curves</i>	175
Glenn Richard Luecke, <i>A note on quasidiagonal and quasitriangular operators</i>	179
Paul Milnes, <i>On the extension of continuous and almost periodic functions</i>	187
Hidegoro Nakano and Kazumi Nakano, <i>Connector theory</i>	195
James Michael Osterburg, <i>Completely outer Galois theory of perfect rings</i>	215
Lavon Barry Page, <i>Compact Hankel operators and the F. and M. Riesz theorem</i>	221
Joseph E. Quinn, <i>Intermediate Riesz spaces</i>	225
Shlomo Vinner, <i>Model-completeness in a first order language with a generalized quantifier</i>	265
Jorge Viola-Prioli, <i>On absolutely torsion-free rings</i>	275
Philip William Walker, <i>A note on differential equations with all solutions of integrable-square</i>	285
Stephen Jeffrey Willson, <i>Equivariant maps between representation spheres</i>	291