

# Pacific Journal of Mathematics

## **A PRODUCT INTEGRAL SOLUTION OF A RICCATI EQUATION**

BURRELL WASHINGTON HELTON

# A PRODUCT INTEGRAL SOLUTION OF A RICCATI EQUATION

BURRELL W. HELTON

*In Memory of Professor H. S. Wall*

Product integrals are used to show that, if  $dw, G, H$  and  $K$  are functions from number pairs to a normed complete ring  $N$  which are integrable and have bounded variation on  $[a, b]$  and  $v^{-1}$  exists and is bounded on  $[a, b]$ , then the integral equation

$$\beta(x) = w(x) + (LRLR) \int_a^x (\beta H + G\beta + \beta K\beta)$$

has a solution  $\beta(x) = v^{-1}(x)u(x)$  on  $[a, b]$ , where  $u$  and  $v$  are defined by the matrix equation

$$[u(x), v(x)] = [w(a), 1] {}_a\prod^x \left( I + \begin{bmatrix} H & -K \\ dw & -G \end{bmatrix} \right)$$

The above results are used to show that if  $p, q, h$  and  $r$  are quasicontinuous functions from the numbers to  $N$  such that  $h$  is left continuous and has bounded variation and  $p, q$  and  $h$  commute, then the solution on  $[a, b]$  of the differential-type equation  $f^{**} + f^*p + fq = r$  is

$$f(x) = f(a) {}_a\prod^x (1 - \beta dh) + (R) \int_a^x dz {}_z\prod^x (1 - \beta dh),$$

where  $f(x) - f(a) = (L) \int_a^x f^* dh$ ,  $\beta$  is the solution of

$$\beta(x) = (L) \int_a^x q dh + (LL) \int_a^x \beta(-p dh) + (LR) \int_a^x \beta dh \beta,$$

and  $z$  is defined in terms of  $p, q, r, h$  and  $\beta$ .

**1. Introduction.** Adam [1] introduced the concept of continuous continued fractions and showed that the solution of  $y' = g'y^2 - f'$  could be given as a continuous continued fraction, provided  $f'$  and  $g'$  are continuous and positive. Wall [11] [12] showed that, if  $F_{11}, F_{12}, F_{21}$  and  $F_{22}$  are continuous functions of bounded variation from the real numbers to the complex numbers and  $|b - a|$  is sufficiently

small, then the solution of

$$(1) \quad w(x) = z + \int_b^x w^2 dF_{21} + \int_b^x wd(F_{22} - F_{11}) - \int_b^x dF_{12}$$

is  $w(x) = [M_{11}(x, b)z + M_{12}(x, b)][M_{21}(x, b)z + M_{22}(x, b)]^{-1}$ , where  $F = \begin{bmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{bmatrix}$  and  $\begin{bmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{bmatrix}$  is the function such that  $M(x, y) = 1 + \int_x^y M(x, s)dF(s)$ . MacNerney, using the Stieltjes integral in [7] and the subdivision-refinement-type mean integral in [8], extended Wall's results to some types of quasicontinuous linear transformations and showed that the solution of Equation (1) can also be expressed as a continuous continued fraction [8, Theorem 5.3]. In this paper the product integral theory developed by MacNerney [8] [9] and the author [3] is used to find and express (in §3) the solution of

$$\beta(x) = w(x) + (LRLR) \int_a^x (\beta H + G\beta + \beta K\beta)$$

and to find and express (in §4) the solution of

$$f^{**} + f^*p + fq = r,$$

where  $w, p, q, r, G, H, K$  are quasicontinuous functions from numbers or pairs of numbers to a normed complete ring  $N$ .

**2. Definitions and notations.** The symbol  $R$  denotes the set of real numbers and  $N$  is a ring which has an identity element  $1$  and a norm  $|\cdot|$  with respect to which  $N$  is complete and  $|1| = 1$  (henceforth, the symbol  $1$  will be used for this identity element). Functions from  $R$  to  $N$  and from  $R \times R$  to  $N$  will be represented by lower case letters and upper case letters, respectively. All sum and product integrals are subdivision-refinement-type limits. If  $G$  is a function from  $R \times R$  to  $N$ , the product integral of  $G$  exists on  $[a, b]$  iff there exists  $A \in N$  such that if  $\epsilon$  is a positive number then there is a subdivision  $D$  of  $[a, b]$  such that if  $\{x_i\}_0^n$  is a refinement of  $D$  then  $|A - G_1G_2 \cdots G_n| < \epsilon$ , where  $G_i = G(x_{i-1}, x_i)$  for  $i = 1, 2, \dots, n$ . The symbol  ${}_a\Pi^bG$  will be used to represent the limit  $A$ . A similar definition holds for the sum integral. Upper case letters preceding an integral symbol show how the integrand is to be evaluated: i.e.,  $(LRLR) \int_a^b (fH + Gf + fKf) = \int_a^b M$ , where for  $x < y$

$$M(x, y) = f(x)H(x, y) + G(x, y)f(y) + f(x)G(x, y)f(y).$$

Also,  $G \in OA^0$  on  $[a, b]$  iff  $\int_a^b G$  exists and  $\int_a^b |G - fG| = 0$ ;  $G \in OM^0$  on  $[a, b]$  iff  ${}_x\Pi^y(1 + G)$  exists for  $a \leq x \leq y \leq b$  and  $\int_a^b |(1 + G) - \Pi(1 + G)| = 0$ ;  $G \in OB^0$  on  $[a, b]$  iff there is a number  $M$  and a subdivision  $D$  of  $[a, b]$  such that, if  $\{x_i\}_0^n$  is a refinement of  $D$ , then  $\sum_1^n |G(x_{i-1}, x_i)| \leq M$ ; the function  $v^{-1}$  exists on  $[a, b]$  means  $v(x)v(x)^{-1} = v(x)^{-1}v(x) = 1$  for  $x \in [a, b]$ . The function  $G^{-1}$  exists on  $[a, b]$  means there is a subdivision  $\{x_i\}_0^n$  of  $[a, b]$  such that if  $0 < i \leq n$  and  $x_{i-1} \leq x < y \leq x_i$ , then  $G(x, y)^{-1}G(x, y) = G(x, y)G(x, y)^{-1} = 1$ . If  $\{x_i\}_0^n$  is a subdivision, the symbols  $f_{i-1}$ ,  $f_i$ , and  $G_i$  will be used as shorthand notations for  $f(x_{i-1})$ ,  $f(x_i)$  and  $G(x_{i-1}, x_i)$ , respectively. For additional details pertaining to these definitions, see [3], [4], and [9]. The main results of this paper will be designated as theorems; the supporting theorems will be labeled as lemmas.

**3. A Riccati integral equation.** In this section we derive a solution for the integral equation

$$f(x) = w(x) + (LRLR) \int_a^x (fH + Gf + fKf).$$

Since the  $OA^0$  property plays an important role in this paper, please note that the function  $G \in OA^0$  if at least one of the following conditions is satisfied:

- (1) there is a function  $g$  such that

$$G(x, y) = g(y) - g(x);$$

- (2) if  $G(x, y) = f(x)H(x, y)$ , where  $f$  is quasicontinuous and  $H \in OA^0$  and  $OB^0$ , [4, Theorem 2];

- (3) if  $G$  is an integrable function from number pairs to a real Hilbert space which is finite dimensional, [2, Theorem 2].

Also note that, if  $H, K, W, G$  are functions from  $R \times R$  to  $N$  which belong to  $OA^0$  and  $OB^0$ , then  $\begin{bmatrix} H & K \\ W & G \end{bmatrix}$  represents a matrix  $Q$  such that  $Q \in OA^0$  and  $OB^0$  and, by Lemma 3.1,  $Q \in OM^0$ .

**LEMMA 3.1.** *If  $G$  is a function from  $R \times R$  to a normed complete ring and  $G \in OB^0$ , then the following statements are equivalent:*

- (1)  $G \in OA^0$  on  $[a, b]$  and
- (2)  $G \in OM^0$  on  $[a, b]$ .

This is Theorem 3.4 of [3].

**THEOREM 3.2.** *Given. (1)  $[a, b]$  is a number interval. (2)  $w$  is a function from  $R$  to  $N$  and  $H, G$  and  $K$  are functions from  $R \times R$  to  $N$  such that each of  $dw, H, G$  and  $K$  belongs to  $OA^0$  and  $OB^0$ .*

*(3)  $u$  and  $v$  are functions from  $R$  to  $N$  such that if  $x \in [a, b]$  then  $u(x)$  and  $v(x)$  are defined by the matrix equation*

$$[u(x), v(x)] = [w(a), 1] {}_a\prod^x \left( I + \begin{bmatrix} H & -K \\ dw & -G \end{bmatrix} \right);$$

*and  $v^{-1}$  exists and is bounded.*

*(4)  $f$  is a bounded function from  $R$  to  $N$ ,  $f(a) = w(a)$  and  $f(x) = v(x)^{-1}u(x)$  for  $x \in [a, b]$ .*

**Conclusion.** If  $x \in [a, b]$ , then

$$f(x) = w(x) + (LRLR) \int_a^x (fH + Gf + fKf).$$

Furthermore, if  $w$  is a constant function, then

$$f(x) = \left[ {}_a\prod^x (1 - G) - w(a)(LR) \int_a^x {}_a\prod^t (1 + H)K {}_a\prod^x (1 - G) \right]^{-1} \left[ w(a) {}_a\prod^x (1 + H) \right].$$

**Proof.** Let  $Q$  be the function such that  $Q = \begin{bmatrix} 1 + H & -K \\ dw & 1 - G \end{bmatrix}$ ; then  $Q - I \in OA^0$  and  $OB^0$  and, by Lemma 3.1,  $Q - I \in OM^0$ . Suppose  $x \in (a, b]$  and  $\{x_i\}_0^n$  is a subdivision of  $[a, x]$ . If  $0 < i \leq n$ , then there exist  $a_i$  and  $b_i \in N$  such that

$$\begin{aligned} [v(x_i)f(x_i), v(x_i)] &= [u(x_i), v(x_i)] \\ &= [w(a), 1] {}_a\prod^{x_{i-1}} Q_{x_{i-1}} \prod^{x_i} Q \\ &= [u(x_{i-1}), v(x_{i-1})] {}_{x_{i-1}}\prod^{x_i} \begin{bmatrix} 1 + H & -K \\ dw & 1 - G \end{bmatrix} \\ &= [u_{i-1}, v_{i-1}] \begin{bmatrix} 1 + H_i & -K_i \\ \Delta w_i & 1 - G_i \end{bmatrix} + [a_i, b_i] \\ &= v_{i-1} [f_{i-1}, 1] \begin{bmatrix} 1 + H_i & -K_i \\ \Delta w_i & 1 - G_i \end{bmatrix} + [a_i, b_i] \end{aligned}$$

$$= v_{i-1} [f_{i-1}(1 + H_i) + \Delta w_i, -f_{i-1} K_i + (1 - G)] + [a_i, b_i].$$

Therefore,

$$(v^{-1}_{i-1} v_i) f_i = f_{i-1}(1 + H_i) + \Delta w_i + v^{-1}_{i-1} a_i$$

and

$$v^{-1}_{i-1} v_i = -f_{i-1} K_i + 1 - G_i + v^{-1}_{i-1} b_i;$$

hence,

$$(-f_{i-1} K_i + 1 - G_i + v^{-1}_{i-1} b_i) f_i = f_{i-1}(1 + H_i) + \Delta w_i + v^{-1}_{i-1} a_i$$

and

$$f_i - f_{i-1} = \Delta w_i + f_{i-1} H_i + G_i f_i + f_{i-1} K_i f_i - v^{-1}_{i-1} b_i f_i + v^{-1}_{i-1} a_i.$$

Since  $f, u, v$  and  $v^{-1}$  are bounded and since  $\sum_i^n (|a_i| + |b_i|)$  can be made arbitrarily small with an appropriate choice of a subdivision (since  $Q \in OM^0$ ), then the following integral exists and

$$f(x) - f(a) = w(x) - f(a) + (LRLR) \int_a^x (fH + Gf + fKf).$$

Since

$$\prod_i^n \begin{bmatrix} p_i & q_i \\ 0 & r_i \end{bmatrix} = \begin{bmatrix} p & q \\ 0 & r \end{bmatrix},$$

where  $p = \prod_i^n p_i$ ,  $q = \sum_{j=1}^n (\prod_{i=1}^{j-1} p_i) q_j (\prod_{i=j+1}^n r_i)$  and  $r = \prod_{i=1}^n r_i$ , and since all the following integrals and product integrals exist, then

$$[w(a), 1] {}_a \prod \begin{bmatrix} 1 + H & -K \\ 0 & 1 - G \end{bmatrix} = [w(a), 1] \begin{bmatrix} A & B \\ 0 & D \end{bmatrix},$$

where  $A = {}_a \Pi (1 + H)$ ,  $B = (LR) \int_a^x [{}_a \Pi' (+H)](1 - K)[{}_l \Pi^x (1 - G)]$  and  $D = {}_a \Pi (1 - G)$ ; hence, if  $w$  is a constant function, then

$$f(x) = [w(a)B + D]^{-1} [w(a)A].$$

**THEOREM 3.3.** *Given. (1)  $[a, b]$  is a number interval;  
(2)  $w$  is a function from  $R$  to  $N$  and  $H, G$  and  $K$  are functions from*

$R \times R$  to  $N$  such that each of  $dw, H, G$  and  $K$  belongs to  $OA^0$  and  $OB^0$ ;

(3)  $u$  and  $v$  are functions from  $R$  to  $N$  such that, if  $x \in [a, b]$ , then  $u(x)$  and  $v(x)$  are defined by the matrix equation

$$[u(x), v(x)] = [w(a), 1] {}_a\prod^x \left( I + \begin{bmatrix} H & -K \\ dw & -G \end{bmatrix} \right)$$

and  $v(x)^{-1}$  exists;

(4)  $f$  is a bounded function from  $R$  to  $N$ ,  $f(a) = w(a)$ ,  $(1 - G_i - f_{i-1} K_i)^{-1}$  exists and

$$f(x) = w(x) + (LRLR) \int_a^x (fH + Gf + fKf)$$

for  $x \in [a, b]$ .

*Conclusion.* If  $x \in [a, b]$ , then  $f(x) = v(x)^{-1} u(x)$ .

*Proof.* Suppose  $x \in [a, b]$  and  $\{x_i\}_0^n$  is a subdivision of  $[a, b]$ . If  $0 < i \leq n$ , then there exists  $\epsilon_i \in N$  such that

$$\begin{aligned} f(x_i) &= w(x_i) + (LRLR) \int_a^{x_i} (fH + Gf + fKf) \\ &= \Delta w_i + f_{i-1} + f_{i-1} H_i + G_i f_i + f_{i-1} K_i f_i + \epsilon_i \end{aligned}$$

and  $f_i = b_i^{-1} a_i$ , where  $b_i = 1 - G_i - f_{i-1} K_i$  and  $a_i = f_{i-1} (1 + H_i) + (\Delta w_i + \epsilon_i)$ . For  $i = 1, 2, 3, \dots, n$ , let  $R_i$  be the  $2 \times 2$  matrix  $R_i = \begin{bmatrix} 1 + H_i & -K_i \\ \Delta w_i + \epsilon_i & 1 - G_i \end{bmatrix}$ ; let  $a_0 = w(a)$  and  $b_0 = 1$ ; then  $\{a_i\}_0^n$  and  $\{b_i\}_0^n$  are elements of  $N$  such that, if  $0 < i \leq n$ , then  $f_i = b_i^{-1} a_i$  and

$$[a_i, b_i] = [f_{i-1}, 1] R_i = [b_{i-1}^{-1} a_{i-1}, 1] R_i = b_{i-1}^{-1} [a_{i-1}, b_{i-1}] R_i.$$

Therefore

$$[a_n, b_n] = \left( \prod_{i=n}^1 b_{i-1}^{-1} \right) [f_0, 1] \prod_{i=1}^n R_i$$

and

$$(1) \quad \left( \prod_{i=1}^n b_{i-1} \right) b_n [f_n, 1] = \prod_{i=1}^n b_{i-1} [a_n, b_n] = [f_0, 1] \prod_{i=1}^n R_i.$$

Let  $Q$  be the function from  $R \times R$  to the set of  $2 \times 2$  matrices such that  $Q = \begin{bmatrix} 1+H & -K \\ dw & 1-G \end{bmatrix}$ . Since  $f$  is quasicontinuous and since each of  $dw, H, G$  and  $K$  belong to  $OA^0$  and  $OB^0$ , then  $Q - I$  and  $-G - fK \in OA^0$  and  $OB^0$  and it follows from Lemma 3.1 that  $Q - I$  and  $-G - fK$  belong to  $OM^0$ , the corresponding product integrals exist,  $\int_a^b |Q - \Pi Q| = 0$  and  $\int_a^b |(1 - G - fK) - \Pi(1 - G - fK)| = 0$ . For each subdivision  $\{x_i\}_0^n$  of  $[a, x]$ , there exist elements  $d_1, d_2$ , and  $d_3$  such that Equation (1) can be rewritten

$$\left\{ (L) {}_a \prod^x (1 - G - fK) + d_1 \right\} [f_n, 1] = [f_0, 1] \left( {}_a \prod^x Q + d_2 + d_3 \right),$$

where  $1 - G_i - f_{i-1} K_i$  is playing the role of  $b_i$  and

$$d_1 = \prod_{i=1}^n (1 - G_i - f_{i-1} K_i) - (L) {}_a \prod^x (1 - G - fK),$$

$$d_2 = \prod_{i=1}^n Q_i - {}_a \prod^x Q$$

and

$$d_3 = \prod_{i=1}^n R_i - \prod_{i=1}^n Q_i = \sum_{i=1}^n \left( \prod_{j=1}^{i-1} Q_j \right) (R_i - Q_i) \prod_{j=i+1}^n R_j.$$

Since  $R_i - Q_i = \begin{bmatrix} 0 & 0 \\ \epsilon_i & 0 \end{bmatrix}$ , it follows from the  $OM^0$  and  $OA^0$  properties that each of  $|d_1|$ ,  $|d_2|$  and  $|d_3|$  can be made arbitrarily small; hence  $(L) {}_a \Pi^x (1 - G - fK)[f(x), 1] = [f_0, 1] {}_a \Pi^x Q = [u(x), v(x)]$ . It follows from the meaning of equality for matrices that  $(L) {}_a \Pi^x (1 - G - fK) = v(x)$ ,  $v(x)f(x) = u(x)$  and  $f(x) = v(x)^{-1}u(x)$ .

**LEMMA 3.4.** *If  $G \in OB^0$  on  $[a, b]$  and  $\epsilon > 0$ , then there is a number  $p \in (a, b]$  such that, if  $\{x_i\}_0^n$  is a subdivision of  $[a, p]$ , then  $\sum_2^n |G_i| < \epsilon$ .*

**THEOREM 3.5.** *Given.  $H, W, K$  and  $G$  are functions from  $R \times R$  to  $N$  such that each of  $H, W, K$  and  $G$  belongs to  $OA^0$  and  $OB^0$  on  $[a, b]$  and  $u$  and  $v$  are functions from  $R$  to  $N$  and are defined by the matrix equation*

$$[u(x), v(x)] = [u(a), v(a)] {}_a \prod^x \left( I + \begin{bmatrix} H & W \\ K & G \end{bmatrix} \right)$$



for  $x \in [a, b]$ . *Conclusion.* (1) If  $p \in (a, b]$  and  $0 < k < 1$  and  $|v(a) - 1| + \sum_1^n |u_{i-1} W_i + v_{i-1} G_i| < k$  for each subdivision  $\{x_i\}_0^n$  of  $[a, p]$ , then  $v^{-1}$  exists and is bounded on  $[a, p]$ . (2) If  $|v(a) - 1| + |u(a)W(a, a^+) + v(a)G(a, a^+)| < 1$ , then there exists  $p \in (a, b]$  such that  $v^{-1}$  exists and is bounded on  $[a, p]$ .

*Proof.* Since  $H, W, K$  and  $G \in OA^0$  and  $OB^0$  on  $[a, b]$ , then  $\begin{bmatrix} H & W \\ K & G \end{bmatrix} \in OA^0$  and  $OB^0$  on  $[a, b]$  and, by Lemma 3.1,  $\begin{bmatrix} H & W \\ K & G \end{bmatrix} \in OM^0$  on  $[a, b]$ ; also,  $u$  and  $v$  are quasicontinuous and bounded on  $[a, b]$ .

We now prove Conclusion 1. Let  $x \in [a, p]$  and let  $\{x_i\}_1^n$  be a subdivision of  $[a, x]$ . For  $i = 1, 2, \dots, n$ , there exist  $a_i$  and  $b_i \in N$  such that

$$\begin{aligned} [u(x_i), v(x_i)] &= [u(a), v(a)] {}_a\prod^{x_i} \left( I + \begin{bmatrix} H & W \\ K & G \end{bmatrix} \right) \\ &= [u_{i-1}, v_{i-1}] {}_{x_{i-1}}\prod^{x_i} \left( I + \begin{bmatrix} H & W \\ K & G \end{bmatrix} \right) \\ &= [u_{i-1}, v_{i-1}] \left[ \begin{array}{cc} 1 + H_i & W_i \\ K_i & 1 + G_i \end{array} \right] + [a_i, b_i] \\ &= [u_{i-1}(1 + H_i) + v_{i-1}K_i, u_{i-1}W_i + v_{i-1} + v_{i-1}G_i] + [a_i, b_i] \end{aligned}$$

and

$$v_i - 1 = (v_{i-1} - 1) + u_{i-1}W_i + v_{i-1}G_i + b_i;$$

hence, by iteration and the norm properties,

$$\begin{aligned} |v(x) - 1| &= |v_n - 1| \leq |v_0 - 1| + \sum_1^n |u_{i-1}W_i + v_{i-1}G_i| + \sum_1^n |b_i| \\ &< k + \sum_1^n |b_i|. \end{aligned}$$

Let  $r = (k + 1)/2$ . Since  $\begin{bmatrix} H & W \\ K & G \end{bmatrix} \in OM^0$  and  $u$  and  $v$  are bounded on  $[a, b]$ , then there is a subdivision  $\{x_i\}_0^n$  of  $[a, x]$  such that  $\sum_1^n |b_i| < r - k$  and, hence,  $|v(x) - 1| < r < 1$ . Let  $v$  denote  $v(x)$ ; then  $v = 1 + (v - 1)$ ,  $v^{-1}$  exists, and

$$v^{-1} = 1 - (v - 1) + (v - 1)^2 - (v - 1)^3 + \dots$$

and

$$|v^{-1}| \leq (1 - |v - 1|)^{-1} \leq (1 - r)^{-1}.$$

Therefore,  $v^{-1}$  exists and is bounded by  $[1 - (k + 1)/2]^{-1}$  on  $[a, p]$ .

Since  $u$  and  $v$  are bounded and  $G$  and  $W \in OB^0$  on  $[a, b]$ , then there exist numbers  $p$  and  $k$  satisfying Conclusion 1, provided  $|v(a) - 1| + |u(a)W(a, a^+) + v(a)G(a, a^+)| < 1$ ; hence, Conclusion 2 follows as a corollary to Conclusion 1.

**LEMMA 3.6.** *If  $G$  is a function from  $R \times R$  to  $N$  such that  $G \in OA^0$  and  $OB^0$ , then  $|G| \in OA^0$ .*

A proof for this lemma is given in [6].

**LEMMA 3.7.** *If  $G$  is a function from  $R \times R$  to  $N$ , and  $G \in OA^0$  and  $OB^0$ , then  $\left| \int_a^b G \right| \leq \int_a^b |G|$ .*

Outline of proof.

$$\left| \int_a^b G \right| \leq \sum_1^n \left| \int_{x_{i-1}}^{x_i} G - G_i \right| + \sum_1^n |G_i|.$$

**LEMMA 3.8.** *Given.  $H$  and  $G$  are functions from  $R \times R$  to  $R$  and  $c$  is a number such that  $H \geq 0$ ,  $G \geq 0$ ,  $1 - G \geq c > 0$ , and  $H$  and  $G \in OA^0$  and  $OB^0$  on  $[a, b]$ ;  $f$  is a bounded function from  $R$  to  $R$  and  $k$  is a number such that  $f(x) \leq k + (LR) \int_a^x (fH + fG)$  for  $x \in [a, b]$ .*

*Conclusion.* If  $x \in [a, b]$ , then  $f(x) \leq k_a \Pi^x (1 + H)(1 - G)^{-1}$ . This is Theorem 4 of [4].

**LEMMA 3.9.** *If  $G \in OA^0$  and  $OB^0$  and  $f$  is quasicontinuous on  $[a, b]$ , then  $fG$  and  $Gf \in OA^0$  on  $[a, b]$ .*

This is a special case of [4, Theorem 2].

**THEOREM 3.10.** *Given. (1)  $[a, b]$  is a number interval;*

*(2)  $w$  is a function from  $R$  to  $N$  and  $H, G$  and  $K$  are functions from  $R \times R$  to  $N$  such that each of  $dw, H, G$  and  $K$  belongs to  $OA^0$  and  $OB^0$  on  $[a, b]$ ;*

*(3)  $f$  and  $g$  are bounded functions from  $R$  to  $N$  and  $c$  is a number such that  $1 - |B| \geq c > 0$ , where  $B(x, y) = G(x, y) + g(x)K(x, y)$  and on  $[a, b]$  each of  $f$  and  $g$  is a solution of the integral equation*

$$f(x) = w(x) + (LRLR) \int_a^x (fH + Gf + fKf).$$

*Conclusion.* If  $x \in [a, b]$ , then  $f(x) = g(x)$ .

*Proof.* Since  $f$  and  $g$  are bounded and since  $dw, H, G$  and  $K \in OA^0$  and  $OB^0$ , then each of  $f, g$  and  $|f - g|$  is a quasicontinuous function. Let  $A$  be the function  $A(x, y) = H(x, y) + K(x, y)f(y)$  for  $a \leq x < y \leq b$ ; then it follows from Lemmas 3.6 and 3.9 that  $A, B, |A|$  and  $|B| \in OA^0$  and  $OB^0$  and that  $(LR) \int_a^b [|f - g| |A| + |B| |f - g|]$  exists. If  $x \in [a, b]$ , then

$$\begin{aligned} |f(x) - g(x)| &= \left| (LR) \int_a^x [(f - g)A + B(f - g)] \right| \\ &\leq 0 + (LR) \int_a^x [|f - g| |A| + |B| |f - g|] \quad (\text{Lemma 3.7}). \end{aligned}$$

It follows from Lemma 3.8 that

$$|f(x) - g(x)| \leq 0 \cdot {}_a\prod^x (1 + |A|)(1 - |B|)^{-1} = 0.$$

Therefore, if  $x \in [a, b]$ , then  $f(x) = g(x)$ .

The restrictions  $1 - |B| \geq c > 0$  and  $(1 - G_i - f_{i-1}K_i)^{-1}$  cannot be deleted from the hypothesis of Theorem 3.10 and Theorem 3.3, respectively. Consider the following example. Let  $u, v$ , and  $g$  be functions from  $R$  to  $R$  such that  $u(x) = 0$  for  $x \in [0, 2]$ ,  $v(x) = g(x) = 0$  for  $x \in [0, 1]$  and  $v(x) = g(x) = 1$  for  $x \in (1, 2]$ . Each of  $u$  and  $v$  is a solution on  $[0, 2]$  for the equation  $f(x) = (R) \int_0^x fdg$ . See [5] for solutions of equations in which the restriction  $1 - |B| \geq c > 0$  does not hold.

Theorems similar to Theorems 3.2, 3.3 and 3.10 can be proved for  $f(x) = u(x)v(x)^{-1}$ ,

$$f(x) = w(x) + (RLRL) \int_a^x (fG + Hf + fKf),$$

and

$$\begin{bmatrix} u(x) \\ v(x) \end{bmatrix} = {}_a\prod^x Q \begin{bmatrix} w(a) \\ 1 \end{bmatrix},$$

where  $Q = \begin{bmatrix} 1 + H & dw \\ -K & 1 - G \end{bmatrix}$  and

$${}_a\prod^x Q = \lim Q(x_{n-1}, x_n) \cdots Q(x_1, x_2)Q(x_0, x_1).$$

We will now compare the Riccati equation for Riemann-Stieltjes integrals with the Riccati equation for the  $(LRLR)$ -integral. In this and the next paragraph,  $G$  is continuous at  $p$  means  $G(p^-, p) = 0 = G(p, p^+)$ ; also, the symbol  $(RS) \int_a^b E(f)$  is used to denote a Riemann-Stieltjes-type integral: i.e., for each subdivision  $\{x_i\}_0^n$  of  $[a, b]$ , the approximating sum has the form  $\sum_1^n E[f(c_i)]$ , where  $x_{i-1} \leq c_i \leq x_i$  for  $i = 1, 2, \dots, n$ . Suppose that  $w, H, G$  and  $K$  satisfy the hypothesis of Theorem 3.2. If  $f$  is the solution of the Riccati equation

$$f(x) = w(x) + (RS) \int_a^x fH + (RS) \int_a^x Gf + (RS) \int_a^x fKf$$

on  $[a, b]$ , then  $f$  is the solution of

$$(1) \quad f(x) = w(x) + (LRLR) \int_a^x (fH + Gf + fKf)$$

on  $[a, b]$ . If  $f$  is a solution of

$$(2) \quad f(x) = w(x) + (RS) \int_a^x (fH + Gf + fKf)$$

on  $[a, b]$  and either  $f$  is continuous on  $[a, b]$  or each of  $H, G$  and  $K$  is continuous on  $[a, b]$ , then  $f$  is the solution of Equation 1 on  $[a, b]$ . Equation 2 can have a solution  $f$  on  $[a, b]$  even though each of  $f, w, H, G$  and  $K$  has a discontinuity.

EXAMPLE. Suppose that  $N$  is a field,  $a < p \leq b$ , and  $g$  is a function of bounded variation which is continuous on  $[a, p)$  and on  $[p, b]$ ;  $f$  is the function such that

$$f(x) = 1 + (LRLR) \int_a^x (fdg + dgf + fdgf)$$

for  $x \in [a, p)$  and

$$f(x) = -2 - f(p^-) + (LRLR) \int_p^x (fdg + dgf + fdgf)$$

for  $x \in [p, b]$ ; also,

$$g(p) - g(p^-) = -2[1 + f(p^-)]/f(p^-)[f(p^-) + 2].$$

The function  $f$  is the solution on  $[a, b]$  of Equation (2) with  $dg = H = G = K$ ; however,  $f$  is not the solution of Equation (1) unless  $f(p^-) = -1$ . Furthermore, if  $g(p)$  is defined differently, then Equation (2) has no solution on  $[a, p]$ .

In order for the Riemann-Stieltjes equation to have a solution which is not a solution of the  $(LRLR)$ -equation, there must be an interdependence between the functions  $w, H, G$  and  $K$ . The following discussion illustrates this. Suppose that  $N$  is a field and that  $w, H, G$  and  $K$  are functions that satisfy the hypothesis of Theorem 3.2 and that on  $[a, b]$  the function  $f$  is a solution of Equation (2) but is not a solution of Equation (1); then there is a number  $p \in [a, b]$  such that  $f$  is not continuous at  $p$ . For convenience suppose that  $f(p^-) \neq f(p)$  and, in the following manipulations, let  $f_1, f_2, \Delta w, H, G$  and  $K$  denote  $f(p^-)$ ,  $f(p)$ ,  $w(p) - w(p^-)$ ,  $H(p^-, p)$ ,  $G(p^-, p)$  and  $K(p^-, p)$ , respectively. Then

$$f(p) = f(p^-) + \Delta w + (RS) \int_{p^-}^p (fH + Gf + fKf),$$

$$f_2 = f_1 + \Delta w + f_1H + Gf_1 + f_1Kf_1,$$

$$= f_1 + \Delta w + f_2H + Gf_2 + f_2Kf_2,$$

$$f_2H + Gf_2 + f_2Kf_2 = f_1H + Gf_1 + f_1Kf_1$$

and

$$(f_2 - f_1)(H + Kf_2) + (G + f_1K)(f_2 - f_1) = 0.$$

Since  $f_2 - f_1 \neq 0$  and  $N$  is a field, then

$$H + G + Kf_2 + f_1K = 0.$$

Substituting for  $f_2$  and simplifying, we obtain

$$(3) \quad K^2f_1^2 + (2 + H + G)Kf_1 + (H + G + \Delta wK) = 0.$$

Since  $f_1 = f(p^-) = w(p^-) + (RS) \int_a^{p^-} (fH + Gf + fKf)$ , then the value of  $f(p^-)$  depends only on the values of  $w, H, G$  and  $K$  on the half open interval  $[a, p)$ ; however, Equation (3) depends on the values of  $w, H, G$  and  $K$  on the closed interval  $[a, p]$ . Hence, these functions cannot be defined independently. For example, if  $K \neq 0$  and a different value is assigned to  $w(p)$ , then Equation (3) is no longer true and the Riemann-Stieltjes equation has no solution on  $[a, p]$  unless compensating values are assigned to  $H(p^-, p)$ ,  $G(p^-, p)$  and  $K(p^-, p)$ . However, the new  $(LRLR)$ -Riccati equation will have a solution on  $[a, p]$ .

**4. A differential-type equation.** In this section we find the solution of  $f^{**} + f^*p + fq = r$ , where  $f^*$  and  $f^{**}$  are defined as follows. If  $[a, b]$  is a number interval and  $h$  is a left continuous function from  $R$  to  $N$  such that  $dh \in OB^0$ , then  $D(h, a, b)$  denotes the set of ordered pairs of functions such that  $(f, g) \in D(h, a, b)$  iff  $g$  is a quasicontinuous function from  $R$  to  $N$  such that  $f(x) - f(a) = (L) \int_a^x g dh$  for  $x \in [a, b]$ . If  $(f, g) \in D(h, a, b)$ , then  $g$  is denoted by  $f^*$ . Also,

$f^{**} = (f^*)^*$  and  $f \cong w$  iff  $(L) \int_a^x f dh = (L) \int_a^x w dh$  for  $x \in [a, b]$ . In this section all integrals and product integrals are Cauchy-left-type integrals unless indicated otherwise.

**LEMMA 4.1.** If  $(f, f^*)$  and  $(g, g^*) \in D(h, a, b)$ , then  $(f + g, f^* + g^*) \in D(h, a, b)$ .

**LEMMA 4.2.** If  $(f, f^*)$  and  $(g, g^*) \in D(h, a, b)$ ,  $g^*, h$  and  $g$  commute and  $z$  is the function such that  $z(x) = g(x^+) - g(x)$  for  $x \in [a, b]$ , then  $(fg, f^*g + fg^* + f^*z) \in D(h, a, b)$ .

*Indication of proof.* Since  $(g, g^*)$  and  $(f, f^*) \in D(h, a, b)$ , then  $g$  is left continuous and  $df \in OB^0$ ; hence,

$$\begin{aligned} \int_a^x df dg &= (L) \int_a^x (df)z, \\ (L) \int_a^x (df)g &= (R) \int_a^x [(df)g - (df)(dg)] \end{aligned}$$

and

$$\begin{aligned} (L) \int_a^x (f^*g + fg^* + f^*z)dh &= (LLL) \int_a^x [(df)g + fdg + (df)z] \\ &= (RLL) \int_a^x [(df)g + fdg - (df)dg + (df)z] \\ &= (RL) \int_a^x [(df)g + fdg] \\ &= f(x)g(x) - f(a)g(a) \end{aligned}$$

**LEMMA 4.3.** Given.  $[a, b]$  is a number interval;  $f$  and  $h$  are functions from  $R$  to  $N$  such that  $f(a) = h(a)$  and  $dh \in OB^0$ ;  $G$  is a function from  $R \times R$  to  $N$  such that  $G \in OB^0$  and  $OA^0$

*Conclusion.* The following statements are equivalent:

- (1) if  $x \in [a, b]$ , then  $f(x) = h(x) + (L) \int_a^x fG$ ; and
- (2) if  $x \in [a, b]$ , then

$$f(x) = f(a) {}_a\prod^x (1 + G) + (R) \int_a^x dh {}_t\prod^x (1 + G).$$

This lemma is a special case of Theorem 5.1 of [3].

**THEOREM 4.4.** *Given. (1)  $[a, b]$  is a number interval; (2)  $h, p, q, u, v, \beta$  and  $s$  are functions from  $R$  to  $N$  such that  $h$  is left continuous,  $dh \in OB^0$ ,  $p$  and  $q$  are quasicontinuous on  $[a, b]$  and, if  $x \in [a, b]$ , then  $u(x)$  and  $v(x)$  are defined by the matrix equation*

$$[u(x), v(x)] = [0, 1](\dot{L}) {}_a\prod^x \left( I + \begin{bmatrix} -p & -1 \\ q & 0 \end{bmatrix} dh \right),$$

*$v(x)^{-1}$  exists,  $\beta(x) = v(x)^{-1}u(x)$  and  $s(x) = \beta(x^+) - \beta(x)$ ; also,  $v^{-1}$  is bounded on  $[a, b]$ ; (3) if  $a \leq x \leq y \leq b$ , then  $p(x)$ ,  $p(y)$ ,  $q(x)$ ,  $q(y)$ ,  $h(x)$  and  $h(y)$  commute; (4)  $f$  and  $r$  are functions from  $R$  to  $N$  and  $r$  is quasicontinuous.*

*Conclusion.* The following statements are equivalent.

- (1) There exist functions  $f^*$  and  $f^{**}$  such that  $(f, f^*)$  and  $(f^*, f^{**}) \in D(h, a, b)$  and such that on  $[a, b]$

$$f^{**} + f^*p + fq = r.$$

- (2) If  $x \in [a, b]$ , then

$$f(x) = f(a)(L) {}_a\prod^x (1 - \beta dh) + (R) \int_a^x dz(L) {}_t\prod^x (1 - \beta dh),$$

where  $\alpha = p - \beta - s$ ,  $z(x) = f(a) + (L) \int_a^x wdh$ ,  $g(x) = f^*(a) + (L) \int_a^x rdh$  and

$$w(x) = f^*(a)(L) {}_a\prod^x (1 - \alpha dh) + (R) \int_a^x dg(L) {}_t\prod^x (1 - \alpha dh).$$

*Proof.* Since  $dh \in OB^0$  and  $h$  is left continuous and since  $p$  and  $q$  are quasicontinuous, then  $u$  and  $v$  are left continuous and

quasicontinuous. Since  $v^{-1}$  is bounded and  $\beta = v^{-1}u$ , then  $\beta$  is left continuous, quasicontinuous and commutes with  $h$ . If  $x \in [a, b]$ , it follows from Theorem 3.2 that

$$\beta(x) = (L) \int_a^x q dh + (LL) \int_a^x \beta(-p dh) + (LR) \int_a^x \beta dh \beta.$$

Let  $\alpha, s$  and  $k$  be the functions such that  $s(t) = \beta(t^+) - \beta(t)$ ,  $\alpha = p - \beta - s$ ,  $k(a) = 0$ , and  $k = q + \beta^2 - \beta p + \beta s$ ; then, for  $x \in [a, b]$ ,

$$\begin{aligned} (L) \int_a^x k dh &= (L) \int_a^x (q + \beta^2 - \beta p + \beta s) dh \\ &= (L) \int_a^x q dh + \left[ (LR) \int_a^x \beta dh \beta - (L) \int_a^x \beta dh d\beta \right] \\ &\quad + (LL) \int_a^x \beta(-p dh) + (LL) \int_a^x \beta s dh. \end{aligned}$$

Since  $\beta$  is left continuous, then

$$(L) \int_a^x \beta dh d\beta = (LL) \int_a^x \beta s dh,$$

$\int_a^x k dh = \beta(x) - \beta(a)$  and  $(\beta, k) \in D(h, a, b)$ ;  $k$  will be denoted by  $\beta^*$ . Note that  $\beta, \alpha, \beta^*, p, q$  and  $h$  commute on  $[a, b]$  and that  $q = \beta^* + \beta\alpha$ .

*Proof of  $1 \rightarrow 2$ .* Since the triple  $(f, f^*), (\beta, \beta^*), s$  satisfies the hypothesis of Lemma 4.2, then  $(f\beta, f^*\beta + f\beta^* + f^*s) \in D(h, a, b)$ . Hence,

$$\begin{aligned} (f^* + f\beta)^* + (f^* + f\beta)\alpha &\equiv f^{**} + f^*\beta + f\beta^* + f^*s + f^*\alpha + f\beta\alpha \\ &= f^{**} + f^*(\beta + s + \alpha) + f(\beta^* + \beta\alpha) \\ &= f^{**} + f^*p + fq = r \end{aligned}$$

and

$$(f^* + f\beta)^* \equiv r - (f^* + f\beta)\alpha.$$

If we integrate each member of the preceding equation with respect to  $h$  and recall that  $\beta(a) = 0$ , we obtain



$$(f^* + f\beta)(x) = g(x) + (L) \int_a^x (f^* + f\beta)(-\alpha dh),$$

where  $g(x) = f^*(a) + (L) \int_a^x r dh$ . It follows from Lemma 4.3,  $1 \rightarrow 2$ , that

$$(f^* + f\beta)(x) = f^*(a) {}_a\prod^x (1 - \alpha dh) + (R) \int_a^x dg, {}_a\prod^x (1 - \alpha dh)$$

for  $x \in [a, b]$ . Let  $w(x)$  represent the right member in the preceding equation. If  $x \in [a, b]$ , then  $f^*(x) = w(x) - f(x)\beta(x)$  and by integrating both members we obtain

$$f(x) = z(x) + (L) \int_a^x f(-\beta dh),$$

where  $z(x) = f(a) + (L) \int_a^x w dh$  and  $z(a) = f(a)$ . It follows from Lemma 4.3,  $1 \rightarrow 2$ , that

$$f(x) = f(a) {}_a\prod^x (1 - \beta dh) + (R) \int_a^x dz, {}_a\prod^x (1 - \beta dh).$$

*Proof of  $2 \rightarrow 1$ .* Functions  $f^{**}$  and  $f^*$  will be defined such that  $(f, f^*)$  and  $(f^*, f^{**}) \in D(h, a, b)$  and such that on  $[a, b]$   $f^{**} + f^*p + fq = r$ .

Let  $f^* = w - f\beta$ . Since  $f$  satisfies the second statement of the conclusion, it follows from Lemma 4.3,  $2 \rightarrow 1$ , that for  $x \in [a, b]$

$$\begin{aligned} f(x) &= z(x) + (L) \int_a^x f(-\beta dh) \\ &= f(a) + (L) \int_a^x w dh + (L) \int_a^x f(-\beta dh) \\ &= f(a) + (L) \int_a^x f^* dh \end{aligned}$$

and  $(f, f^*) \in D(h, a, b)$ .

Let  $f^{**}$  be the function such that

$$f^{**} = r - (f^* + f\beta)\alpha - (f^*\beta + f\beta^* + f^*s).$$

Since  $\beta(a) = 0$  and

$$\begin{aligned}(f^* + f\beta)(x) &= w(x) \\ &= f^*(a) {}_a\prod^x (1 - \alpha dh) + (R) \int_a^x dg {}_t\prod^x (1 - \alpha dh)\end{aligned}$$

for  $x \in [a, b]$ , it follows from Lemma 4.3,  $2 \rightarrow 1$ , that

$$(f^* + f\beta)(x) = g(x) + (L) \int_a^x (f^* + f\beta)(-\alpha dh)$$

and, hence,

$$f^*(x) = g(x) + (L) \int_a^x (f^* + f\beta)(-\alpha dh) - f(x)\beta(x).$$

Since  $(f\beta, f^*\beta + f\beta^* + f^*s) \in D(h, a, b)$  and  $\beta(a) = 0$ , it follows from the definition of  $f^{**}$  that

$$\begin{aligned}(L) \int_a^x f^{**} dh &= (L) \int_a^x [r - (f^* + f\beta)\alpha - (f^*\beta + f\beta^* + f^*s)] dh \\ &= -f^*(a) + \left[ g(x) + (L) \int_a^x (f^* + f\beta)(-\alpha dh) \right. \\ &\quad \left. - f(x)g(x) \right] \\ &= f^*(x) - f^*(a)\end{aligned}$$

for  $x \in [a, b]$ ; hence,  $(f^*, f^{**}) \in D(h, a, b)$ .

Since

$$\begin{aligned}f^{**} + f^*p + fq &= [r - (f^* + f\beta)\alpha - (f^*\beta + f\beta^* + f^*s)] \\ &\quad + f^*(\alpha + \beta + s) + f(\beta^* + \alpha\beta) = r,\end{aligned}$$

then the triple  $f, f^*, f^{**}$  satisfies the given equation.

Suppose that on  $[a, b]$  the functions  $h, p$  and  $q$  are defined as in Theorem 4.4 except for the restrictions pertaining to  $v^{-1}$ . If  $h \in C^0$ , it follows from Theorem 3.5 that there is a subdivision  $\{x_i\}_0^n$  of  $[a, b]$  and functions  $\{\beta_i\}_1^n$ ,  $\{u_i\}_1^n$  and  $\{v_i\}_1^n$  such that for  $i = 1, 2, \dots, n$  and  $x \in [x_{i-1}, x_i]$

$$[u_i(x), v_i(x)] = [0, 1] {}_{x_{i-1}}\prod^x \left( I + \begin{bmatrix} -p & -1 \\ q & 0 \end{bmatrix} dh \right),$$

$\beta_i(x) = v_i(x)^{-1} u_i(x)$ , and  $v_i^{-1}$  exists and is bounded on  $[x_{i-1}, x_i]$ . Hence, for  $i = 1, 2, \dots, n$ , Theorem 4.4 gives the solution of  $f^{**} + f^*p + fq = r$  on  $[x_{i-1}, x_i]$  which is unique for a given pair  $f^*(x_{i-1})$  and  $f(x_{i-1})$ . Therefore, Theorem 4.4 can be used to find a unique solution on  $[a, b]$  for given values of  $f(a)$  and  $f^*(a)$ .

A theorem similar to Theorem 4.4 can be stated and proved for the equation  $f^{**} + pf^* + qf = r$ ; however, Theorem 5.2 of [3] would be used in the proof instead of Lemma 4.3. If  $(f, f^*)$  means  $f(x) - f(a) = (R) \int_a^x f^* dh$  and  $h$  is right continuous, a theorem similar to Theorem 4.4 can be stated and proved.

## REFERENCES

1. P. Puig Adam, *Los fracciones continuas de cocientes incompletos diferenciales y sus aplicaciones*, Revista Mat. Hisp. Amer. II, (1951), 180–190.
2. W. D. L. Appling, *Interval functions and real Hilbert spaces*, Rend.-Circl. Mat. Palermo, Ser. 2, **11** (1962), 154–156.
3. B. W. Helton, *Integral equations and product integrals*, Pacific J. Math. **16** (1966), 297–322.
4. ———, *A product integral representation for a Gronwall inequality*, Proc. Amer. Math. Soc., **23** (1969), 493–500.
5. ———, *Solutions of  $f(x) = f(a) + (RL) \int_a^x (fH + fG)$  for rings*, Proc. Amer. Math. Soc., **25** (1970), 735–742.
6. Jon C. Helton, *An existence theorem for sum and product integrals*, Proc. Amer. Math. Soc., **39** (1973), 149–154.
7. J. S. MacNerney, *Stieltjes integrals in linear spaces*, Ann. of Math., (2) **61** (1955), 354–367.
8. ———, *Continuous products in linear spaces*, J. Elisha Mitchell Sci. Soc., **71** (1955), 185–200.
9. ———, *Integral equations and semigroups*, Illinois J. Math., **7** (1963), 148–173.
10. Alfred Stohr, *Über einen integralartigen Grenzübergang bei Kettenbrüchen*, Math. Nachr., **6** (1951), 103–107.
11. H. S. Wall, *Concerning continuous continued fractions and certain systems of Stieltjes integral equations*, Rend. Circ. Mat. Palermo, (2), **2** (1953), 73–84.
12. ———, *Concerning harmonic matrices*, Archiv. der Math., **V** (1954), 160–167.

Received September 12, 1973 and in revised form June 11, 1974.

SOUTHWEST TEXAS STATE UNIVERSITY



Shimshon A. Amitsur, <i>Central embeddings in semi-simple rings</i> .....	1
David Marion Arnold and Charles Estep Murley, <i>Abelian groups, <math>A</math>, such that <math>\text{Hom}(A, - - -)</math> preserves direct sums of copies of <math>A</math></i> .....	7
Martin Bartelt, <i>An integral representation for strictly continuous linear operators</i> .....	21
Richard G. Burton, <i>Fractional elements in multiplicative lattices</i> .....	35
James Alan Cochran, <i>Growth estimates for the singular values of square-integrable kernels</i> .....	51
C. Martin Edwards and Peter John Stacey, <i>On group algebras of central group extensions</i> .....	59
Peter Fletcher and Pei Liu, <i>Topologies compatible with homeomorphism groups</i> .....	77
George Gasper, Jr., <i>Products of terminating <math>{}_3F_2(1)</math> series</i> .....	87
Leon Gerber, <i>The orthocentric simplex as an extreme simplex</i> .....	97
Burrell Washington Helton, <i>A product integral solution of a Riccati equation</i> .....	113
Melvyn W. Jeter, <i>On the extremal elements of the convex cone of superadditive <math>n</math>-homogeneous functions</i> .....	131
R. H. Johnson, <i>Simple separable graphs</i> .....	143
Margaret Humm Kleinfeld, <i>More on a generalization of commutative and alternative rings</i> .....	159
A. Y. W. Lau, <i>The boundary of a semilattice on an <math>n</math>-cell</i> .....	171
Robert F. Lax, <i>The local rigidity of the moduli scheme for curves</i> .....	175
Glenn Richard Luecke, <i>A note on quasidiagonal and quasitriangular operators</i> .....	179
Paul Milnes, <i>On the extension of continuous and almost periodic functions</i> .....	187
Hidegoro Nakano and Kazumi Nakano, <i>Connector theory</i> .....	195
James Michael Osterburg, <i>Completely outer Galois theory of perfect rings</i> .....	215
Lavon Barry Page, <i>Compact Hankel operators and the F. and M. Riesz theorem</i> .....	221
Joseph E. Quinn, <i>Intermediate Riesz spaces</i> .....	225
Shlomo Vinner, <i>Model-completeness in a first order language with a generalized quantifier</i> .....	265
Jorge Viola-Prioli, <i>On absolutely torsion-free rings</i> .....	275
Philip William Walker, <i>A note on differential equations with all solutions of integrable-square</i> .....	285
Stephen Jeffrey Willson, <i>Equivariant maps between representation spheres</i> .....	291