

Pacific Journal of Mathematics

CLASSIFICATION OF SINGULAR INTEGRALS OVER A LOCAL FIELD

CHARLES P. DOWNEY

CLASSIFICATION OF SINGULAR INTEGRALS OVER A LOCAL FIELD

CHARLES DOWNEY

The singular integral operators over a local field K whose kernels are multiplicative characters of the unit sphere of K are shown to be precisely those continuous operators on $\mathcal{L}_2(K)$ which commute with translation and dilation, anti-commute with an appropriately defined rotation, and whose multipliers satisfy a smoothness condition. The characterization is analogous to that of the Hilbert transform over the real numbers.

1. Classically, the Hilbert transform over \mathbf{R} is, up to a constant multiple, the only continuous operator on $\mathcal{L}_2(\mathbf{R})$ which commutes with translation and (positive) dilation and anti-commutes with reflection. See [9], page 55. The Hilbert transform is a singular integral operator with kernel the only (nontrivial) multiplicative character of the unit sphere of \mathbf{R} .

Singular integrals over a local field have been developed. (See, for example, Phillips [6], Phillips-Taibleson [7], and Chao [1].) Those with kernel a multiplicative character of the unit sphere satisfy a classification similar to that of the classical Hilbert transform.

The classification theorem is in § 4. The main results are Theorems 4.1 and 4.2. Section 3 contains the necessary results regarding the character group of the unit sphere of a local field; § 2 contains other preliminary results, notation, and definitions.

2. Let $Z, Z^+, \mathbf{Q}, \mathbf{R}$, and \mathbf{C} denote the integers, the positive integers, the rational number, the real numbers, and the complex numbers, respectively. F_{p^n} will denote the (unique) field with p^n elements. The symbols \mathbf{Q}_p and Z_p will denote the p -adic numbers and the p -adic integers, respectively. For any set S , ξ_S will denote the characteristic function of S . The complement of S will be written S_c .

The necessary analysis on local fields is stated without proof below. Most of it may be found in Chapters I and II of Weil [11].

A local field is a nondiscrete, locally compact, zero-dimensional topological (commutative) field. These have been completely classified. Those of characteristic $p \neq 0$ can be identified as the fields of formal power series over a finite field. Those of characteristic 0 are either the p -adic numbers of finite extensions of the p -adic numbers. See [11], page 11.

Let K be a local field with λ Haar measure for $(K, +)$. The modular function for K , $|\cdot|$, is given by $|x| = \lambda(xS)/\lambda(S)$ for $0 < \lambda(S) < \infty$. Haar measure for the multiplicative group $K^\times = K \sim \{0\}$ is $\lambda/|\cdot|$.

Let R be the ring of integers of the local field K and P be the unique maximal ideal of R . Then $\text{ord}(R/P) = q$, the module of K , a prime power. The ideal P has a generator π , so that $\pi R = P$. We have $|\pi| = q^{-1}$, and, in fact, any $x \in K$ with $|x| = q^{-1}$ will generate P . Those elements of modulus q^{-1} will be called primes in K .

For $n \in \mathbb{Z}$ we define

$$P^n = \{x \in K: |x| \leq q^{-n}\}; D^n = \{x \in K: |x| = q^{-n}\}.$$

Then $P^1 = P, P^0 = R$, and $R \sim P = D^0$. The set $\{P^n\}_{n=0}^\infty$ is a neighborhood base at 0 of open and closed subgroups of $(K, +)$. The set $\{1 + P^n\}_{n=1}^\infty$ is a neighborhood base at 1 of open and closed subgroups for the topological group (K^\times, \cdot) .

We define the operators τ_δ for $\delta \neq 0$ on functions by $\tau_\delta f(x) = f(\delta x)$. Regarding the prime π as fixed, we single out a set of such operators, the dilation operators, \mathcal{D}_j , defined by $\mathcal{D}_j f(x) = f(\pi^j x) j \in \mathbb{Z}$. A function f is homogeneous degree zero if $\mathcal{D}_j f = f$ for all $j \in \mathbb{Z}$. For $x \in K$, translation operators T_x are defined on functions by $T_x f(y) = f(x + y)$.

There is a character χ of the additive group of K which is identically one on R and nontrivial on P^{-1} . Then for any $y \in K$, $\chi_y(x) = \chi(xy)$ defines a character of K . In fact, the mapping $y \rightarrow \chi_y$ is a topological isomorphism of $(K, +)$ onto its dual. We thus identify K with its dual.

The Fourier transform for K is initially defined on $\mathcal{L}_1(K)$ by

$$\mathcal{F} f(x) = \hat{f}(x) = \int_K f(y) \overline{\chi(xy)} dy.$$

[The integral is taken with respect to λ . Here and elsewhere the λ will be suppressed.] The transform \mathcal{F}^{-1} is defined by $\mathcal{F}^{-1} f(x) = \check{f}(x) = \int_K f(y) \chi(xy) dy$. Both \mathcal{F} and \mathcal{F}^{-1} extend uniquely to \mathcal{L}_2 . It is easy to see that, as \mathcal{L}_2 operators, $\tau_\delta \mathcal{F} = |\delta|^{-1} \mathcal{F} \tau_{\delta^{-1}}$ and $\tau_\delta \mathcal{F}^{-1} = |\delta|^{-1} \mathcal{F}^{-1} \tau_{\delta^{-1}}$.

The following result will be used extensively in the sequel: Let L be a continuous linear operator from $\mathcal{L}_2(K)$ to $\mathcal{L}_2(K)$. Then a necessary and sufficient condition that L commute with translation is that there exist a function m , in $\mathcal{L}_\infty(K)$, such that $\mathcal{F}(Lf) = m \mathcal{F} f$ for all $f \in \mathcal{L}_2(K)$. See [5], pp. 92-94.

The space \mathcal{L} of test functions on K and its topological dual \mathcal{L}' , the space of distributions, are defined as in [8]. Both are

complete linear spaces. The action of a $\mu \in \mathcal{F}'$ on an $f \in \mathcal{F}$ will be denoted (μ, f) .

The space \mathcal{F} is contained densely in $\mathcal{L}_p, 1 \leq p < \infty$. The Fourier transform is thus well-defined on \mathcal{F} . The Fourier transform on \mathcal{F}' is given by $(\hat{\mu}, f) = (\mu, \hat{f})$. Thus defined, the Fourier transform is a linear topological isomorphism on both \mathcal{F} and \mathcal{F}' .

Functions and measures will be identified with the distributions they induce. Convolution of a distribution and a test function is defined by $\mu * f(x) = (\mu, T_x \tilde{f})$, where $\tilde{f}(x) = f(-x)$.

Let $\mu \in \mathcal{F}'$, and let σ be a (not necessarily unitary) multiplicative character of $K^\times (= K \sim \{0\})$. Then, as in [8], we say μ is homogeneous of degree σ if for all $t \in K^\times, \mu_t = \sigma(t)\mu$, where μ_t is that distribution defined by $(\mu_t, \phi) = (\mu, |t|^{-1}\tau_{t^{-1}}\phi)$.

We take M to be $M^\times \cup \{0\}$, where M^\times is the group of roots of unity in K of order prime to p . Then M^\times is the unique cyclic group of order $q-1$ ([11], p. 16). Let g be a generator of M^\times . Then each $0 \neq x \in K$ may be written uniquely as $x = \pi^j g^k (1 + p_x)$, where $k, j \in \mathbb{Z}, 0 \leq k \leq q-2, p_x \in P$. A multiplicative character of K^\times is given by its values at π, g , and on $1 + P$.

Let ω be a multiplicative character of K^\times . There is some $n \in \mathbb{Z}$ such that ω is trivial on $1 + P^n$. If ω is trivial on $1 + P^n$ but not on $1 + P^{n-1}, n \geq 1$, we say ω is ramified of degree n . If ω is trivial on D^0 , we say ω is unramified. Given a character ω of $1 + P$, ω is the restriction of a character of K^\times , say ω' . The ramification degree of ω' depends only on ω , and we define the ramification degree of ω to be that of ω' .

We define the local field gamma function on ramified characters of K^\times by

$$\Gamma(\omega) = \text{p.v.} \int_K \frac{\chi(x)\omega(x)dx}{|x|},$$

where

$$\text{p.v.} \int_K f(x)dx = \lim_{n \rightarrow \infty} \int_{P^{-1} \cap (P^{+n})^c} f(x)dx.$$

See [8] for details and further definition of Γ .

3. LEMMA 3.1. *Let K be a local field of characteristic $p \neq 0$ with module $q = p^f$. Let $\{\alpha_i, \dots, \alpha_f\}$ be a basis for F_q over F_p . Then given $x \in P$ and $N \in \mathbb{Z}^+$,*

(a) *there are unique integers $a_{i,j}, n_j, \nu_j$, with $0 \leq a_{i,j} < p, (n_j, p) = 1$ for $1 \leq i \leq f, 1 \leq j \leq N$, such that $1 + x = \prod_{j=1}^N \prod_{i=1}^f (1 + \alpha_i \pi^{n_j})^{a_{i,j} p^{\nu_j}} (p^{N+1})$, and*

(b) $1 + x \in (1 + P^N) \sim (1 + P^{N+1})$ if and only if $\alpha_{i,j} = 0$ for $1 \leq i \leq f, 1 \leq j \leq N$ and at least one of the $\alpha_{i,N} \neq 0, 1 \leq i \leq f$.

Proof. The proof is similar to that of Proposition 10, page 34 of [11], and is omitted.

Given $N \in \mathbb{Z}^+$ we establish the following notation to be used in the following lemma and theorem. For each $j, 1 \leq j \leq N$, write $j = n_j P^{\nu_j}$, where $(n_j, p) = 1$; define m_j as the smallest integer such that $m_j \geq \log_p((N+1)/n_j)$ then define β_j as a primitive p^{m_j} th root of 1 in C .

LEMMA 3.2. *With the above notation, $m_N = \nu_N + 1$.*

Proof. The proof is a direct computation and is omitted.

THEOREM 3.1. *Let K be a local field of characteristic $p \neq 0$ and ω a character of $1 + P \subset K$ ramified degree $N + 1$. Then for $x \in 1 + P$, ω is given by*

$$\omega(x) = \prod_{j=1}^N \prod_{i=1}^f \beta_j^{k_{i,j} a_{i,j} p^{\nu_j}},$$

where

$$(*) \quad x \equiv \prod_{j=1}^N \prod_{i=1}^f (1 + \alpha_i \pi^{n_j})^{a_{i,j} p^{\nu_j}} (P^{N+1})$$

for some unique $k_{i,j}, 0 \leq k_{i,j} < p^{m_j}$ with at least one of $k_{i,N}, 1 \leq i \leq f$, relatively prime to p .

Proof. Since ω is constant on cosets of P^{N+1} it suffices to consider $x \bmod P^{N+1}$. For any $x \in 1 + P$, the numbers $a_{i,j}, n_j, \nu_j$ are determined as in Lemma 3.1 so that (*) holds. Clearly ω will be completely determined by its values on $\{1 + \alpha_i \pi^{n_j}\}, 1 \leq i \leq f, 1 \leq j \leq N$, and the range of ω is contained in the p^{th} power roots of unity.

The definition of m_j as the smallest integer greater than or equal to $\log_p(N+1)/n_j$ makes m_j the smallest integer such that

$$(1 + \alpha_i \pi^{n_j})^{p^{m_j}} \in 1 + P^{N+1}.$$

Thus $(\omega(1 + \alpha_i \pi^{n_j}))^{p^{m_j}} = 1$, and $\omega(1 + \alpha_i \pi^{n_j}) = \beta_j^{k_{i,j}}$ for some unique $k_{i,j}, 0 \leq k_{i,j} < p^{m_j}$. Thus ω has the form required. The remainder of the theorem follows easily from the fact that β_N is a p^{m_N} th root of unity and ω must be nontrivial on P^N .

From Proposition 9 of Chapter II, § 3 of [11], we have:

PROPOSITION. *Let K be a d -dimensional extension of \mathbf{Q}_p . Then there is an integer $m \geq 0$ such that $1 + P$, as a multiplicative group is isomorphic to the additive group $Z_p^d \times F_{p^m}$, where m is the largest integer such that K contains a primitive $p^{m\text{th}}$ root of unity. For proof see [11].*

Let $\{u_i\}_{i=1}^d$ be those elements of $1 + P$ which map to the vectors with 1 in the i^{th} coordinate and zeros elsewhere by the isomorphism in the proposition. Let u_{d+1} be a primitive p^{th} power root of unity in K of maximal order, say p^m . Then any $x \in 1 + P$ is given uniquely by $x = \prod_{i=1}^{d+1} u_i^{a_i}$, where $a_i \in Z_p$, $1 \leq i \leq d$ and $a_{d+1} \in Z$, $0 \leq a_{d+1} < p^m$.

LEMMA 3.3. *Let K be a d -dimensional extension of \mathbf{Q}_p . Then given nonnegative integers k_i , $1 \leq i \leq d$, each $x \in 1 + P \subset K$ has a representation as*

$$x = u_{d+1}^{a_{d+1}} \prod_{i=1}^d u_i^{n_i} u_i^{b_i}, \quad \text{where}$$

$b_i \in Z_p$ with $|b_i|_{Z_p} < p^{-k_i}$ and n_i is a nonnegative integer. If n_i is picked to be as small as possible, this representation is unique.

Proof. The proof is direct from the above proposition and the density of Z^+ in Z_p .

Given $N \in Z^+$, define, for $1 \leq i \leq d + 1$, \mathcal{L}_i to be the smallest integer such that $u_i^{p^{\mathcal{L}_i}} \in 1 + P^{N+1}$ and β_i to be a fixed primitive $p^{\mathcal{L}_i\text{th}}$ root of 1 in C . With this notation we have the following:

THEOREM 3.2. *Let K be a local field of characteristic 0 and ω a character of $1 + P \subset K$ ramified of degree $N + 1$. Then for $x \in 1 + P$, ω is given for some unique k_i , $0 \leq k_i < p^{\mathcal{L}_i}$, $1 \leq i \leq d + 1$, by*

$$\omega(x) = \prod_{i=1}^d \beta_i^{k_i n_i} \beta_{d+1}^{k_{d+1} a_{d+1}} \quad \text{for } x = \prod_{i=1}^d u_i^{n_i} u_i^{b_i} u_{d+1}^{a_{d+1}},$$

where for $1 \leq i \leq d$, $b_i \in Z_p$ with $|b_i|_{Z_p} < p^{-\mathcal{L}_i}$ and $n_i \in Z^+$.

Proof. The density of Z in Z_p shows that an (additive) character of Z_p is determined by its value at 1. Thus a (multiplicative) character $1 + P$ will be determined by its values at the u_i , $1 \leq i \leq d + 1$. Here $\omega(u_i)^{p^{\mathcal{L}_i}} = 1$ since $u_i^{p^{\mathcal{L}_i}} \in 1 + P^{N+1}$ and ω is ramified of degree $N + 1$. Thus $\omega(u_i) = \beta_i^{k_i}$ for some (unique) k_i , $0 \leq k_i < p^{\mathcal{L}_i}$.

This characterization of the character group of K depends on

the p^{th} roots of unity in K . Since K is a finite dimensional extension of \mathbb{Q}_p , we look for a relationship between the degree d of K over \mathbb{Q}_p and the existence of p^{th} roots of unity in K .

THEOREM 3.3. *Let K be a local field of characteristic 0. If K is the p -adic field \mathbb{Q}_p for some prime $p \neq 2$, then K has no nontrivial p^{th} roots of unity. If K is an extension of \mathbb{Q}_p , $p \neq 2$, let the degree of ramification (see [11]) of K over \mathbb{Q}_p be e ; then,*

- (a) K has no p^{th} roots of 1 if $(p - 1)$ does not divide e ,
- (b) K may or may not have p^{th} roots of 1 if $p - 1$ divides e .

Proof. For the proof of (a) see [2]. Part (b) follows from [2] and the fact that the extension of \mathbb{Q}_p by a root of $x^{p-1} - p$ is fully ramified of degree $p - 1$ and has no p^{th} roots of unity.

LEMMA 4.1. *Let ω be a homogeneous degree zero multiplicative character of K^* , ramified of degree $k > 0$. Then ω is a kernel for a singular integral operator. The multiplier m for the singular integral operator T with kernel ω satisfies*

$$m(x) = \omega(-1)\Gamma(\omega)\omega^{-1}(x).$$

Proof. The operator T is defined for $f \in \mathcal{L}_p$, $1 \leq p < \infty$ by

$$Tf(x) = \lim_{k \rightarrow \infty} \int_{(p^k)_c} \frac{\omega(y)}{|y|} (f(x - y)dy).$$

Theorem 3.1 of [7] gives sufficient conditions on the kernel ω for the limit to exist (in \mathcal{L}_p). That ω satisfies those conditions is easily verified. Then from [7] we know T is bounded on \mathcal{L}_p , $1 < p < \infty$ and weak type $(1, 1)$.

The remainder of the lemma is done by Chao [1] for the case ω ramified of degree 1. The same proof establishes the result stated here.

Note. Chao [1] uses Theorem 4 of [8] to establish the conclusion of Lemma 4.1 for the case ω ramified of degree 1. However, he fails to compensate for the fact that he defines the Fourier transform as herein, i.e., $\mathcal{F}f(y) = \int f(x)\overline{\chi(xy)}dx$, while in [8] it is defined as $\int f(x)\chi(xy)dx$. Thus the result of [1] which corresponds to the conclusion of Lemma 4.4 above does not contain the necessary factor of $\omega(-1)$.

With notation as in Theorem 3.1, we define rotation operators $S_{i,j}$ for functions on a p -series field as follows:

$$S_{1,0}f(x) = f(gx),$$

where g is a fixed primitive $(q-1)^{\text{st}}$ root of unity in K ; and

$$S_{i,j}f(x) = f((1 + \alpha_i \pi^n x))$$

for $1 \leq i \leq f, j \geq 1$.

Given N we determine $\beta_j, 1 \leq j \leq n$ as in Theorem 3.1, and let β_0 be a $(q-1)^{\text{st}}$ root of unity in \mathcal{C} . Also as in that theorem, note that given N the choice of integers $k_{i,j}, 0 \leq k_{i,j} < p^{mj}, 1 \leq i \leq f, 1 \leq j \leq N$ determines a character of $1 + P$. If we also pick a $k_{1,0}, 0 \leq k_{1,0} < q-1$, and set $\omega(g) = \beta^{k_{1,0}}$, then the set $\{k_{i,j}\}$ determines character of D^0 . That character will be called the character determined by $\{k_{i,j}\}$. As it may be used as a kernel for a singular integral operator, that operator will be identified as the one determined by $\{k_{i,j}\}$.

We can now state

THEOREM 4.1. *Let K be a p -series field and L a continuous linear operator from $\mathcal{L}_2(K)$ to $\mathcal{L}_2(K)$ which satisfies*

- (a) \mathcal{L} commutes with translation and dilation,
- (b) there is some $N \geq 0$ such that the multiplier corresponding to L is constant on cosets of P^{N+1} ,
- (c) L anti-commutes with the rotations $S_{i,j}, 1 \leq i \leq f, 1 \leq j \leq N$, and $S_{1,0}$ in the sense that

$$LS_{i,j} = \beta_j^{-k_{i,j}} S_{i,j}L,$$

for some $k_{i,j}$. Then L is a constant multiple of the singular integral operator determined by $\{k_{i,j}\}$.

Before proving Theorem 4.1, we consider the p -adic case. Let $u_i, 1 \leq i \leq d+1$ be as in Theorem 3.2, and let $u_0 = g$, the fixed $(q-1)^{\text{st}}$ root of $1 \in K$. In this case we define rotation operators as: $S_i f(x) = f(u_i x), 0 \leq i \leq d+1$. Given N , we determine $\beta_i, 0 \leq i \leq d+1$ by: β_0 is a primitive $(q-1)^{\text{st}}$ root of $1 \in \mathcal{C}$; $\beta_i, 1 \leq i \leq d+1$, is a primitive p^{l_i} root of $1 \in \mathcal{C}$, where l_i is the smallest integer such that $u_i^{p^{l_i}} \in 1 + P^{N+1}$. Also, for each i we consider integers k_i such that $0 \leq k_0 < q-1, 0 \leq k_i < p^{l_i}, 1 \leq i \leq d+1$.

By Theorem 3.2 and the fact that $D^0 = M^{\times} x(1 + P)$, the set $\{k_i\}_{i=0}^{d+1}$ determines a unique character of D^0 by $\omega(u_i) = \beta_i^{k_i}$. The character ω will be called the character determined by the $\{k_i\}$. It is clearly constant on $1 + P^{N+1}$.

THEOREM 4.2. *Let K be a local field of characteristic 0, and let L be a continuous linear operator from $\mathcal{L}_2(K)$ to $\mathcal{L}_2(K)$ which*

satisfies

- (a) L commutes with translation and dilation,
- (b) there is some N such that the multiplier for L is constant on cosets of P^{N+1} ,
- (c) L anti-commutes with the rotations $S_i, 0 \leq i \leq d + 1$ in the sense that

$$LS_i = \beta_i^{-k_i} S_i L .$$

Then L is a constant multiple of the singular integral transform determined by the $\{k_i\}$. The proof of Theorems 4.1 and 4.2 will utilize the following Lemma.

LEMMA 4.2. *Let K be a local field. If characteristic $K = 0$, let L satisfy the hypothesis of Theorem 4.2. If characteristic $K = p \neq 0$, let L satisfy the hypothesis of Theorem 4.1. Then for $f \in \mathcal{L}$, L is given by convolution with a unique distribution μ , homogeneous of degree $\omega/|\cdot|$, where ω is the character of D^0 determined the $\{k_{i,j}\}$ or $\{k_i\}$ in the characteristic $p \neq 0$ and characteristic 0 case, respectively.*

Proof. Since L is a bounded linear operator from \mathcal{L}_2 to \mathcal{L}_2 which commutes with translation, by Theorem 9 of [10], it is given, on \mathcal{L} , by convolution with a unique distribution μ . We need only to show μ homogeneous of degree ν , where $\nu(x) = \omega(x)/|x|, x \neq 0$.

There is a function m in $\mathcal{L}_\infty(K)$ so that for $f \in \mathcal{L}_2(K), (Lf)^\wedge = m\hat{f}$. For $f \in \mathcal{L}, \hat{f} \in \mathcal{L}$, thus $m\hat{f} \in \mathcal{L}_1(K)$ since $m \in \mathcal{L}_\infty(K)$. Then $Lf = (m\hat{f})^\vee$ is continuous since it is the inverse Fourier transform of an \mathcal{L}_1 function.

Let $\gamma \in 1 + P^{N+1}$. Then $\gamma^{-1} \in 1 + P^{N+1}$, and, since m is constant on cosets of P^{N+1} , we have:

$$\begin{aligned} (L\tau_\gamma f)^\wedge(x) &= m(\tau_\gamma f)^\wedge(x) = m(x)\hat{f}(\gamma^{-1}x) \\ &= m(\gamma^{-1}x)\hat{f}(\gamma^{-1}x) = \tau_\gamma^{-1}m\hat{f}(x) . \end{aligned}$$

Thus

$$L\tau_\gamma f = \tau_\gamma Lf \quad \text{in } \mathcal{L}_2 .$$

Fix $t \in K^\times$. By (a) and (c) of Theorems 4.1 and 4.2 and the above equality, we have:

$$L\tau_t f = \omega^{-1}(t)\tau_t Lf \quad \text{in } \mathcal{L}_2$$

and

$$L\tau_t f(x) = \omega^{-1}(t)\tau_t Lf(x) \quad \text{a.e.}$$

But since both $L\tau_i f$ and $\tau_i Lf$ are continuous, we have the above equality everywhere.

For $f \in \mathcal{L}$,

$$\mu * \tau_i f(0) = \omega^{-1}(t)(\mu * f)(t \cdot 0),$$

and

$$(\mu, \tau_i \tilde{f}) = \omega^{-1}(t)(\mu, \tilde{f}).$$

Thus

$$\begin{aligned} (\mu_t, f) &= (\mu, |t|^{-1} \tau_{i-1} f) \\ &= |t|^{-1} (\mu, \tau_{i-1} f) \\ &= \frac{\omega(t)}{|t|} (\mu, f) = \nu(t)(\mu, f). \end{aligned}$$

Since this holds for all $f \in \mathcal{L}$, $\mu_t = \nu(t)\mu$.

Now we are ready for the

Proof of Theorems 4.1 and 4.2. By Lemma 4.2 for $f \in \mathcal{L}$, $Lf = \mu * f$, where μ is homogeneous of degree $\omega/|\cdot|$. But by Lemma 5 of [8], the only distributions which are homogeneous of degree σ , σ multiplicative character of K^\times such that $\sigma(x)$ is not identically $|x|^{-1}$, are constant multiples of σ . Thus $\mu = c\omega/|\cdot|$, and $Lf = (c\omega)/(|\cdot|) * f$, $f \in \mathcal{L}$. Thus, on the test functions, a dense subset of \mathcal{L}_2 , L agrees with L' , the singular integral operator defined by $L'f(x) = c \int (\omega(y)/(|y|) f(x-y) dy$. But since L and L' are continuous, $L = L'$ on \mathcal{L}_2 .

5. **Example.** The conclusions of Theorems 4.1 and 4.2 may be obtained by direct calculation. We indicate the method in the case $q = 3$ and ω ramified of degree 1. Here $M^\times = \{1, -1\}$ and ω will assume only the values ± 1 . [This is the "exact" analog of the Hilbert transform for the reals.]

Let H be the singular integral operator with ω as kernel. Both theorems then have the form: **Theorem:** Let K be local field with module $q = 3$ and L be a continuous operator on $\mathcal{L}_2(K)$ which satisfies:

- (a) L commutes with translation and dilation;
- (b) the multiplier, m , for L is constant on $1 + P$;
- (c) L anti-commutes with the rotation τ_{-1} by $L\tau_{-1} = -\tau_{-1}L$.

Then L is a constant multiple of H .

Proof. From the relation $(Lf)^\wedge = m\hat{f}$ it follows as in the real

case (see [9]) that $m(-x) = -m(x)$. Since any $x \in K^\times$ may be written $x = \pm\pi^j(1 + \rho_x)$, $\rho_x \in P$, $m(x) = \pm m(1) = \omega^{-1}(x)m(1)$. The theorem then follows from Lemma 4.1.

Lemma 4.1 may also be shown directly. For the case above we may even evaluate the multiplier m_H explicitly. Taking the fundamental character χ to be that given in [1] (a variation of that given in [6]), and the form of m_H from [7], we obtain $m_H(x) = (i)/(\sqrt{3})\omega(x)$. As in [1], a similar easy calculation gives $\Gamma(\omega) = -i/\sqrt{3}$, exemplifying Lemma 4.1. In further analogy with the real case, it is apparent from the multiplier that $H^2 = -(1/3)I$.

REFERENCES

1. J. A. Chao, *H^p -spaces of conjugate systems on local fields*, Stucka Math., **49** (1974), 267-287.
2. H. Hasse, *Zalentheorie*, Berlin Akademie-Verlag, 1963.
3. E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis II*, Springer-Verlag, 1970.
4. P. J. McCarthy, *Algebraic Extensions of Field*, Blaisdell, 1966.
5. R. Larsen, *An Introduction to the Theory of Multipliers*, Springer-Verlag, 1971.
6. K. Phillips, *Hilbert transforms for the p -adic and p -series fields*, Pacific J. Math., **23** (1967), 329-347.
7. K. Phillips and M. H. Taibleson, *Singular integrals in several variable over a local field*, Ibid., **30** (1969), 209-231.
8. P. J. Sally and M. H. Taibleson, *Special functions on locally compact fields*, Acta Math., **116** (1966) 279-309.
9. E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
10. M. H. Taibleson, *Harmonic analysis on n -dimensional vector spaces over local fields. I*, Math. Annalen, **176** (1968), 191-207.
11. A. Weil, *Basic Number Theory*, Springer-Verlag, 1967.

Received October 14, 1974 and in revised form April 12, 1975.

UNIVERSITY OF NEBRASKA AT OMAHA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

| | |
|---|-----|
| Aharon Atzmon, <i>A moment problem for positive measures on the unit disc</i> | 317 |
| Peter W. Bates and Grant Bernard Gustafson, <i>Green's function inequalities for two-point boundary value problems</i> | 327 |
| Howard Edwin Bell, <i>Infinite subrings of infinite rings and near-rings</i> | 345 |
| Grahame Bennett, Victor Wayne Goodman and Charles Michael Newman, <i>Norms of random matrices</i> | 359 |
| Beverly L. Brechner, <i>Almost periodic homeomorphisms of E^2 are periodic</i> | 367 |
| Beverly L. Brechner and R. Daniel Mauldin, <i>Homeomorphisms of the plane</i> | 375 |
| Jia-Arng Chao, <i>Lusin area functions on local fields</i> | 383 |
| Frank Rimi DeMeyer, <i>The Brauer group of polynomial rings</i> | 391 |
| M. V. Deshpande, <i>Collectively compact sets and the ergodic theory of semi-groups</i> | 399 |
| Raymond Frank Dickman and Jack Ray Porter, <i>θ-closed subsets of Hausdorff spaces</i> | 407 |
| Charles P. Downey, <i>Classification of singular integrals over a local field</i> | 417 |
| Daniel Reuven Farkas, <i>Miscellany on Bieberbach group algebras</i> | 427 |
| Peter A. Fowler, <i>Infimum and domination principles in vector lattices</i> | 437 |
| Barry J. Gardner, <i>Some aspects of T-nilpotence. II: Lifting properties over T-nilpotent ideals</i> | 445 |
| Gary Fred Gruenhagen and Phillip Lee Zenor, <i>Metritzation of spaces with countable large basis dimension</i> | 455 |
| J. L. Hickman, <i>Reducing series of ordinals</i> | 461 |
| Hugh M. Hilden, <i>Generators for two groups related to the braid group</i> | 475 |
| Tom (Roy Thomas Jr.) Jacob, <i>Some matrix transformations on analytic sequence spaces</i> | 487 |
| Elyahu Katz, <i>Free products in the category of k_w-groups</i> | 493 |
| Tsang Hai Kuo, <i>On conjugate Banach spaces with the Radon-Nikodým property</i> | 497 |
| Norman Eugene Liden, <i>K-spaces, their antispace and related mappings</i> | 505 |
| Clinton M. Petty, <i>Radon partitions in real linear spaces</i> | 515 |
| Alan Saleski, <i>A conditional entropy for the space of pseudo-Menger maps</i> | 525 |
| Michael Singer, <i>Elementary solutions of differential equations</i> | 535 |
| Eugene Spiegel and Allan Trojan, <i>On semi-simple group algebras. I</i> | 549 |
| Charles Madison Stanton, <i>Bounded analytic functions on a class of open Riemann surfaces</i> | 557 |
| Sherman K. Stein, <i>Transversals of Latin squares and their generalizations</i> | 567 |
| Ivan Ernest Stux, <i>Distribution of squarefree integers in non-linear sequences</i> | 577 |
| Lowell G. Sweet, <i>On homogeneous algebras</i> | 585 |
| Lowell G. Sweet, <i>On doubly homogeneous algebras</i> | 595 |
| Florian Vasilescu, <i>The closed range modulus of operators</i> | 599 |
| Arthur Anthony Yanushka, <i>A characterization of the symplectic groups $P\text{Sp}(2m, q)$ as rank 3 permutation groups</i> | 611 |
| James Juei-Chin Yeh, <i>Inversion of conditional Wiener integrals</i> | 623 |