

Pacific Journal of Mathematics

A CHARACTERIZATION OF THE SYMPLECTIC GROUPS $\mathrm{PSp}(2m, q)$ AS RANK 3 PERMUTATION GROUPS

ARTHUR ANTHONY YANUSHKA

A CHARACTERIZATION OF THE SYMPLECTIC GROUPS $PSp(2m, q)$ AS RANK 3 PERMUTATION GROUPS

ARTHUR YANUSHKA

In this paper the following characterization of the symplectic groups $PSp(2m, q)$ for $m > 2$ as rank 3 permutation groups is obtained:

THEOREM. Let G be a transitive rank 3 group of permutations of a finite set X such that the orbit lengths for G_b , the stabilizer of a point b in X are $1, q(q^{r-2} - 1)/(q - 1)$ and q^{r-1} for integers $q > 1$ and $r > 4$. Let b^\perp denote the union of b and the G_b orbit of length $q(q^{r-2} - 1)/(q - 1)$. Let $R(bc)$ denote $\cap \{z^\perp : b, c \in z^\perp\}$. Assume $R(bc) \neq \{b, c\}$, for all distinct pairs of points, b and c . Assume that the pointwise stabilizer of b^\perp is transitive on the points unequal to b of $R(bc)$ for $c \in b^\perp$. Then r is even, q is a prime power and $G \cong H$, a group of symplectic collineations of projective $r - 1$ space over the finite field of q elements and $PSp(r, q) \trianglelefteq H$.

The rank of a transitive permutation group is the number of orbits of the stabilizer of a point. The projective classical groups of symplectic type $PSp(2m, q)$ for $m \geq 2$ and for a prime power q are transitive groups of rank 3 when considered as groups of permutations of the absolute points of the corresponding projective space. Indeed, the pointwise stabilizer of $PSp(2m, q)$ has 3 orbits of lengths $1, q(q^{2m-2} - 1)/(q - 1)$ and q^{2m-1} .

Let G be any rank 3 group of permutations of a set X such that the pointwise stabilizer has orbit lengths of $1, q(q^{r-2} - 1)/(q - 1)$ and q^{r-1} for any integers $r \geq 4$ and $q \geq 2$. The characterization problem is to impose some restrictions on G and on X to force the conclusion that X is a projective space and that G is a group of symplectic collineations. Let b^\perp denote the union of b and the G_b -orbit of length $q(q^{r-2} - 1)/(q - 1)$. There are several rank 3 characterizations of the symplectic groups. Assume that q is a prime power and that $r \geq 6$. Kantor [5] proved that G can be regarded as an automorphism group of a symplectic geometry, acting on the set of singular points. Next assume that q is any integer, that $r = 4$ and that the pointwise stabilizer of b^\perp contains at least q elements. D. G. Higman [4] proved that G can be regarded as a group of symplectic collineations of projective 3-space over the finite field of q elements and that G contains $PSp(4, q)$. Later Tsuzuku [6] extended Higman's theorem to $r \geq 4$ under the additional assumption that q is a prime

power. This paper essentially generalizes Higman's theorem to all higher dimensions, without the assumption that q is a prime power.

A brief outline of the proof follows. The assumption that the pointwise stabilizer in G of b^\perp is transitive on the points unequal to b of the "hyperbolic line" $R(bc)$ for $c \notin b^\perp$ yields that G_a , the stabilizer of the points a and b of X , is transitive on the points of $a^\perp \cap b^\perp - R(ab)$. This fact implies that G_a is a rank 3 permutation group on $\{R(ab): b \in a^\perp - a\}$, and the set of "totally singular lines" carry $q + 1$ points. We then show that X together with its totally singular lines forms a nondegenerate Shult space [1] of rank ≥ 3 . Next we use a theorem of Buekenhout and Shult [1] to conclude that X is isomorphic to the set of points of a classical geometry of symplectic type. Therefore G is a group of symplectic collineations. Finally we show that the nontrivial elements of the pointwise stabilizer of b^\perp correspond to symplectic elations with center b and that G contains $PSp(r, q)$.

In §2 we collect the necessary facts about rank 3 groups from the basic papers of D. G. Higman [3], [4]. We refer the reader to a paper of Buekenhout and Shult [1] for the definition of Shult space and a brief introduction to polar spaces. In §3 we prove the characterization theorem. Finally the author wishes to thank Donald Higman for making him aware of the work of Buekenhout and Shult whose theorem makes the proof of the characterization of $PSp(2m, q)$ given here considerably shorter than the original version.

2. Rank 3 permutation groups. In this section we collect the necessary facts about rank 3 permutation groups which will be used in the proof of the characterization theorem.

Let G be a finite transitive group of permutations of a finite set X . Then the *rank* of G is the number of orbits of the stabilizer of a point. Rank 3 means that for $b \in X$ the stabilizer of b , G_b , has exactly 3 orbits on X , denoted b , $D(b)$ and $C(b)$. Choose the notation in such a way that $g(D(a)) = D(g(a))$ and $g(C(a)) = C(g(a))$ for all $a \in X$, $g \in G$. Let $|Y|$ denote the number of elements in a set Y . Set

$$|X| = n, |D(b)| = k \text{ and } |C(b)| = l$$

so that $n = 1 + k + l$. Set

$$|D(a) \cap D(b)| = \begin{cases} \lambda & \text{for } b \in D(a) \\ \mu & \text{for } b \in C(a) \end{cases}$$

The *parameters* of G are the triple (n, k, l) .

LEMMA 2.1. *Let G be a rank 3 permutation group. Then*

- (i) $\mu l = k(k - \lambda - 1)$.
- (ii) G is primitive iff $0 < \mu < k$. If G is primitive, then $(l, k) > 1$ where (l, k) denotes the greatest common divisor of l and k .
- (iii) If G is imprimitive, then either $(l + 1) | k$ or $(k + 1) | l$ where $a | b$ denotes that a divides b .
- (iv) If $|G|$ is odd, $k = l$.
- (v) If $|G|$ is even, then $D(\lambda - \mu)^2 + 4(k + \mu)$ is a square.
- (vi) If $|G|$ is even, then $a \in D(b)$ iff $b \in D(a)$.

Proof. See [3] and [4].

Assume $|G|$ is even. Define the "lines" of X as follows: for $a \neq b$ in X define

$$R(ab) = \bigcap \{z^\perp : a, b \in z^\perp\}$$

where $z^\perp = z \cup D(z)$. Call $R(ab)$ totally singular (resp. hyperbolic) if $a \in b^\perp$ (resp. $a \notin b^\perp$).

LEMMA 2.2. Let G be a rank 3 group of even order. Then

- (i) $g(R(ab)) = R(g(a)g(b))$ for all $a, b \in X, g \in G$.
- (ii) If $x \in R(ab)$ and $x \neq a$, then $R(ax) = R(ab)$ if $b \in D(a)$ or if $b \in C(a)$ and $\mu > \lambda + 1$.
- (iii) $x \in R(ab) - \{a\}$ iff $a^\perp \cap x^\perp = a^\perp \cap b^\perp$.
- (iv) $|R(ab)| - 1$ divides k if $b \in D(a)$.

Proof. See [4].

Let $T(a)$ denote the pointwise stabilizer of a^\perp . Then $T(a)$ is a normal subgroup of G_a .

LEMMA 2.3. Let G be a primitive rank 3 group of even order such that $\mu > \lambda + 1$. Then

- (i) $T(a)$ fixes all lines through a .
- (ii) $T(a)_x = 1$ for $x \in C(a)$.
- (iii) $|T(a)|$ divides $|R(ab)| - 1$, if $b \in C(a)$.

NOTATION. If $Y \subseteq X$, let G_Y denote the global stabilizer of Y . If $Y, Z \subseteq X$, then $G_{Y,Z}$ denotes $G_Y \cap G_Z$.

If $Y \subseteq X$, let $X - Y$ denote the set of elements of X which do not belong to Y .

For a natural number r , let v_r denote $(q^r - 1)/(q - 1)$.

3. The proof of the theorem. We now begin the proof of the characterization theorem. Assume that G is a rank 3 permutation

group of a set X which satisfies the hypotheses of the theorem.

LEMMA 3.1. (i) G is primitive of even order.

(ii) $\mu = \lambda + 2 = v_{r-2}$.

(iii) $a^\perp \cap b^\perp \neq R(ab)$ for $b \in D(a)$.

Proof. (i) Assume G is imprimitive. By Lemma 2.1 (iii) either $(k+1)|l$ or $(l+1)|k$. The first case does not occur because $k+1 = v_{r-1}$, $l = q^{r-1}$ and $(v_{r-1}, q^{r-1}) = 1$. The second case does not occur because $l+1 > k$. So G is primitive. Since $k \neq l$, $|G|$ is even by Lemma 2.1 (iv).

(ii) By Lemma 2.1 (i), $\mu q^{r-1} = qv_{r-2}(qv_{r-2} - \lambda - 1)$. By Lemma 2.1 (ii), $\mu > 0$. Since $(q^{r-2}, v_{r-2}) = 1$, there is a natural number t such that $v_{r-2}t = \mu$. So $\lambda + 1 = q(v_{r-2} - tq^{r-3})$ and $v_{r-2} - tq^{r-3} \geq 1$. If $t > 1$, then

$$v_{r-2} - 1 = qv_{r-3} \geq tq^{r-3} \geq 2q^{r-3}$$

which implies $2q^{r-4} \geq q^{r-3} + 1$, a contradiction because $q \geq 2$. So $t = 1$, $\mu = v_{r-2}$ and $\lambda - 1 = qv_{r-3}$.

(iii) Assume $a^\perp \cap b^\perp = R(ab)$ for $b \in D(a)$. Let $|R(ab)| = s + 1$. So $\lambda + 2 = s + 1 = \mu$. Since $s|k = q\mu$ by Lemma 2.2 (iv) and $(q, \mu) = 1$, there is a natural number t such that $st = q$. Then $\mu - 1 = qv_{r-3} = s$ implies $tv_{r-3} = 1$ and $r = 4$, a contradiction. This completes the proof of the lemma.

LEMMA 3.2. (i) $|a^\perp \cap C(b)| = q^{r-2}$ for $b \in D(a)$.

(ii) G_{ab} is transitive on the points of $a^\perp \cap C(b)$ for $b \in D(a)$.

Proof. (i) Since $a^\perp \cap C(b) = a^\perp - (a^\perp \cap b^\perp)$, by Proposition 3.1

(iii) $|a^\perp \cap C(b)| = k + 1 - (\lambda + 2) = q^{r-2}$.

(ii) Let $b \in D(a)$ and let $d \in a^\perp \cap C(b)$. Now

$$\begin{aligned} qv_{r-2} \cdot |G_{ab} : G_{abd}| &= |G_b : G_{ab}| \cdot |G_{ab} : G_{abd}| = |G_b : G_{bd}| |G_{bd} : G_{abd}| \\ &= q^{r-1} \cdot |G_{bd} : G_{abd}|. \end{aligned}$$

Let $x = |G_{ab} : G_{abd}|$. Since $(v_{r-2}, q^{r-2}) = 1$, it follows that $q^{r-2} | x$. But $x \leq q^{r-2}$ because

$$d^{G_{ab}} \subseteq a^\perp \cap C(b).$$

So $x = q^{r-2}$ and the proof is complete.

PROPOSITION 3.3. G_{ab} is transitive on the points of $a^\perp \cap b^{\perp_{\frac{1}{2}}} - R(ab)$ for $b \in D(a)$.

Proof. Let c and e be distinct points of $a^\perp \cap b^\perp - R(ab)$. Since

$c \notin R(ab)$, by Lemma 2.2 (iii) $c^\perp \not\subseteq a^\perp \cap b^\perp$. There is $u \in a^\perp \cap b^\perp \cap C(c)$. Since $e \notin R(ab)$, there is $v \in a^\perp \cap b^\perp \cap C(e)$. There are 4 possible cases to consider: (1) $u \in C(e)$, (2) $v \in C(c)$, (3) $u = e$ or $v = c$ and (4) $u \in D(e)$ and $v \in D(c)$.

(1) $u \in a^\perp \cap b^\perp \cap C(c) \cap C(e)$. Since $|R(uc)| > 2$, there is $y \in R(uc) - \{u, c\}$. By Proposition 3.1 (ii) and Lemma 2.2 (ii) it follows that $R(y) = R(uc) \subseteq a^\perp \cap b^\perp$. Because $R(uc)$ is a hyperbolic line and $T(y)$ is transitive on the points unequal to y of $R(y)$, there exists t in $T(y)$ such that $t(c) = u$. Since $a, b \in y^\perp$, it follows that $t \in G_{ab}$. Similarly there is $z \in R(ue) - \{u, e\}$ and then $R(z) = R(ue) \subseteq a^\perp \cap b^\perp$. Because $R(ue)$ is a hyperbolic line and $T(z)$ is transitive on the points unequal to z of $R(z)$, there exists s in $T(z) \leq G_{ab}$ such that $s(u) = e$. Thus $st(c) = e$ and $st \in G_{ab}$.

(2) $v \in a^\perp \cap b^\perp \cap C(c) \cap C(e)$. This case has a proof similar to that of case (1).

(3) If $u = e$ or $v = c$, then $R(ce)$ is a hyperbolic line in $a^\perp \cap b^\perp$. Pick $z \in R(c, e) - \{c, e\}$. There exists t in $T(z) \leq G_{ab}$ such that $t(c) = e$.

(4) $u \in a^\perp \cap b^\perp \cap C(c) \cap D(e)$ and $v \in a^\perp \cap b^\perp \cap D(c) \cap C(e)$. Since $|R(ce)| > 2$, there is $w \in R(ce) - \{c, e\}$. Note that $w \in C(u)$, for if $w \in u^\perp$, then $c \in R(ce) = R(w) \subseteq u^\perp$, a contradiction. Now $w \in R(ce) \subseteq a^\perp \cap b^\perp$. But $w \notin R(ab)$ because $u \in a^\perp \cap b^\perp \cap C(w) \cap C(c)$. By case (1) there exists $g \in G_{ab}$ such that $g(c) = w$. Note that $w \in C(v)$ for if $w \in v^\perp$, then $e \in R(ce) = R(w) \subseteq v^\perp$, a contradiction. Now $v \in a^\perp \cap b^\perp \cap C(w) \cap C(e)$. By case (1) there exists $h \in G_{ab}$ such that $h(w) = e$. So $hg(c) = e$ and $hg \in G_{ab}$. This completes the proof of the proposition.

PROPOSITION 3.4. *The group G_a is a rank 3 permutation group on the set of totally singular lines through a .*

Proof. Clearly G_a is transitive on the set of totally singular lines through a since $D(a)$ is an orbit of G_a . For $b \in D(a)$ define the sets $D(R(ab))$ and $C(R(ab))$ as follows:

$$\begin{aligned} D(R(ab)) &= \{R(ac) : c \in a^\perp \cap b^\perp - R(ab)\} \\ C(R(ab)) &= \{R(ac) : c \in a^\perp \cap C(b)\}. \end{aligned}$$

We claim that these sets are well-defined, form a partition of the set of totally singular lines through a unequal to $R(ab)$ and are nontrivial orbits of $G_{aR(ab)}$.

These sets are well-defined. Indeed suppose $R(ab) = R(ad)$ for $b, d \in D(a)$. By Lemma 2.2 (iii), $a^\perp \cap b^\perp = a^\perp \cap d^\perp$ and so $a^\perp \cap C(b) = a^\perp \cap C(d)$. Thus $D((ab)) = D(R(ad))$ and $C(R(ab)) = C(R(ad))$.

Let $R(az)$ be a totally singular line. Either $z \in C(b)$ in which case $R(az) \in C(R(ab))$ or $z \in b^\perp$. If $z \in b^\perp$, then either $z \in R(ab)$ in which

case $R(az) = R(ab)$ or $z \notin R(ab)$ in which case $R(az) \in D(R(ab))$. Thus

$$R(ab) \cup D(R(ab)) \cup C(R(ab))$$

is a partition of the set of totally singular lines through a .

If $R(ac)$ and $R(ae)$ belong to $D(R(ab))$, then c and e are elements of $a^\perp \cap b^\perp = R(ab)$. By Proposition 3.3 there is $g \in G_{ab}$ such that $g(c) = e$. So $g(R(ab)) = R(ab)$ and $g(R(ac)) = R(ae)$. Thus $D(R(ab))$ is an orbit of $G_{aR(ab)}$.

If $R(ac)$ and $R(ae)$ belong to $C(R(ab))$, then c and e are elements of $a^\perp \cap C(b)$. By Lemma 3.2 (ii) there is $g \in G_{ab}$ such that $g(c) = e$. Thus $C(R(ab))$ is an orbit of $G_{aR(ab)}$ and G_a is a rank 3 group on the set of totally singular lines through a , as desired.

PROPOSITION 3.5. *Totally singular lines carry $q + 1$ points.*

Proof. Let $|R(ab)| = s + 1$. We will show that $s = q$ by determining the rank 3 parameters of G_a on the set of totally singular lines through a . Let $k_2 = |D(R(ab))|$ and $l_2 = |C(R(ab))|$. Then by Lemma 3.1

$$k_2 = (\lambda + 2 - (s + 1))/s = (qv_{r-3}/s) - 1$$

and

$$l_2 = (k + 1 - (\lambda + 2))/s = q^{r-2}/s.$$

So there is a natural number t such that $st = q$. We claim that $t = 1$.

Now G_a is a rank 3 group with $k_2 = tv_{r-3} - 1$ and $l_2 = tq^{r-3}$. We claim that G_a is primitive. If G_a is imprimitive, then by Lemma 2.1 (iii) either $k_2 + 1 = tv_{r-3}$ divides $l_2 = tq^{r-3}$, a contradiction since $(v_{r-3}, q^{r-3}) = 1$ or $l_2 + 1$ divides k_2 , a contradiction since $k_2 < l_2 + 1$. So G_a is primitive. By Lemma 2.1 (ii), $(k_2, l_2) > 1$. Let $z = (k_2, l_2)$. We claim that $z = q + t - 1$.

Since $(1 - q)v_{r-3} + q^{r-3} = 1$, it follows that $(1 - q)k_2 + l_2 = q + t - 1$ and that there is a natural number u such that $zu = q + t - 1$. By Lemma 2.1 (i) $\mu_2 l_2 = k_2(k_2 - \lambda_2 - 1)$. So there is a natural number w such that $wk_2/z = \mu_2$. Then $wl_2/z = k_2 - \lambda_2 - 1$ and

$$2 \leq \lambda_2 + 2 = tv_{r-3} - tq^{r-3}w/z.$$

Now $(z, t) = 1$ for if the prime $p \mid (z, t)$, then $p \mid k_2 = tv_{r-3} - 1$ and $p \mid t$, a contradiction. So $v_{r-3} - q^{r-3}w/z$ is a natural number. From the substitution of $z = (q + t - 1)/u$ into $v_{r-3} - q^{r-3}w/z \geq 1$, it follows that

$$(uw - 1)q^{r-4} + 1 \leq tv_{r-4} \leq (q - 1)v_{r-4}$$

because $t \leq q - 1$ as $s > 1$. Then $0 \geq (uw - 2)q^{r-4} + 2$ which forces $u = w = 1$. So $z = q + t - 1$, $\mu_2 = k_2/z$ and $\lambda_2 + 2 = t\mu_2$.

Now $|G_a|$ is even. For if $|G_a|$ is odd, then $k_2 = l_2$ and $tv_{r-3} - 1 = tq_{r-3}$, which is impossible. By Lemma 2.1 (v)

$$\begin{aligned} D &= (t\mu_2 - 2 - \mu_2)^2 + 4(z\mu_2 - \mu_2) = (t - 1)^2\mu_2^2 + 4(q - 1)\mu_2 + 4 \\ &= ((t - 1)\mu_2 + b + 2)^2 \end{aligned}$$

for some nonnegative integer b . If $b = 0$, then $t = q$ and $s = 1$, a contradiction. So $b \geq 1$ and

$$4(q - 1)\mu_2 + 4 = 2(t - 1)\mu_2(b + 2) + (b + 2)^2$$

implies $b = 2c$ for some natural number c . It follows that

$$((q - 1) - (t - 1)(c + 1))\mu_2 = c(c + 2).$$

Assume $t > 1$. Since $(q - 1) - (t - 1)(c + 1) > 0$, it follows that $q \geq (t - 1)c + t \geq c + 2$ and $\mu_2 \leq c(c + 2) < q^2$. But

$$\mu_2 = k_2/z \geq (2v_{r-3} - 1)/2q.$$

If $r \geq 7$, then $\mu_2 \geq (2v_4 - 1)/2q > q^2$, a contradiction.

If $r = 5$, then $\mu_2 = t - (t - 1)^2/(q + t - 1)$. Since $t > 1$, there is a natural number f such that $(q + t - 1)f = (t - 1)^2$. Since $q = st$, it follows that $stf = (t - 1)(t - 1 - f)$ and that $t|(t - 1 - f)$, a contradiction.

If $r = 6$, then

$$\mu_2 = (tv_3 - 1)/(q + t - 1) = (t - 1)q^2/(q + t - 1) + q + 1.$$

Note $t > 2$ for $t = 2$ implies $q^2/(q + 1)$ is a natural number. So $\mu_2 \geq 2q^2/2q + q + 1 = 2q + 1$. Since μ_2 divides $c(c + 2)$ and $(c, c + 2) = 1$ or 2 , it follows that $\mu_2 \leq 2(c + 2)$. But $c + 2 \leq q$ and so $\mu_2 \leq 2q$, a contradiction. Therefore $t = 1$ and $s = q$ for all $r \geq 5$.

LEMMA 3.6. If $b \in D(a)$, then $X = \bigcup \{c^\perp : c \in R(ab)\}$.

Proof. We know $|R(ab)| = q + 1$. Let $R(ab) = \{d_1, d_2, \dots, d_{q+1}\}$. Let $R = \bigcup \{d_i^\perp : 1 \leq i \leq q + 1\}$. Express R as a pairwise disjoint union of $q + 1$ subsets of X .

$$R = d_1^\perp \cup \bigcup_{i=2}^{q+1} \left(d_i^\perp \cap \bigcap_{j=1}^{i-1} C(d_j) \right).$$

We claim that

$$d_i^\perp \cap \bigcap_{j=1}^{i-1} C(d_j) = d_i^\perp \cap C(d_1).$$

This is true for $i = 2$. Let $i > 2$. Then

$$d_i^\perp \cap C(d_i) = \left(d_i^\perp \cap C(d_i) \cap \bigcap_{j=2}^{i-1} C(d_j) \right) \cup \left(d_i^\perp \cap C(d_i) \cap \left(\bigcup_{j=2}^{i-1} d_j^\perp \right) \right).$$

Suppose there is $x \in d_i^\perp \cap C(d_i) \cap \left(\bigcup_{j=2}^{i-1} d_j^\perp \right)$. Then $x \in d_i^\perp \cap d_j^\perp \cap C(d_i)$ for $i \neq j$. So $d_i \in R(d_j) \subseteq x^\perp$, a contradiction. So

$$d_i^\perp \cap C(d_i) \cap \left(\bigcup_{j=2}^{i-1} d_j^\perp \right) = \emptyset$$

and the claim holds for $2 \leq i \leq q+1$. So

$$R = d_1^\perp \cup \bigcup_{i=2}^{q+1} \left(d_i^\perp \cap C(d_i) \right)$$

and this union is pairwise disjoint. Now

$$|R| = k + 1 + q(k + 1 - (\lambda + 2)) = v_r = |X|.$$

Thus $R = X$ and the proof of the lemma is complete.

PROPOSITION 3.7. *X together with the totally singular lines of X forms a nondegenerate Shult space of finite rank ≥ 3 in which lines carry $q+1$ points.*

Proof. It suffices to show that if $x \in R(ab)$ for $b \in D(a)$, then x is adjacent to either one point or all points of $R(ab)$. By definition two distinct points are adjacent if they determine a totally singular line. By Lemma 3.6, there exists $c \in R(ab)$ such that $x \in c^\perp$. If $x \in d^\perp$ for $d \in R(ab) - \{c\}$, then $R(ab) = R(cd) \subseteq x^\perp$ and $x \in e^\perp$ for all $e \in R(ab)$. Thus X is a nondegenerate Shult space in which lines carry $q+1$ points.

It remains to show that X has rank ≥ 3 . For $b \in D(a)$, there is $c \in a^\perp \cap b^\perp - R(ab)$ by Lemma 3.1 (iii). Define the “plane” $R(abc)$ by

$$R(abc) = \bigcap \{z^\perp : a, b, c \in z^\perp\}.$$

We claim that $R(abc)$ is a subspace of the Shult space X . If so, then X has rank ≥ 3 since

$$a \subset R(ab) \subset R(abc)$$

is a chain of subspaces of X . To prove that $R(abc)$ is a subspace, we need the following lemma.

LEMMA 3.8. *$w \in R(abc)$ iff $w^\perp \supseteq a^\perp \cap b^\perp \cap c^\perp$.*

Proof. Let $w \in R(abc)$. If $u \in a^\perp \cap b^\perp \cap c^\perp$, then $a, b, c \in u^\perp$ and $w \in u^\perp$ since $w \in R(abc)$. So $u \in w^\perp$ and $a^\perp \cap b^\perp \cap c^\perp \subseteq w^\perp$. Conversely,

assume $a^\perp \cap b^\perp \cap c^\perp \subseteq w^\perp$. Let $a, b, c \in z^\perp$. Then $z \in a^\perp \cap b^\perp \cap c^\perp \subseteq w^\perp$ and $w \in z^\perp$. By definition of "plane," $w \in R(abc)$ and the lemma is proved.

By definition $R(abc)$ is a subspace if any two points of $R(abc)$ are adjacent and if any line meeting $R(abc)$ in more than one point is contained in $R(abc)$. Let $d, e \in R(abc)$. Since $a, b, c \in a^\perp \cap b^\perp \cap c^\perp$, it follows that $R(abc) \subseteq a^\perp \cap b^\perp \cap c^\perp$. By Lemma 3.8.

$$d \in R(ab) \subseteq a^\perp \cap b^\perp \cap c^\perp \subseteq e^\perp.$$

So any two points of $R(abc)$ are adjacent. Let the line $R(xy)$ meet $R(abc)$ in $\{u, v\}$. Then $R(xy) = R(uv)$ and $x^\perp \cap y^\perp = u^\perp \cap v^\perp$ by Lemma 2.2. If $z \in R(xy)$, then

$$z^\perp \supseteq x^\perp \cap y^\perp = u^\perp \cap v^\perp \supseteq a^\perp \cap b^\perp \cap c^\perp$$

since $u, v \in R(abc)$. By Lemma 3.8, $z \in R(abc)$. Thus $R(xy) \subseteq R(abc)$ and $R(abc)$ is a subspace of the Shult space X , as desired.

PROPOSITION 3.9. (i) q is a prime power and r is even.

(ii) Either X is isomorphic to the polar space S associated with an alternating form f defined on a projective space P of dimension $r - 1$ over $GF(q)$ or X is isomorphic to the polar space S associated with a symmetric form f defined on a projective space P of dimension r over $GF(q)$ for q odd.

Proof. By Proposition 3.7 and Theorem 4 of Buekenhout and Shult [1], X is a polar space of rank ≥ 3 in which lines carry $q + 1 \geq 3$ points. Since $|X| = v_r$ is finite, by Theorem 1 of Buekenhout and Shult [1], X is isomorphic to the set of singular points of a classical symplectic, unitary or orthogonal geometry. Because a line of X carries $q + 1$ points and corresponds to a totally singular line of a classical geometry, it follows that q is a prime power. Note that $|X| = v_r$ equals the number of singular points of a classical geometry. It follows that either the geometry is symplectic or orthogonal and that $r = 2m$ for some $m \geq 3$ since X has rank ≥ 3 . Statement (ii) now follows.

PROPOSITION 3.10. (i) G is isomorphic to a subgroup of $PGU(f)$, the group of collineations of P which preserve the form f .

(ii) For $x \in X$, $\varphi(x^\perp) = \{w \in P: f(w, w) = 0, f(w, \varphi(x)) = 0\}$ where $\varphi: X \rightarrow S$ is a polar space isomorphism.

(iii) For $x, y \in X$, $\varphi(R(x, y))$ is the set of singular points of the projective line determined by $\varphi(x)$ and $\varphi(y)$.

(iv) X is isomorphic to a symplectic geometry.

Proof. (i) The group G is a subgroup of the group of automorphisms of the polar space X , which we denote by $\text{Aut}(X)$. If $\varphi: X \rightarrow S$ is a polar space isomorphism, then define a map

$$\begin{aligned}\psi: \text{Aut}(X) &\longrightarrow \text{Aut}(S) \quad \text{by} \\ \psi(s) &= \varphi s \varphi^{-1}\end{aligned}$$

for $s \in \text{Aut}(X)$. It follows that ψ is a group isomorphism. Now $P\Gamma U(f) \cong \text{Aut}(S)$ by a natural map defined by

$$\begin{aligned}P\Gamma U &\longrightarrow \text{Aut}(S) \\ u &\longrightarrow \text{the restriction of } u \text{ to } S.\end{aligned}$$

See Dieudonné [2] pp. 82–84. So $\psi(G)$ is a subgroup of $P\Gamma U(f)$.

(ii) This statement claims that $\varphi(x \cup D(x))$ is the hyperplane of singular points of S which are perpendicular to $\varphi(x)$. Denote this hyperplane by $\varphi(x)^\perp$, where $^\perp$ is the polarity determined by the form f .

Since $x^\perp = \bigcup \{R(xb): b \in D(x)\}$, it follows that

$$\varphi(x^\perp) = \bigcup \{\varphi(R(xb)): b \in D(x)\} \subseteq \varphi(x)^\perp$$

because $\varphi(R(xb))$ is a totally singular line of P . So $\varphi(x^\perp) \subseteq \varphi(x)^\perp$.

Conversely for $z \in \varphi(x)^\perp$, there exists $b \in X$ such that $z = \varphi(b)$ and $\varphi(b)^\perp \varphi(x)$. Suppose $b \notin x^\perp$. Then $b \in C(x)$, an orbit of G_x . For $c \in C(x)$ there exists $g \in G_x$ such that $g(b) = c$. Then $\psi(g) \in \text{Aut}(S)$ and $\psi(g)$ preserves the polarity $^\perp$. Since $\varphi(b)^\perp \varphi(x)$, it follows that $(\psi(g)(\varphi(b)))^\perp (\psi(g)(\varphi(x)))$ and $\varphi(c)^\perp \varphi(x)$. So $\varphi(c) \in \varphi(x)^\perp$ for all $c \in C(x)$. Since $\varphi(x^\perp) \subseteq \varphi(x)^\perp$, it follows that $\varphi(X) = S \subseteq \varphi(x)^\perp$, a contradiction. Thus $b \in x^\perp$, $\varphi(b) = z \in \varphi(x^\perp)$ and $\varphi(x)^\perp \subseteq \varphi(x^\perp)$.

(iii) Since $R(xy) = \bigcap \{u^\perp: x, y \in u^\perp\}$, it follows from (ii) that

$$\varphi(R(xy)) = \bigcap \{v^\perp: v \in S \text{ and } \varphi(x), \varphi(y) \in v^\perp\}.$$

So $\varphi(R(xy))$ is the set of singular points of the projective line determined by $\varphi(x)$ and $\varphi(y)$.

(iv) Assume X is an orthogonal geometry. If $y \in C(x)$, then $\varphi(R(xy))$ is a hyperbolic line in an orthogonal geometry and so carries just 2 singular points. But $|\varphi(R(xy))| = |R(xy)| > 2$, by hypothesis of the theorem. This contradiction shows that X must be a symplectic geometry. So the proposition is established.

PROPOSITION 3.11. (i) *The nontrivial elements of $T(x)$ correspond to elations of P .*

(ii) *$\psi(G)$ contains $P\text{Sp}(2m, q)$ as a normal subgroup.*

Proof. (i) Because X is symplectic, all points of P are singular and $S = P$. Because $|R(ax)| > 2$ by Lemma 2.1 (iii), there exists a nontrivial element t of $T(x)$. Then t fixes x^\perp pointwise and t fixes no point outside x^\perp by Lemma 2.3 (ii). It easily follows from Proposition 3.10 (ii) that $\psi(t)$ fixes the hyperplane $\varphi(x)^\perp$ pointwise and $\psi(t)$ fixes no point outside this hyperplane. Thus $\psi(t)$ is an elation of P . Since $|T(x)| \mid (|R(xy)| - 1)$ for $y \in C(x)$, since hyperbolic lines of S carry $q + 1$ points and since $T(x)$ is transitive on $R(xy) - \{x\}$, it follows that $|T(x)| = q$.

(ii) $\psi(G)$ contains q elations for each point v of P . Since these elations generate $PSp(2m, q)$, (ii) holds.

REFERENCES

1. F. Buekenhout and E. Shult, *On the foundations of polar geometry*, Geometriae Dedicata, **3** (1974), 155-170.
2. J. Dieudonné, *La géométrie des groupes classiques*, second edition, Springer-Verlag, Berlin, 1963.
3. M. Hestenes and D. Higman, *Rank 3 subgroups and strongly regular graphs*, Computers in number theory and algebra, SIAM-AMS Proceedings, American Mathematical Society, Providence, 1971.
4. D. Higman, *Finite permutation groups of rank 3*, Math. Z., **86** (1964), 145-156.
5. W. Kantor, *Rank 3 characterizations of classical geometries*, to appear.
6. T. Tsuzuku, *On a problem of D. G. Higman*, to appear.

Received April 21, 1975.

UNIVERSITY OF MICHIGAN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Aharon Atzmon, <i>A moment problem for positive measures on the unit disc</i>	317
Peter W. Bates and Grant Bernard Gustafson, <i>Green's function inequalities for two-point boundary value problems</i>	327
Howard Edwin Bell, <i>Infinite subrings of infinite rings and near-rings</i>	345
Grahame Bennett, Victor Wayne Goodman and Charles Michael Newman, <i>Norms of random matrices</i>	359
Beverly L. Brechner, <i>Almost periodic homeomorphisms of E^2 are periodic</i>	367
Beverly L. Brechner and R. Daniel Mauldin, <i>Homeomorphisms of the plane</i>	375
Jia-Arng Chao, <i>Lusin area functions on local fields</i>	383
Frank Rimi DeMeyer, <i>The Brauer group of polynomial rings</i>	391
M. V. Deshpande, <i>Collectively compact sets and the ergodic theory of semi-groups</i>	399
Raymond Frank Dickman and Jack Ray Porter, <i>θ-closed subsets of Hausdorff spaces</i>	407
Charles P. Downey, <i>Classification of singular integrals over a local field</i>	417
Daniel Reuven Farkas, <i>Miscellany on Bieberbach group algebras</i>	427
Peter A. Fowler, <i>Infimum and domination principles in vector lattices</i>	437
Barry J. Gardner, <i>Some aspects of T-nilpotence. II: Lifting properties over T-nilpotent ideals</i>	445
Gary Fred Gruenhagen and Phillip Lee Zenor, <i>Metritzation of spaces with countable large basis dimension</i>	455
J. L. Hickman, <i>Reducing series of ordinals</i>	461
Hugh M. Hilden, <i>Generators for two groups related to the braid group</i>	475
Tom (Roy Thomas Jr.) Jacob, <i>Some matrix transformations on analytic sequence spaces</i>	487
Elyahu Katz, <i>Free products in the category of k_w-groups</i>	493
Tsang Hai Kuo, <i>On conjugate Banach spaces with the Radon-Nikodým property</i>	497
Norman Eugene Liden, <i>K-spaces, their antispace and related mappings</i>	505
Clinton M. Petty, <i>Radon partitions in real linear spaces</i>	515
Alan Saleski, <i>A conditional entropy for the space of pseudo-Menger maps</i>	525
Michael Singer, <i>Elementary solutions of differential equations</i>	535
Eugene Spiegel and Allan Trojan, <i>On semi-simple group algebras. I</i>	549
Charles Madison Stanton, <i>Bounded analytic functions on a class of open Riemann surfaces</i>	557
Sherman K. Stein, <i>Transversals of Latin squares and their generalizations</i>	567
Ivan Ernest Stux, <i>Distribution of squarefree integers in non-linear sequences</i>	577
Lowell G. Sweet, <i>On homogeneous algebras</i>	585
Lowell G. Sweet, <i>On doubly homogeneous algebras</i>	595
Florian Vasilescu, <i>The closed range modulus of operators</i>	599
Arthur Anthony Yanushka, <i>A characterization of the symplectic groups $\mathrm{PSp}(2m, q)$ as rank 3 permutation groups</i>	611
James Juei-Chin Yeh, <i>Inversion of conditional Wiener integrals</i>	623