

Pacific Journal of Mathematics

**FUNDAMENTAL UNITS AND CYCLES IN THE PERIOD OF
REAL QUADRATIC NUMBER FIELDS. I**

LEON BERNSTEIN

FUNDAMENTAL UNITS AND CYCLES IN THE PERIOD OF REAL QUADRATIC NUMBER FIELDS

LEON BERNSTEIN

PART I

0. Introduction. In this paper we introduce the concept of "Cycles in the Period" of the simple continued fraction expansion of a real quadratic irrational. This is expressed in the

DEFINITION. Let M, D, d be positive rational integers, M square free, $M = D^2 + d, d \leq 2D$. Let k, a, s be nonnegative rational integers, $0 \leq a \leq k - 1$; let $f = f(k, a, s; d, D)$ be a polynomial with rational integral coefficients. For a fixed s , the finite sequence of polynomials

$$(0.1) \quad F(s) = f(k, a, s; d, D), f(k, a + 1, s; d, D), \dots, \\ f(k, a + k - 1, s; d, D)$$

will be called "Cycle in the Period" of the simple continued fraction expansion of \sqrt{M} if, for $s_0 \geq 1$, this expansion has the form

$$(0.2) \quad \sqrt{M} = [\overline{b_0, b_1, \dots, F(0), \dots, F(s_0 - 1), f(k, a, s_0; d, D), \dots,} \\ \overline{f(k, a + b, s_0; d, D), \dots, f(k, a, s_0; d, D), F'(s_0 - 1), \dots,} \\ \overline{F'(0), f(k, a - 1, 0; d, D), \dots, b_1, 2b_0}]$$

$b \geq 1; b \leq k - 1; k$ is the length of the cycle; $F'(s)$ means that the order of the $f - s$ must be reversed.

In the first part of this paper, the main result is the construction of infinitely many classes of quadratic fields $Q(\sqrt{M})$, each containing infinitely many M of a simple structure. Among the various classes thus constructed, there are a few in whose expansion of \sqrt{M} cycles in the period surprisingly have the length ≤ 12 . Functions $f(k, a, s; d, D), f(k, a + 1, s; d, D), \dots$ are of course stated explicitly; hence we are able to construct numbers \sqrt{M} such that the primitive period of their expansion has any given length m which is a function of the parameter k .

Expansions of \sqrt{M} which have the structure of cycles in the period were generally not known up to now. In a recent paper Y. Yamamoto [6] has given a few numerical examples of expansions of

real quadratic irrationals in which cycles of length two appear. The present paper, which generalizes this concept, was, however, developed independently of Yamoto's beautiful discovery.

It is an old dream of mathematicians to find infinite classes of real quadratic number fields $Q(\sqrt{M})$ for which the continued fraction expansion can be stated explicitly as a function of M . Very little knowledge in this direction was available up to now. The bit of it that was there is based on a theorem by Th. Muir [4], and in his "encyclopedia" of continued fractions O. Perron [5] has given a few demonstrations of Muir's theorem which the author does not believe to be of great practical significance. The most that can be achieved with Muir's theorem, is the explicitly stated expansion of certain classes of \sqrt{M} with a primitive period up to length six. In a recent paper [2] the author has given the following infinite classes of \sqrt{M} whose continued fraction expansion has a primitive period of lengths 10, 12 and 8 respectively; these are

$$\begin{aligned}
 & \sqrt{D^2 + 4d} \\
 (0.3) \quad & = [D, \overline{(2d)^{-1}(D-d), 1, 1, 2^{-1}(D-1), 2d^{-1}D, 2^{-1}(D-1)}, \\
 & \quad \overline{1, 1, (2d)^{-1}(D-d), 2D}] \\
 & \quad d \mid D; d > 1, D \text{ odd}; D > d
 \end{aligned}$$

$$\begin{aligned}
 & \sqrt{D^2 - 4d} \\
 (0.4) \quad & = [D-1, \overline{1, (2d)^{-1}(D-3d), 2, 2^{-1}(D-3), 1, 2d^{-1}(D-d)}, \\
 & \quad \overline{1, 2^{-1}(D-3), 2, (2d)^{-1}(D-3d), 1, 2b_0}] \\
 & \quad d \mid D; d > 1, D \text{ odd}; b_0 = D-1; D^2 - 4d \text{ squarefree}.
 \end{aligned}$$

$$\begin{aligned}
 & \sqrt{9d^2 - 4d} \\
 (0.5) \quad & = [3d-1, \overline{3, 3(d-1)2^{-1}, 1, 4, 1, 3(d-1)2^{-1}, 3, 2(3d-1)}] \\
 & \quad d > 1, d \text{ odd}.
 \end{aligned}$$

This is a special case of (0.9)

For $d = 1$, the primitive period in the above expansions has length at most half of the original one, as the reader verify easily. Then author constructed these expansions in order to find the well-known fundamental unit of G. Degert [3] in the corresponding quadratic field by an approach different from Degert's.

We shall recall the basic rules of expanding \sqrt{M} for further references:

$$\begin{aligned}
 (0.6) \quad & \sqrt{M} = w = x_0 = (w + P_0)Q_0^{-1} = b_0 + x^{-1}; P_0 = 0; Q_0 = 1; b_0 \\
 & = [w].
 \end{aligned}$$

$$(0.7) \quad \begin{aligned} x_k &= (w + P_k)Q_k^{-1} = b_k + x_{k+1}^{-1}; P_k = b_{k-1}Q_{k-1} - P_{k-1}; Q_{k-1}Q_k \\ &= w^2 - P_k^2; b_k = [x_k]; (k = 1, 2, \dots). \end{aligned}$$

From $P_k + P_{k-1} = b_{k-1}Q_{k-1}$, $w^2 - P_k^2 = Q_{k-1}Q_k$, we derive the formula

$$(0.8) \quad \begin{aligned} (w + P_{k-1})(w + P_k) &= Q_{k-1}(Q_k + b_{k-1}(w + P_k)) \\ &\quad (k = 1, 2, \dots). \end{aligned}$$

With m denoting the length of the primitive period of \sqrt{M} as a periodic continued fraction expansion, the following rules hold:

$$(0.9) \quad \begin{aligned} \text{Let } m = 2k; \text{ then } P_k &= P_{k+1}, \text{ and vice versa; also } b_{k-i} = \\ b_{k+i}; (i = 1, \dots, k); b_{2k} &= 2b_0^k. \end{aligned}$$

$$(0.10) \quad \begin{aligned} \text{Let } m = 2k + 1; \text{ then } Q_k &= Q_{k+1}, \text{ and vice versa; also} \\ b_k &= b_{k+1}; b_{k-i} = b_{k+i+1} (i = 1, \dots, k-1); b_{2k+1} = 2b_0. \end{aligned}$$

$$(0.11) \quad \begin{aligned} \text{If } m = 2k, \text{ then } P_{k-i} &= P_{k+i+1} (i = 0, \dots, k-1); Q_{k-i} = \\ Q_{k+i} (i = 1, \dots, k). \end{aligned}$$

$$(0.12) \quad \begin{aligned} \text{If } m = 2k + 1, \text{ then } P_{k-i} &= P_{k+i+2} (i = 0, \dots, k-1); \\ Q_{k-i} &= Q_{k+i+1} (i = 0, \dots, k). \end{aligned}$$

$$(0.13) \quad \begin{aligned} Q_v > 2 (v = 1, 2, \dots). \text{ Only if } m = 2k, Q_k = 2 \text{ is possible} \\ \text{(then } P_k = P_{k+1}). \end{aligned}$$

In Part II (later in this volume) the fundamental unit of $Q(\sqrt{M})$, M square free, is stated explicitly, being calculated from the periodic expansion of \sqrt{M} .

The explicit representation of this expansion is therefore a primary issue. Units of algebraic number fields of any degree have been recently investigated by H. Zassenhaus [7]. This author also thinks that the calculation of units from the periodic expansion of a basis of the field, generally by Jacobi-Perron algorithm, is a most suitable tool.

In the following chapters w^2 is squarefree.

1. Expansion of $w = \sqrt{[(2a+1)^k + a]^2 + 2a+1}$; $a, k \geq 1$. The formula holds

$$(1.1) \quad w^2 = A^{2k} + 2aA^k + (a+1)^2 = (A^k + a)^2 + A; A = 2a+1.$$

$$(1.2) \quad [w] = A^k + a; w = A^k + a + r; \quad 0 < r < 1.$$

The reader will easily verify the following expansion for $k \geq 2$.

$$(1.3) \quad w = A^k + a + \frac{1}{x_1}; P_1 = A^k + a; Q_1 = A;$$

$$(1.4) \quad \frac{w + A^k + a}{A} = 2A^{k-1} + \frac{1}{x_2}; P_2 = A^k - a; Q_2 = 4aA^{k-1} + 1.$$

$$(1.5) \quad \frac{w + A^k - a}{4aA^{k-1} + 1} = 1 + \frac{1}{x_3}; P_3 = (A - 2)A^{k-1} + a + 1; Q_3 = 2A^{k-1}.$$

$$(1.6) \quad \frac{w + (A - 2)A^{k-1} + a + 1}{2A^{k-1}} = A - 1 + \frac{1}{x_4}; P_4 = A^k - (a + 1);$$

$$Q_4 = A^2.$$

$$(1.7) \quad \frac{w + A^k - (a + 1)}{A^2} = 2A^{k-2} - 1 + \frac{1}{x_5}.$$

We now prove the formulas

$$(1.8) \quad \begin{cases} \text{(i)} & P_{3s-1} = A^k - [A^s - (a + 1)]; Q_{3s-1} = 2(A^s - 1)A^{k-s} \\ & \quad - [A^s - 2(a - 1)]; b_{3s-1} = 1 \\ \text{(ii)} & P_{3s} = (A^s - 2)A^{k-s} + (a + 1); Q_{3s} = 2A^{k-s}; b_{3s} = A^s - 1 \\ \text{(iii)} & P_{3s+1} = A^k - (a + 1); Q_{3s+1} = A^{s+1}; b_{3s+1} = 2A^{k-(s+1)} - 1. \end{cases}$$

Proof by idduction. Formulas (1.8) are correct for $s = 1$, as can be easily verified from (1.3) to (1.7). We prove: (1.8) is true for s being replaced by $s + 1$. We obtain from (1.8), (iii), presuming $1 \leq s + 1 \leq k$,

$$(1.9) \quad \begin{cases} \frac{w + A^k - (a + 1)}{A^{s+1}} = 2A^{k-(s+1)} - 1 + \frac{1}{x_{3s+2}}; P_{3s+2} \\ \quad = A^k - [A^{s+1} - (a + 1)] \\ Q_{3s+2} = 2(A^{s+1} - 1)A^{k-(s+1)} - [A^{s+1} - 2(a + 1)]. \end{cases}$$

Since

$$\begin{aligned} 1 &< 1 + \frac{2A^{k-(s+1)} - 1 + r}{Q_{3s+2}} = x_{3s+2} < 1 + \frac{2A^{k-2}}{2A^k - 2A^{k-1} - A^k} \\ &= 1 + \frac{2}{(2a - 1)A} < 2 \end{aligned}$$

we obtain

$$(1.10) \quad \begin{cases} \frac{w + A^k - A^{s+1} + (a + 1)}{2A^k - 2A^{k-(s+1)} - A^{s+1} + 2(a + 1)} = 1 + \frac{1}{x_{3s+3}}; \\ P_{3s+3} = (A^{s+1} - 2)A^{k-(s+1)} + (a + 1); Q_{3s+3} = 2A^{k-(s+1)}. \end{cases}$$

Since, for $k \geq 2$,

$$A^{s+1} - 1 < A^{s+1} - 1 + \frac{A + r}{2A^{k-(s+1)}} = x_{3s+3} < A^{s+1} - 1 + \frac{2A}{2A^{k-(s+1)}} \leq A^{s+1},$$

we obtain

$$(1.11) \quad \begin{cases} \frac{w + A^k - 2A^{k-(s+1)} + a + 1}{2A^{k-(s+1)}} = A^{s+1} - 1 + \frac{1}{x_{3s+4}}; \\ P_{3s+4} = A^k - (a + 1); Q_{3s+4} = A^{s+2}. \end{cases}$$

Since

$$2A^{k-(s+2)} - 1 < 2A^{k-(s+2)} - 1 + \frac{A^{s+2} - 1 + r}{A^{s+2}} = x_{3s+4} < 2A^{k-(s+2)} - 1 + 1,$$

we have

$$(1.12) \quad b_{3s+4} = 2A^{k-(s+2)} - 1.$$

With formulas (1.9)-(1.12) formulas (1.8), with s replaced by $s + 1$ are verified. Since, as can easily be verified from (1.8), no P_v equals P_{v+1} , and no Q_v equals Q_{v+1} in the cycle $\{P_{3s-1}, P_{3s}, P_{3s+1}\}$ and $\{Q_{3s-1}, Q_{3s}, Q_{3s+1}\}$. We look for the possibility that some $Q_v = 2$. This happens for

$$(1.13) \quad Q_{3k} = 2.$$

We now obtain from (1.8), (ii), and from (1.2)

$$(1.14) \quad \begin{cases} \frac{w + A^k + a - 1}{2} = \frac{2A^k + 2a - r'}{2} = A^k + a - 1 + \frac{1}{x_{3k+1}} \\ P_{3k+1} = A^k + a - 1 = b_0 - 1 = P_{3k}. \end{cases}$$

The length of the primitive period of the expansion of w thus equals $1 + 6(k - 1) + 3 + 2 = 6k$, and we can state.

THEOREM 1. *Let a, k be natural numbers, $a \geq 1, k \geq 2, w^2 = (A^k + a)^2 + A, A = 2a + 1, w^2$ squarefree. The expansion of w as a periodic continued fraction has a primitive period of length $6k$ and the form*

$$(1.15) \quad \begin{cases} w = [\overline{b_0, b_1, \dots, b_{3s-1}, b_{3s}, b_{3s+1}, \dots, b_{3k-1}, b_{3k}, b_{3k-1}, \dots, b_1, 2b_0}] \\ b_0 = A^k + a; b_1 = 2A^{k-1}; b_{3k-1} = 1; b_{3k} = b_0 - 1 \\ b_{3s-1} = 1; b_{3s} = A^s - 1; b_{3s+1} = 2A^{k-s-1} - 1; s = 1, \dots, k - 1. \end{cases}$$

For $k = 1$, we obtain by a special calculation that

$$(1.16) \quad \sqrt{9a^2 + 8a + 2} = [3a + 2, 1, 3a, 1, 2, 6a + 2].$$

2. Expansion of $w = \sqrt{(A^k - a)^2 + A}; A = 2a + 1; a, k \geq 1$.
The formulas hold

$$(2.1) \quad w^2 = A^{2k} - 2aA^k + (a+1)^2.$$

$$(2.2) \quad [w] = A^k - a; w = A^k - a + r, 0 < r < 1.$$

The reader will easily verify the following

THEOREM 2. *Let a, k be natural numbers, $a \geq 1, k \geq 2, w^2 = (A^k - a)^2 + A, A = 2a + 1, w^2$ squarefree. The expansion of w as a periodic continued fraction has a primitive period of length $6k - 2$ and the form*

$$(2.3) \quad w = [b_0, \overline{\dots, b_{3s-2}, b_{3s-1}, b_{3s}, \dots, b_{3k-2}, b_{3k-1}, b_{3k-2}, \dots, b_1, 2b_0}]$$

$$(2.4) \quad \begin{cases} b_0 = A^k - a; b_{3k-1} = b_0 - 1; b_{3k-2} = 1 \\ b_{3s-2} = 2A^{k-s} - 1; b_{3s-1} = A^s - 1; b_{3s} = 1; s = 1, \dots, k-1. \end{cases}$$

For $k = 1$, we obtain the expansion

$$(2.5) \quad \sqrt{a^2 + 4a + 2} = [a + 1, \overline{1, a, 1, 2(a+1)}].$$

(2.6) is obtained from $\sqrt{a^2 + 4a + 2} = \sqrt{(a+2)^2 - 2}$, for which the author has found (2.6) in [1].

3. Expansion of $w = \sqrt{(A^k + a + 1)^2 - A}; A = 2a + 1; a, k \geq 1$.
The formulas hold

$$(3.1) \quad w^2 = A^{2k} + 2(a+1)A^k + a^2;$$

$$(3.2) \quad [w] = A^k + a; w = A^k + a + r; 0 < r < 1.$$

The reader will easily verify the following

THEOREM 3. *Let a, k be natural numbers, $a \geq 1, k \geq 2; w^2 = (A^k + a + 1)^2 - A; A = 2a + 1, w^2$ squarefree. The expansion of w as a periodic continued fraction has a primitive period of length $4k + 2$ and the form*

$$(3.3) \quad \begin{cases} w = [b_0, \overline{\dots, b_{2s-1}, b_{2s}, \dots, b_{2k+1}, b_{2k}, \dots, b_1, 2b_0}] \\ b_0 = b_{2k+1} = A^k + a; b_{2s-1} = A^{s-1}; b_{2s} = 2A^{k-s} \\ s = 1, \dots, k. \end{cases}$$

For $k = 1$ we obtain the expansion

$$\sqrt{9a^2 + 10a + 3} = [3a + 1, \overline{1, 1, 2, 3a + 1, 2, 1, 6a + 2}].$$

4. Expansion of $w = \sqrt{[A^k - (a+1)]^2 - A}; A = 2a + 1, a, k \geq 2$.
The formulas hold

$$(4.1) \quad w^2 = A^{2k} - 2(a+1)A^k + a^2$$

$$(4.2) \quad \begin{aligned} [w] &= A^k - (a + 2); \\ w &= A^k - a - 2 + r; 0 < r < 1. \end{aligned}$$

The reader will easily verify the following:

THEOREM 4. *Let $a, k \geq 1$ be natural numbers, $w^2 = [A^k - (a + 1)]^2 - A$, $A = 2a + 1$, w^2 squarefree. The expansion of w , as a periodic continued fraction, has a primitive period of length $4(2k - 1)$ and the form*

$$(4.3) \quad \left\{ \begin{aligned} w &= [b_0, \overline{b_1, b_2, \dots, b_{4s-1}, b_{4s}, b_{4s+1}, b_{4s+2}, \dots,} \\ &\quad \overline{b_{4k-5}, b_{4k-4}, b_{4k-3}, b_{4k-2}, b_{4k-1}, \dots, b_2, b_1, 2b_0}] \\ b_0 &= A^k - (a + 2); b_1 = 1, b_2 = 2A^{k-1} - 3 \\ b_{4s-1} &= 1; b_{4s} = A^s - 2; b_{4s+1} = 1; b_{4s+2} = 2(A^{k-s-1} - 1); \\ s &= 1, 2, \dots, k - 2, k \geq 3; a \geq 2; \\ b_{4k-5} &= 1; b_{4k-4} = A^{k-1} - 2; b_{4k-3} = 2; b_{4k-2} = b_0. \end{aligned} \right.$$

For $k = 2$, $a \geq 1$ the expansion holds

$$(4.4) \quad \begin{aligned} &\sqrt{(4a^2 + 3a)^2 - 2a + 1} \\ &= [4a^2 + 3a - 1, \overline{1, 4a - 1, 1, 2a - 1, 2, 4a^2 + 3a - 1, 2,} \\ &\quad \overline{2a - 1, 1, 4a - 1, 1, 8a^3 + 6a - 2}]. \end{aligned}$$

For $k = 1$, $a \geq 4$, the expansion holds

$$(4.5) \quad \sqrt{a^2 - 2a - 1} = [a - 2, \overline{1, a - 3, 1, 2(a - 2)}].$$

5. Expansion of $w = \sqrt{[A^k + (A - 1)]^2 + 4A}$; $A = 2^d b$, b odd ≥ 1 ; $d \geq 1$; $k \geq 2$. The formulas hold (d and b not both equal 1)

$$(5.1) \quad \begin{aligned} w^2 &= A^{2k} + 2(A - 1)A^k + (A + 1)^2; [w] \\ &= A^k + A - 1; w = [w] + r, 0 < r < 1. \end{aligned}$$

The reader will easily verify the following:

THEOREM 5. *Let $A = 2^d b$, b odd ≥ 1 , $d \geq 2$, $k \geq 2$ be natural numbers, $w^2 = [A^k + (A - 1)]^2 + 4A$ squarefree. If $k \geq 3$ is odd, then the expansion of w , as a periodic continued fraction has a primitive period of length $5k - 6$. The expansion has the form*

$$(5.2) \quad \left\{ \begin{aligned} w &= [b_0, \overline{b_1, \dots, b_{5s-3}, b_{5s-2}, b_{5s-1}, b_{5s}, b_{5s+1}, \dots,} \\ &\quad \overline{b_{1/2(5k-11)}, b_{1/2(5k-9)}, b_{1/2(5k-7)}, b_{1/2(5k-5)}, b_{1/2(5k-3)}, \dots, b_1, 2b_0}] \\ (i) \quad b_0 &= A^k + A - 1; b_1 = 2^{d-1}bA^{k-2} \\ (ii) \quad b_{5s-3} &= 2; b_{5s-2} = 2^{d-1}bA^{s-1} - 1; b_{5s-1} = b_{5s} = 1; b_{5s+1} \\ &= 2^{d-1}bA^{k-s-2} - 1. \end{aligned} \right.$$

$$\left\{ \begin{array}{l} s = 1, 2, \dots, \frac{1}{2}(k-3), k \geq 5. \\ \text{(iii)} \quad \begin{cases} b_{1/2(5k-11)} = 2, \\ b_{1/2(5k-9)} = 2^{d-1}bA^{1/2(k-3)} - 1, \\ b_{1/2(5k-7)} = 1. \end{cases} \end{array} \right.$$

For $k = 3$, the expansion has the form

$$(5.3) \quad \left\{ \begin{array}{l} \sqrt{[A^3 + (A-1)]^2 + 4A} \\ = [A^3 + a - 1, \overline{2^{d-1}bA}, 2, \overline{2^{d-1}b} - 1, 1, 1, \overline{2^{d-1}b} \\ - 1, 2, \overline{2^{d-1}bA}, 2(A^3 + A - 1)] \end{array} \right.$$

If $k \geq 2$ is even, the expansion of w , as a periodic continued fraction has a primitive period of length $5k - 6$. The expansion has the form

$$(5.4) \quad \left\{ \begin{array}{l} w = [b_0, \overline{b_1, \dots, b_{5s-3}, b_{5s-2}, b_{5s-1}, b_{5s}, b_{5s+1}, \dots, 2, \dots,} \\ \overline{b_{5s+1}, b_{5s}, b_{5s-1}, b_{5s-2}, b_{5s-3}, \dots, b_1, 2b_0}] \\ s = 1, 2, \dots, \frac{1}{2}(k-2); k \geq 4. \\ b_0, b_1, b_{5s-3}, b_{5s-2}, b_{5s-1}, b_{5s}, b_{5s+1} \text{ as in (5.17), (i), (ii).} \end{array} \right.$$

For $k = 2$, the expansion has the form

$$(5.5) \quad \left\{ \begin{array}{l} \sqrt{[A^2 + (A-1)]^2 + 4A} \\ = [A^2 + A - 1, \overline{2^{d-1}b}, 2, \overline{2^{d-1}b}, 2(A^2 + A - 1)] \end{array} \right.$$

EXAMPLE 1. For $d = 2, b = 1, A = 4$, we obtain from (5.18)

$$\sqrt{4505} = [67, \overline{8, 2, 1, 1, 1, 1, 2, 8, 134}].$$

EXAMPLE 2. For $d = 2, b = 1, A = 4$, we obtain from (5.20)

$$\sqrt{377} = [19, \overline{2, 2, 2, 38}].$$

The reader should note that for $k = 1$, we obtain the known expansion $\sqrt{4A^2 + 1} = [2A, \overline{4A}]$.

As is known in the case of an odd period of length $2r + 1$, the formulas hold

$$(5.6) \quad w^2 = P_{r+1}^2 + Q_{r+1}^2.$$

In our case $5k - 6 = 2r + 1$, hence

$$(5.7) \quad r + 1 = \frac{5}{2}(k - 1),$$

and so we obtain

$$(5.8) \quad Q_{5/2(k-1)}^2 + P_{5/2(k-1)}^2 = [A^k + (A-1)]^2 + 4A.$$

We obtain from (5.8), (iv), for $s = 1/2(k-1)$,

$$(5.9) \quad \begin{aligned} Q_{5/2(k-1)} &= (A^{1/2(k-1)} + 1)A^{k-1/2(k-1)} - [A^{1/2(k-1)+1} + 1(A+1)], \\ P_{5/2(k-1)} &= A^{k-(k-1)/2} + A^{(k-1)/2+1}, \\ P_{5/2(k-1)} &= 2A^{(k+1)/2}; \quad Q_{5/2(k-1)} = A^k - (A+1). \end{aligned}$$

Indeed:

$$\begin{aligned} P_{5/2(k-1)}^2 + Q_{5/2(k-1)}^2 &= 4A^{k+1} + A^{2k} - 2(A+1)A^k + (A+1)^2 \\ &= A^{2k} + 2(A-1)A^k + (A+1)^2 = w^2, \end{aligned}$$

by (5.1).

6. Expansion of $w = \sqrt{[A^k - (A-1)]^2 + 4A}$, $A = 2^d b$, $d \geq 1$, $b \geq 1$ odd, $k \geq 2$. The formulas hold (d and b not both equal 1)

$$(6.1) \quad \begin{aligned} w^2 &= A^{2k} - 2(A-1)A^k + (A+1)^2; \quad [w] = A^k - (A-1); \\ w &= [w] + r, \quad 0 < r < 1. \end{aligned}$$

The reader will easily verify the following

THEOREM 6. *Let $A = 2^d b$, b odd ≥ 1 , $d \geq 2$, be natural numbers, $w^2 = [A^k - (A-1)]^2 + 4A$ squarefree. If $k \geq 3$ is odd, then the expansion of w as a periodic continued fraction has a primitive period of length $5(k-1)$. The expansion has the form*

$$(6.2) \quad \left\{ \begin{aligned} w &= [\overline{b_0, \dots, b_{5s+1}, b_{5s+2}, b_{5s+3}, b_{5s+4}, b_{5s+5}, \dots, b_{1/2(5k-13)}, b_{1/2(5k-11)}}, \\ &\quad \overline{b_{1/2(5k-9)}, b_{1/2(5k-7)}, b_{1/2(5k-5)}, b_{1/2(5k-3)}, b_{1/2(5k-1)}, \dots, 2b_0}] \\ b_{5s+1} &= 2^{d-1}bA^{k-s-2} - 1; \\ b_{5s+2} &= 1; \\ b_{5s+3} &= 1; \\ b_{5s+4} &= 2^{d-1}bA^s - 1; \\ b_{5s+5} &= 2; \quad s = 0, 1, \dots, \frac{1}{2}(k-5); \\ b_{1/2(5k-13)} &= 2^{d-1}bA^{1/2(k-1)} - 1; \quad b_{1/2(5k-11)} = b_{1/2(5k-9)} = 1; \\ b_{1/2(5k-7)} &= 2^{d-1}bA^{1/2(k-3)} - 1; \quad b_{1/2(5k-5)} = 2. \\ b_0 &= A^k - (A-1). \end{aligned} \right.$$

For $k-3$, w has the expansion

$$(6.3) \quad \left\{ \begin{aligned} &\sqrt{[A^3 - (A-1)]^2 + 4A} \\ &= [\overline{A^3 - A + 1, 2^{d-1}bA - 1, 1, 1, 2^{d-1}b - 1, 2, 2^{d-1}b} \\ &\quad \overline{-1, 1, 1, 2^{d-1}bA - 1, 2(A^3 - A + 1)}]. \end{aligned} \right.$$

If $k \geq 4$ is even then the expansion of w as a periodic continued fraction has a primitive period of length $5k - 5$. The expansion has the form

$$(6.4) \quad \begin{cases} w = [\overline{b_0, \dots, b_{5s+1}, b_{5s+2}, b_{5s+3}, b_{5s+4}, b_{5s+5}, \dots}, \\ \quad \overline{b_{1/2(5k-8)}, 1, 1, b_{1/2(5k-8)}, \dots, 2b_0}] \\ b_0 = A^k - (A - 1); \\ b_{5s+1}, b_{5s+2}, b_{5s+3}, b_{5s+4}, b_{5s+5} \text{ as in (6.12); } s = 0, 1, \dots, \frac{1}{2}(k-4), \\ b_{1/2(5k-8)} = 2^{d-1}bA^{1/2(k-2)} - 1. \end{cases}$$

For $k = 2$, the expansion of w is

$$(6.5) \quad \begin{cases} \sqrt{[A^2 - (A - 1)]^2 + 4A} \\ = [A^2 - A + 1, \overline{2^{d-1}b - 1, 1, 1, 2^{d-1}b - 1, 2(A^2 - A + 1)}]. \end{cases}$$

By formula (5.21) we obtain here

$$(6.6) \quad \begin{cases} P_{5s+3}^2 + Q_{5s+3}^2 = w^2, s = \frac{1}{2}(k-2); P_{1/2(5k-4)} = 2A^{1/2k}; \\ A_{1/2(5k-4)} = A^k - (A + 1). \end{cases}$$

Indeed:

$$(2A^{2k})^2 + [A^k - (A + 1)]^2 = A^{2k} - 2(A - 1)A^k + (A + 1)^2 = w^2.$$

7. Expansion of $w = \sqrt{[A^k + (A + 1)]^2 - 4A}$, $A = 2^db$, b odd ≥ 1 ; $d \geq 1$, $k \geq 1$. We can eliminate the case $k = 1$, for then

$$w = \sqrt{4A^2 + 1} = [2A, \bar{4A}].$$

The formulas hold (d and b not both equal 1)

$$(7.1) \quad \begin{cases} w^2 = A^{2k} + 2(A + 1)A^k + (A - 1)^2; [w] = A^k + A; \\ w = A^k + A + r, 0 < r < 1. \end{cases}$$

The reader will easily verify the following

THEOREM 7. *Let $A = 2^db$, b odd ≥ 1 , $d \geq 2$ be natural numbers, $w^2 = [A^k + (A + 1)]^2 - 4A$, $k \geq 1$, w^2 squarefree. The length of the primitive period in the expansion of w as a periodic continued fraction equals $6(k - 1) + 1$. If k is even ≥ 4 , the expansion has the form*

$$(7.2) \quad \begin{cases} w = [\overline{b_0, \dots, b_{6s+1}, b_{6s+2}, b_{6s+3}, b_{6s+4}, b_{6s+5}, b_{6s+6}, \dots}, \\ \quad \overline{b_{3k-5}, b_{3k-4}, 1, 1, b_{3k-4}, b_{3k-5}, \dots, 2b_0}] \\ b_0 = A^k + A, \end{cases}$$

$$\left\{ \begin{array}{l} b_{6s+1} = 1; b_{6s+2} = 2^{d-1}bA^{k-s-2} - 1; b_{6s+3} = b_{6s+4} = 1; \\ b_{6s+5} = 2^{d-1}bA^s - 1; b_{6s+6} = 1; s = 0, 1, \dots, \frac{1}{2}(k-4) \\ b_{3k-5} = 1; b_{3k-4} = 2^{d-1}bA^{1/2(k-2)}. \end{array} \right.$$

If k is odd ≥ 5 , the expansion has the form

$$(7.3) \quad \left\{ \begin{array}{l} w = [b_0, \overline{\dots, b_{6s+1}, b_{6s+2}, b_{6s+3}, b_{6s+4}, b_{6s+5}, b_{6s+6}, \dots}, \\ \overline{b_{3k-8}, b_{3k-7}, b_{3k-6}, b_{3k-5}, b_{3k-4}, 1, 1, b_{3k-4}, b_{3k-5}}, \\ \overline{b_{3k-6}, b_{3k-7}, b_{3k-8}, \dots, 2b_0}] \\ b_0 = A^k + A, \\ b_{6s+1} = 1; b_{6s+2} = 2^{d-1}bA^{k-s-2} - 1; b_{6s+3} = b_{6s+4} = 1; \\ b_{6s+5} = 2^{d-1}bA^s - 1; b_{6s+6} = 1; s = 0, 1, \dots, \frac{1}{2}(k-5). \\ b_{3k-8} = 1; b_{3k-7} = 2^{d-1}bA^{1/2(k-1)} - 1; b_{3k-6} = b_{3k-5} = 1; \\ b_{3k-4} = 2^{d-1}bA^{1/2(k-3)} - 1. \end{array} \right.$$

For $k = 2$ we obtain the expansion

$$(7.4) \quad \left\{ \begin{array}{l} \sqrt{(A^2 + A + 1)^2 - 4A} \\ = [A^2 + A, \overline{1, 2^{d-1}b - 1, 1, 1, 2^{d-1}b - 1, 1, 2(A^2 + A)}] \end{array} \right.$$

For $k = 3$ we obtain the expansion

$$(7.5) \quad \left\{ \begin{array}{l} \sqrt{(A^3 + A + 1)^2 - 4A} \\ = [A^3 + A, \overline{1, 2^{d-1}bA - 1, 1, 1, 2^{d-1}b - 1, 1, 2^{d-1} - 1, 1, 1}, \\ \overline{2^{d-1}bA - 1, 1, 2(A^3 + A)}] \end{array} \right.$$

The formula $P_{r+1}^2 + Q_{r+1}^2 = w^2$ is also verified easily, with $2r + 1 = 6(k-1) + 1$.

For $d = 2, b = 1, A = 4$, formula (7.14) does not hold, for in this case $w^2 = 425$ and is not squarefree.

8. Expansion of $w = \sqrt{[A^k + (A+1)]^2 - 4A}$; $A = 2^db, b$ odd $\geq 1; d \geq 1; k \geq 2$. The formulas hold (for b and d not both equal 1)

$$(8.1) \quad \left\{ \begin{array}{l} w^2 = A^{2k} - 2(A+1)A^k + (A-1)^2; [w] = A^k - A - 2; \\ w = A^k - A - 2 + r; 0 < r < 1. \end{array} \right.$$

The reader will verify easily the following

THEOREM 8. *Let $A = 2^db, b$ odd $\geq 1, d \geq 1$, be natural numbers, $w^2 = [A^k - A + 1]^2 - 4A$ squarefree. If $k \geq 4$ is even, the length of the primitive period in the expansion of w as a periodic continued fraction equals $4k - 2$, and the expansion has the form*

$$(8.2) \quad \left\{ \begin{array}{l} w = [\overline{b_0, b_1, b_2, \dots, b_{4s-1}, b_{4s}, b_{4s+1}, b_{4s+2}, \dots, 2, \dots, b_{2(k+1)-4s}}, \\ \quad \overline{b_2(k+1)-4s-1, b_{2(k+1)-4s-2}, b_{2(k+1)-4s-3}, \dots, b_2, b_1, 2b_0}] ; \\ b_0 = A^k - A - 2; b_1 = 1; b_2 = 2^{d-1}bA^{k-2} - 2 ; \\ b_{4s-1} = 2; b_{4s} - 2^{d-1}bA^{s-1} - 1; b_{4s+1} = 2; b_{4s+2} = 2^{d-1}bA^{k-s-2} - 1 ; \\ s = 1, 2, \dots, \frac{1}{2}(k-2) . \end{array} \right.$$

For $k = 2$, we obtain the expansion

$$(8.3) \quad \left\{ \begin{array}{l} \sqrt{(A^2 - A - 1)^2 - 4A} \\ = [\overline{A^2 - A - 2, 1, 2^{d-1}b - 2, 2, 2^{d-1}b - 2, 1, 2(A^2 - A - 2)}] \end{array} \right. \quad (A > 4) .$$

If $k \geq 5$ is odd, the length of the primitive period in the expansion of w as a periodic continued fraction also equals $4k - 2$, and the expansion has the form

$$(8.4) \quad \left\{ \begin{array}{l} w = [\overline{b_0, b_1, b_2, \dots, b_{4s-1}, b_{4s}, b_{4s+1}, b_{4s+2}, \dots, b_{2k-3}, b_{2k-2}, 2, b_{2k-2}}, \\ \quad \overline{b_{2k-3}, b_2, b_1, 2b_0}] \\ b_0, b_1, b_2, b_{4s-1}, b_{4s}, b_{4s+1}, b_{4s+2} \text{ from (8.13); } s = 1, 2, \dots, \frac{1}{2}(k-3) ; \\ b_{2k-3} = 2; b_{2k-2} = 2^{d-1}bA^{1/2(k-3)} - 1 . \end{array} \right.$$

For $k = 3$, we obtain the expansion

$$(8.5) \quad \left\{ \begin{array}{l} \sqrt{(A^3 - A - 1)^2 - 4A} \\ = [\overline{A^3 - A - 2, 1, 2^{d-1}bA - 2, 2, 2^{d-1}b - 1, 2, 2^{d-1}b - 1, 2,} \\ \quad \overline{2^{d-1}bA - 2, 1, 2(A^3 - A - 2)}] . \end{array} \right.$$

9. Expansion of $w = \sqrt{[(4A)^k + (A-1)]^2 + 4A}$; $A = 2^ab$, b odd; $d \geq 2$. Though, at a first glance, the structure of w looks similar to that of the §§5-8, there are surprising restrictions on the choice of A , and k . The reader will verify easily the following expansion and formulas.

$$(9.1) \quad \left\{ \begin{array}{l} w^2 = (4A)^{2k} + 2(A-1)(4A)^k + (A+1)^2 ; \\ [w] = (4A)^k + A - 1; w = (4A)^k + A - 1 + r; 0 < r < 1 . \end{array} \right.$$

$$(9.2) \quad w = (4A)^k + A - 1 = \frac{1}{x_1}; P_1 = (4A)^{k-1} + A - 1; Q_1 = 4A .$$

$$(9.3) \quad \left\{ \begin{array}{l} \frac{w + (4A)^k + A - 1}{4A} = 2(4A)^{k-1} + \frac{1}{x_2} ; \\ P_2 = (4A)^k - (A - 1); Q_2 = 4(A - 1)(4A)^{k-1} + 1 . \end{array} \right.$$

$$(9.4) \quad \begin{cases} \frac{w + (4A)^k - (A - 1)}{4(A - 1)(4A)^{k-1} + 1} = 2 + \frac{1}{x_3}; \\ P_3 = 4(A - 2)(4A)^{k-1} + (A + 1); Q_3 = 16(4A)^{k-1}. \end{cases}$$

$$(9.5) \quad \begin{cases} \frac{w + 4(A - 2)(4A)^{k-1} + (A + 1)}{16(4A)^{k-1}} 2^{d-1}b - 1 + \frac{1}{x_4}; \\ P_4 = 4(A - 2)(4A)^{k-1} - (A + 1); Q_4 = 4(A - 1)(4A)^{k-1} \\ + [A^2 - (A + 1)]. \end{cases}$$

$$(9.6) \quad \begin{cases} \frac{w + (4A)^k - 8(4A)^{k-1} - (A + 1)}{(4A)^k - 4(4A)^{k-1} + [A^2 - (A + 1)]} = 1 + \frac{1}{x_5}; \\ P_5 = 4(4A)^{k-1} + A^2; Q_5 = 4(A + 1)(4A)^{k-1} - [A^2 + (A + 1)]. \end{cases}$$

$$(9.7) \quad \begin{cases} \frac{w + 4(4A)^{k-1} + A^2}{(4A)^k + 4(4A)^{k-1} - [A^2 + (A + 1)]} = 1 + \frac{1}{x_6}; \\ P_6 = (4A)^k - [2A^2 + (A + 1)]; Q_6 = 4A^2 \end{cases}$$

$$(9.8) \quad \frac{w + (4A)^k - [2A^2 + (A + 1)]}{4A^2} = 8(4A)^{k-2} - 1 + \frac{1}{x_7}.$$

The reader will now verify by induction the following formulas.

$$(9.9) \quad \left\{ \begin{array}{l} \text{(i)} \quad P_{5s+2} = (4A)^k - [2A^{s+1} - (A + 1)]; \\ \quad \quad Q_{5s+2} = 4^{s+1}(A^{s+1} - 1)(4A)^{k-s-1} - [A^{s+1} - (A + 1)]; \\ \quad \quad b_{5s+2} = 2; \\ \text{(ii)} \quad P_{5s+3} = 4^{s+1}(A^{s+1} - 2)(4A)^{k-s-1} + (A + 1); \\ \quad \quad Q_{5s+3} = 4^{s+2}(4A)^{k-s-1}; b_{5s+2} = 2^{d-1}bA^s - 1; \\ \text{(iii)} \quad P_{5s+4} = 4^{s+1}(A^{s+1} - 2)(4A)^{k-s-1} - (A + 1); \\ \quad \quad Q_{5s+4} = 4^{s+1}(A^{s+1} - 1)(4A)^{k-s-1} + [A^{s+2} - (A + 1)]; \\ \quad \quad b_{5s+4} = 1; \\ \text{(iv)} \quad P_{5s+5} = 4^{s+1}(4A)^{k-s-1} + A^{s+2}; \\ \quad \quad Q_{5s+5} = 4^{s+1}(A^{s+1} + 1)(4A)^{k-s-1} - [A^{s+2} + (A + 1)]; \\ \quad \quad b_{5s+5} = 1; \\ \text{(v)} \quad P_{5s+6} = (4A)^k - [2A^{s+2} + (A + 1)]; Q_{5s+6} = 4A^{s+2}; b_{5s+6} \\ \quad \quad = 2 \cdot 4^{s+1}(4A)^{k-s-2} - 1. \\ \quad \quad s = 0, 1, \dots \end{array} \right.$$

Formulas (9.9) are correct for $s = 0$, in virtue of formulas (9.3)–(9.8); then it proved that they are correct if s is replaced by $s + 1$. Comparing successive $P_v - s$, we see that the only possibility of equality is

$$(9.10) \quad (I) \quad P_{5s+2} = P_{5s+3}.$$

This implies

$$(9.11) \quad 4^k = A^{2(s+1)-k}; A = 2^d, b = 1.$$

From (9.11) we obtain $2k = d[2(s+1) - k]$, hence

$$(9.12) \quad k(d+2) = 2d(s+1).$$

Solving the Diophantine equation (9.12), we obtain all possible solutions as follows.

$$(9.13) \quad \left\{ \begin{array}{l} \text{(a)} \quad k = 2ud; s = u(d+2) - 1; u = 1, 2, \dots; \\ \text{(b)} \quad k = 2ut; d = 2t; s = (t+1)u - 1; t, u = 1, 2, \dots; \\ \text{(c)} \quad k = (2u+1)d; s = \frac{1}{2}(d+2)(2u+1) - 1; d \equiv 0 \pmod{2}; \\ \text{(d)} \quad 2k = (2u+1)d; 4s = (d+2)(2u+1) - 4; d \equiv 2 \pmod{4}. \end{array} \right.$$

The reader should note the following procedure: after k and d have been chosen from (9.13), (a)-(d), s is a function of k, d ; for constant k_0, d_0 we shall denote

$$(9.14) \quad s_0 = F(k_0, d_0).$$

The length of the primitive period in the expansion of w for any choice of (k, d) from (9.13) then becomes $m = 10s_0 + 4$.

Comparing successive $Q_v - s$, we see that the only possibility of equality is

$$(9.15) \quad \text{(II)} \quad Q_{5s+4} = Q_{5s+5}.$$

This implies

$$(9.16) \quad 4^k = A^{2s+3-k}; A = 2^d; b = 1.$$

From (9.16) we obtain $2k = d(2s+3-1)$, hence

$$(9.17) \quad k(d+2) = d(2s+3).$$

Solving the Diophantine equation (9.17), we obtain all possible solutions as follows

$$(9.18) \quad \left\{ \begin{array}{l} \text{(a)} \quad k = ud, 2s+3 = u(2+d), u, d \equiv 1(2); \\ \text{(b)} \quad k = ut; d = 2t; 2s+3 = u(t+1); u \equiv 1(2); t \equiv 0(2). \end{array} \right.$$

We again denote $s_0 = F(k_0, d_0)$, for any choice of fixed k and d from (9.18). The length of the primitive period in the expansion of w for any choice of (k, d) from (9.18) then becomes $m = 10s_0 + g$. For $d = 1, (b = 1)$, we obtain $w = \sqrt{(8^k + 1)^2 + 8}$, which is easily expanded and is left to the reader. We can now state.

THEOREM 9. *Let $A = 2^d$, $d \geq 2$, $w^2 = [(4A)^k + (A - 1)]^2 + 4A$ square-free. If (k_0, d_0, s_0) is any solution vector of the Diophantine equation (9.12), given by the value Table (9.13), (a)–(d), then the primitive period in the expansion of w as a periodic continued fraction has length $m = 10s_0 + 4$, ($s_0 \geq 1$), and the form*

$$(9.19) \quad \begin{cases} w = [\overline{b_0, b_1, \dots, b_{5s+2}, b_{5s+3}, b_{5s+4}, b_{5s+5}, b_{5s+6}, \dots, 2, \dots, b_{5(s_0-s)+1}}, \\ \quad \overline{b_{5(s_0-s)}, b_{5(s_0-s)-1}, b_{5(s_0-s)-2}, b_{5(s_0-s)-3}, \dots, b_1, 2b_0}] \\ b_0 = (4A)^{k_0} + A - 1; b_1 = 2(4A)^{k_0-1}; \\ b_{5s+2} = 2; b_{5s+3} = 2^{d_0-1}A^s - 1; b_{5s+4} = b_{5s+5} = 1; \\ b_{5s+6} = 2 \cdot 4^{s+1}(4A)^{k_0-s-2} - 1. \quad s = 0, 1, \dots, s_0 - 1. \end{cases}$$

If (k_0, s_0, d_0) is any solution vector of the Diophantine equation (9.17), given by the value table (9.18), (a), (b), then the primitive period in the expansion of w as a periodic continued fraction has length $m = 10s_0 + 9$, and the form

$$(9.20) \quad \begin{cases} w = [\overline{b_0, b_1, \dots, b_{5s+2}, b_{5s+3}, b_{5s+4}, b_{5s+5}, b_{5s+6}, \dots, 2, 2^{d_0-1}bA^{s_0} - 1}, \\ \quad \overline{1, 1, 2^{d_0-1}bA^{s_0} - 1, 2, \dots, b_1, 2b_0}] \\ \text{where the } b_0, b_1, b_{5s+2}, \dots, b_{5s+6} \text{ are the same as in (9.19),} \\ s = 0, 1, \dots, s_0 - 1. \end{cases}$$

10. Expansion of $w = \sqrt{[(4A)^k + A + 1]^2 - 4A}$; $A = 2^db$, b odd, $d \geq 2$. The reader will verify easily the following formulas and expansions

$$(10.1) \quad \begin{cases} w^2 = (4A)^{2k} + 2(A + 1)(4A)^k + (A - 1)^2; [w] = (4A)^k + A; \\ w = (4A)^k + A + r; 0 < r < 1. \end{cases}$$

$$(10.2) \quad w = (4A)^k + A + \frac{1}{x_1}; P_1 = (4A)^k + A; Q_1 = 2(4A)^k - (2A - 1).$$

$$(10.3) \quad \frac{w + (4A)^k + A}{2(4A)^k - (2A - 1)} = 1 + \frac{1}{x_2}; P_2 = (4A)^k - (3A - 1); Q_2 = 4A.$$

$$(10.4) \quad \begin{cases} \frac{w + (4A)^k - (3A - 1)}{4A} = 2(4A)^{k-1} - 1 + \frac{1}{x_3}; \\ P_3 = (4A)^k - (A + 1); Q_3 = 4(A + 1)(4A)^{k-1} - 1. \end{cases}$$

$$(10.5) \quad \begin{cases} \frac{w + (4A)^k - (A + 1)}{(4A)^k + 4(4A)^{k-1} - 1} = 1 + \frac{1}{x_4}; \\ P_4 = 4(4A)^{k-1} + A; Q_4 = 4(A - 1)(4A)^{k-1} + (2A - 1). \end{cases}$$

$$(10.6) \quad \begin{cases} \frac{w + 4(4A)^{k-1} + A}{(4A)^k - 4(4A)^{k-1} + 2A - 1} = 1 + \frac{1}{x_5}; \\ P_5 = 4(A - 2)(4A)^{k-1} + A - 1; Q_5 = 16(4A)^{k-1}. \end{cases}$$

$$(10.7) \quad \begin{cases} \frac{w + 4(A - 2)(4A)^{k-1} + A - 1}{16(4A)^{k-1}} = 2^{d-1}b - 1 + \frac{1}{x_6}; \\ P_6 = 4(A - 2)(4A)^{k-1} - (A - 1); Q_6 = 4(A - 1)(4A)^{k-1} \\ + [A^2 - (A - 1)]. \end{cases}$$

$$(10.8) \quad \begin{cases} \frac{w + (4A)^k - 8(4A)^{k-1} - (A - 1)}{(4A)^k - 4(4A)^{k-1} + [A^2 - (A - 1)]} = 1 + \frac{1}{x_7} \\ P_7 = 4(4A)^{k-1} + A^2; Q_7 = 4(A + 1)(4A)^{k-1} - [A^2 + (A - 1)]. \end{cases}$$

$$(10.9) \quad \begin{cases} \frac{w + 4(4A)^{k-1} + A^2}{(4A)^k + 4(4A)^{k-1} - [A^2 + (A - 1)]} = 1 + \frac{1}{x_8} \\ P_8 = (4A)^k - [2A^2 + (A - 1)]; Q_8 = 4A^2. \end{cases}$$

We now prove by induction the following formulas as before; they are correct for the parameter $s = 0$, and it is then proved that they are correct for substituting s by $s + 1$:

$$(10.10) \quad \left\{ \begin{array}{ll} \text{(i)} & P_{6s+2} = (4A)^k - [2A^{s+1} + (A - 1)]; Q_{6s+2} = 4A^{s+1}; b_{6s+2} \\ & = 2 \cdot 4^s(4A)^{k-s-1} - 1; \\ \text{(ii)} & P_{6s+3} = (4A)^k - [2A^{s+1} - (A - 1)]; \\ & Q_{6s+3} = 4^{s+1}(A^{s+1} + 1)(4A)^{k-s-1} - [A^{s+1} - (A - 1)]; \\ & b_{6s+3} = 1; \\ \text{(iii)} & P_{6s+4} = 4^{s+1}(4A)^{k-s-1} + A^{s+1}; \\ & Q_{6s+4} = 4^{s+1}(A^{s+1} - 1)(4A)^{k-s-1} + [A^{s+1} + (A - 1)]; \\ & b_{6s+4} = 1; \\ \text{(iv)} & P_{6s+5} = 4^{s+1}(A^{s+2} - 2)(4A)^{k-s-1} + A - 1; \\ & Q_{6s+5} = 4^{s+2}(4A)^{k-s-1}; b_{6s+5} = 2^{d-1}bA^s - 1; \\ \text{(v)} & P_{6s+6} = 4^{s+1}(A^{s+1} - 2)(4A)^{k-s-1} - (A - 1); \\ & Q_{6s+6} = 4^{s+1}(A^{s+1} - 1)(4A)^{k-s-1} + [A^{s+2} - (A - 1)]; \\ & b_{6s+6} = 1; \\ \text{(vi)} & P_{6s+7} = 4^{s+1}(4A)^{k-s-1} + A^{s+2}; \\ & Q_{6s+7} = 4^{s+1}(A^{s+1} + 1)(4A)^{k-s-1} - [A^{s+2} + (A - 1)]; \\ & b_{6s+7} = 1; s = 0, 1, \dots \end{array} \right.$$

Comparing successive $P_v - s$ and $Q_v - s$, we obtain the cases

$$(10.11) \quad \text{(I)} \quad Q_{6s+3} = Q_{6s+4}.$$

This implies

$$(10.12) \quad 4^k = A^{2(s+1)-k}; A = 2^d; b = 1.$$

From (10.12) we obtain $2k = d[2(s + 1) - k]$,

$$(10.13) \quad (d + 2)k = 2d(s + 1) .$$

(10.13) is the same equation as (9.12) and its solutions are given by (9.13). If (k_0, d_0, s_0) is a solution vector of (10.13), then the length of the primitive period in the expansion of w equals $m = 10s_0 + 7$.

$$(10.14) \quad (II) \quad Q_{6s+6} = Q_{6s+7} .$$

This implies

$$(10.15) \quad 4^k = A^{2s+3-k}, A = 2^d; b = 1 .$$

From (10.15) we obtain $2k = d(2s + 3 - k)$,

$$(10.16) \quad k(d + 2) = d(2s + 3) .$$

Equation (10.16) is equation (9.17), and its solutions are given by (9.18), (a), (b). If (k_0, d_0, s_0) is a solution vector of (10.16), then the length of the primitive period in the expansion of w equals $m = 10s_0 + 13$. The restriction on $d \geq 2$ results from the value of $b_5 = 2^{d-1} - 1$ in (10.7). If $d = 1$, we would have $b_5 = 0$, which is impossible. The case $d = 1, A = 2$, yields $w = \sqrt{(8^k + 3)^2 - 8}$, and the expansion of this w is left to the reader. We can now state.

THEOREM 10. *Let $A = 2^d, d \geq 2, w^2 = [(4A)^k + (A + 1)]^2 - 4A$ squarefree. If (k_0, d_0, s_0) is any solution vector of the Diophantine equation (10.13), given by the value table (9.13), (a)-(d), then the primitive period in the expansion of w as a periodic continued fraction has length $m = 10s_0 + 7$, and the form*

$$(10.17) \quad \left\{ \begin{array}{l} w = [b_0, \overline{b_1, \dots, b_{6s+2}, b_{6s+3}, b_{6s+4}, b_{6s+5}, b_{6s+6}, b_{6s+7}, \dots}, \\ \quad \overline{2 \cdot 4^{s_0}(4A)^{k_0-s_0-1} - 1, 1, 1, 2 \cdot 4^{s_0}(4A)^{k_0-s_0-1}, \dots, b_1, 2b_0}] \\ b_0 = (4A)^{k_0} + A; b_1 = 1, \\ b_{6s+2} = 2 \cdot 4^s(4A)^{k_0-s-1} - 1; b_{6s+3} = b_{6s+4} = 1; \\ b_{6s+5} = 2^{d_0-1}A^s - 1; b_{6s+6} = b_{6s+7} = 1. \end{array} \right.$$

If (k_0, d_0, s_0) is any solution vector of the Diophantine equation (10.16), given by the value table (9.18), (a), (b), then the primitive period in the expansion of w as a periodic continued fraction has length $m = 10s_0 + 13$, and the form

$$(10.18) \quad \left\{ \begin{array}{l} w = [b_0, \overline{b_1, \dots, b_{6s+2}, b_{6s+3}, b_{6s+4}, b_{6s+5}, b_{6s+6}, b_{6s+7}, \dots}, \\ \quad \overline{2 \cdot 4^{s_0}(4A)^{k_0-s_0-1} - 1, 1, 1, 2^{d_0-1}A^{s_0} - 1, 1, 1, 2^{d_0-1}A^{s_0} - 1, \\ \quad 1, 1, 2 \cdot 4^{s_0}(4A)^{k_0-s_0-1} - 1, \dots, b_1, 2b_0}] \\ b_0, b_1, b_{6s+2}, \dots, b_{6s+7} \text{ from (10.17), } s = 0, 1, \dots, s_0 - 1. \end{array} \right.$$

11. Expansion of $w = \sqrt{[A^k + (A - 1)]^2 + 4A}; A = 2^d b + 1; d \geq$

1; b odd. This expansion is remarkable in the sense that the cycles of the period are of length eleven; so are the cycles in the next two sections, while in the last section the length of the cycle is even twelve.

We obtain the formulas

$$(11.1) \quad \begin{aligned} w^2 &= A^{2k} + 2(A-1)A^k + (A+1)^2; [w] = A^k + A - 1; w \\ &= A^k + A - 1 + r; 0 < r < 1. \end{aligned}$$

The reader will easily verify the following:

THEOREM 11. *Let $A = 2^d b + 1$, $d \geq 1$, b odd, $w^2 = [A^k + (A-1)]^2 + 4A$ squarefree; let further $k \equiv 4 \pmod{6}$, $k \geq 4$, $s_0 = (1/6)(k-4)$. Then the length of the primitive period in the expansion of w as a periodic continued fraction equals $1/3(11k-14)$, and the expansion has the form, for $k \geq 10$,*

$$(11.2) \quad \left\{ \begin{aligned} w &= [\overline{b_0, b_1, \dots, b_{11s+2}, b_{11s+3}, \dots, b_{11s+12}, \dots, b_{11s_0+2}, b_{11s_0+3}, b_{11s_0+4}}, \\ &\quad \overline{2, b_{11s_0+4}, b_{11s_0+3}, b_{11s_0+2}, \dots, b_1, 2b_0}] \\ b_0 &= A^k + A - 1; b_1 = 2^{d-1}bA_{k-2}; b_{11s_0+2} = 1; b_{11s_0+3} = 2A^{1/2(k-2)} - 1; \\ b_{11s_0+4} &= 2^{d-1}bA_{1/2(k-2)}; b_{11s+2} = 1; b_{11s+3} = 2A^{3s+1} - 1; \\ b_{11s+4} &= 2^{d-1}bA_{k-3(s+1)}; b_{11s+5} = 2; b_{11s+6} = 2^{d-1}bA_{3s+1}; \\ b_{11s+7} &= 2A^{k-3(s+1)} - 1; b_{11s+8} = 1; b_{11s+9} = 2^{d-1}bA_{3s+2}; \\ b_{11s+10} &= b_{11s+1} - 1; b_{11s+12} = 2^{d-1}bA_{k-3(s+1)-2} - 2; \\ s &= 0, 1, \dots, s_0 - 1; A_u = (A-1)^{-1}(A^{u+1} - 1); u = 0, 1, \dots; \\ A_0 &= 1. \end{aligned} \right.$$

If $k = 4$, the expansion of w has the form

$$(11.3) \quad \left\{ \begin{aligned} \sqrt{[A^4 + (A-1)]^2 + 4A} &= [A^4 + A - 1, \overline{2^{d-1}b(A^2 + A + 1), 1}, \\ 2A - 1, 2^{d-1}b, 2, 2^{d-1}b, 2A - 1, 1, 2^{d-1}b(A^2 + A + 1), \\ &\quad \overline{2(A^4 + A - 1)}] \end{aligned} \right.$$

If $k = 3$, the expansion has the form

$$(11.4) \quad \left\{ \begin{aligned} \sqrt{[A^3 + (A-1)]^2 + 4A} &= [b_0, b_1, \dots, b_{12}, b_{13}, b_{14}, b_{12}, \dots, b_1, 2b_0] \\ b_0 &= A^3 + A - 1; b_1 = 2^{d-1}b(A+1); b_2 = 1; b_3 = 2A - 1; \\ b_4 &= 2^{d-1}b; b_5 = 2; b_6 = 2^{d-1}b(A+1); b_7 = b_8 = 1; \\ b_9 &= 2^{d-1}b(A^2 + A) + A - 1; b_{10} = 2A^2 + 1; b_{11} = 1; \\ b_{12} &= 2^{d-1}b - 1; b_{13} = b_{14} = 1. \end{aligned} \right.$$

The length of the primitive period in the expansion of w for $k = 3$ equals 27. Let further be $k \equiv 1 \pmod{6}$, $k \geq 7$, $s_0 = (1/6)(k-7)$. Then the length of the period in the expansion of w as a periodic continued fraction equals $1/3(11k-14)$, and the expansion has the form, for $k \geq 13$,

$$(11.5) \left\{ \begin{array}{l} w = [b_0, \overline{b_1, \dots, b_{11s+2}, b_{11s+3}, \dots, b_{11s+12}, \dots, b_{11s_0+2}, b_{11s_0+3}, \dots,} \\ \quad \overline{b_{11s_0+9}, 1, 1, b_{11s_0+9}, b_{11s_0+8}, \dots, b_{11s_0+2}, \dots, b_1, 2b_0}] \\ b_0 = A^{k+}A - 1; b_1 = 2^{d-1}bA_{k-2}; b_{11s_0+2} = 1; b_{11s_0+3} = 2A^{1/2(k-s)} - 1; \\ b_{11s_0+4} = 2^{d-1}bA_{1/2(k+1)}; b_{11s_0+5} = 2; b_{11s_0+6} = 2^{d-1}bA_{1/2(k-5)}; \\ b_{11s_0+7} = 2A^{1/2(k+1)} - 1; b_{11s_0+8} = 1; \\ b_{11s_0+9} = 2^{d-1}bA_{1/2(k-3)} - 1; \\ b_{11s+2}, b_{11s+3}, \dots, b_{11s+12} \text{ as in (11.19); } s = 0, 1, s_0 - 1. \end{array} \right.$$

It is left to the reader to find the expansion of w for the cases $k = 5, 6, 7$.

12. Expansion of $w = \sqrt{[A^k - (A-1)]^2 + 4A}$; $A = 2^db + 1$; $d \parallel v$; $1; b$ odd. We use the notation, as before,

$$(12.1) \quad \begin{aligned} A^{u+1} - 1 &= (A-1)A_u; u = 0, 1, \dots; A_0 \\ &= 1; \frac{1}{2}(A^{u+1} - 1) = 2^{d-1}bA_u. \end{aligned}$$

The following formulas hold:

$$(12.2) \quad \begin{cases} w^2 = A^{2k} - 2(A-1)A^k + (A+1)^2; [w] = A^k - A + 1; \\ w = A^k - A + 1 + r, 0 < r < 1. \end{cases}$$

The reader will easily verify the following

THEOREM 12. *Let $A = 2^db + 1$, $d \geq 1$, b odd, $w^2 = [A^k - (A-1)]^2 + 4A$ squarefree; let further $k \equiv 1 \pmod{6}$, $k \geq 7$, $s_0 = (1/6)(k-7)$. Then the length of the primitive period in the expansion of w as a periodic continued fraction equals $11/3(k-1)$, and the expansion has the form, for $k \geq 13$,*

$$(12.3) \left\{ \begin{array}{l} w = [b_0, \overline{b_1, \dots, b_{11s+1}, b_{11s+2}, \dots, b_{11s+11}, \dots, b_{11s_0+1}, b_{11s_0+2}, \dots,} \\ \quad \overline{b_{11s_0+10}, 2, b_{11s_0+10}, b_{11s_0+9}, \dots, b_{11s_0+1}, \dots, 2b_0}] \\ b_{11s+1} = 2^{d-1}bA_{k-3s-2}; b_{11s+2} = 2A^{3s+1} - 1; \\ b_{11s+3} = b_{11s+5} = b_{11s+6} = b_{11s+8} = 1; b_{11s+4} = 2^{d-1}bA_{k-3(s+1)} - 1; \\ b_{11s+7} = 2^{d-1}bA_{3s+1} - 1; b_{11s+9} = 2A^{k-3(s+1)} - 1; \\ b_{11s+10} = 2^{d-1}bA_{3s+2}; b_0 = A^k - A + 1; b_{11s_0+1} = 2^{d-1}bA_{1/2(k+3)}; \\ b_{11s_0+2} = 2A^{1/2(k-5)} - 1; b_{11s_0+5} = b_{11s_0+6} = b_{11s_0+8} = 1; \\ b_{11s_0+4} = 2^{d-1}bA_{1/2(k+1)} - 1; b_{11s_0+7} = 2^{d-1}bA_{1/2(k-5)} - 1; \\ b_{11s_0+9} = 2A^{1/2(k+1)} - 1; b_{11s_0+10} = 2^{d-1}bA_{1/2(k-3)}; b_{11s+11} = 2. \\ s = 0, 1, \dots, s_0 - 1; A_u = (A-1)^{-1}(A^{u+1} - 1), u = 0, 1, \dots; \\ A_0 = 1. \end{array} \right.$$

For $k = 7$, the expansion has the form

$$(12.4) \quad \left\{ \begin{array}{l} \sqrt{[A^7 - (A-1)]^2 + 4A} = [b_0, \overline{b_1, b_2, \dots, b_{10}, 2, b_{10}, \dots, b_2, b_1, 2b_0}], \\ b_0 = A^7 - A + 1; b_1 = 2^{d-1}bA_5; b_2 = 2A - 1; \\ b_3 = b_5 = b_6 = b_8 = 1; b_4 = 2^{d-1}bA_4 - 1; b_7 = 2^{d-1}bA_1 - 1; \\ b_9 = 2A^4 - 1; b_{10} = 2^{d-1}bA_2. \end{array} \right.$$

Let further be $k \equiv 4 \pmod{6}$; $s_0 = (1/6)(k-4)$; $k \geq 4$. Then the length of the primitive period in the expansion of w equals $11/3(k-1)$, and the expansion has the form, for $k \geq 10$,

$$(12.5) \quad \left\{ \begin{array}{l} w = [b_0, \overline{b_{11s+1}, b_{11s+2}, \dots, b_{11s+11}, \dots, b_{11s_0+1}, b_{11s_0+2}, \\ b_{11s_0+3}, b_{11s_0+4}, 1, 1, b_{11s_0+4}, b_{11s_0+3}, b_{11s_0+2}, b_{11s_0+1}, \dots, 2b_0}] \\ b_0, b_{11s+1}, b_{11s+2}, \dots, b_{11s+11} \text{ as in (12.18);} \\ b_{11s_0+1} = 2^{d-1}bA_{1/2k}; b_{11s_0+2} = 2A^{1/2(k-2)} - 1; \\ b_{11s_0+3} = 1; b_{11s_0+4} = 2^{d-1}bA_{1/2(k-2)} - 1. \\ s = 0, 1, \dots, s_0 - 1; A_u \text{ as in (12.18).} \end{array} \right.$$

For $k = 4$, the expansion has the form

$$(12.6) \quad \left\{ \begin{array}{l} \sqrt{[A^4 - (A-1)]^2 + 4A} = [b_0, \overline{b_1, b_2, b_3, b_4, 1, 1, b_4, b_3, b_2, b_1, 2b_0}] \\ b_0 = A^4 - A + 1; b_1 = 2^{d-1}bA_2; b_2 = 2A - 1; b_3 = 1; \\ b_4 = 2^{d-1}bA_1 - 1. \end{array} \right.$$

13. Expansion of $w\sqrt{(A^k + A + 1)^2 - 4A}$; $A = 2^db + 1$; $d \geq 1$; b odd.

THEOREM 13. *Let $A = 2^db + 1$, $d \geq 1$, b odd, $w^2 = (A^k + A + 1)^2 - 4A$ squarefree; let further $k \equiv 4 \pmod{6}$, $k \geq 4$; $s_0 = 1/6(k-4)$. Then the length of the primitive period in the expansion of w as a continued fraction equals $1/3(10k-7)$, and the expansion has the form, for $k \geq 10$,*

$$(13.1) \quad \left\{ \begin{array}{l} w = [b_0, \overline{b_1, \dots, b_{10s+2}, b_{10s+3}, \dots, b_{10s+11}, \dots, b_{10s_0+2}, b_{10s_0+3}, b_{10s_0+4}, \\ 1, 1, b_{10s_0+4}, b_{10s_0+3}, b_{10s_0+2}, \dots, b_1, 2b_0}] \\ b_0 = A^k + A; b_1 = 1; b_{11s_0+2} = 2^{d-1}bA_{1/2k}; b_{10s_0+3} = 2A^{1/2(k-2)}; \\ b_{10s_0+4} = 2^{d-1}bA_{1/2(k-2)}; b_{10s+2} = 2^{d-1}bA_{k-3s-2}; \\ b_{10s+3} = 2A^{3s+1}; b_{10s+4} = 2^{d-1}bA_{k-3(s+1)}; \\ b_{10s+5} = b_{10s+6} = b_{10s+10} = b_{10s+11} = 1; b_{10s+7} = 2^{d-1}bA_{3s+1}; \\ b_{10s+8} = 2A^{k-3(s+1)}; b_{10s+9} = 2^{d-1}bA_{3s+2}; \\ s = 0, 1, \dots, s_0 - 1; A^{u+1} - 1 = (A-1)A_u; u = 1, 2, \dots. \end{array} \right.$$

For $k = 4$, the expansion has the form

$$(13.2) \left\{ \begin{aligned} & \sqrt{(A^4 + A + 1)^2 - 4A} = [b_0, \overline{b_1, b_2, b_3, b_4, 1, 1, b_4, b_3, b_2, b_1, 2b_0}] , \\ & b_0 = A^4 + A; b_1 = 1; b_2 = 2^{d-1}bA_2; b_3 = 2A; b_5 = 2^{d-1}b(A + 1) . \end{aligned} \right.$$

Let further $k \equiv 1 \pmod{6}$, $k \geq 7$, $s_0 = (1/6)(k - 7)$. Then the length of the primitive period in the expansion of w as a periodic continued fraction equals $1/3(10k - 7)$, and the expansion has the form, for $k \geq 13$,

$$(13.3) \left\{ \begin{aligned} & w = [b_0, \overline{b_1, \dots, b_{10s_0+2}, b_{10s_0+3}, \dots, b_{10s_0+11}, \dots, b_{10s_0+2}, b_{10s_0+3}, \dots,} \\ & \quad \overline{b_{10s_0+9}, 1, 1, b_{10s_0+9}, b_{10s_0+8}, \dots, b_{10s_0+2}, \dots, b_1, 2b_0}] \\ & b_0, b_1, b_{10s_0+2}, b_{10s_0+3}, \dots, b_{10s_0+11} \text{ as in (13.17), } s = 0, 1, \dots, \\ & s_0 - 1; b_{10s_0+2} = 2^{d-1}bA_{1/2(k+3)}; b_{10s_0+3} = 2A^{1/2(k-5)}; \\ & b_{10s_0+4} = 2^{d-1}bA_{1/2(k+1)}; b_{10s_0+5} = b_{10s_0+6} = 1; \\ & b_{10s_0+7} = 2^{d-1}bA_{1/2(k-5)}; b_{10s_0+8} = 2A^{1/1(k+1)}; \\ & b_{10s_0+9} = 2^{d-1}bA_{1/2(k-3)} . \end{aligned} \right.$$

For $k = 7$, the expansion of w has the form

$$(13.4) \left\{ \begin{aligned} & \sqrt{(A^7 + A + 1)^2 - 4A} = [b_0, b_1, \overline{b_2, \dots, b_9, 1, 1, b_9, \dots, b_2, b_1, 2b_0}] \\ & b_0 = A^7 + A; b_1 = 1; b_2 = 2^{d-1}bA_5; b_3 = 2A; b_4 = 2^{d-1}bA_4; \\ & b_5 = b_6 = 1; b_7 = 2^{d-1}bA_1; b_8 = 2A^4; b_9 = 2^{d-1}bA_2 . \end{aligned} \right.$$

14. Expansion of $w = \sqrt{[A^k - (A + 1)]^2 - 4A}$; $A = 2^db + 1$; $d \geq 1$; $b \geq 1$ odd. This case is the most interesting of all treated in this part of the paper, since the length of the cycle is greater than any previous one, namely 12. With the previous notation, $A^{u+1} - 1 = (A - 1)A_u$, $u = 0, 1, \dots$; $A_0 = 1$, the reader will easily verify the following expansions and formulas.

$$(14.1) \left\{ \begin{aligned} & w^2 = A^{2k} - 2(A + 1)A^k + (A - 1)^2; [w] = A^k - A - 2; \\ & w = A^k - A - 2 + r, 0 < r < 1 . \end{aligned} \right.$$

$$(14.2) \quad w = A^k - A - 2 + \frac{1}{x_1}; P_1 = A^k - (A + 2); Q_1 = 2A^k + 3(2A + 1) .$$

$$(14.3) \quad \frac{w + A^k - A - 2}{2A^k - 6A - 3} = 1 + \frac{1}{x_2}; P_2 = A^k - (5A + 1); Q_2 = 4A .$$

$$(14.4) \left\{ \begin{aligned} & \frac{w + A^k - 5A - 1}{4A} = 2^{d-1}bA_{k-2} - 2 + \frac{1}{x_3}; P_3 = A^k - (5A - 1); \\ & Q_3 = (2A - 1)A^{k-1} - 2(3A - 1) . \end{aligned} \right.$$

$$(14.5) \left\{ \begin{aligned} & \frac{w + A^k - 5A + 1}{2A^k - A^{k-1} - 6A + 2} = 1 + \frac{1}{x_4}; P_4 = (A - 1)A^{k-1} - (A - 1); \\ & Q_4 = A^{k-1} . \end{aligned} \right.$$

$$(14.6) \quad \frac{w + A^k - A^{k-1} - A + 1}{A^{k-1}} 2 = (A - 1) + \frac{1}{x_5};$$

$$P_5 = (A - 1)A^{k-1} + (A - 1); Q_5 = (2A - 1)A^{k-1} - 2[2A^2 - (A - 1)].$$

$$(14.7) \quad \begin{cases} \frac{w + A^k - A^{k-1} + A - 1}{2A^k - A^{k-1} - 2[2A^2 - (A - 1)]} = 1 + \frac{1}{x_6}; \\ P_6 = A^k - [4A^2 - (A - 1)]; Q_6 = 4A^2. \end{cases}$$

$$(14.8) \quad \begin{cases} \frac{w + A^k - 4A^2 + A - 1}{4A^2} = 2^{d-1}bA_{k-3} - 1 + \frac{1}{x_7}; \\ P_7 = A^k - [2A^2 + (A - 1)]; Q_7 = (A^2 - 1)A^{k-2} - [A^2 + (A - 1)]; \end{cases}$$

$$(14.9) \quad \begin{cases} \frac{w + A^k - 2A^2 - A + 1}{A^k - A^{k-2} - A^2 - A + 1} = 2 + \frac{1}{x_8}; \\ P_8 = (A^2 - 2)A^{k-2} - (A - 1); Q_8 = 4A^{k-2}; \end{cases}$$

$$(14.10) \quad \begin{cases} \frac{w + A^k - 2A^{k-2} - (A - 1)}{4A^{k-2}} \\ = 2^{d-1}bA_1 - 1 + \frac{1}{x_9}; P_9 = (A^2 - 4)A^{k-2} + (A - 1); Q_9 \\ = 2(A^2 - 2)A^{k-2} - [A^3 - 2(A - 1)]; \end{cases}$$

$$(14.11) \quad \begin{aligned} & \frac{w + A^k - 4A^{k-2} + (A - 1)}{2A^k - 4A^{k-2} - A^3 + 2(A - 1)} \\ & = 1 + \frac{1}{x_{10}}; P_{10} = A^k - [A^3 - (A - 1)]; Q_{10} = A^3; \end{aligned}$$

$$(14.12) \quad \begin{cases} \frac{w + A^k - A^3 + A - 1}{A^3} = 2A^{k-3} - 2 + \frac{1}{x_{11}}; \\ P_{11} = A^k - [A^3 + (A - 1)]; \\ Q_{11} = 2(A^3 - 2)A^{k-3} - [A^3 + 2(A - 1)]; \end{cases}$$

$$(14.13) \quad \begin{cases} \frac{w + A^k - A^3 - (A - 1)}{2A^k - 4A^{k-3} - A^3 - 2(A - 1)} = 1 + \frac{1}{x_{12}}; \\ P_{12} = (A^3 - 4)A^{k-3} - (A - 1); Q_{12} = 4A^{k-3}; \end{cases}$$

$$(14.14) \quad \begin{cases} \frac{w + A^k - 4A^{k-3} - (A - 1)}{4A^{k-3}} = 2^{d-1}bA_2 - 1 + \frac{1}{x_{13}}; \\ P_{13} = (A^3 - 2)A^{k-3} + (A - 1); \\ Q_{13} = (A^3 - 1)A^{k-3} - [A^4 - (A - 1)]; \end{cases}$$

$$(14.15) \quad \begin{cases} \frac{w + A^k - 2A^{k-3} + (A - 1)}{A^k - A^{k-3} - A^4 + (A - 1)} = 2 + \frac{1}{x_{14}}; \\ P_{14} = A^k - [2A^4 - (A - 1)]; Q_{14} = 4A^4; \end{cases}$$

$$(14.16) \quad \frac{w + A^k - 2A^4 + (A - 1)}{4A^4} 2d^{-1}bA_{k-5} - 1 + \frac{1}{x_{15}}.$$

We now prove the formulas:

$$(14.17) \quad \left\{ \begin{array}{l} P_{12s+3} = A^k - [4A^{3s+1} + (A - 1)]; \\ Q_{12s+3} = (2A^{3s+1} - 1)A^{k-3s-1} - 2[2A^{3s+1} + (A - 1)]; b_{12s+3} = 1; \\ P_{12s+4} = (A^{3s+1} - 1)A^{k-3s-1} - (A - 1); \\ Q_{12s+4} = A^{k-3s-1}; b_{12s+4} = 2A^{3s+1} - 2; \\ P_{12s+5} = (A^{3s+1} - 1)A^{k-3s-1} + (A - 1); \\ Q_{12s+5} = (2A^{3s+1} - 1)A^{k-3s-1} - 2[2A^{3s+2} - (A - 1)]; b_{12s+5} = 1; \\ P_{12s+6} = A^k - [4A^{3s+2} - (A - 1)]; \\ Q_{12s+6} = 4A^{3s+2}; b_{12s+6} = 2^{d-1}bA_{k-3(s+1)} - 1; \\ P_{12s+7} = A^k - [2A^{3s+2} + (A - 1)]; \\ Q_{12s+7} = (A^{3s+2} - 1)A^{k-3s-2} - [A^{3s+2} + (A - 1)]; b_{12s+7} = 2; \\ P_{12s+8} = (A^{3s+2} - 2)A^{k-3s-2} - (A - 1); \\ Q_{12s+8} = 4A^{k-3s-2}; b_{12s+8} = 2^{d-1}bA_{3s+1} - 1; \\ P_{12s+9} = (A^{3s+2} - 4)A^{k-3s-2} + (A - 1); \\ Q_{12s+9} = 2(A^{3s+2} - 2)A^{k-3s-2} - [A^{3(s+1)} - 2(A - 1)]; b_{12s+9} = 1; \\ P_{12s+10} = A^k - [A^{3(s+1)} - (A - 1)]; \\ Q_{12s+10} = A^{3(s+1)}; b_{12s+10} = 2A^{k-3(s+1)} - 2; \\ P_{12s+11}^s = A^k - [A^{3(s+1)} + (A - 1)]; \\ Q_{12s+11} = 2(A^{3(s+1)} - 2)A^{k-3(s+1)} - [A^{3(s+1)} + 2(A - 1)]; \\ b_{12s+11} = 1; \\ P_{12s+12} = (A^{3(s+1)} - 4)A^{k-3(s+1)} - (A - 1); \\ Q_{12s+12} = 4A^{k-3(s+1)}; b_{12s+12} = 2^{d-1}bA_{3s+2} - 1; \\ P_{12s+13} = (A^{3(s+1)} - 2)A^{k-3(s+1)} + (A - 1); \\ Q_{12s+13} = (A^{3(s+1)} - 1)A^{k-3(s+1)} - [A^{3(s+1)+1} - (A - 1)]; \\ b_{12s+13} = 2; \\ P_{12s+14} = A^k - [2A^{3(s+1)+1} - (A - 1)]; Q_{12s+14} = 4A^{3(s+1)+1}; \\ b_{12s+14} = 2^{d-1}bA_{k-3(s+1)-2} - 1. \\ s = 0, 1, \dots \end{array} \right.$$

Comparison of successive $P_v - s$ and $Q_v - s$ shows that equality takes place in the following cases:

$$(14.18) \quad \left\{ \begin{array}{l} P_{12s+7} = P_{12s+8} \text{ implying } k = 2(3s + 2); k \equiv 4(\text{mod } 6); \\ s_0 = \frac{1}{6}(k - 4). \end{array} \right.$$

$$(14.19) \quad \begin{cases} P_{12s+13} = P_{12s+14} \text{ implying } k = 6(s+1) + 1; k \equiv 1 \pmod{6}; \\ s_0 = \frac{1}{6}(k-7). \end{cases}$$

The length of the primitive period in the expansion of w equals, in case (14.18), $m = 4 + 24 \cdot 1/6(k-4) + 8 + 2 = 4k - 2$; in case (14.19) the length of the primitive period equals $m = 4 + 24 \cdot (1/6)(k-7) + 20 + 2 = 4k - 2$. We can now state.

THEOREM 14. *Let $A = 2^d b + 1$, $d \geq 1$, b odd, $w^2 = [A^k - (A+1)]^2 - 4A$ squarefree; let further $k \equiv 4 \pmod{6}$, $k \geq 4$, $s_0 = (1/6)(k-4)$. Then the length of the primitive period in the expansion of w as a periodic continued fraction equals $4k - 2$, and the expansion has the form $k \geq 10$,*

$$(14.20) \quad \left\{ \begin{array}{l} w = [b_0, \overline{b_1, b_2, \dots, b_{12s+3}, b_{12s+4}, \dots, b_{12s+14}, \dots, b_{12s_0+3}, b_{12s_0+4}}, \\ \quad \overline{b_{12s_0+5}, b_{12s_0+6}, 2, b_{12s_0+6}, b_{12s_0+5}, b_{12s_0+4}, b_{12s_0+3}, \dots, b_2, b_1, 2b_0}] \\ b_0 = A^k - A - 2; b_1 = 1; b_2 = 2^{d-1}bA_{k-2} - 2; b_{12s_0+3} = b_{12s_0+5} = 1; \\ b_{12s_0+4} = 2A^{1/2(k-2)} - 2; b_{12s_0+6} = 2^{d-1}bA_{1/2(k-2)}; \\ b_{12s+3} = b_{12s+5} = \frac{1}{2}b_{12s+7} = b_{12s+9} = b_{12s+11} = \frac{1}{2}b_{12s+13} = 1; \\ b_{12s+4} = 2A^{3s+1} - 2; b_{12s+6} = 2^{d-1}bA_{k-3(s+1)} - 1; \\ b_{12s+8} = 2^{d-1}bA_{3s+1} - 1; b_{12s+10} = 2A^{k-3(s+1)} - 2; \\ b_{12s+12} = 2^{d-1}bA_{3s+2} - 1; b_{12s+14} = 2^{d-1}bA_{k-3s-5} - 1. \\ s = 0, 1, \dots, s_0 - 1; A^{u+1} - 1 = (A-1)A_u; u = 0, 1, \dots; \\ A_0 = 1. \end{array} \right.$$

For $k = 4$, we obtain the expansion

$$(14.21) \quad \left\{ \begin{array}{l} \sqrt{[A^4 - (A+1)]^2 - 4A} \\ = [A^4 - A - 2, \overline{1, 2^{d-1}bA_2 - 2, 1, 2A - 2, 1, 2^{d-1}bA_1, 2,} \\ \quad \overline{2^{d-1}bA_1, 1, 2A - 2, 1, 2^{d-1}bA_2 - 2, 1, 2(A^4 - A - 2)}]. \end{array} \right.$$

Let further $k \equiv 1 \pmod{6}$; $k \geq 7$, $s_0 = 1/6(k-7)$. Then the length of the period in the expansion of w as a periodic continued fraction equals $4k - 2$, and the expansion has the form

$$\left\{ \begin{array}{l} w = [b_0, \overline{b_1, b_2, \dots, b_{12s+3}, b_{12s+4}, \dots, b_{12s+14}, \dots, b_{12s_0+3}}, \\ \quad \overline{b_{12s_0+4}, \dots, b_{12s_0+12}, 2, b_{12s_0+12}, \dots, b_{12s_0+4}, b_{12s_0+3}, \dots, b_2, b_1, 2b_0}] \\ b_0, b_1, b_2, b_{12s+3}, b_{12s+4}, \dots, b_{12s+14} \text{ as in (14.20), } s = 0, 1, \dots, \\ \quad s_0 - 1. \\ b_{12s_0+3} = b_{12s_0+5} = \frac{1}{2}b_{12s_0+7} = b_{12s_0+9} = b_{12s_0+11} = 1; \end{array} \right.$$

$$\begin{cases} b_{12s_0+4} = 2A^{1/2(k-5)} - 2; b_{12s_0+6} = 2^{d-1}bA_{1/2(k+1)} - 1; \\ b_{12s_0+8} = 2^{d-1}bA_{1/2(k-5)} - 1; b_{12s_0+10} = 2A^{1/2(k+1)} - 2; \\ b_{12s_0+12} = 2^{d-1}bA_{1/2(k-3)} - 1. \end{cases}$$

The reader will have no difficulty to formulate the theorem for the case $k = 7$.

REFERENCES

1. L. Bernstein, *The Jacobi-Perron Algorithm, Its Theory and Application*, Lecture Notes in Mathematics 207, Springer-Verlag, 1971, 1-160.
2. ———, *Periodicity and Units in Quadratic Number Fields*, Submitted.
3. G. Degert, *Ueber die Bestimmung der Grundeinheit gewisser real-quadratischer Zahlkoerper*, Hamburger Math. Abh., Bd. XXII, (1958), 92-97.
4. Th. Muir, (i) *The expression of a quadratic surd as a continued fraction*, Glasgow, 1874; (ii) *A New Special Class of Determinants*, Proc. Ed. 8, 1874.
5. O. Perron, *Die Lehre von den Kettenbruechen*, Zweite verbesserte Auflage, Chelsea Publ. Co., New York, I-XII + 1-524.
6. Y. Yamamoto, *Real quadratic number fields with large fundamental units*, Osaka J. Math., **8**, no. 2 (1971) 261-271.
7. H. Zassenhaus, *On the units of orders*, J. Algebra, **20** (1972), 368-395.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Ralph Artino, <i>Gevrey classes and hypoelliptic boundary value problems</i>	1
B. Aupetit, <i>Caractérisation spectrale des algèbres de Banach commutatives</i> . . .	23
Leon Bernstein, <i>Fundamental units and cycles in the period of real quadratic number fields. I</i>	37
Leon Bernstein, <i>Fundamental units and cycles in the period of real quadratic number fields. II</i>	63
Robert F. Brown, <i>Fixed points of automorphisms of compact Lie groups</i>	79
Thomas Ashland Chapman, <i>Concordances of noncompact Hilbert cube manifolds</i>	89
William C. Connett, V and Alan Schwartz, <i>Weak type multipliers for Hankel transforms</i>	125
John Wayne Davenport, <i>Multipliers on a Banach algebra with a bounded approximate identity</i>	131
Gustave Adam Efroymson, <i>Substitution in Nash functions</i>	137
John Sollion Hsia, <i>Representations by spinor genera</i>	147
William George Kitto and Daniel Eliot Wulbert, <i>Korovkin approximations in L_p-spaces</i>	153
Eric P. Kronstadt, <i>Interpolating sequences for functions satisfying a Lipschitz condition</i>	169
Gary Douglas Jones and Samuel Murray Rankin, III, <i>Oscillation properties of certain self-adjoint differential equations of the fourth order</i>	179
Takaši Kusano and Hiroshi Onose, <i>Nonoscillation theorems for differential equations with deviating argument</i>	185
David C. Lantz, <i>Preservation of local properties and chain conditions in commutative group rings</i>	193
Charles W. Neville, <i>Banach spaces with a restricted Hahn-Banach extension property</i>	201
Norman Oler, <i>Spaces of discrete subsets of a locally compact group</i>	213
Robert Olin, <i>Functional relationships between a subnormal operator and its minimal normal extension</i>	221
Thomas Thornton Read, <i>Bounds and quantitative comparison theorems for nonoscillatory second order differential equations</i>	231
Robert Horace Redfield, <i>Archimedean and basic elements in completely distributive lattice-ordered groups</i>	247
Jeffery William Sanders, <i>Weighted Sidon sets</i>	255
Aaron R. Todd, <i>Continuous linear images of pseudo-complete linear topological spaces</i>	281
J. Jerry Uhl, Jr., <i>Norm attaining operators on $L^1[0, 1]$ and the Radon-Nikodým property</i>	293
William Jennings Wickless, <i>Abelian groups in which every endomorphism is a left multiplication</i>	301