

# Pacific Journal of Mathematics

**SETS OF UNIQUENESS AND MULTIPLICITY FOR  $L^p$**

ANNA MARIA MANTERO

# SETS OF UNIQUENESS AND MULTIPLICITY FOR $L^p$

ANNA MARIA MANTERO

**In the first part of this paper, it is proved that: if  $1 < q < p \leq 2$  and  $G$  is a nondiscrete, locally compact abelian (LCA) group with character group  $\Gamma$ , there exists a subset of positive measure  $E \subset G$  which is a set of uniqueness for  $L^q(\Gamma)$  and, at the same time, a set of multiplicity for  $L^p(\Gamma)$ .**

**This is followed by some results of the same type concerning the spaces  $L^{p,\alpha}(\Gamma)$ ,  $\alpha \neq 0$ , when  $G$  is the Cantor group.**

I. I. Hirschman, Jr., and Y. Katznelson prove all the results of this paper in the case when  $G = T$  (the circle group) and  $\Gamma = Z$  (the integer) (see [5]). They also prove results concerning the spaces  $L^{p,\alpha}(Z)$ . The core of the present work consists in proving our theorem for the case in which  $G$  is a compact group the elements of which have bounded order. To obtain the theorem for a general LCA group, we use the latter result, the theorem of Hirschman and Katznelson, and the structure theory for LCA groups. The existence of sets of uniqueness for  $L^p(\Gamma)$ ,  $1 \leq p < 2$ , which are of positive measure (and therefore are sets of multiplicity for  $L^2(\Gamma)$ ) was proved by Y. Katznelson ([6]; [7], p. 101), for the case  $G = T$ , and by A. Figà-Talamanca and G. I. Gaudry for the general case (see [2]). The results contained in this paper appeared in part in our thesis for the "laurea" in Mathematics at the University of Genova. This thesis was prepared under the guidance of Prof. A. Figà-Talamanca, to whom we are grateful for advice and assistance.

1. Preliminaries. Let  $G$  be a nondiscrete LCA group and  $\Gamma$  its character group. If  $f$  is an integrable function on  $G$ ,

$$\hat{f}(\gamma) = \int_G f(x) \cdot \gamma(-x) dx \quad \text{for any } \gamma \in \Gamma$$

denotes the Fourier transform of  $f$ .

Let us define now a uniqueness set and a multiplicity set for  $L^p(\Gamma)$  as follows:

**DEFINITION.** Let  $E$  be a measurable subset of  $G$ ; then  $E$  is called a *set of uniqueness* for  $L^p(\Gamma)$  (or set of  $p$ -uniqueness), if no non-zero integrable function  $f$ , carried by  $E$ , satisfies the condition  $\hat{f} \in L^p(\Gamma)$ .

A subset which is not a set of uniqueness is called a *set of multiplicity* for  $L^p(\Gamma)$  (or set of  $p$ -multiplicity).

Throughout this paper we indicate the Haar measure of a set  $S$ , by  $m(S)$ .

2. The main result. We can now state the first result of this paper.

**THEOREM 1.** *For every real numbers  $p, q$  such that  $1 < q < p \leq 2$  there exists a subset  $E \subset G$ , of positive measure, which is a set of  $q$ -uniqueness and  $p$ -multiplicity.*

*Proof.* From a theorem of structure [8, p. 40] we know that every LCA group has an open subgroup which is the direct sum of a compact group  $K$  and a Euclidean space  $\mathbf{R}^n (n \geq 0)$ . Thus it is sufficient to prove the theorem for groups of the form

$$G = K \oplus \mathbf{R}^n \quad \text{for } n \geq 0.$$

We must distinguish several cases:

*Case (1).*  $G$  is a compact group with Haar measure normalized ( $n = 0$ ). In this case the proof of the theorem can be obtained from the following two lemmas:

**LEMMA 1.** *Given a real numbers  $\varepsilon, 0 < \varepsilon < 1$ , and  $p, q \in (1, 2]$  such that  $q < p$ , we can find a measurable set  $E \subset G$  and two functions  $F$  and  $\phi$ , defined on  $G$ , satisfying the following properties:*

- (a)  $m(E) \geq 1 - \varepsilon$
- (b)  $F(x) = 1$  for any  $x \in E$
- (c)  $\|\hat{F}\|_{q'} \leq \varepsilon$  if  $1/q + 1/q' = 1$
- (d)  $\phi(x) \geq 0$  for any  $x \in G$
- (e)  $m\{x \in G: \phi(x) = 1\} \geq 1 - 2\varepsilon$
- (f)  $\log \|\hat{\phi}\|_p \leq \varepsilon$
- (g)  $\{x \in G: \phi(x) \neq 0\} \subset E$ .

*Proof of the Lemma 1.* We consider three cases according to the order of the elements of  $\Gamma$ .

*Case 1 (a).*  $\Gamma$  is a torsion group of bounded order. Let  $\mathcal{A}$  be an independent set of generators for  $\Gamma$ . Let  $\{\Gamma_n\}_{n=1}^\infty$  be a sequence of subgroups of  $\Gamma$ , all having the same finite order  $r$  and which are generated by elements of  $\mathcal{A}$  in such a way that, if  $(\Gamma_1, \Gamma_2, \dots)$  denotes the group generated by  $\Gamma_1, \Gamma_2, \dots$ , then

$$\Gamma_{n+1} \cap (\Gamma_1, \Gamma_2, \dots, \Gamma_n) = \{0\} \quad \text{for } n = 1, 2, \dots.$$

Now, for any (large) positive integer  $N$ , we can define—on the group

$G$ —two sets of  $N$  functions,  $F_1, F_2, \dots, F_N$  and  $\phi_1, \phi_2, \dots, \phi_N$  in the following way:

$$F_i(x) = 1 - \sum_{\gamma \in \Gamma_i} \gamma(x) \quad \text{for } i = 1, \dots, N;$$

$$\phi_i(x) = 1 - \frac{1}{r} \sum_{\gamma \in \Gamma_i} \gamma(x) \quad \text{for } i = 1, \dots, N.$$

Since  $F_i$  and  $\phi_i$  ( $1 \leq i \leq N$ ) are trigonometric polynomials, we can trivially deduce that:

$$(I) \quad \|\hat{F}_i\|_{q'} \leq r^{1/q'} \quad \text{for any } i = 1, \dots, N.$$

$$(II) \quad \log \|\hat{\phi}_i\|_p \leq r^{1-p} \quad \text{for any } i = 1, \dots, N.$$

Moreover:

$$(III) \quad m\{x \in G: F_i(x) = 1\} = 1 - 1/r \quad \text{for } i = 1, \dots, N.$$

$$(IV) \quad m\{x \in G: \phi_i(x) = 1\} = 1 - 1/r \quad \text{for } i = 1, \dots, N.$$

Now, let us define, for every integer  $N$ :

$$E = \bigcap_{i=1}^N \{x \in G: F_i(x) = 1\},$$

$$F(x) = \frac{1}{N} \sum_{i=1}^N F_i(x) \quad \text{for } x \in G$$

$$\phi(x) = \prod_{i=1}^N \phi_i(x) \quad \text{for } x \in G.$$

It follows that:

$$(i) \quad m(E) = m\{x \in G: F(x) = 1\} \geq 1 - N/r.$$

Because the  $F_i$  are supported by disjoint subgroups and  $\hat{F}_i(0) = 0$  ( $1 \leq i \leq N$ ), then (I) yields:

$$(ii) \quad \|\hat{F}\|_{q'} = (1/N) (\sum_{i=1}^N \|\hat{F}_i\|_{q'}^{q'})^{1/q'} \leq N^{-1+1/q'} \cdot r^{1/q'}.$$

Since  $\|\hat{\phi}_i\|_p = \|\hat{\phi}_j\|_p$  for any  $i, j \in \{1, 2, \dots, N\}$ , and by the definition of  $\Gamma_n$  there exists at most one way to write  $\gamma \in \Gamma$  as the sum of elements  $\gamma_1 \in \Gamma_1, \gamma_2 \in \Gamma_2, \dots, \gamma_N \in \Gamma_N$ , then

$$(iii) \quad \log \|\hat{\phi}\|_p = \log (\prod_{i=1}^N \|\hat{\phi}_i\|_p) \leq N \cdot r^{1-p}.$$

Since  $p > q$ , it is possible to choose a positive integer  $N$  and a large positive integer  $r$  such that the conditions

$$\begin{cases} \|\hat{F}\|_{q'} \leq N^{-1/q} \cdot r^{1/q'} \leq \varepsilon \\ \log \|\hat{\phi}\|_p \leq N \cdot r^{1-p} \leq \varepsilon, \end{cases}$$

are simultaneously satisfied.

Since  $\Gamma$  is of bounded order (and therefore it has infinitely many elements of the same order) we can choose  $r$  arbitrarily large so that it is the common order of infinitely many disjoint subgroups  $\Gamma_1, \Gamma_2, \dots$ .

To complete the proof of the lemma in Case 1 (a), we note that:

$$(i) \quad \phi(x) = \prod_{i=1}^N (1 - \chi_{M_i})(x) \geq 0 \quad \text{for } x \in G$$

where—for every  $i = 1, 2, \dots, N$ — $M_i = \{x \in G: \gamma(x) = 1, \gamma \in \Gamma_i\}$  and

$\chi_{M_i}$  is the characteristic function of  $M_i$ .

$$(ii) \quad m(E) \geq 1 - N/r \geq 1 - N \cdot r^{1-p} \geq 1 - \varepsilon$$

because  $r > 1$  and  $(p - 1) \leq 1$ .

$$(iii) \quad m\{x \in G: \phi(x) = 1\} = m(E) \geq 1 - \varepsilon.$$

*Case 1 (b).*  $\Gamma$  contains an element of infinite order. Let  $\gamma'$  be an element of  $\Gamma$  such that  $n \cdot \gamma' \neq 0$ , for every  $n \in \mathbb{N}$ . We call  $\Gamma'$  the subgroup of  $\Gamma$  generated by  $\gamma'$ . Thus  $\Gamma'$  is isomorphic to  $\mathbb{Z}$  and its character group  $\hat{\Gamma}'$  is a compact group isomorphic to  $G/M$ , where

$$M = \{x \in G: \gamma'(x) = 1\}$$

is the annihilator of  $\Gamma'$  [8, p. 35]. But  $\hat{\Gamma}'$  is also isomorphic to  $T$ . Thus we can consider—given a positive real number  $\varepsilon$ —the construction of the functions which satisfy the Lemma 1 in the case  $G = T$ , given by Hirschman and Katznelson in [5, p. 226]. We call these functions  $f$  and  $\varphi$ . Since  $f$  and  $\varphi$  belong to  $L^2(T)$ , we can write:

$$f(\vartheta) = \sum_{k \in \mathbb{Z}} a_k \cdot e^{ik\vartheta} \quad \text{for any } \vartheta \in (0, 2\pi]$$

$$\varphi(\vartheta) = \sum_{k \in \mathbb{Z}} b_k \cdot e^{ik\vartheta} \quad \text{for any } \vartheta \in (0, 2\pi] .$$

Now we define:

$$F(x) = \sum_{k \in \mathbb{Z}} a_k \cdot \gamma'(x)^k \quad \text{for } x \in G$$

and

$$\phi(x) = \sum_{k \in \mathbb{Z}} b_k \cdot \gamma'(x)^k \quad \text{for } x \in G .$$

From the definition of  $M$  it follows that such functions are constant on the sets  $x + M$ , where  $x \in G$ , and therefore we can consider  $F$  and  $\phi$  as defined on  $G/M$ . Hence for every  $x \in G/M$  we have

$$F(x) = \sum_{\gamma \in \Gamma'} \hat{F}(\gamma) \cdot x(\gamma) = \sum_{k \in \mathbb{Z}} \hat{f}(k) e^{ik\vartheta} = f(\vartheta) ,$$

where  $\vartheta \in (0, 2\pi]$  is such that  $e^{i\vartheta} \in T$  is the image of  $x \in G/M$  under the isomorphism mapping  $G/M$  onto  $T$ .

Similarly:

$$\phi(x) = \varphi(\vartheta) .$$

It follows that [4, p. 91, vol. II]:

$$m\{x \in G: F(x) = 1\} = m\{\vartheta \in (0, 2\pi]: f(\vartheta) = 1\}$$

and

$$m\{x \in G: \phi(x) = 1\} = m\{\vartheta \in (0, 2\pi]: \varphi(\vartheta) = 1\}.$$

Thus, by observing that:

$$\|\hat{F}\|_{q'} = \|\hat{f}\|_{q'}; \quad \|\hat{\phi}\|_p = \|\hat{\varphi}\|_p;$$

and recalling [5], the proof of the lemma is easily completed in this case.

*Case 1 (c).*  $\Gamma$  is a torsion group of infinite order. Suppose that  $\gamma' \in \Gamma$  has order  $n$ . We call  $\Gamma'$  the cyclic subgroup of  $\Gamma$  generated by  $\gamma'$ . The discrete group  $\Gamma'$  is isomorphic to  $\mathbb{Z}(n)$ . Its character group  $\hat{\Gamma}'$  is a cyclic compact subgroup of  $T$  which is also isomorphic to  $G/M$ , if  $M$  is the annihilator of  $\Gamma'$ .

Similarly to the Case 1 (b), if

$$f(\vartheta) = \sum_{k \in \mathbb{Z}} A_k \cdot e^{ik\vartheta} \quad \text{for } \vartheta \in (0, 2\pi]$$

and

$$\varphi(\vartheta) = \sum_{k \in \mathbb{Z}} B_k \cdot e^{ik\vartheta} \quad \text{for } \vartheta \in (0, 2\pi]$$

are the functions satisfying the Lemma 1 when  $G = T$  and  $\varepsilon/2$  replaces  $\varepsilon$ , we pose:

$$\begin{aligned} F(x) &= \sum_{k=-\lfloor (n-1)/2 \rfloor}^{\lfloor n/2 \rfloor} \left( \sum_{i=-\infty}^{+\infty} A_{k+ni} \right) \cdot \gamma'(x)^k & \text{for } x \in G, \\ \phi(x) &= \sum_{k=-\lfloor (n-1)/2 \rfloor}^{\lfloor n/2 \rfloor} \left( \sum_{i=-\infty}^{+\infty} B_{k+ni} \right) \cdot \gamma'(x)^k & \text{for } x \in G. \end{aligned}$$

Since these functions are constant on  $x + M \subset G (x \in G)$ , we can consider them as defined on  $G/M$ . Thus, for every  $x \in G/M$ , if  $e^{i\vartheta}$  is the image of  $x$  under the isomorphism mapping  $G/M$  onto  $T$ ,

$$F(x) = \sum_{\gamma \in \Gamma'} \hat{F}(\gamma)x(\gamma) = \sum_{m \in \mathbb{Z}} \hat{f}(m)e^{im\vartheta} = f(\vartheta).$$

Using the same reasoning we get, for any  $x \in G/M$  and for any  $\vartheta$  defined as above,

$$\phi(x) = \varphi(\vartheta).$$

In order that the functions  $F$  and  $\phi$  satisfy the Lemma 1, they must verify these non-obvious conditions:

- (1)  $\|\hat{F}\|_{q'} \leq \varepsilon$ ,
- (2)  $\log \|\hat{\phi}\|_p \leq \varepsilon$ ,
- (3)  $m\{x \in G: F(x) = 1\} \geq 1 - \varepsilon$ ,
- (4)  $m\{x \in G: \phi(x) = 1\} \geq 1 - 2\varepsilon$ .

We will show that these conditions are satisfied if the order of

the element  $\gamma'$  is large enough. Indeed, given  $\hat{f} \in L^1(Z)$  and  $\varepsilon > 0$  there exists  $n'$  so large that for  $n \geq n'$  we have:

$$\sum_{|k| > \lfloor (n-1)/2 \rfloor} |A_k| \leq \varepsilon/2.$$

Consequently, by Minkowski's inequality and recalling that if  $1 \leq s$  then  $L^1(Z) \subseteq L^s(Z)$  we have:

$$\begin{aligned} \|\hat{F}\|_{q'} &= \left( \sum_k \left| A_k + \sum_{i \neq 0} A_{k+ni} \right|^{q'} \right)^{1/q'} \\ &\leq \left( \sum_k |A_k|^{q'} \right)^{1/q'} + \left( \sum_k \left| \sum_{i \neq 0} A_{k+ni} \right|^{q'} \right)^{1/q'} \\ &\leq \|\hat{f}\|_{q'} + \sum_k \left| \sum_{i \neq 0} A_{k+ni} \right| \\ &\leq \|\hat{f}\|_{q'} + \sum_{\substack{k \neq 0 \\ i \neq 0}} |A_{k+ni}| \\ &\leq \|\hat{f}\|_{q'} + \varepsilon/2 \leq \varepsilon. \end{aligned}$$

Similarly, given  $\hat{\varphi} \in L^1(Z)$  and  $\varepsilon > 0$ , there exists  $n''$  so large that for  $n \geq n''$  we have

$$\sum_{|k| > \lfloor (n-1)/2 \rfloor} |B_k| \leq \|\hat{\varphi}\|_p \cdot \varepsilon/2.$$

Thus if  $\gamma'$  has order  $n \geq n''$ ,

$$\begin{aligned} \lg \|\hat{\varphi}\|_p &= \lg \left( \sum_k \left| B_k + \sum_{i \neq 0} B_{k+ni} \right|^p \right)^{1/p} \\ &\leq \lg \left( \left( \sum_k |B_k|^p \right)^{1/p} + \left( \sum_k \left| \sum_{i \neq 0} B_{k+ni} \right|^p \right)^{1/p} \right) \\ &\leq \lg \left( \|\hat{\varphi}\|_p + \sum_{\substack{k \neq 0 \\ i \neq 0}} |B_{k+ni}| \right) \\ &\leq \lg (\|\hat{\varphi}\|_p \cdot (1 + \varepsilon/2)) \leq \varepsilon. \end{aligned}$$

Finally, if  $\eta$  is the measure of the subset of  $(0, 2\pi]$  where  $f(\vartheta) \neq 1$  and the order  $n$  of  $\gamma'$  satisfies

$$n > (\eta/\varepsilon)^{1/2}$$

we have:

$$\begin{aligned} m\{x \in G: F(x) = 1\} &\geq 1 - \varepsilon, \\ m\{x \in G: \phi(x) = 1\} &\geq 1 - 2\varepsilon. \end{aligned}$$

Therefore, to fulfill conditions (1) to (4) and to prove the lemma it is enough to choose  $\gamma' \in \Gamma$  with order  $n$  satisfying

$$n > \max(n', n'', (\eta/\varepsilon)^{1/2}).$$

**LEMMA 2.** *Given a sequence of positive real numbers  $\{\varepsilon_k\}_{k=1}^\infty$  such that  $2 \sum_k \varepsilon_k < 1$ , it is possible—for every  $k \in N$  to find  $E^{(k)}, F^{(k)}, \phi^{(k)}$  satisfying Lemma 1 with  $\varepsilon = \varepsilon_k$  such that for all  $N \in N$*

$$(1) \quad \left\| \left( \prod_{k=1}^N \phi^{(k)} \right)^\wedge \right\|_p \leq \exp \left[ \sum_{k=1}^{N-1} \varepsilon_k \right] \cdot \prod_{k=1}^N \|\hat{\phi}^{(k)}\|_p \leq e .$$

*Proof.* Inequality (1) can be proved easily enough by induction. Suppose that (1) holds for  $N = 1, 2, \dots, R-1$ . We want to show that:

$$(2) \quad \left\| \left( \prod_{k=1}^R \phi^{(k)} \right)^\wedge \right\|_p \leq \exp(\varepsilon_{R-1}) \left\| \left( \prod_{k=1}^{R-1} \phi^{(k)} \right)^\wedge \right\|_p \cdot \|\hat{\phi}^{(R)}\|_p .$$

We must distinguish two cases:

(1°)  $G$  is as in the Case 1(a) of Lemma 1. Then we must choose  $\phi^{(R)}$  such that

$$\text{supp } \hat{\phi}^{(R)} \cap (\text{supp } \hat{\phi}^{(1)}, \dots, \text{supp } \hat{\phi}^{(R-1)}) = \{0\} .$$

(2°)  $G$  is as the Cases 1 (b) and 1 (c) of Lemma 1. Then we define  $\phi^{(R)}$  starting from the function  $\varphi_R$  which satisfy the analogous inequality in [5]. Using condition (f) of Lemma 1, we see that (1) is true in both cases.

*Conclusion of the proof of Theorem 1 in Case (1).* Choose a sequence of real numbers  $\varepsilon_k > 0, k = 1, 2, \dots$ , such that  $2 \sum_k \varepsilon_k < 1$ . We define

$$E = \bigcap_{k=1}^\infty E^{(k)} .$$

Since we assume that  $2 \sum_k \varepsilon_k < 1$ , then clearly:

$$m(E) = 1 - \sum_{k=1}^\infty \varepsilon_k > 0 .$$

We assert now that  $E$  is a set of uniqueness for  $L^q(I)$ . Let  $g$  be an integrable function with support included in  $E$  for which  $\|\hat{g}\|_q < +\infty$ . Since  $F^{(k)}(x) = 1$  for  $x \in E, k \in N$  we have, for  $\gamma$  fixed:

$$\begin{aligned} \hat{g}(\gamma) &= \int_G g(x) \gamma(-x) dx = \int_G F^{(k)}(x) g(x) \gamma(-x) dx \\ &= F^{(k)} * \hat{g}(\gamma) , \end{aligned}$$

and, from Hölder's inequality:

$$|\hat{g}(\gamma)| \leq \|\hat{g}\|_q \cdot \|\hat{F}^{(k)}\|_{q'} \leq \|\hat{g}\|_q \cdot \varepsilon_k .$$

Letting  $k \rightarrow \infty$  we have that  $\hat{g}(\gamma) = 0$  for all  $\gamma \in I$  and thus that  $g \equiv 0$ .

We also assert that  $E$  is a set of multiplicity for  $L^p(I)$ . Let  $\phi(N, x) = \prod_{k=1}^N \phi^{(k)}(x)$  for  $x \in G$ . Since  $L^p(I)$  is weakly boundedly



compact for  $p \in (1, 2]$ , it follows from (1) that there exists  $\lambda(\gamma) \in L^p(\Gamma)$  which is a weak limit point of  $\hat{\phi}(N, \gamma)$ ,  $N = 1, 2, \dots$ . By (e) of Lemma 1,

$$m\{x \in G: \phi(N, x) = 1\} \geq 1 - 2 \sum_k \varepsilon_k > 0,$$

and since, for every  $x \in G$ ,  $\phi(N, x) \geq 0$ , by (d) of Lemma 1, we see that:

$$\hat{\phi}(N, 0) \geq 1 - 2 \sum \varepsilon_k, \quad N = 1, 2, \dots,$$

and hence

$$\lambda(0) = 1 - 2 \sum \varepsilon_k > 0.$$

Therefore  $\lambda(\gamma) \neq 0$ .

Let  $h$  be a function defined on  $G$  such that  $\hat{h}(\gamma) = \lambda(\gamma)$ . Moreover,  $h \in L^1(G)$  since it has support of finite measure and, by the Hausdorff-Young theorem,  $h \in L^{p'}(G)$ . Then  $h$  is a nonzero function supported in the intersection of the supports of  $\phi(N, x)$ ,  $x \in G$  and  $N \in N$ , and therefore  $(\text{supp } h) \subset E$ .

*Case (2).*  $G$  is a Euclidean space  $R^n$  for some  $n > 0$ . We can identify  $T$  with the real interval  $[-1/2, 1/2]$  and therefore we can identify  $T^n$  to be the vector product  $[-1/2, 1/2]^n$ . Choose now a sequence of real numbers  $\{\varepsilon_k\}_{k=1}^\infty$  such that  $4 \sum_k \varepsilon_k < 1$ .

For every  $\varepsilon_k$  we consider the corresponding function  $f_k$  which satisfies Lemmas 1 and 2 in the case of the torus. Thus we can extend periodically the function  $f_k$  to all the real line and we call  $f_k$  also this (periodic) extension. Now we extend the domain of definition of  $f_k$  from  $R$  to  $R^n$  as follows:

$$f_k^\dagger(x_1, \dots, x_n) = f_k(x_1)$$

and we choose a positive, continuous function  $P$  defined on  $R^n$  which satisfies:

- (1)  $0 \leq P(x_1, \dots, x_n) \leq 1$  for any  $(x_1, \dots, x_n) \in R^n$
- (2)  $P(x_1, \dots, x_n) = 1$  on  $[-1/2 + 1/8, 1/2 - 1/8]^n$
- (3)  $P(x_1, \dots, x_n) = 0$  out of  $[-1/2, 1/2]^n$
- (4)  $\hat{P} \in C^\infty(\hat{R}^n)$ .

By defining  $F_k$  as

$$F_k(x_1, \dots, x_n) = f_k^\dagger(x_1, \dots, x_n) \cdot P(x_1, \dots, x_n)$$

we have a function which is obtained from  $f_k^\dagger$  by dropping to zero its value outside of  $[-1/2, 1/2]^n$  and which equals  $f_k^\dagger$  on a smaller cube.

Now we show that the range of the operator  $T$ ,

$$T: \hat{g} \longrightarrow (g^{\sharp} P)^{\wedge}$$

where we define  $g^{\sharp}(x_1, \dots, x_n) = g(x_1)$  is contained in  $L^s(\hat{R}^n)$ ,  $1 \leq s \leq \infty$ , for all continuous  $g$  on  $T$  with  $\hat{g} \in L^1(Z)$ , that is for all  $g \in A(T)$ . Moreover, we show that  $T$  is continuous from  $L^1(Z)$ -equipped with  $L^s(Z)$ -norm into  $L^s(\hat{R}^n)$ ,  $1 \leq s \leq \infty$ .

To keep the notation simple we only prove the case when  $n = 1$ . We also make use of the following formula for  $\alpha \in \hat{R}$ :

$$(gP)^{\wedge}(\alpha) = \sum_{n=-\infty}^{+\infty} \hat{g}(n) \hat{P}(\alpha - n).$$

By the Riesz-Thorin theorem, it is sufficient to prove that  $T$  is continuous for  $p = 1$  and  $p = \infty$ .

If  $p = 1$ ,

$$\begin{aligned} \|Tg\|_1 &= \int_{\mathbf{R}} |(gP)^{\wedge}(\alpha)| d\alpha \leq \sum_{n=-\infty}^{+\infty} |\hat{g}(n)| \left( \sup_{\alpha} \int_{\mathbf{R}} |\hat{P}(\alpha - n)| d\alpha \right) \\ &\leq \|\hat{g}\|_1 \cdot \|\hat{P}\|_{L^1(\hat{R})}. \end{aligned}$$

If  $p = \infty$ ,

$$\begin{aligned} \|Tg\|_{\infty} &= \sup_{\alpha} |(gP)^{\wedge}(\alpha)| \leq \sup_{\alpha} \sum_{n=-\infty}^{+\infty} |\hat{g}(n)| \cdot |\hat{P}(\alpha - n)| \\ &\leq \|\hat{g}\|_{\infty} \cdot \sup_{0 \leq \alpha < 1} \sum_n |\hat{P}(\alpha - n)|; \end{aligned}$$

the sum  $\sum_n |\hat{P}(\alpha - n)|$  is bounded independently of  $\alpha$  and therefore

$$\|Tg\|_{\infty} \leq C_1 \cdot \|\hat{g}\|_{\infty},$$

where  $C_1$  depends only on  $P$ .

Therefore,

$$\|\hat{F}^{(k)}\|_{L^{q'}(\hat{R})} \leq C_2 \cdot \|\hat{f}_k\|_{1q},$$

where  $C_2$  depends only on  $P$ .

We claim that

$$E = \bigcap_{k=1}^{\infty} \{x \in \mathbf{R}^n: F_k(x) = 1\}$$

is a set of positive measure, of  $q$ -uniqueness and  $p$ -multiplicity.

It is easy to check that

$$m(E) \geq (1 - \sum_k \varepsilon_k - 1/4)(3/4)^{n-1} > 0.$$

The proof for  $E$  that is a set of  $q$ -uniqueness is the same as for Theorem 1, Case (1).

To prove the  $p$ -multiplicity of  $E$ , let  $h^{\sharp}$  denote the periodic

extension to  $\mathbf{R}^n$ —defined in the same way as for  $f_k^\chi$  in Case (2)—of the function  $h$  found in Case (1), where  $G = T$ .

Let  $P$  be defined on  $\mathbf{R}^n$  as above. It remains only to show that the function

$$H(x_1, \dots, x_n) = h^\chi(x_1, \dots, x_n)P(x_1, \dots, x_n)$$

is a nonzero function supported by  $E$  (obvious), and satisfying  $H \in L^p(\hat{\mathbf{R}}^n)$ .

This last fact follows from the boundness of the operator  $T: \hat{g} \rightarrow (g^\chi P)^\wedge$  from  $L^1(Z)$  to  $L^p(\hat{\mathbf{R}}^n)$ , which we have proved above.

Thus the theorem is completed also in Case (2).

*Case (3).*  $G$  is a direct sum of a compact group  $K$  and a Euclidean space  $\mathbf{R}^n$  for  $K \neq \{0\}$  and  $n > 0$ . We can consider the set  $E \subset \mathbf{R}^n$  found in Case (2) and, by choosing the Haar measure on  $K$  such that  $m(K) = 1$ , it is easy to check that

$$E \oplus K$$

is a set which resolves our theorem. We simply recall that the dual group of a direct sum is a sum of the corresponding dual groups [8, p. 36].

3. A variant. We consider the case when  $G$  is the Cantor group.

**DEFINITION.** We call Cantor group  $D$ , the complete direct sum of  $Z(2)_n$  for  $n \in N$  where each  $Z(2)_n$  is the group of order two. Thus  $D$  is the group of all sequences  $\{\xi_n\}$ ,  $\xi_n = 0$  or  $\xi_n = 1$  with coordinatewise addition modulo 2, and with the topology that makes the mapping

$$\{\xi_n\} \rightsquigarrow 2 \sum_{n=1}^{\infty} \xi_n \cdot 3^{-n}$$

a homeomorphism of the group onto the classical Cantor set of the real line. It will be convenient to identify  $D$  with the interval  $[0, 1]$ . This is done by considering the mapping

$$t: D \longrightarrow [0, 1]$$

$$\{\xi_n\}_{n=1}^{\infty} \rightsquigarrow \sum_{n=1}^{\infty} \xi_n \cdot 2^{-n}$$

which is continuous, invertible a.e. and Haar measure preserving. Because the dual of a group of order two is again the group of order two, the dual of the group  $D$  is the direct sum of a sequence of groups of order two. In other words, the generators of  $\Gamma$  (the dual

of  $D$ ), are the functions

$$r_n(x) = (-1)^{\varepsilon_n},$$

and every element of  $\Gamma$  has the form

$$w(x) = \prod_{i=1}^k r_{n_i}(x) = (-1)^{\varepsilon_{n_1} + \dots + \varepsilon_{n_k}},$$

If  $t(x)$  is not a dyadic rational number, we can identify  $x$  with  $t(x)$  and then we can identify the  $r_n$  functions with the classical Rademacher functions on  $[0, 1]$ . It is clear that the harmonic analysis on  $L^p(D)$  is the same as on  $L^p([0, 1])$ , if we identify  $D$  with the real interval  $[0, 1]$ , its Haar measure with the Lebesgue measure, and its characters with the Walsh functions (finite products of Rademacher functions).

We recall now the Paley order for the Walsh functions and hence for the characters of  $D$ :

$$\begin{aligned} w_0 &= r_0 \equiv 1, \\ w_{2^{n-1}} &= r_n, & \text{for } n \geq 1, \\ w_n &= r_{n_1} \cdot \dots \cdot r_{n_k} & \text{where } n = 2^{n_1-1} + \dots + 2^{n_k-1}. \end{aligned}$$

Let us set, for  $f$  defined on  $D$ :

$$\|\hat{f}\|_{p,\alpha} = \left( \sum_{n \in N} |\hat{f}(w_n)|^p (1+n)^{p\alpha} \right)^{1/p}$$

where  $1 < p < \infty$  and  $0 < \alpha < 1 - 1/p$ .

If we make the obvious generalization of a set of uniqueness and a set of multiplicity for  $L^p(\Gamma)$ —given in §1—to the case  $L^{p,\alpha}(\Gamma)$ ,  $\alpha \neq 0$ , we can prove:

**THEOREM 2.** *For every real numbers  $p, q$  such that  $1 < p, q < \infty$ , if  $p'\alpha < \beta q'$ , where  $\alpha, \beta \in \mathbf{R}$ ,  $0 < \alpha < 1/p'$  and  $0 < \beta < 1/q'$ , there exists a subset  $E \subset D$  of positive measure which is a set of uniqueness for  $L^{q,\beta}(\Gamma)$  and a set of multiplicity for  $L^{p,\alpha}(\Gamma)$ .*

*Proof.* Given a real number  $\varepsilon > 0$  we must find a function  $F$  such that

$$(1) \quad \|\hat{F}\|_{q',-\beta} \leq \varepsilon \quad \text{where } 1/q + 1/q' = 1,$$

$$(2) \quad m\{x \in D: F(x) = 1\} \geq 1 - \varepsilon,$$

and a function  $\phi$  such that

$$(3) \quad \phi(x) \geq 0, \quad \text{for any } x \in D,$$

$$(4) \quad m\{x \in D: \phi(x) = 1\} \geq 1 - \varepsilon,$$

$$(5) \quad \log \|\hat{\phi}\|_{p,\alpha} \leq \varepsilon.$$

To this end we define a sequence  $\{F_m\}_{m=0}^\infty$  of functions:

$$F_m(x) = 1 - \sum_n w_n \quad \text{for } m = 0, 1, 2, \dots,$$

where the sum is taken over the finite set

$$\{0, 1 \cdot 2^{km}, 2 \cdot 2^{km}, 3 \cdot 2^{km}, \dots, (2^k - 1)2^{km}\}$$

and  $k$  is a positive integer to be specified later.

We can give the following equivalent definition for  $F_0$ :

$$F_0(x) = \begin{cases} 1 - 2^k & x \in (0, 1/2^k) \\ 1 & \text{elsewhere} . \end{cases}$$

If we extend  $F_0$  to a periodic function  $F_0^*$  of period one, defined on all of  $\mathbb{R}$  and equal to  $F_0$  on  $[0, 1]$  we can also define:

$$F_m(x) = F_0^*(2^{km}x) \quad \text{for } 0 \leq x \leq 1 .$$

It is clear that:

$$\int F_m(x) dx = 0 \quad \text{and} \quad m\{x \in D: F_m(x) = 1\} = 1 - 1/2^k .$$

Moreover, from the following relation:

$$\hat{F}_m(w_n) = \begin{cases} 1 & \text{if } n = 0, 1 \cdot 2^{km}, 2 \cdot 2^{km}, 3 \cdot 2^{km}, \dots, (2^k - 1)2^{km} , \\ 0 & \text{otherwise} , \end{cases}$$

we deduce:

$$\|\hat{F}_m\|_{q', -\beta} = \left( \sum_n (1 + n)^{-\beta q'} \right)^{1/q'} \leq A 2^{k(-\beta + 1/q')} \cdot 2^{km(-\beta)} .$$

Analogously we define a sequence  $\{\phi_m\}_{m=0}^\infty$  of functions such that

$$\phi_0(x) = 1 ,$$

$$\phi_m(x) = 1 - \frac{1}{2^k} \sum_n w_n(x) , \quad \text{for } m = 1, 2, \dots ,$$

where the sum is taken over the set

$$\{0, 1 \cdot 2^{k(m-1)}, 2 \cdot 2^{k(m-1)}, \dots, (2^k - 1)2^{k(m-1)}\} .$$

Since the  $\phi_m$  are trigonometric polynomials, it is trivial that:

$$\hat{\phi}_m(w_n) = \begin{cases} 1 - 1/2^k & \text{if } n = 0 \\ 1/2^k & \text{if } n = 2^{k(m-1)}, \dots, (2^k - 1)2^{k(m-1)} \\ 0 & \text{otherwise} , \end{cases}$$

and so, an easy calculation yields:

$$\log \|\hat{\phi}_m\|_{p,\alpha} \leq B \cdot 2^{k(1-p)} \cdot 2^{km(p\alpha)}.$$

Moreover, we have

$$m\{x: \phi_m(x) = 1\} \geq 1 - 1/2^k.$$

Indeed we can give the following equivalent definition for  $\phi_1$ :

$$\phi_1(x) = \begin{cases} 0 & \text{for } x \in (0, 1/2^k) \\ 1 & \text{elsewhere} \end{cases}$$

and notice that, if  $\phi_1^*$  is a periodic extension of  $\phi_1$  on all the real line, the following relation holds:

$$\phi_m(x) = \phi_1^*(2^{k(m-1)}x) \quad \text{for any } x, 0 \leq x \leq 1.$$

Now, if we choose  $k$  and  $m$  such that are solutions of the system:

$$(2) \quad \begin{cases} A2^{k(1/q'-\beta)} & 2^{km(-\beta)} \leq \varepsilon \\ B2^{k(1-p)} & 2^{km(p\alpha)} \leq \varepsilon \end{cases}$$

(we observe that  $A$  and  $B$  are constants depending on  $p, q, \alpha, \beta$ ),  $F_m$  and  $\phi_m$  satisfy properties (1), (2), (3), (4), and (5).

These solutions exist because  $p'\alpha < \beta q'$ , by hypothesis.

Let us define:

$$E = \{x \in D: F_m(x) = 1\}.$$

Choose a sequence of real numbers  $\varepsilon_i > 0$ ,  $i = 1, 2, \dots$ , such that  $2 \sum_i \varepsilon_i < 1$ , and repeat the preceding construction for every  $\varepsilon = \varepsilon_i$ . If we also choose the solutions  $k$  and  $m$  of (2) in such a way that:

$$k_i > k_{i-1}m_{i-1},$$

we can show, by induction, that, for every  $N \in \mathbb{N}$ :

$$\left\| \left( \prod_{i=1}^N \phi_i \right)^\wedge \right\|_{p,\alpha} \leq \prod_{i=1}^N \|\hat{\phi}_i\|_{p,\alpha}.$$

We assert now that our set of  $(p, \alpha)$ -multiplicity and  $(q, \beta)$ -uniqueness is

$$E = \bigcap_{i=1}^{\infty} E_i.$$

The proof of the theorem is now the same as that for Theorem 1, Case (1).

## REFERENCES

1. R. E. Edwards, *Fourier Analysis: A Modern Introduction*, Volumes I and II, Holt, Rinehart and Winston, New York, 1967 and 1968.

2. A. Figà-Talamanca and G. I. Gaudry, *Multipliers and sets of uniqueness of  $L^p$* , Michigan Math. J., **17** (1970), 179-191.
3. P. R. Halmos, *Measure Theory*, Van Nostrand, New York, 1950.
4. E. Hewitt and K. Ross, *Abstract Harmonic Analysis*, Volumes I and II, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963.
5. I. I. Hirschman, Jr., and Y. Katznelson, *Sets of uniqueness and multiplicity for  $l^{p,\alpha}$* , Israel J. Math., **3** (1965), 221-231.
6. Y. Katznelson, *An Introduction to Harmonic Analysis*, Wiley, New York, 1968.
7. ———, *Sets of uniqueness for some classes of trigonometrical series*, Bull. Amer. Math. Soc., **70** (1964), 722-723.
8. W. Rudin, *Fourier Analysis on Groups*, Interscience, New York-London, 1962.

Received April 5, 1972. This paper was written in part while the author held an undergraduate fellowship of the Consiglio Nazionale delle Ricerche.

UNIVERSITA DI GENOVA, ITALY

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan



Joseph Anthony Ball and Arthur R. Lubin, <i>On a class of contractive perturbations of restricted shifts</i> .....	309
Joseph Becker and William C. Brown, <i>On extending higher derivations generated by cup products to the integral closure</i> .....	325
Andreas Blass, <i>Exact functors and measurable cardinals</i> .....	335
Joseph Eugene Collison, <i>A variance property for arithmetic functions</i> .....	347
Craig McCormack Cordes, <i>Quadratic forms over nonformally real fields with a finite number of quaternion algebras</i> .....	357
Freddy Delbaen, <i>Weakly compact sets in <math>H^1</math></i> .....	367
G. D. Dikshit, <i>Absolute Nörlund summability factors for Fourier series</i> .....	371
Edward Richard Fadell, <i>Nielsen numbers as a homotopy type invariant</i> .....	381
Josip Globevnik, <i>Analytic extensions of vector-valued functions</i> .....	389
Robert Gold, <i>Genera in normal extensions</i> .....	397
Solomon Wolf Golomb, <i>Formulas for the next prime</i> .....	401
Robert L. Griess, Jr., <i>The splitting of extensions of <math>SL(3, 3)</math> by the vector space <math>F_3^3</math></i> .....	405
Thomas Alan Keagy, <i>Matrix transformations and absolute summability</i> .....	411
Kazuo Kishi, <i>Analytic maps of the open unit disk onto a Gleason part</i> .....	417
Kwangil Koh, Jiang Luh and Mohan S. Putcha, <i>On the associativity and commutativity of algebras over commutative rings</i> .....	423
James C. Lillo, <i>Asymptotic behavior of solutions of retarded differential difference equations</i> .....	431
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues</i> .....	445
Anna Maria Mantero, <i>Sets of uniqueness and multiplicity for <math>L^p</math></i> .....	467
J. F. McClendon, <i>Embedding metric families</i> .....	481
L. Robbiano and Giuseppe Valla, <i>Primary powers of a prime ideal</i> .....	491
Wolfgang Ruess, <i>Generalized inductive limit topologies and barrelledness properties</i> .....	499
Judith D. Sally, <i>Bounds for numbers of generators of Cohen-Macaulay ideals</i> ....	517
Helga Schirmer, <i>Mappings of polyhedra with prescribed fixed points and fixed point indices</i> .....	521
Cho Wei Sit, <i>Quotients of complete multipartite graphs</i> .....	531
S. Sznaider and Zbigniew Zielezny, <i>Solvability of convolution equations in <math>\mathcal{H}'_p</math>, <math>p &gt; 1</math></i> .....	539
Mitchell Herbert Taibleson, <i>The existence of natural field structures for finite dimensional vector spaces over local fields</i> .....	545
William Yslas Vélez, <i>A characterization of completely regular fields</i> .....	553
P. S. Venkatesan, <i>On right unipotent semigroups</i> .....	555
Kenneth S. Williams, <i>A rational octic reciprocity law</i> .....	563
Robert Ross Wilson, <i>Lattice orderings on the real field</i> .....	571
Harvey Eli Wolff, <i>V-localizations and V-monads. II</i> .....	579