

Pacific Journal of Mathematics

PACKING SPHERES IN ORLICZ SPACES

CHARLES EDWARD CLEAVER

PACKING SPHERES IN ORLICZ SPACES

CHARLES E. CLEAVER

A collection of open balls of radius r can be packed in the unit ball U of a Banach space provided each ball is a subset of U and the intersection of any two is empty. In an infinite dimensional Banach space, it is possible to find a largest number Λ so that if $r \leq \Lambda$ then an infinite number of spheres of radius r can be packed in U . In this paper, upper and lower bounds are found for this number in Orlicz spaces.

For the space l_2 , this number was found by Rankin [7] to be $1/(1 + \sqrt{2})$ and this result was extended in [1] to show that the number in $l_p (1 \leq p < \infty)$ is $1/(1 + 2^{1-1/p})$. In 1970 Kottman [4] showed that $1/3 \leq \Lambda \leq 1/2$ for any Banach space. More recently, Wells and Williams [10] used a generalized Riesz-Thorin interpolation theorem to obtain the exact value of Λ in the $L^p(\mu) (1 \leq p < \infty)$ spaces with some restrictions on the measure space when $2 < p < \infty$. The results in this paper include all the above and also show that all restrictions can be removed in the L^p case. Recent results have demonstrated that the structure of Orlicz spaces is quite different from L^p spaces and very little seems to be known in the Orlicz case. The packing criteria lead to some results on isometric embeddings of subspaces and to notions of noncompactness.

2. Preliminaries. An Orlicz function M will be a continuous convex nondecreasing function defined for $x \geq 0$ and such that $M(0) = 0$, $M(\infty) = \infty$ and $M(x) > 0$ for $x > 0$. The Orlicz space $L_M(X, \mathcal{A}, \mu) (= L_M)$ is the set of measurable scalar-valued functions defined on the measure space (X, \mathcal{A}, μ) such that $f \in L_M$ if and only if $\|f\|' < \infty$ where

$$\|f\|'_M = \inf \left\{ k > 0: \int_X M\left(\frac{|f|}{k}\right) d\mu \leq 1 \right\}.$$

For each Orlicz function M , a complementary function N is defined by

$$N(x) = \sup \{xy - M(y): 0 < y < \infty\}.$$

If $M(x) = \int_0^x p(t)dt$ where p is a right continuous nondecreasing function, then $N(p(x)) = xp(x) - M(x)$ (cf [5]). Using this function, another norm can be defined on L_M

$$\|f\|_M = \sup \left\{ \int_X |fg| d\mu : \|g\|'_N \leq 1 \right\}.$$

These norms are equivalent if every set of positive μ -measure contains a subset of positive finite μ -measure and in this paper the latter will be used. In the case of $M(x) = x^p$, $p > 1$, it follows that $\|f\|_p = \|f\|'_M = K \|f\|_M$ where K is independent of f (cf [11]). It will be assumed in the remainder of the paper that M is chosen so that the simple functions are dense in L_M .

If M_1 and M_2 are two Orlicz functions then M_s will denote the inverse of $M_s^{-1} = (M_1^{-1})^{1-s}(M_2^{-1})^s$ for $0 \leq s \leq 1$, where M^{-1} is the unique inverse of the Orlicz function M . The function M_s is an Orlicz function and satisfies most of properties of M_1 and M_2 including the fact that the simple functions are dense in L_{M_s} if the same is true in L_{M_1} and L_{M_2} . The complementary function to M_s is not always the same as the inverse of $N_s^{-1} = (N_1^{-1})^{1-s}(N_2^{-1})^s$ where N_1 and N_2 are the respective complements of M_1 and M_2 . However, the complement of M_s and the inverse of N_s^{-1} generate the same Orlicz space with equivalent norms (cf [8]). Since the complementary function is the one of interest in this paper, N_s will denote the complement of M_s .

One condition which guarantees the separability of L_M is the Δ_2 -condition. An Orlicz function is said to satisfy the Δ_2 -condition at ∞ if $\lim_{x \rightarrow \infty} \sup M(2x)/M(x) < \infty$. In the case of sequence spaces, separability occurs if and only if the Δ_2 -condition holds at 0. A necessary and sufficient condition that M satisfy the Δ_2 -condition is that $\lim_{x \rightarrow \infty} \sup xM'(x)/M(x) = \alpha < \infty$ where $M'(x)$ is the derivative of M (cf [5], p. 24). If M' and N' are both continuous where N is the complement of M , then this condition is equivalent to

$$\liminf_{x \rightarrow \infty} \frac{xN'(x)}{N(x)} > \alpha/\alpha - 1.$$

This and elementary calculus lead to a lemma that will be useful in later sections.

LEMMA 2.1. *Let M and N be complementary functions with M' and N' continuous. If*

$$\alpha = \limsup_{x \rightarrow \infty} \frac{xM'(x)}{M(x)}$$

then

$$\liminf_{x \rightarrow \infty} \frac{N^{-1}(x)}{N^{-1}(2x)} \geq \frac{1}{2^{(\alpha-1)/\alpha}}.$$

3. Interpolation. In this section a generalized interpolation theorem is described and then applied to obtain inequalities that will be useful in next section. This theorem generalizes Theorem 1 in [8] and follows the development in [10] of the L_p case.

Let $(X_1, \mu_1), (X_2, \mu_2), \dots, (X_n, \mu_n)$ be measure spaces and $M = (M_1, M_2, \dots, M_n)$ be an n -tuple of Orlicz functions. Define the direct sum $\bigoplus L_{M_k}(\mu_k)$ by

$\bigoplus L_{M_k}(\mu_k) = \{f = (f_1, f_2, \dots, f_n) \mid f_k \in L_{M_k}(\mu_k), k = 1, 2, \dots, n\}$ with usual addition and scalar multiplication. For each $r, 1 \leq r \leq \infty$ and each n -tuple $\lambda = (\lambda_1, \dots, \lambda_n)$ of positive weights, introduce the following norm on $\bigoplus L_{M_k}(\mu_k)$,

$$\|f\|_{M,r} = \begin{cases} \left\{ \sum_{k=1}^n \|f_k\|_{M_k}^r \lambda_k \right\}^{1/r} & 1 \leq r < \infty \\ \max_{1 \leq k \leq n} \|f_k\|_{M_k} & r = \infty. \end{cases}$$

The space of all f such that $\|f\|_{M,r} < \infty$ is a Banach space and will be denoted by $L_M^\lambda(\lambda)$.

For two n -tuples $M_1 = (M_{11}, M_{12}, \dots, M_{1n})$ and $M_2 = (M_{21}, M_{22}, \dots, M_{2n})$ define $M_s = (M_{s1}, M_{s2}, \dots, M_{sn}), 0 \leq s \leq 1$, where M_{sk} is the inverse of the function $M_{sk}^{-1} = (M_{1k}^{-1})^{1-s} (M_{2k}^{-1})^s, k = 1, 2, \dots, n$.

Now let $(Y_1, \nu_1), (Y_2, \nu_2), \dots, (Y_m, \nu_m)$ be another collection of measure spaces, $\eta = (\eta_1, \dots, \eta_m)$ and define m -tuples Q_1, Q_2 in the same manner as M_1 and M_2 . Letting $X = (X_1, X_2, \dots, X_n)$ and $Y = (Y_1, Y_2, \dots, Y_m)$, the following interpolation theorem was proved in [2].

THEOREM 3.1. *Let $1 \leq r_i, t_i \leq \infty, i = 1, 2, 0 \leq s \leq 1$ with $1/r = 1 - s/r_1 + s/r_2, 1/t = 1 - s/t_1 + s/t_2$ and suppose M_i and $Q_i, i = 1, 2$, are defined on X and Y respectively. If T is a linear transformation from $L_{M_i}^{t_i}(\lambda)$ into $L_{Q_i}^{r_i}(\eta), i = 1, 2$, with bounds K_1 and K_2 respectively, then T takes $L_{M_s}^t$ into $L_{Q_s}^r$ and*

$$\|Tf\|_{Q_s,r} \leq K_1^{1-s} K_2^s \|f\|_{M_s,t}.$$

This result is quite useful in establishing inequalities as the following theorem demonstrates.

THEOREM 3.2. *Let M be an Orlicz function, $M_0(x) = x^2$ and $M_s^{-1} = (M^{-1})^{1-s} (M_0^{-1})^s, 0 \leq s \leq 1$. Then for any collection of positive numbers c_1, c_2, \dots, c_n such that $\sum_{i=1}^n c_i = 1$, the inequality*

$$\sum_{i,j=1}^n c_i c_j \|f_i - f_j\|_{V_s}^{2/(2-s)} \leq 2\gamma^{2(1-s)/(2-s)} \sum_{i=1}^n c_i \|f_i\|_{M_s}^{2/(2-s)}$$

holds wherever $f_1, f_2, \dots, f_n \in L_{M_s}$ and $\gamma = \max_{1 \leq i \leq n} (1 - c_i)$.

Proof. Let $M_i, i = 1, 2$ be the constant n -tuple with each component M and $Q_i, i = 1, 2$, the constant n^2 -tuple with each component M . Setting $t_1 = r_1 = 1, t_2 = r_2 = 2, c = (c_1, c_2, \dots, c_n)$ and $c^2 = (c_i c_j)_{i,j=1}^n$ define T from $L_{M_i}^i(c)$ into $L_{Q_i}^i(c^2)$ by $T(f_1, f_2, \dots, f_n) = (f_i - f_j)_{i,j=1}^n$. Now

$$\begin{aligned} \|Tf\|_{M,1} &= \sum_{i,j=1}^n c_i c_j \|f_i - f_j\|_M \\ &\leq \sum_{i,j=1}^n c_i c_j (\|f_i\|_M + \|f_j\|_M) - 2 \sum_{i=1}^n c_i^2 \|f_i\|_M \\ &= 2 \sum_{i=1}^n \|f_i\|_M (1 - c_i) c_i \leq 2\gamma \sum_{i=1}^n c_i \|f_i\|_M = 2\gamma \|f\|_{M,1}. \end{aligned}$$

It follows from properties of Hilbert space that $\|Tf\|_{M_0,2} \leq \sqrt{2} \|f\|_{M_0,2}$. According to Theorem 3.1, T takes $L_{M_s}^{2/(2-s)}(c^2)$ into $L_{M_s}^{2/(2-s)}(c^2)$ and

$$\|Tf\|_{M_s, 2/(2-s)} \leq (2\gamma)^{1-s} (\sqrt{2})^s \|f\|_{M_s, 2/(2-s)}.$$

This says

$$\left\{ \sum_{i,j=1}^n c_i c_j \|f_i - f_j\|_{M_s}^{2/(2-s)} \right\}^{(2-s)/2} \leq (2\gamma)^{1-s} (\sqrt{2})^s \left\{ \sum_{i=1}^n c_i \|f_i\|_{M_s}^{2(2-s)} \right\}^{(2-s)/2}.$$

Raising both sides to the $2/(2-s)$ power, the desired inequality is obtained.

The above theorem reduces to the results found in [10] for the L_p case.

COROLLARY 3.3. *Let $1 < p < \infty$ and c_1, c_2, \dots, c_n be any collection of positive numbers such that $\sum_{i=1}^n c_i = 1$. Then for any f_1, f_2, \dots, f_n in L_p ,*

$$(i) \quad \sum_{i,j=1}^n c_i c_j \|f_i - f_j\|_p^p \leq 2\gamma^{2-p} \sum_{i=1}^n c_i \|f_i\|_p^p, \quad 1 \leq p \leq 2$$

and

$$(ii) \quad \sum_{i,j=1}^n c_i c_j \|f_i - f_j\|_{p'}^{p'} \leq 2\gamma^{2-p'} \sum_{i=1}^n c_i \|f_i\|_{p'}^{p'}, \quad 2 < p < \infty$$

where $p' = p/p - 1$.

Proof. To prove (i), choose l so that $1 < l < p \leq 2$ and let $M(x) = x^l$. If we set $s = 2/p((p-l)/(p-2))$, $M_s(x) = x^p$ and let $l \rightarrow 1$, then $2/(2-s)$ approaches p . Similarly one can show (ii) by choosing $l > p$ and allowing $l \rightarrow \infty$.

4. Packing. The main object of this section is to find bounds on the number A_M where A_M satisfies the property that for $r \leq A_M$, infinite packing is possible and for $r > A_M$, only a finite number of balls of radius r can be packed in the unit ball of the Orlicz space L_M . It has been shown by Kottman [4] that $1/3 \leq A_M \leq 1/2$. These bounds are improved below in the spaces $L_M[0, 1]$ but it is clear that the techniques apply to a wider class of spaces.

DEFINITION 4.1. A family of balls $\{B_r(f_j)\}_{j \in I}$ of radius r and centers $\{f_j\}_{j \in I}$ can be packed in the unit ball B_1 of L_M provided

- (i) $B_r(f_j) \subset B_1$ for each $j \in I$
- (ii) $\text{int}(B_r(f_j)) \cap \text{int}(B_r(f_k)) = \emptyset, j \neq k$.

If a family of balls $\{B_r(f_j)\}_{j \in I}$ can be packed in B_1 then it is clear that

$$(4.1) \quad \|f_j\| \leq 1 - r, j \in I$$

$$(4.2) \quad \|f_j - f_k\| \geq 2r, j \neq k$$

must be satisfied. Thus to find an example to serve as lower bound one needs to find vectors f_1, f_2, \dots , satisfying these inequalities.

Given an Orlicz function M with complement N , choose a sequence of disjoint measurable sets $\{E_j\}_{j=1}^\infty$ in $[0, 1]$ and define

$$g_k = \frac{1}{N^{-1}\left(\frac{1}{\mu(E_k)}\right)\mu(E_k)} \chi_{E_k}, \quad k = 1, 2, \dots$$

Each g_k has the property that $\|g_k\|_M = 1$ (cf [5]). To compute the norm of the difference of two of these, consider the function

$$h \equiv N^{-1}\left(\frac{1}{2\mu(E_k)}\right)\chi_{E_k} + N^{-1}\left(\frac{1}{2\mu(E_n)}\right)\chi_{E_n}.$$

Then

$$\int_X N(h) = \int_{E_k} \frac{1}{2\mu(E_k)} \chi_{E_k} + \int_{E_n} \frac{1}{2\mu(E_n)} \chi_{E_n} = 1$$

and hence $\|h\|'_N \leq 1$. Now

$$\begin{aligned} \|g_k - g_n\|_M &= \sup_{\|f\|'_N \leq 1} \int_X |g_k - g_n| |f| d\mu \\ &\geq \int_X |g_k - g_n| |h| = \frac{N^{-1}\left(\frac{1}{2\mu(E_k)}\right)}{N^{-1}\left(\frac{1}{\mu(E_k)}\right)} + \frac{N^{-1}\left(\frac{1}{2\mu(E_n)}\right)}{N^{-2}\left(\frac{1}{\mu(E_n)}\right)}. \end{aligned}$$

By choosing a subsequence we obtain

$$\|g_k - g_n\|_M \geq 2 \liminf_{x \rightarrow \infty} \frac{N^{-1}(x)}{N^{-1}(2x)}.$$

Putting $f_k = (1 - r)g_k$, $k = 1, 2, \dots$, it follows that $\|f_k\| = 1 - r$ and

$$\|f_k - f_n\| \geq (1 - r)2 \liminf_{x \rightarrow \infty} \frac{N^{-1}(x)}{N^{-1}(2x)}.$$

Setting

$$\beta = \liminf_{x \rightarrow \infty} \frac{N^{-1}(x)}{N^{-1}(2x)},$$

the inequalities (4.1) and (4.2) will be satisfied provided $(1 - r)2\beta \geq 2r$ or $r \leq 1/(1 + 1/\beta)$. This example shows that $A_M \geq 1/(1 + 1/\beta)$ and leads to the following theorem.

THEOREM 4.2. *$L_M[0, 1]$ be an Orlicz space with N the complement of M and set $M_s^{-1} = (M^{-1})^{1-s}(M_0^{-1})^s$ where $M_0(x) = x^2$, $0 \leq s \leq 1$. Then with*

$$(4.3) \quad \beta = \liminf_{x \rightarrow \infty} \frac{N_s^{-1}(x)}{N_s^{-1}(2x)},$$

$$\frac{1}{1 + 1/\beta} \leq A_{M_s} \leq \frac{1}{1 + 2^{s/2}}.$$

Furthermore, if $1/(1 + 2^{s/2}) < r < 1$ then at most a finite number $\Gamma_{M_s}(r)$ of balls of radius r can be packed in B_1 and that number satisfies

$$(4.4) \quad \Gamma_{M_s}(r) \leq \left[1 - 1/2 \left(\frac{1 - r}{r} \right)^{2/s} \right]^{-1}.$$

Proof. It remains to show (4.4) and the right hand side of (4.3). Suppose there are n disjoint balls of radius r with centers f_1, f_2, \dots, f_n packed in B_1 . Then by Theorem 3.2,

$$(4.5) \quad \sum_{i,j=1}^n c_i c_j \|f_i - f_j\|_{M_s}^{2/(2-s)} \leq 2\gamma \frac{2(1-s)}{2-s} \sum_{i=1}^n c_i \|f_i\|_{M_s}^{2/(2-s)}$$

for any collection c_1, c_2, \dots, c_n of positive numbers such that $\sum_{i=1}^n c_i = 1$. In particular, if $c_i = 1/n$, $i = 1, 2, \dots, n$ then $\gamma = 1 - 1/n$ and (4.5) reduces to

$$(4.6) \quad \sum_{i,j=1}^n \frac{1}{n^2} \|f_i - f_j\|_{M_s}^{2/(2-s)} \leq 2 \left(1 - \frac{1}{n}\right)^{2(1-s)/(2-s)} \sum_{i=1}^n \frac{1}{n} \|f_i\|_{M_s}^{2/(2-s)}$$

since the balls are disjoint, $\|f_i - f_j\| \geq 2r$, $i \neq j$, and $\|f_i\| \leq 1 - r$. Hence (4.6) implies

$$\frac{1}{n^2} n(n-1)(2r)^{2/(2-s)} \leq 2 \left(\frac{n-1}{n}\right)^{2(1-s)/(2-s)} \frac{1}{n} \cdot n(1-r)^{2/(2-s)}.$$

This inequality then reduces to

$$(4.7) \quad r \leq \frac{1}{1 + 2^{s/2} \left(\frac{n-1}{n}\right)^{s/2}}.$$

If we allow $n \rightarrow \infty$, the right hand side of (4.3) is obtained. The inequality (4.4) follows by solving (4.7) for n .

In the case when M and N have continuous derivatives, proposition 2.1 gives a lower bound in terms of M_s .

COROLLARY 4.3. *Let M and N be complementary Orlicz functions with M satisfying the Δ_2 -condition. If M and N have continuous derivatives and $M_s^{-1} = (M^{-1})^{1-s}(M_0^{-1})^s$, $0 \leq s \leq 1$, then*

$$(4.8) \quad \frac{1}{1 + 2^{(\alpha-1)/\alpha}} \leq A_{M_s} \leq \frac{1}{1 + 2^{s/2}}$$

where

$$\alpha = \limsup_{x \rightarrow \infty} \frac{xM'_s(x)}{M_s(x)}.$$

If we set $M(x) = x^p$ and use a proof similar to Corollary 3.3, the exact value $A_M \equiv A_p$ is obtained for L^p , $1 \leq p \leq 2$.

COROLLARY 4.4. *Let $1 \leq p \leq 2$. Then $A_p = 1/(1 + 2^{1-1/p})$ for the space $L_p(\mu)$.*

This holds for any measure space because, for $M(x) = x^p$ in the example preceding Theorem 4.2, $N^{-1}(2x)/N^{-1}(x) = 2^{1-1/p}$ for all x .

The upper bounds are independent of the measure space but not the lower. Corollary 4.3 does not give the exact number for $2 < p < \infty$ but gives a lower bound which was shown in [1] to be exact for l_p . However, it is demonstrated in [10] that the number in $L_p[0, 1]$, $2 < p < \infty$ is $1/(1 + 2^{1/p})$. A simple generalization of this gives us new lower bounds in Orlicz spaces.

For each positive integer n and each integer j , $0 < j \leq 2^n$, define $E_{nj} = ((j-1)/2^n, j/2^n)$. Now for each integer n , define the function g_n by

$$g_n = \frac{1}{N^{-1}(1)} \sum_{k=1}^{2^n} (-1)^{k+1} \chi_{E_{nk}}$$

where N is the complementary function of the Orlicz function M and $\chi_{E_{nk}}$ is the characteristic function of the set E_{nk} . Then $\|g_n\|_M = 1$ for each n and $\|g_n - g_m\| = N^{-1}(2)/N^{-1}(1)$, $n \neq m$. Consider the spheres $S_r(f_j)$, $j = 1, 2, \dots$ with centers $f_j = (1-r)g_j$. Thus $\|f_i\| = 1-r$ and $\|f_j - f_k\| = (1-r)N^{-1}(2)/N^{-1}(1)$. The inequalities (4.1) and (4.2) will be satisfied provided $(1-r)N^{-1}(2)/N^{-1}(1) \geq 2r$ or $r \leq 1/(1 + 2N^{-1}(1)/N^{-1}(2))$.

THEOREM 4.4. *Let $L_M[0, 1]$ be an Orlicz space and set $M_s^{-1} = (M^{-1})^{1-s}(M_s^{-1})^s$ where $\phi_s(x) = x^2$ and $0 \leq s \leq 1$. If N_s is the complementary function to M_s , then*

$$\frac{1}{1 + \frac{2N_s^{-1}(1)}{N_s^{-1}(2)}} \leq A_{M_s} \leq \frac{1}{1 + 2^{s/2}}.$$

The example constructed above does not depend on $[0, 1]$ but rather on being able to find sets E_{nj} with the same properties. However, for the L_p spaces the construction on $[0, 1]$ is enough and theorem 16.2 in [10] generalizes to the following.

COROLLARY 4.5. *Let $2 \leq p < \infty$ and μ be any measure which is not purely atomic. Then for $L_p(\mu)$,*

$$A_p(\mu) = \frac{1}{1 + 2^{1/p}}.$$

Proof. For the space $L_p[0, 1]$, the usual argument gives the result. It is known ([3] or [9]) that if μ is not purely atomic, $L_p(\mu)$ has a subspace isometric to $L_p[0, 1]$. Suppose there are infinitely many balls of radius r in $L_p[0, 1]$ then there is a sequence of points satisfying inequalities (4.1) and (4.2) in the subspace and hence in $L_p(\mu)$. Thus the lower bound for A in $L_p(\mu)$ is greater than or equal to A_p and since the upper bound is independent of the measure, the result follows.

The problem of embedding $L_p[0, 1]$ into $L_r[0, 1]$ has been studied extensively and it has been shown [cf 3] that for $1 \leq r \leq p < 2$, $L_p[0, 1]$ is isometric to a subspace of $L_r[0, 1]$. More recently Nielsen [6] has given conditions under which $L_p(0, \infty)$ is isomorphic to a

subspace $L_p[0, 1]$. Also the Khintchin inequality implies l_2 is isomorphic to a subspace of $L_M[0, 1]$ for every Orlicz function M and furthermore l_2 is actually isometric to a subspace of $L_p[0, 1]$ for every $p, 1 \leq p < \infty$. Consistent with these results is the following.

THEOREM 4.6. *Let M_1 and M_2 be Orlicz functions and suppose L_{M_2} is isometric to a subspace of L_{M_1} . Then $\Lambda_{M_2} \geq \Lambda_{M_1}$. In particular if l_2 is isometric to a subspace of L_M then $1/(1 + \sqrt{2}) \leq \Lambda_M \leq 1/2$.*

A converse to this theorem would be of interest. A reasonable conjecture might be to try to show that if $[\alpha_{M_1}, \alpha_{M_2}] < [\alpha_{M_1}, \alpha_{M_2}] < 2$ (see [3] for definitions) and $\Lambda_{M_2} \geq \Lambda_{M_1}$ then $L_{M_1}[0, 1]$ is isometric to a subspace of $L_{M_2}[0, 1]$.

For the sequence case, the situation is different. Using the example preceding Theorem 4.2 with each E_n a singleton, it follows that $\lambda_M \geq 1/(1 + (N^{-1}(1)/N^{-1}(1/2)))$. The proof in [1] for l_p depends on the strong property

$$(4.9) \quad M^{-1}\left(\sum_{j=1}^{\infty} M(x_j)\right) = \|X\|_M$$

where $M(x) = x^p/p$. If we mimic their proof the following is obtained.

THEOREM 4.7. *Let M and N be complementary functions both satisfying the Λ_2 -condition at 0, and M satisfies (4.9). Then for the space l_M ,*

$$\frac{1}{1 + \frac{N^{-1}(1)}{N^{-1}\left(\frac{1}{2}\right)}} \leq \lambda_M \leq \frac{1}{1 + \frac{2}{M^{-1}(2M(1))}}.$$

Furthermore, if $N^{-1}(1/2) \leq 1/2 N^{-1}(1)M^{-1}(2M(1))$, then for

$$\frac{1}{1 + \frac{2}{M^{-1}(2M(1))}} < r \leq \frac{1}{1 + 2\frac{N^{-1}(1)}{N^{-1}(2)}},$$

any finite number of spheres of radius r can be packed in the unit ball of l_M but not an infinite number.

Proof. The Λ_2 -condition on both M and N is equivalent to reflexivity. Now suppose there are an infinite number of balls of radius r in l_M with centers y_1, y_2, \dots satisfying inequalities (4.1)

and (4.2). Assume y is the weak limit point of $\{y_j\}$, then $y \in l_M$ and $\|y\| \leq 1 - r$. Let $\varepsilon > 0$ and fix a positive integer n . Then there exists N such that $\|\tilde{y}_k\| < \varepsilon(1 - r)$ where

$$\tilde{y}_{kj} = \begin{cases} 0 & j \leq N \\ y_{kj} & j > N \end{cases}.$$

Then

$$\begin{aligned} M\left(\frac{2r}{1-r}\right) &\leq M\left(\frac{\|y_n - y_m\|}{1-r}\right) = \sum_{j=1}^{\infty} M\left(\frac{|y_{nj} - y_{mj}|}{1-r}\right) \\ &= \sum_{j=1}^N M\left(\frac{|y_{nj} - y_{mj}|}{1-r}\right) + \sum_{j>N} M\left(\frac{|y_{nj} - y_{mj}|}{1-r}\right). \end{aligned}$$

Now

$$\begin{aligned} M^{-1}\left(\sum_{j>N} M\left(\frac{|y_{nj} - y_{mj}|}{1-r}\right)\right) &= \left\| \frac{y_n - \tilde{y}_m}{1-r} \right\| \\ &\leq \frac{\|y_m\|}{1-r} + \frac{\|\tilde{y}_n\|}{1-r} \leq 1 + \varepsilon. \end{aligned}$$

Thus

$$M\left(\frac{2r}{1-r}\right) \leq \sum_{j=1}^N M\left(\frac{|y_{nj} - y_{mj}|}{1-r}\right) + M(1 + \varepsilon).$$

This argument is independent of m and hence

$$M\left(\frac{2r}{1-r}\right) - M(1 + \varepsilon) \leq \sum_{j=1}^N M\left(\frac{|y_{nj} - y_j|}{1-r}\right).$$

Letting $N \rightarrow \infty$ and $\varepsilon \rightarrow 0$, this becomes

$$M\left(\frac{2r}{1-r}\right) - M(1) \leq \sum_{j=1}^{\infty} M\left(\frac{|y_{nj} - y_j|}{1-r}\right).$$

Repeat the argument using y in place of y_n obtaining

$$M\left(\frac{2r}{1-r}\right) - M(1) \leq \sum_{j=1}^{N'} M\left(\frac{|y_{nj} - y_j|}{1-r}\right) + M(1 + \varepsilon).$$

Now, letting $n \rightarrow \infty$ and $\varepsilon \rightarrow 0$, it follows that

$$M\left(\frac{2r}{1-r}\right) \leq 2M(1)$$

and

$$r \leq \frac{1}{1 + \frac{2}{M^{-1}(2M(1))}}.$$

The last statement follows by constructing the example preceding Theorem 4.4 on the set $(1, 2, \dots, 2^n)$ in place of $[0, 1]$.

COROLLARY 4.8. *For $1 \leq p \leq \infty$, $\lambda_p = 1/(1 + 2^{1-1/p})$ for the spaces l_p . Furthermore if $2 \leq p < \infty$ then for*

$$\frac{1}{1 + 2^{1-1/p}} < r \leq \frac{1}{1 + 2^{1/p}},$$

any finite number of spheres of radius r can be packed in l_p , but not an infinite number.

REFERENCES

1. J. A. C. Burlack, R. A. Rankin, A. P. Robertson, *The packing of spheres in the space l_p* , Proc. M. A. Glasgow, **4** (1958), 145-146.
2. C. E. Cleaver, *On the Extension of Lipschitz-Holder maps on Orlicz Spaces*, Studia Mathematica, **27** (1972), 195-204.
3. J. Lindenstrauss, and L. Tzafriri, *Classical Banach Spaces*, Lecture Notes in Mathematics 338, Springer-Verlag 1973.
4. C. A. Kottman, *Packing and reflexivity in Banach spaces*, TAMS, **150** (1970), 565-576.
5. M. A. Kransnosel'skii, and Ya. B., Rutickii, *Convex Functions and Orlicz Spaces*, (Translation), 1961.
6. N. J. Nielsen, *On the Orlicz Function Spaces $L_M(0, \infty)$* , Preprint Series, ISBN 82-553-0166-6, No. 4, (1974), University of Oslo.
7. R. A. Rankin, *On packings of spheres in Hilbert space*, Proc. M. A. Glasgow, **2** (1955), 139-144.
8. M. M. Rao, *Interpolation, ergodicity, and Martingales*, J. of Math. and Mech., **16** (1966), 543-568.
9. H. P. Rosenthal, *On the subspaces of $L_p(p > 2)$ spanned by sequences of independent random variables*, Israel J. Math., **8** (1970), 273-303.
10. J. H. Wells and Lynn R. Williams. *Imbedding and Extension Problems in Analysis*, Springer-Verlag, 1975.
11. A. C. Zaananen, *Linear Analysis*, Amsterdam, 1953.

Received November 19, 1974 and in revised form April 12, 1976. The results in this paper were obtained while the author was a visiting professor at Georgia Institute of Technology.

KENT STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Andrew Adler, <i>Weak homomorphisms and invariants: an example</i>	293
Howard Anton and William J. Pervin, <i>Separation axioms and metric-like functions</i>	299
Ron C. Blei, <i>Sidon partitions and p-Sidon sets</i>	307
T. J. Cheatham and J. R. Smith, <i>Regular and semisimple modules</i>	315
Charles Edward Cleaver, <i>Packing spheres in Orlicz spaces</i>	325
Le Baron O. Ferguson and Michael D. Rusk, <i>Korovkin sets for an operator on a space of continuous functions</i>	337
Rudolf Fritsch, <i>An approximation theorem for maps into Kan fibrations</i>	347
David Sexton Gilliam, <i>Geometry and the Radon-Nikodym theorem in strict Mackey convergence spaces</i>	353
William Hery, <i>Maximal ideals in algebras of topological algebra valued functions</i>	365
Alan Hopenwasser, <i>The radical of a reflexive operator algebra</i>	375
Bruno Kramm, <i>A characterization of Riemann algebras</i>	393
Peter K. F. Kuhfittig, <i>Fixed points of locally contractive and nonexpansive set-valued mappings</i>	399
Stephen Allan McGrath, <i>On almost everywhere convergence of Abel means of contraction semigroups</i>	405
Edward Peter Merkes and Marion Wetzel, <i>A geometric characterization of indeterminate moment sequences</i>	409
John C. Morgan, II, <i>The absolute Baire property</i>	421
Eli Aaron Passow and John A. Roulier, <i>Negative theorems on generalized convex approximation</i>	437
Louis Jackson Ratliff, Jr., <i>A theorem on prime divisors of zero and characterizations of unmixed local domains</i>	449
Ellen Elizabeth Reed, <i>A class of T_1-compactifications</i>	471
Maxwell Alexander Rosenlicht, <i>On Liouville's theory of elementary functions</i>	485
Arthur Argyle Sagle, <i>Power-associative algebras and Riemannian connections</i>	493
Chester Cornelius Seabury, <i>On extending regular holomorphic maps from Stein manifolds</i>	499
Elias Sai Wan Shiu, <i>Commutators and numerical ranges of powers of operators</i>	517
Donald Mark Topkis, <i>The structure of sublattices of the product of n lattices</i>	525
John Bason Wagoner, <i>Delooping the continuous K-theory of a valuation ring</i>	533
Ronson Joseph Warne, <i>Standard regular semigroups</i>	539
Anthony William Wickstead, <i>The centraliser of $E \otimes_{\lambda} F$</i>	563
R. Grant Woods, <i>Characterizations of some C^*-embedded subspaces of $\beta\mathbb{N}$</i>	573