

Pacific Journal of Mathematics

VARIATION OF GREEN'S POTENTIAL

JANG-MEI GLORIA WU

VARIATION OF GREEN'S POTENTIAL

JANG-MEI G. WU

Let u be the Green's potential of a nonnegative mass distribution μ on unit disk D , defined by

$$(1.1) \quad u(z) = \int_D G(z, w) d\mu(w),$$

where G is the Green's function on D . The following will be proved.

THEOREM 1. *Suppose $0 < \rho < 1$ and*

$$(1.2) \quad \int_D (1 - |w|) \log \frac{1}{1 - |w|} d\mu(w) < \infty.$$

Then for almost all θ , $0 \leq \theta < 2\pi$, u has finite variation on the line segment joining $\rho e^{i\theta}$ and $e^{i\theta}$.

THEOREM 2. *Fix θ , $0 \leq \theta < 2\pi$ and suppose*

$$(1.3) \quad \int_D \frac{1 - |w|}{|e^{i\theta} - w|} d\mu(w) < \infty.$$

If L is a circular arc in D centered at $e^{i\theta}$, then for all a on L except a set of capacity zero, u has finite variation on the line segment joining a and $e^{i\theta}$.

It is known [4; 150] that if μ satisfies

$$(1.4) \quad \int_D (1 - |w|) d\mu(w) < \infty,$$

then u is finite everywhere except a set of capacity zero in D . Carleson [2; 32] has implicitly proved the following result.

THEOREM A. *Suppose $0 < \rho < 1$ and*

$$(1.5) \quad \int_D (1 - |w|)^\alpha d\mu(w) < \infty$$

for some fixed α , $0 \leq \alpha < 1$. Then for all θ , $0 \leq \theta < 2\pi$, except a set E of

α -dimensional Hausdorff measure zero if $0 < \alpha < 1$, or of capacity zero if $\alpha = 0$, u has finite variation on the line segment joining $pe^{i\theta}$ and $e^{i\theta}$.

Some ideas involved in Theorems 1 and 2 are borrowed from Rudin [6] and Cargo [1], who studied the variation of Blaschke products on line segments.

In §4 we shall extend the above three theorems to curves with certain differentiability properties, which need not be line segments.

For convenience, we shall use $l(z, w)$ to denote the line segment joining z, w and $V(F, \gamma)$ to denote the total variation of the function F on the curve γ whenever this is meaningful.

2. Lemmas.

LEMMA 1. *If l is a Jordan rectifiable curve in \bar{D} , then*

$$V(u, l) \leq \int_D V(G(\cdot, w), l) d\mu(w).$$

The proof follows directly from the definition of variation and (1.1).

LEMMA 2. *If $z, w \in D$ and $|z - w| \geq (1 - |w|)/2$ then $|(z - w)/(1 - \bar{w}z)| \geq 1/5$.*

Proof. For fixed w , $|(z - w)/(1 - \bar{w}z)|$ attains minimum $1/(3|w| + 2)$ on the circle $|z - w| = (1 - |w|)/2$. The lemma follows from minimal modulus principle.

LEMMA 3. *Let $w \in D$, A be a subset of D and $m = \min\{(1 - |z|)/|1 - z| : z \in A\}$. If the distance from w to A is less than $(1 - |w|)/2$, then $(1 - |w|)/|1 - w| \geq m/3$.*

Proof. Let $z \in A$ and $|z - w| \leq (1 - |w|)/2$. It is easy to check $\frac{1}{2}(1 - |w|) \leq 1 - |z| \leq \frac{2}{3}(1 - |w|)$. Therefore

$$\frac{1 - |w|}{|1 - w|} \geq \frac{1 - |w|}{|1 - z| + |z - w|} \geq \frac{\frac{2}{3}(1 - |z|)}{|1 - z| + 1 - |z|} \geq \frac{m}{3}.$$

LEMMA 4. (Cargo [1; 145]) *Suppose $a, w \in D$, $0 \leq \theta < 2\pi$ and $z(r) = a + r(e^{i\theta} - a)$ for $0 \leq r \leq 1$. Then*

$$\int_0^1 \frac{1 - |w|^2}{|1 - \bar{w}z(r)|^2} dr < \frac{8}{(1 - |a|)\sin\left[\frac{\pi}{2}(1 + |a|)\right]} \frac{1 - |w|}{|e^{i\theta} - w|}.$$

LEMMA 5. Let $w, a \in D$, $0 \leq \theta < 2\pi$ and b be the projection of a on $l(w, e^{i\theta})$. If $0 < m \leq \min\{(1 - |z|)/(|e^{i\theta} - z|): z \in l(a, e^{i\theta})\}$, we have

$$(2.1) \quad V(G(\cdot, w), l(a, e^{i\theta})) \leq \left(\frac{40}{\sin \frac{\pi}{2}(1 + |a|)} + \frac{12}{m} \log \left| \frac{1}{b - a} \right| \right) \frac{1}{1 - |a|} \frac{1 - |w|}{|e^{i\theta} - w|}.$$

Proof. We assume that w is not on the line through $a, e^{i\theta}$, otherwise $b = a$ and (2.1) is trivial. Also we may assume $\theta = 0$ by a suitable rotation. Let $z(r) = a + r(1 - a)$, $0 \leq r \leq 1$, be a representation of $l(a, 1)$ and $\text{grad } G$ be the gradient of Green's function with respect to the first variable, that is,

$$(2.2) \quad |\text{grad } G(z, w)| = \frac{1 - |w|^2}{|1 - \bar{w}z| |z - w|}$$

if $z \neq w$. Let $S_1 = \{r: |z(r) - w| \geq (1 - |w|)/2\} \cap [0, 1]$ and $S_2 = [0, 1] \setminus S_1$. Thus

$$\begin{aligned} V(G(\cdot, w), l(a, 1)) &\leq |1 - a| \int_{S_1 + S_2} \frac{1 - |w|^2}{|1 - \bar{w}z(r)| |z(r) - w|} dr \\ &= |1 - a| (I_1 + I_2). \end{aligned}$$

From Lemma 2 and Lemma 4 we have

$$I_1 < \frac{40}{(1 - |a|) \sin \left[\frac{\pi}{2}(1 + |a|) \right]} \frac{1 - |w|}{|1 - w|}.$$

Assume S_2 is nonempty and let α, β ($\alpha < \beta$) be the two real numbers so that the points $z(\alpha) = a + \alpha(1 - a)$ and $z(\beta) = a + \beta(1 - a)$ have distance exactly $(1 - |w|)/2$ from w . We observe that

$$\text{Im} \frac{z(\alpha) - w}{1 - a} = \text{Im} \frac{z(\beta) - w}{1 - a} = -\text{Im} \frac{w - a}{1 - a}$$

and $|\text{Im}(w - a)/(1 - a)|$ is the distance from w to $l(a, 1)$, which is $|(b - a)/(1 - a)| |1 - w|$. Therefore

$$-\text{Re} \frac{z(\alpha) - w}{1 - a} = \text{Re} \frac{z(\beta) - w}{1 - a} > 0.$$

With the aid of Lemma 3, we estimate I_2 as follows:

$$\begin{aligned}
 I_2 &\leq 2 \int_{\alpha}^{\beta} \frac{1}{|z(r) - w|} dr \\
 &= \frac{2}{|1-a|} \left[\log \left(\left| \frac{z(r) - w}{1-a} \right| + \operatorname{Re} \frac{z(r) - w}{1-a} \right) \right]_{\alpha}^{\beta} \\
 &= \frac{2}{|1-a|} \log \frac{\left| \frac{z(\beta) - w}{1-a} \right| + \operatorname{Re} \frac{z(\beta) - w}{1-a}}{\left| \frac{z(\beta) - w}{1-a} \right| - \operatorname{Re} \frac{z(\beta) - w}{1-a}} \\
 &\leq \frac{2}{|1-a|} \log \frac{4 \left| \frac{z(\beta) - w}{1-a} \right|^2}{\left(\operatorname{Im} \frac{z(\beta) - w}{1-a} \right)^2} \\
 &\leq \frac{4}{|1-a|} \log \frac{(1 - |w|)}{|b-a| |1-w|} \\
 &\leq \frac{12}{m|1-a|} \frac{1 - |w|}{|1-w|} \log \frac{1}{|b-a|}.
 \end{aligned}$$

Thus (2.1) follows from the estimations for I_1 and I_2 .

3. Proofs and remarks.

Proof of Theorem 1. For a fixed $w \in D$, let T_1 be $\{z: \rho \leq |z| < 1, |z - w| \geq (1 - |w|)/2\}$ and $T_2 = \{z: \rho \leq |z| < 1\} \setminus T_1$. We consider

$$\int_{|z| \geq \rho} |\operatorname{grad} G(z, w)| d\sigma_z = \int_{T_1} \cdot d\sigma_z + \int_{T_2} \cdot d\sigma_z = I_1 + I_2$$

where $d\sigma_z = r dr d\theta$, $z = re^{i\theta}$. From (2.2), Lemma 2 and Lemma 4 we obtain

$$I_1 \leq C \int_0^{2\pi} \frac{1 - |w|}{|e^{i\theta} - w|} d\theta$$

where C is a constant depending on ρ only. It is easy to verify that there exists a constant C_1 so that

$$I_1 \leq C_1 (1 - |w|) \log \frac{1}{1 - |w|} \quad \text{when } |w| > \frac{1}{2}$$

and

$$I_1 \leq C_1(1 - |w|) \quad \text{when} \quad |w| \leq \frac{1}{2}.$$

Noticing that $|1 - \bar{w}z| \geq 1 - |w|$, we have

$$I_2 \leq \int_{|z-w| < (1-|w|)/2} \frac{2}{|z-w|} d\sigma_z = 2\pi(1 - |w|).$$

From (1.2), (1.4) and the estimations above, we see that

$$\int_0^{2\pi} V(u, l(\rho e^{i\theta}, e^{i\theta})) d\sigma \leq \frac{1}{\rho} \int_D \int_{|z| \geq \rho} |\text{grad } G(z, w)| d\sigma_z d\mu(w) < \infty.$$

Theorem 1 follows.

Proof of Theorem 2. We may assume $\theta = 0$. Let α be the distance from 1 to L , L_0 be a closed subarc of L and $m = \text{minimum of the numbers in } \{1 - |a| : a \in L_0\} \cup \{(1 - |z|)/(1 - |z|) : z \in l(a, 1) \text{ and } a \in L_0\}$. It is clear that $m > 0$.

Let E be the set $\{a \in L_0 : V(u, l(a, 1)) = \infty\}$. We shall show E is of capacity zero. If we assume otherwise, there exist $M > 0$ and a finite positive measure $\nu(a)$ on E such that

$$\int_E \log \frac{1}{|z - a|} d\nu(a) < M$$

for every complex z . With the aid of Lemma 5 we obtain

$$\begin{aligned} & \int_E V(G(\cdot, w), l(a, 1)) d\nu(a) \\ & \leq \int_E \left(\frac{40}{\sin \frac{\pi}{2}(2 - m)} + \frac{12}{m} \log \frac{1}{|b - a|} \right) \frac{1}{m} \frac{1 - |w|}{|1 - w|} d\nu(a) \\ & < \left[\frac{40}{\sin \frac{\pi}{2}(2 - m)} \nu(E) + \frac{12M}{m} \right] \frac{1}{m} \frac{1 - |w|}{|1 - w|}. \end{aligned}$$

From Lemma 1 and the assumption of Theorem 2 we see that $\int_E V(u, l(a, 1)) d\nu(a) < \infty$ which is a contradiction. Hence E is of capacity zero. Theorem 2 is proved.

REMARK 1. The conclusion in Theorem 2 is best possible, because

for any compact set E of capacity zero in D , there is a mass distribution μ on E such that the Green's potential of μ is $+\infty$ on E and finite on $D \setminus E$ [4; 152 and 271].

REMARK 2. Carleson [2; 26] showed that (1.5) implies (1.3) for all θ except a set of α -dim Hausdorff measure zero for $0 < \alpha < 1$ or of capacity zero for $\alpha = 0$. Frostman [3] showed that (1.2) implies (1.3) for almost all θ . But we note that Theorems A and 1 do not follow from Theorem 2 for any obvious reason.

REMARK 3. If B is the Blaschke product with zeros at z_n , then $-\log |B|$ is the Green's potential given by $\nu = \sum \delta_{z_n}$ where δ_{z_n} is the unit mass at z_n . If l is a line segment in \bar{D} not passing through any z_n , then $\log(z - z_n)/(1 - \bar{z}_n z)$ is analytic in a neighborhood of l . Thus we have

$$\begin{aligned} (3.1) \quad V(\arg B, l) &\leq \sum V\left(\arg \frac{z - z_n}{1 - \bar{z}_n z}, l\right) \\ &\leq \sum \int_l \left| \operatorname{grad} \arg \frac{z - z_n}{1 - \bar{z}_n z} \right| |dz| \\ &= \int_D \int_l |\operatorname{grad} G(z, w)| |dz| d\nu(w) \end{aligned}$$

and

$$\begin{aligned} (3.2) \quad V(|B|, l) &\leq \int_l |B'(z)| |dz| \\ &\leq \int_D \int_l |\operatorname{grad} G(z, w)| |dz| d\nu(w). \end{aligned}$$

In the above theorems, when we dealt with variation of u on line segment l , we in fact studied an upper bound of it, which is $\int_D \int_l |\operatorname{grad} G(z, w)| |dz| d\mu(w)$. Therefore from (3.1) and (3.2) we see that whenever ν satisfies the same condition for μ in Theorem j , $j = 1, 2$ or A, B has the same variation property for u stated in Theorem j . With the aid of a theorem by F. Riesz [5; 401] on variation of analytic functions, we can conclude the following result of Cargo [1; 147]: if $\sum_{n=1}^{\infty} (1 - |z_n|)/|e^{i\theta} - z_n| < \infty$ then all the subproducts of B have finite variation on each line segment in D ending at $e^{i\theta}$.

4. Variation on curves. Let $0 < \eta < 1$, $0 < m < M$ and f be a one to one map from $\{\eta \leq |z| \leq 1\}$ to $\bar{D} \setminus \{0\}$ mapping $|z| = 1$ onto $|z| = 1$ and satisfying

$$(4.1) \quad m|\alpha - \beta| \leq |f(\alpha) - f(\beta)| \leq M|\alpha - \beta|$$

whenever $\eta \leq |\alpha| \leq 1, \eta \leq |\beta| \leq 1$. We have the following result.

THEOREM 3. *Let μ satisfy the condition in Theorem $j, j = 1, 2$ or A , and assume that the line segments we considered in Theorem j are short enough to be in $\{\eta < |z| \leq 1\}$. Then the variations of u on the images under f of the corresponding line segments are finite outside an exceptional set of the same size as in Theorem j .*

We note that when $j = A$ or 1 , Theorem 3 can be considered as a generalization of a theorem in [7], in the sense of the size of the exceptional set, of variation vs. limit and of the more generalized curves.

Proof. Let S be the image of f and l be any fixed closed line segment in $\{\eta < |z| \leq 1\}$. We may assume μ has support on S . In fact, if v is the Green's potential given by $\mu \mid D \setminus S$, then $v(z)$ is harmonic in the interior of S and tends to zero uniformly as z tends to the unit circle; thus $v(z)$ has finite variation on $f(l)$.

Let $g = f^{-1}$ and $\text{grad } G$ be the gradient with respect to the first variable of G . Thus from (2.2) we have

$$(4.2) \quad \frac{|\text{grad } G(f(z), w)|}{|\text{grad } G(z, g(w))|} = \frac{1 - |w|^2}{1 - |g(w)|^2} \frac{|1 - \overline{g(w)}z|}{|1 - \overline{w}f(z)|} \frac{|g(w) - z|}{|w - f(z)|}.$$

We observe that

$$(4.3) \quad \begin{aligned} \left| \frac{1 - \overline{g(w)}z}{1 - \overline{w}f(z)} \right| &\leq \frac{1 - |g(w)|^2 + |g(w)| |g(w) - z|}{|1 - \overline{w}f(z)|} \\ &\leq 2 \frac{1 - |g(w)|}{1 - |w|} + \frac{|g(w) - z|}{|w - f(z)|}. \end{aligned}$$

Combining (4.1), (4.2) and (4.3), we have

$$|\text{grad } G(f(z), w)| \leq M_1 |\text{grad } G(z, g(w))|$$

for some constant M_1 . Because $f(l)$ is Lipschitz, for fixed w , we have

$$\begin{aligned}
 (4.4) \quad V(G(z, w), f(l)) &\leq \int_{f(l)} |\operatorname{grad} G(z, w)| |dz| \\
 &\cong \int_l |\operatorname{grad} G(f(z), w)| M |dz| \\
 &\cong MM_1 \int_l |\operatorname{grad} G(z, g(w))| |dz|.
 \end{aligned}$$

From (4.4) and the proof of Theorem *j*, we see that it is enough to show

$$(4.5) \quad \int_D (1 - |g(w)|)^\alpha d\mu(w) < \infty,$$

$$(4.6) \quad \int_D (1 - |g(w)|) \log \frac{1}{1 - |g(w)|} d\mu(w) < \infty,$$

$$(4.7) \quad \int_D \frac{1 - |g(w)|}{|g(e^{i\theta}) - g(w)|} d\mu(w) < \infty,$$

for Theorems A, 1, 2 respectively. (4.5) and (4.7) are simple consequences of (4.1). We shall show (4.6). Let δ be the distance from 0 to S and assume $w \in S$. We have

$$\log \frac{1}{1 - |g(w)|} - \log \frac{1}{1 - |w|} < \log M,$$

thus

$$\log \frac{1}{1 - |g(w)|} < \left(1 + \frac{\log M}{\log \frac{1}{1 - \delta}}\right) \log \frac{1}{1 - |w|},$$

therefore

$$\begin{aligned}
 (4.8) \quad (1 - |g(w)|) \log \frac{1}{1 - |g(w)|} \\
 < \left(1 + \frac{\log M}{\log \frac{1}{1 - \delta}}\right) \frac{1}{m} (1 - |w|) \log \frac{1}{1 - |w|}.
 \end{aligned}$$

Since we assume μ has support on S , (1.2) and (4.8) imply (4.6). The proof of Theorem 3 is complete.

REFERENCES

1. G. T. Cargo, *The segmental variation of Blaschke products*, Duke Math. J., **30** (1963), 143–149.
2. L. Carleson, *On a class of meromorphic functions and its associated exceptional sets*, (1950), Thesis, University of Uppsala.
3. O. Frostman, *Sur les produits de Blaschke*, Kungl. Fysiografiska Sällskapets i Lund Förhandlingar, **12** (1942), 169–182.
4. L. L. Helms, *Introduction to potential theory*, (1969), Wiley-Interscience, New York.
5. F. Riesz, *Eine Ungleichung für harmonische Funktionen*, Monatsh. Math. Phys., **43** (1936), 401–406.
6. W. Rudin, *The radial variation of analytic functions*, Duke Math. J., **22** (1955), 235–242.
7. Jang-Mei G. Wu, *Boundary limits of Green's potentials along curves*, Studia Math., (to appear).

Received November 10, 1975 and in revised form June 1, 1976.

UNIVERSITY OF ILLINOIS—URBANA

Helen Elizabeth Adams, <i>Factorization-prime ideals in integral domains</i>	1
Patrick Robert Ahern and Robert Bruce Schneider, <i>The boundary behavior of Henkin's kernel</i>	9
Daniel D. Anderson, Jacob R. Matijevic and Warren Douglas Nichols, <i>The Krull intersection theorem. II</i>	15
Efraim Pacillas Armendariz, <i>On semiprime P.I.-algebras over commutative regular rings</i>	23
Robert H. Bird and Charles John Parry, <i>Integral bases for bicyclic biquadratic fields over quadratic subfields</i>	29
Tae Ho Choe and Young Hee Hong, <i>Extensions of completely regular ordered spaces</i>	37
John Dauns, <i>Generalized monofrom and quasi injective modules</i>	49
F. S. De Blasi, <i>On the differentiability of multifunctions</i>	67
Paul M. Eakin, Jr. and Avinash Madhav Sathaye, <i>R-endomorphisms of $R[[X]]$ are essentially continuous</i>	83
Larry Quin Eifler, <i>Open mapping theorems for probability measures on metric spaces</i>	89
Garret J. Etgen and James Pawlowski, <i>Oscillation criteria for second order self adjoint differential systems</i>	99
Ronald Fintushel, <i>Local S^1 actions on 3-manifolds</i>	111
Kenneth R. Goodearl, <i>Choquet simplexes and σ-convex faces</i>	119
John R. Graef, <i>Some nonoscillation criteria for higher order nonlinear differential equations</i>	125
Charles Henry Heiberg, <i>Norms of powers of absolutely convergent Fourier series: an example</i>	131
Les Andrew Karlovitz, <i>Existence of fixed points of nonexpansive mappings in a space without normal structure</i>	153
Gangaram S. Ladde, <i>Systems of functional differential inequalities and functional differential systems</i>	161
Joseph Michael Lambert, <i>Conditions for simultaneous approximation and interpolation with norm preservation in $C[a, b]$</i>	173
Ernest Paul Lane, <i>Insertion of a continuous function</i>	181
Robert F. Lax, <i>Weierstrass points of products of Riemann surfaces</i>	191
Dan McCord, <i>An estimate of the Nielsen number and an example concerning the Lefschetz fixed point theorem</i>	195
Paul Milnes and John Sydney Pym, <i>Counterexample in the theory of continuous functions on topological groups</i>	205
Peter Johanna I. M. De Paepe, <i>Homomorphism spaces of algebras of holomorphic functions</i>	211
Judith Ann Palagallo, <i>A representation of additive functionals on L^p-spaces, $0 < p < 1$</i>	221
S. M. Patel, <i>On generalized numerical ranges</i>	235
Thomas Thornton Read, <i>A limit-point criterion for expressions with oscillatory coefficients</i>	243
Elemer E. Rosinger, <i>Division of distributions</i>	257
Peter S. Shoenfeld, <i>Highly proximal and generalized almost finite extensions of minimal sets</i>	265
R. Sirois-Dumais and Stephen Willard, <i>Quotient-universal sequential spaces</i>	281
Robert Charles Thompson, <i>Convex and concave functions of singular values of matrix sums</i>	285
Edward D. Tymchatyn, <i>Some n-arc theorems</i>	291
Jang-Mei Gloria Wu, <i>Variation of Green's potential</i>	295