

Pacific Journal of Mathematics

A CHARACTERIZATION OF THE GAUSSIAN DISTRIBUTION IN A HILBERT SPACE

JEFFREY LYNN SPIELMAN

A CHARACTERIZATION OF THE GAUSSIAN DISTRIBUTION IN A HILBERT SPACE

JEFFREY L. SPIELMAN

In this paper we consider the case in which random variables X_j take values in a real, separable Hilbert space \mathcal{H} . We look at a linear form $\sum A_j X_j$ where each A_j is a bounded linear operator in \mathcal{H} . We then assume that this linear form is identically distributed with a monomial and form conditions under which it is possible to deduce that the common distribution of the random variables is the Gaussian distribution.

The study of identically distributed linear forms of independent and identically distributed random variables has been undertaken by several authors. J. Marcinkiewicz studied linear forms in which all moments of the random variables are assumed to exist. He then proved that the common distribution of the random variables was the Normal distribution. R. G. Laha and E. Lukacs have considered the case where one of the linear forms is a monomial. They have obtained characterizations of the Normal distribution for both the case when the variance is assumed finite and when no assumption is made concerning the variance.

1. Statement of the main result. Suppose now that X_1, X_2, \dots is a sequence (possibly finite) of independent, identically distributed, nondegenerate \mathcal{H} -valued random variables, where X_1 has a finite variance (i.e. $\text{Var } X_1 < +\infty$). Let A_1, A_2, \dots be a sequence of 1-1 bounded linear operators in \mathcal{H} , with the following two properties:

$$(1) \quad \sum_j \|A_j\|^2 < +\infty \quad \text{and} \quad \sum_j A_j^* A_j \cong I$$

and

$$(2) \quad \sup_j \|A_j\| < 1.$$

We note that in the above A_j^* represents the adjoint of A_j and that the inequality $\sum_j A_j^* A_j \cong I$ is true in the sense of positive-definiteness. (For example, see page 313 of [7].)

Our goal is to prove the following theorem.

THEOREM 1. *Suppose that $\sum_j A_j X_j$ converges with probability one. If $\sum_j A_j X_j$ has the same distribution as X_1 , then X_1 has a Gaussian distribution.*

In §2 we will prove an important preliminary result (Theorem 2). Then in §3 we will present the proof of Theorem 1.

2. A preliminary result. In this section we will prove the following result.

THEOREM 2. *Let X_1, X_2, \dots be a sequence (possibly finite) of independent, identically distributed, nondegenerate, \mathcal{H} -valued random variables. Suppose that the sum $\sum_j A_j X_j$ exists with probability one, where A_1, A_2, \dots are bounded linear operators in \mathcal{H} , with $\sup_j \|A_j\| < 1$.*

If $\sum_j A_j X_j$ has the same distribution as X_1 , then X_1 has an infinitely divisible distribution.

Note. The hypotheses of Theorem 2 are somewhat weaker than the hypotheses of Theorem 1.

Before beginning the proof of Theorem 2, let us fix some notation.

Let $\varphi(y)$ be the common characteristic functional of X_1, X_2, \dots . Then $\varphi(y) = \mathcal{E}e^{i\langle X_1, y \rangle}$ for all $y \in \mathcal{H}$, where \mathcal{E} denotes mathematical expectation.

The characteristic functional of $A_j X_j$ is then given by:

$$(3) \quad \mathcal{E}e^{i\langle A_j X_j, y \rangle} = \mathcal{E}e^{i\langle X_j, A_j^* y \rangle} = \varphi(A_j^* y)$$

where A_j^* denotes the adjoint operator of A_j .

Now, suppose that $\sum_j A_j X_j$ has the same distribution as X_1 . Then equation (3) gives us:

$$(4) \quad \varphi(y) = \prod_j \varphi(A_j^* y), \quad \text{for all } y \in \mathcal{H},$$

where the product converges uniformly on bounded spheres. (See Theorem 4.4, pg. 171 of [5].)

Since $\sum_j A_j X_j$ converges, then $\sum_{j=n}^\infty A_j X_j$ converges, with probability one, to the origin of \mathcal{H} as $n \rightarrow \infty$. (Of course, if X_1, X_2, \dots is a finite sequence, the preceding statement is unnecessary.)

Thus, it is possible to choose N_0 for any $\epsilon > 0$, such that $P\{\|\sum_{j=N+1}^\infty A_j X_j\| > \epsilon\} < \epsilon$, whenever $N \geq N_0$. Let $\varphi_N(y)$ denote the characteristic functional of $\sum_{j=N+1}^\infty A_j X_j$. Then using equation (4), we have:

$$(5) \quad \varphi(y) = \varphi(A_1^* y) \cdots \varphi(A_N^* y) \varphi_N(y).$$

Proof of Theorem 2. We assume that $\sum_j A_j X_j$ has the same distribution as X_1 . Then equation (5) holds. If we replace y by $A_j^* y$ in equation (5), we obtain:

$$(6) \quad \varphi(A_j^* y) = \varphi(A_1^* A_j^* y) \cdots \varphi(A_N^* A_j^* y) \varphi_N(A_j^* y)$$

for each $j = 1, 2, \dots, N$.

Combining equations (5) and (6) we have:

$$\varphi(y) = \prod_{j=1}^N \varphi((A_j^*)^2 y) \cdot \prod_{j \neq k} \varphi(A_j^* A_k^* y) \prod_{j=1}^N \varphi_N(A_j^* y) \varphi_N(y).$$

If we repeat the above process n times, we get the following result:

$$(7) \quad \varphi(y) = \prod \varphi(A_{j_1}^* \cdots A_{j_n}^* y) \prod_{k=1}^{n-1} \prod \varphi_N(A_{j_1}^* \cdots A_{j_{n-k}}^* y) \varphi_N(y).$$

The product on the right hand side of equation (7) consists of $N^n + N^{n-1} + \dots + N + 1$ factors, where each of the subscripts j_1, \dots, j_n can take any of the values $1, \dots, N$ with repetitions allowed.

Thus, equation (7) says that X_1 is distributed as the sum of $k_n = \sum_{k=0}^n N^k$ independent, \mathcal{H} -valued random variables, $Y_{n,k}$ ($k = 1, 2, \dots, k_n$), for any positive integer n .

We will now show that $Y_{n,k}$ is a *uniformly infinitesimal* collection of random variables. That is, we will show that for any $\epsilon > 0$, $\sup_{1 \leq k \leq k_n} P\{\|Y_{n,k}\| > \epsilon\} \rightarrow 0$ as $n \rightarrow \infty$. Once this has been established, the infinite divisibility of X_1 will follow from Corollary 6.2, page 199 of [5].

Consider the factors on the right hand side of equation (7). Let $\epsilon > 0$ be given. By definition $\varphi_N(y)$ is the characteristic functional of $\sum_{j=N+1}^\infty A_j X_j$ and $P\{\|\sum_{j=N+1}^\infty A_j X_j\| > \epsilon\} < \epsilon$, for all $N \geq N_0$.

Consider now a factor of the form $\varphi_N(A_{j_1}^* A_{j_2}^* \cdots A_{j_{n-k}}^* y)$. This is the characteristic functional of

$$A_{j_{n-k}} \cdots A_{j_2} A_{j_1} \sum_{j=N+1}^\infty A_j X_j.$$

Also,

$$\begin{aligned} P\left\{\left\|A_{j_{n-k}} \cdots A_{j_2} A_{j_1} \sum_{j=N+1}^\infty A_j X_j\right\| > \epsilon\right\} \\ \leq P\left\{\|A_{j_{n-k}}\| \cdots \|A_{j_2}\| \cdot \|A_{j_1}\| \cdot \left\|\sum_{j=N+1}^\infty A_j X_j\right\| > \epsilon\right\} \\ \leq P\left\{\left\|\sum_{j=N+1}^\infty A_j X_j\right\| > \epsilon\right\} < \epsilon \quad \text{whenever } N \geq N_0, \end{aligned}$$

since $\sup_j \|A_j\| < 1$.

Finally, we consider a factor of the form $\varphi(A_{j_1}^* \cdots A_{j_n}^* y)$, which is the characteristic functional of $A_{j_n} \cdots A_{j_1} X_1$. Set $\alpha = \sup_j \|A_j\|$.

Then

$$\begin{aligned} P\{\|A_{j_n} \cdots A_{j_1} X_1\| > \epsilon\} \\ &\leq P\{\|A_{j_n}\| \cdots \|A_{j_1}\| \cdot \|X_1\| > \epsilon\} \\ &\leq P\left\{\|X_1\| > \frac{\epsilon}{\alpha^n}\right\}. \end{aligned}$$

Now choose an integer N' such that $P\{\|X_1\| > \epsilon/\alpha^n\} < \epsilon$, whenever $n \geq N'$. (This is possible because $0 < \alpha < 1$). Set $n_0 = \max\{N_0, N'\}$.

Hence, we have shown that $P\{\|Y_{nk}\| > \epsilon\} < \epsilon$, for all $k = 1, 2, \dots, k_n$, whenever $n \geq n_0$. Therefore, the collection Y_{nk} is uniformly infinitesimal and X_1 is infinitely divisible. This completes the proof of the theorem.

3. Proof of the main result. For convenience, we now will make the assumption that X_1, X_2, \dots are symmetric random variables. Since the common distribution of these random variables is infinitely divisible, the common characteristic functional, $\varphi(y)$, has a unique Levy-Khintchine representation given by:

$$(8) \quad \ln \varphi(y) = -\frac{1}{2} \langle Sy, y \rangle + \int (\cos \langle x, y \rangle - 1) dL(x)$$

where S is an S -operator (a nonnegative, self-adjoint compact operator on \mathcal{H} , with a finite trace), and L is a σ -finite measure with finite mass outside every neighborhood of the origin and with the property that

$$\int_{\|x\| \leq 1} \|x\|^2 dL(x) < +\infty.$$

(see [5], page 181.)

Furthermore, since X_1, X_2, \dots have finite variance, $\varphi(y)$ has a unique Kolmogorov representation, given by:

$$(9) \quad \ln \varphi(y) = -\frac{1}{2} \langle Sy, y \rangle + \int_{\mathcal{H} \setminus \{0\}} \frac{\cos \langle x, y \rangle - 1}{\|x\|^2} dK(x)$$

where S is an S -operator and K is a finite measure on \mathcal{H} . (See [6].)

By equations (4) and (8) we have:

$$\begin{aligned} -\frac{1}{2} \sum_j \langle SA_j^* y, A_j^* y \rangle + \sum_j \int (\cos \langle x, A_j^* y \rangle - 1) dL(x) \\ = -\frac{1}{2} \langle Sy, y \rangle + \int (\cos \langle x, y \rangle - 1) dL(x). \end{aligned}$$

Also,

$$\begin{aligned}
 & -\frac{1}{2} \sum_j \langle SA_j^* y, A_j^* y \rangle + \sum_j \int (\cos \langle x, A_j^* y \rangle - 1) dL(x) \\
 (10) \quad & = -\frac{1}{2} \left\langle \sum_j A_j SA_j^* y, y \right\rangle + \sum_j \int (\cos \langle A_j x, y \rangle - 1) dL(x) \\
 & = -\frac{1}{2} \left\langle \sum_j A_j SA_j^* y, y \right\rangle + \sum_j \int (\cos \langle x, y \rangle - 1) dLA_j^{-1}(x).
 \end{aligned}$$

It is not difficult to show that $\sum_j A_j SA_j^*$ is an S -operator. Also, it is clear that LA_j^{-1} is the σ -finite measure which occurs in the Levy-Khintchine representation of $A_j X_j$, for each j .

We denote by \mathcal{B} , the class of Borel sets in \mathcal{H} . Then the measure K_j , defined by:

$$K_j(D) = \int_D \|x\|^2 dLA_j^{-1}(x), \quad \text{for all } D \in \mathcal{B}$$

is the finite measure which occurs in the Kolmogorov representation of $A_j X_j$, for each j .

Since X_1, X_2, \dots have finite variance,

$$(11) \quad \int \|x\|^2 dL(x) < +\infty. \quad (\text{See [6].})$$

By equation (10), $\ln \Pi_j \varphi(A_j^* y)$

$$(12) \quad = -\frac{1}{2} \left\langle \sum_j A_j SA_j^* y, y \right\rangle + \sum_j \int_{\{x \neq 0\}} \frac{(\cos \langle x, y \rangle - 1)}{\|x\|^2} dK_j(x).$$

We note that

$$\begin{aligned}
 & \sum_j \int_{\{x \neq 0\}} \left| \frac{\cos \langle x, y \rangle - 1}{\|x\|^2} \right| dK_j(x) \leq \sum_j \int_{\{x \neq 0\}} \frac{\|x\|^2 \|y\|^2}{\|x\|^2} dK_j(x) \\
 & = \|y\|^2 \sum_j \int_{\{x \neq 0\}} dK_j(x) = \|y\|^2 \sum_j \int \|x\|^2 dLA_j^{-1}(x) \\
 & = \|y\|^2 \sum_j \int \|A_j x\|^2 dL(x) \leq \left(\sum_j \|A_j\|^2 \right) \|y\|^2 \int \|x\|^2 dL(x) < \infty
 \end{aligned}$$

because of relations (1) and (11).

Thus we may interchange the integral and summation signs in equation (12) to obtain:

$$\ln \prod_j \varphi(A_j^* y) = -\frac{1}{2} \left\langle \sum_j A_j S A_j^* y, y \right\rangle + \int_{\{x \neq 0\}} \frac{\cos \langle x, y \rangle - 1}{\|x\|^2} d\left(\sum_j K_j(x)\right).$$

Then, by the uniqueness of the Kolmogorov representation, we have:

$$\sum_j A_j S A_j^* = S \quad \text{and} \quad \sum_j K_j = K.$$

From the second of these relations, $\sum_j K_j(\mathcal{H}) = K(\mathcal{H})$, which leads to the following sequence of equations.

$$\begin{aligned} \sum_j \int \|x\|^2 dL A_j^{-1}(x) &= \int \|x\|^2 dL(x) \\ \sum_j \int \|A_j x\|^2 dL(x) &= \int \|x\|^2 dL(x) \\ \int \left[\sum_j \|A_j x\|^2 - \|x\|^2 \right] dL(x) &= 0 \\ \int \left[\left\langle \sum_j A_j^* A_j x, x \right\rangle - \langle x, x \rangle \right] dL(x) &= 0. \end{aligned}$$

In view of relation (4), it must then be true that

$$(13) \quad L \left\{ x : \sum_j \|A_j x\|^2 - \|x\|^2 > 0 \right\} = 0.$$

We note that for n a positive integer, $\sum_{j=1}^n A_j X_j$ has characteristic functional $\prod_{j=1}^n \varphi(A_j^* y)$, and

$$(14) \quad \begin{aligned} \ln \prod_{j=1}^n \varphi(A_j^* y) &= -\frac{1}{2} \left\langle \sum_{j=1}^n A_j S A_j^* y, y \right\rangle \\ &+ \int (\cos \langle x, y \rangle - 1) d\left(\sum_{j=1}^n L A_j^{-1}(x)\right). \end{aligned}$$

Thus $\sum_{j=1}^n L A_j^{-1}$ converges weakly, outside closed neighborhoods of $0 \in \mathcal{H}$, to L , as $n \rightarrow \infty$. (See [5], page 189).

It now becomes necessary to state and prove two technical lemmas.

LEMMA 1. For any $\epsilon > 0$,

$$\int_{\|x\| > \epsilon} \|x\|^2 dL(x) = \sum_j \int_{\|A_j x\| > \epsilon} \|A_j x\|^2 dL(x).$$

Proof. Let ϵ_1 and ϵ_2 be positive constants with $\epsilon_1 < \epsilon_2$. Define a function $f(x)$ by:

$$f(x) = \begin{cases} \|x\|^2, & \epsilon_1 < \|x\| \leq \epsilon_2 \\ (\epsilon_2)^2, & \|x\| > \epsilon_2. \end{cases}$$

Then $f(x)$ is bounded and continuous. Thus by comment (14),

$$\int_{\|x\| > \epsilon_1} f(x) dL(x) = \lim_{n \rightarrow \infty} \sum_{j=1}^n \int_{\|x\| > \epsilon_1} f(x) dLA_j^{-1}(x)$$

which implies that

$$\sum_{j=1}^n \int_{\epsilon_1 < \|x\| \leq \epsilon_2} \|x\|^2 dLA_j^{-1}(x) + (\epsilon_2)^2 \sum_{j=1}^n LA_j^{-1}\{\|x\| > \epsilon_2\}$$

converges to

$$\int_{\epsilon_1 < \|x\| \leq \epsilon_2} \|x\|^2 dL(x) + (\epsilon_2)^2 L\{\|x\| > \epsilon_2\}$$

as $n \rightarrow \infty$.

But, again because of comment (14), $\sum_{j=1}^n LA_j^{-1}\{\|x\| > \epsilon_2\}$ converges to $L\{\|x\| > \epsilon_2\}$ as $n \rightarrow \infty$.

Therefore, $\sum_{j=1}^n \int_{\epsilon_1 < \|x\| \leq \epsilon_2} \|x\|^2 dLA_j^{-1}(x)$ converges to

$$\int_{\epsilon_1 < \|x\| \leq \epsilon_2} \|x\|^2 dL(x),$$

whenever we choose $0 < \epsilon_1 < \epsilon_2$.

Let $\epsilon > 0$ be given. Let ϵ_n be a strictly increasing sequence of positive numbers, $\epsilon_n \uparrow +\infty$, with $\epsilon_1 > \epsilon$. For convenience we set $\epsilon = \epsilon_0$.

Then

$$\begin{aligned} \int_{\|x\| > \epsilon_0} \|x\|^2 dL(x) &= \sum_{k=0}^{\infty} \int_{\epsilon_k < \|x\| \leq \epsilon_{k+1}} \|x\|^2 dL(x) \\ &= \sum_{k=0}^{\infty} \sum_j \int_{\epsilon_k < \|x\| < \epsilon_{k+1}} \|x\|^2 dLA_j^{-1}(x) \\ &= \sum_j \sum_{k=0}^{\infty} \int_{\epsilon_k < \|x\| \leq \epsilon_{k+1}} \|x\|^2 dLA_j^{-1}(x) \\ &= \sum_j \int_{\|x\| > \epsilon} \|x\|^2 dLA_j^{-1}(x) = \sum_j \int_{\|A_j x\| > \epsilon} \|A_j x\|^2 dL(x). \end{aligned}$$

This completes the proof.

LEMMA 2. $L(\{x: \|A_k x\|^2 \geq \|x\|^2, \text{ for some } k = 1, 2, \dots\}) = L(\{0\})$.

Proof. Let k be a fixed positive integer.

Set $E_k = \{x : \|A_k x\|^2 = \|x\|^2\}$. Then, using equation (13), $L(E_k) = L(E_k \cap \{x : \sum_j \|A_j x\|^2 = \|x\|^2\})$.

Thus, $L(E_k) = L\{x : \sum_{j \neq k} \|A_j x\|^2 = 0\} = L(\{0\})$, since each operator A_j is 1-1.

$$(15) \quad \text{Similarly, } L\left(\bigcup_k E_k\right) = L(\{0\}).$$

Using the same type of argument, it is easy to show that for all $k = 1, 2, \dots$

$$(16) \quad L\{x : \|A_k x\|^2 > \|x\|^2\} = 0.$$

Combining equations (15) and (16) we are done.

From relations (1) and (13), we see that $L\{x : \|x\|^2 \neq \sum_j \|A_j x\|^2\} = 0$. Hence, referring to Lemma 1, it is true that, for all $\epsilon > 0$,

$$\sum_j \int_{\|x\| > \epsilon} \|A_j x\|^2 dL(x) = \sum_j \int_{\|A_j x\| > \epsilon} \|A_j x\|^2 dL(x),$$

and this implies that

$$(17) \quad \sum_j \left[\int_{\|x\| > \epsilon} \|A_j x\|^2 dL(x) - \int_{\|A_j x\| > \epsilon} \|A_j x\|^2 dL(x) \right] = 0, \text{ for all } \epsilon > 0.$$

But

$$(18) \quad L\{x : x \neq 0 \text{ and } \|A_j x\| \geq \|x\|\} = 0, \text{ for all } j.$$

Thus, each term in the sum of equation (17) must be nonnegative, which yields:

$$\int_{\|x\| > \epsilon} \|A_j x\|^2 dL(x) = \int_{\|A_j x\| > \epsilon} \|A_j x\|^2 dL(x), \text{ for all } \epsilon > 0 \text{ and all } j.$$

Or, using equation (18),

$$\int_{\{\|x\| > \epsilon\} \cap F_j} \|A_j x\|^2 dL(x) = \int_{\{\|A_j x\| > \epsilon\} \cap F_j} \|A_j x\|^2 dL(x),$$

for all $\epsilon > 0$ and all j , where $F_j = \{x : \|A_j x\| < \|x\|\}$, for each $j = 1, 2, \dots$.

The above implies that

$$\int_{\{\|x\|>\epsilon\} \cap \{\|A_j x\| \leq \epsilon\} \cap F_j} \|A_j x\|^2 dL(x) = 0, \quad \text{for all } \epsilon > 0 \text{ and all } j,$$

or,

$$\int_{\{\|x\|>\epsilon \text{ and } \|A_j x\| \leq \epsilon\}} \|A_j x\|^2 dL(x) = 0, \quad \text{for all } \epsilon > 0 \text{ and all } j.$$

So, we must have that

$$(19) \quad L\{x: \|x\| > \epsilon \text{ and } \|A_j x\| \leq \epsilon\} = 0, \quad \text{for all } \epsilon > 0 \text{ and all } j.$$

Consider the set Q^+ of positive rational numbers. Let k be a fixed positive integer.

$$L\left[\bigcup_{r \in Q^+} \{x: \|x\| > r \text{ and } \|A_k x\| \leq r\}\right] = L\{x: \|A_k x\| < \|x\|\} = L[\mathcal{H} \setminus \{0\}].$$

Therefore, $L[\mathcal{H} \setminus \{0\}] \leq \sum_{r \in Q^+} L\{x: \|x\| > r \text{ and } \|A_k x\| \leq r\} = 0$, by equation (19).

This last relation says that L is degenerate at $0 \in \mathcal{H}$, which means that the common characteristic functional of X_1, X_2, \dots is given by:

$$\ln \varphi(y) = -\frac{1}{2} \langle Sy, y \rangle \quad (\text{see Eq. (8)}).$$

Hence X_1, X_2, \dots have a common Gaussian distribution.

Recall that we have assumed X_1, X_2, \dots to be symmetric, but it is now easy to extend our result to the general case by using Cramer's Theorem (see page 141 of [1]).

The proof of Theorem 1 is now completed.

REFERENCES

1. U. Grenander, *Probabilities on Algebraic Structures*, Wiley, New York (1963).
2. R. G. Laha, *On a property of infinitely divisible distributions in a Hilbert space*, Acta Math. Acad. Sci. Hungar. **20** (1969), 143–147.
3. R. G. Laha and E. Lukacs, *On a linear form whose distribution is identical with that of a monomial*, Pacific J. Math., **15** (1965), 207–214.
4. J. Marcinkiewicz, *Sur une propriété de la loi de Gauss*, Math. Zeitschr., **44** (1949), 612–618.
5. K. R. Parthasarathy, *Probability Measures on Metric Spaces*, Academic Press, New York (1967).
6. B. L. S. Prakasa Rao, *Some characterization theorems for Wiener processes in a Hilbert space*, Z. Wahr. verw. Geb., **19** (1971), 103–116.
7. W. Rudin, *Functional Analysis*, McGraw-Hill, New York (1973).

Received April 2, 1976.

CHICAGO STATE UNIVERSITY

CHICAGO, IL 60628

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. A. BEAUMONT
University of Washington
Seattle, WA 98105

R. FINN AND J. MILGRAM
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
• • •
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$ 72.00 a year (6 Vols., 12 issues). Special rate: \$ 36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1977 Pacific Journal of Mathematics
All Rights Reserved

William Allen Adkins, Aldo Andreotti and John Vincent Leahy, <i>An analogue of Oka's theorem for weakly normal complex spaces</i>	297
Ann K. Boyle, M. G. Deshpande and Edmund H. Feller, <i>On nonsingularly k-primitive rings</i>	303
Rolando Basim Chuaqui, <i>Measures invariant under a group of transformations</i>	313
Wendell Dan Curtis and Forrest Miller, <i>Gauge groups and classification of bundles with simple structural group</i>	331
Garret J. Etgen and Willie Taylor, <i>The essential uniqueness of bounded nonoscillatory solutions of certain even order differential equations</i>	339
Paul Ezust, <i>On a representation theory for ideal systems</i>	347
Richard Carl Gilbert, <i>The deficiency index of a third order operator</i>	369
John Norman Ginsburg, <i>S-spaces in countably compact spaces using Ostaszewski's method</i>	393
Basil Gordon and S. P. Mohanty, <i>On a theorem of Delaunay and some related results</i>	399
Douglas Lloyd Grant, <i>Topological groups which satisfy an open mapping theorem</i>	411
Charles Lemuel Hagopian, <i>A characterization of solenoids</i>	425
Kyong Taik Hahn, <i>On completeness of the Bergman metric and its subordinate metrics. II</i>	437
G. Hochschild and David Wheeler Wigner, <i>Abstractly split group extensions</i>	447
Gary S. Itzkowitz, <i>Inner invariant subspaces</i>	455
Jiang Luh and Mohan S. Putcha, <i>A commutativity theorem for non-associative algebras over a principal ideal domain</i>	485
Donald J. Newman and A. R. Reddy, <i>Addendum to: "Rational approximation of e^{-x} on the positive real axis"</i>	489
Akio Osada, <i>On the distribution of a-points of a strongly annular function</i>	491
Jeffrey Lynn Spielman, <i>A characterization of the Gaussian distribution in a Hilbert space</i>	497
Robert Moffatt Stephenson Jr., <i>Symmetrizable-closed spaces</i>	507
Peter George Trotter and Takayuki Tamura, <i>Completely semisimple inverse Δ-semigroups admitting principal series</i>	515
Charles Irvin Vinsonhaler and William Jennings Wickless, <i>Torsion free abelian groups quasi-projective over their endomorphism rings</i>	527
Frank Arvey Wattenberg, <i>Topologies on the set of closed subsets</i>	537
Richard A. Zalik, <i>Integral representation of Tchebycheff systems</i>	553