

Pacific Journal of Mathematics

EQUATIONS OF MEAN CURVATURE TYPE IN 2 INDEPENDENT VARIABLES

LEON M. SIMON

EQUATIONS OF MEAN CURVATURE TYPE IN 2 INDEPENDENT VARIABLES

LEON SIMON

The object of this paper is to develop a regularity theory for equations of mean curvature type in two independent variables. An equation of mean curvature type in two independent variables is defined to be an equation of the form

$$\sum_{i,j=1}^2 a_{ij}(x, u, Du) D_{ij}u = b(x, u, Du)$$

on a domain $\Omega \subset \mathbb{R}^2$, where the functions a_{ij} , b satisfy special structural conditions. Namely, we require that (i) $(1 + |Du|^2)^{-1/2}b(x, u, Du)$ is bounded by a fixed constant (independent of u), and (ii) the quadratic form $\sum_{i,j=1}^2 a_{ij}(x, u, Du)\xi_i\xi_j$ is bounded from above and below in terms of the quadratic form $\sum_{i,j=1}^2 g^{ij}(Du)\xi_i\xi_j$, where $g^{ij}(Du) = \delta_{ij} - D_iu D_ju / (1 + |Du|^2)$, $i, j = 1, 2$, are the coefficients of the minimal surface equation.

R. Finn [2] was the first to consider such equations; he considered the case $b \equiv 0$ and $a_{ij}(x, u, Du) \equiv a_{ij}(Du)$. Later Jenkins [5] and Jenkins-Serrin [6] specialized further to equations which arise as the non-parametric Euler-Lagrange equation of a parametric elliptic functional with integrand independent of the spatial variables (see Appendix 1). The main results in [2] concerned a-priori estimates for the gradient of a solution. In [5], [6] somewhat deeper results were obtained; in particular, pointwise estimates for the principal curvatures of the graph of a solution were established. Recently J. Spruck [11] obtained such a pointwise curvature estimate for the constant mean curvature equation; this was the first such result obtained for a non-homogeneous (i.e. b not identically zero) equation of mean curvature type.

In this paper we intend to use the Hölder estimate established in [8] in order to obtain a strong regularity theory for the entire class of equations of mean curvature type. The plan of the paper is as follows. In §1 we introduce the class of equations of mean curvature type and give a geometric characterization of such equations. In §2 we discuss application of the results of [8] to *homogeneous* equations of mean curvature type; in particular we obtain some a-priori gradient estimates, a Bernstein type theorem, a Bers-type theorem concerning the limiting behaviour of the gradient of solutions defined outside a compact set, a global Hölder continuity estimate for solutions which continuously attain Lipschitz boundary

values on $\partial\Omega$, a pointwise estimate for the principal curvatures of the graph of a solution, and a theorem concerning the removability of isolated singularities. Except for the pointwise curvature estimate, all of these results are obtained without any continuity restrictions on the coefficient functions a_{ij} . In §3 we discuss extensions of the results of §2 to the nonhomogeneous case.

1. Preliminaries. By an equation of mean curvature type, we mean an equation of the form

$$(1.1) \quad \sum_{i,j=1}^2 a_{ij}(x, u, Du) D_{ij}u = b(x, u, Du)$$

on a domain $\Omega \subset \mathbf{R}^2$, where a_{ij} , $i, j = 1, 2$, and b are given real-valued functions on $\Omega \times \mathbf{R} \times \mathbf{R}^2$ with

$$(1.2) \quad |\xi|^2 - \frac{(\xi \cdot p)^2}{1 + |p|^2} \leq \sum_{i,j=1}^2 a_{ij}(x, z, p) \xi_i \xi_j \leq \gamma \left(|\xi|^2 - \frac{(\xi \cdot p)^2}{1 + |p|^2} \right)$$

for all $(x, z, p) \in \Omega \times \mathbf{R} \times \mathbf{R}^2$ and all $\xi = (\xi_1, \xi_2) \in \mathbf{R}^2$; and

$$(1.3) \quad |b(x, z, p)| \leq \mu \sqrt{1 + |p|^2}$$

for all $(x, z, p) \in \Omega \times \mathbf{R} \times \mathbf{R}^2$. Here γ and μ denote fixed constants.

Note that the minimal surface equation can be written in the form (1.1) with $a_{ij}(x, z, p) = \delta_{ij} - p_i p_j / (1 + |p|^2)$ and with $b \equiv 0$. In this case (1.2), (1.3) hold with $\gamma = 1$ and $\mu = 0$. More generally, any equation which arises as the non-parametric Euler-Lagrange equation of a parametric elliptic functional (see Appendix 1) is of the form (1.1), (1.2), (1.3). But quite apart from these examples, the equations of mean curvature type are both natural and interesting in that they are completely characterized as follows:

Suppose u is a $C^2(\Omega)$ function with graph

$$M = \{(x, z) : x \in \Omega, z = u(x)\}.$$

Then there exists real-valued functions α_{ij} , b such that (1.1)-(1.3) hold if and only if the principal curvatures κ_1, κ_2 of M are related at each point of M by an equation of the form

$$(1.1)' \quad \alpha_1 \kappa_1 + \alpha_2 \kappa_2 = \beta,$$

with $\alpha_1, \alpha_2, \beta$ satisfying

$$(1.2)' \quad 1 \leq \alpha_i \leq \gamma, \quad i = 1, 2,$$

$$(1.3)' \quad |\beta| \leq \mu.$$

To demonstrate that this characterization is valid, we let d denote

the distance function of M defined for $X = (x, z) \in \Omega \times \mathbb{R}$ by setting $d(X) = \text{dist}(X, M)$ if $z > u(x)$ and $d(X) = -\text{dist}(X, M)$ if $z < u(x)$. Since d is C^2 and $d(x, u(x)) \equiv 0$, $x \in \Omega$, we then have, by the chain rule, the identities $D_i d(X) + D_i u(x) D_3 d(X) = 0$ and

$$(1.4) \quad \begin{aligned} & D_{ij} d(X) + D_i u(x) D_{3j} d(X) + D_j u(x) D_{3i} d(X) \\ & + D_i u(x) D_j u(x) D_{33} d(X) + D_3 d(X) D_{ij} u(x) = 0, \end{aligned}$$

$i, j = 1, 2$, where $X = (x, u(x))$. Since $D_3 d(X) = v^{-1}$, $v = \sqrt{1 + |Du(x)|^2}$, (1.1) then implies

$$(1.5) \quad \sum_{i,j=1}^3 a_{ij}^*(x) D_{ij} d(X) + b^*(x) = 0,$$

where $b^*(x) = v^{-1}b(x, u(x), Du(x))$ and where the 3×3 matrix $(a_{ij}^*(x))$ is defined by setting $a_{ij}^*(x) = a_{ij}(x, u(x), Du(x))$ for $i, j = 1, 2$ and

$$(1.6) \quad \begin{aligned} \alpha_{33}^*(x) &= \alpha_{3i}^*(x) = \sum_{j=1}^2 D_j u(x) a_{ij}^*(x), \quad i = 1, 2, \\ \alpha_{33}^* &= \sum_{i,j=1}^2 D_i u(x) D_j u(x) a_{ij}^*(x). \end{aligned}$$

Note that these last relations are equivalent to

$$(1.6)' \quad \sum_{j=1}^3 \alpha_{ji}^*(x) \nu_j = \sum_{j=1}^3 \alpha_{ij}^*(x) \nu_j = 0, \quad i = 1, 2, 3,$$

where $\nu = v^{-1}(-Du(x), 1) (= Dd(X))$ is the upward unit normal of M . Next we let Q be the matrix with rows e_1, e_2, ν , where e_1, e_2 are principal directions of M at X , so that $Q(D_{ij} d(X))Q^t = \text{diag}[\kappa_1, \kappa_2, 0]$, where κ_1, κ_2 are principal curvatures of M at X . Thus (1.5) can be written in the form (1.1)', with α_1, α_2 the first two elements on the leading diagonal of $Q(a_{ij}^*(x))Q^t$ and with $\beta = b^*(x)$. (1.3)' is now true by (1.3). To check (1.2)', we first note that, by (1.6),

$$\sum_{i,j=1}^3 a_{ij}^*(x) \xi_i \xi_j = \sum_{i,j=1}^2 \alpha_{ij}^*(x) (\xi_i + \xi_3 D_i u(x)) (\xi_j + \xi_3 D_j u(x)), \quad \xi \in \mathbb{R}^3,$$

and it then follows from (1.2) that

$$|\xi'|^2 \leq \sum_{i,j=1}^3 \alpha_{ij}^*(x) \xi_i \xi_j \leq \gamma |\xi'|^2, \quad \xi' = \xi - (\nu \cdot \xi) \nu,$$

where $\nu = v^{-1}(-Du(x), 1)$. (1.2)' easily follows from this.

To prove the converse implication we suppose that (1.1)', (1.2)', (1.3)' hold at $X = (x, u(x)) \in M$, we let $(a_{ij}^*(x)) = Q^t \text{diag}[\alpha_1, \alpha_2, 0]Q$, where Q is as above, and let $b^*(x) = \beta$. Then (1.5) holds and consequently, since we still have the relations (1.6), (1.6)', an application of (1.4) yields

$$\sum_{i,j=1}^2 a_{ij}^*(x) D_{ij} u + b^*(x) = 0 .$$

We then define, for $i, j = 1, 2$,

$$a_{ij}(x, z, p) = \begin{cases} a_{ij}^*(x) & \text{if } z = u(x) \text{ and } p = Du(x) \\ \delta_{ij} - \frac{p_i p_j}{1 + |p|^2} & \text{otherwise ,} \end{cases}$$

and

$$b(x, z, p) = \begin{cases} b^*(x) & \text{if } z = u(x) \text{ and } p = Du(x) \\ 0 & \text{otherwise .} \end{cases}$$

(1.1) (1.3) are now easily checked.

Notice that if we square each side of (1.1)' and divide by $\alpha_1 \alpha_2$, then we obtain

$$\frac{\alpha_1}{\alpha_2} \kappa_1^2 + \frac{\alpha_2}{\alpha_1} \kappa_2^2 = -2\kappa_1 \kappa_2 + \frac{\beta^2}{\alpha_1 \alpha_2} ,$$

and by (1.2)', (1.3)' this gives

$$(1.7) \quad \kappa_1^2 + \kappa_2^2 \leq A_1 \kappa_1 \kappa_2 + A_2 , \quad A_1 = -2\gamma, A_2 = \gamma \mu^2 .$$

This last inequality asserts precisely that the Gauss map of the graph of M is (A_1, A_2) -quasiconformal in the sense of [8]. (See [8], (1.8), (1.9).) In particular the Gauss map is $(A_1, 0)$ -quasiconformal, with $A_1 = -2\gamma$, in case $b \equiv 0$ (for then we can set $\mu = 0$). These observations are the key in applying the results of [8] to the equations of mean curvature type.

2. The homogeneous case ($b \equiv 0$). Throughout this section it is assumed that u is a $C^2(\Omega)$ solution of (1.1), that $b \equiv 0$ on $\Omega \times \mathbf{R} \times \mathbf{R}^2$, and that (1.2) holds.

M will denote the graph of u ; that is

$$M = \{X = (x, u(x)): x \in \Omega\} .$$

v and $\nu = (\nu_1, \nu_2, \nu_3)$ will denote the functions defined on M by

$$v(X) = \sqrt{1 + |Du(x)|^2} , \quad X = (x, u(x)) \in M$$

and

$$\nu(X) = v^{-1}(-Du(x), 1) , \quad X = (x, u(x)) \in M .$$

(Thus ν is just the upward unit normal of M .)

$\kappa_1(X), \kappa_2(X)$ will denote the principal curvatures of M at X .

x_0 will denote a fixed point of Ω .

X_0 will denote the point $(x_0, u(x_0)) \in M$.

$D_\rho(x_0) = \{x \in \mathbf{R}^2: |x - x_0| < \rho\}$;

$S_\rho(X_0) = \{X \in M: |X - X_0| < \rho\} = M \cap B_\rho(X_0)$,

where

$B_\rho(X_0) = \{X \in \mathbf{R}^3: |X - X_0| < \rho\}$.

We will begin by listing some results which follow directly from [8] §3, 4 (by virtue of the remarks at the end of §1 above).

THEOREM 1. (See Theorem (4.2) of [8].) *If $D_\rho(x_0) \subset \Omega$, then*

$$\sup_{S_{\rho/2}(X_0)} v \leq c \inf_{S_{\rho/2}(X_0)} v,$$

where $c > 0$ depends only on γ .

THEOREM 2. (See Corollary (4.2) of [8].) *If $u \geq 0$ on $D_\rho(x_0)$, then*

$$|Du(x_0)| \leq c_1 \exp \{c_2 u(x_0)/\rho\},$$

where c_1, c_2 depend only on γ .

THEOREM 3. (See Theorem (4.3) of [8].) *If $D_\rho(x_0) \subset \Omega$, then*

$$|\nu(X) - \nu(\bar{X})| \leq c(v(X_0))^{-1} \left\{ \frac{|X - \bar{X}|}{\rho} \right\}^\alpha, \quad X, \bar{X} \in S_{\rho/2}(X_0),$$

where $c > 0$ and $\alpha \in (0, 1)$ depend only on γ .

THEOREM 4. (See Theorem (4.1) of [8].) *If $\Omega = \mathbf{R}^2$, then u is linear.*

(Note that Theorem 4 follows directly from Theorem 3 by letting $\rho \rightarrow \infty$.)

THEOREM 5. (See Theorem (4.4) of [8].) *Suppose u extends continuously to $\bar{\Omega}$, suppose φ is a Lipschitz function on \mathbf{R}^2 with $\sup |\nabla \varphi| \leq L$. Then, if $\lim_{\substack{x \rightarrow y \\ x \in \Omega}} u(x) = \varphi(y)$ at each $y \in \partial\Omega$, we have*

$$|u(x) - u(\bar{x})| \leq c\{M^{1-\alpha} + |x - \bar{x}|^{1-\alpha}\} |x - \bar{x}|^\alpha, \quad x, \bar{x} \in \Omega$$

where $M = \sup_\Omega |u - \varphi|$ and $c > 0, \alpha \in (0, 1)$ are constants depending only on L .

Notice that there is absolutely no dependence on the domain Ω in the estimate above. We should point out also that from Theorem 5 various other continuity estimates follow. (See Theorems 3 and 4 of [10].)

We now wish to mention some additional results which do not quite directly follow from [8]. First we have the following theorem, which is an analogue of a theorem established by Bers [1] for solutions of the minimal surface equation.

THEOREM 6. *Suppose $\Omega = \mathbf{R}^2 \sim K$, where K is compact. Then there is a vector $a \in \mathbf{R}^2$ such that $Du(x) \rightarrow a$ uniformly for $|x| \rightarrow \infty$.*

A somewhat stronger result than Theorem 6 will be established in Theorem 6' of §2; in Theorem 6' the condition that $b \equiv 0$ will be replaced by the requirement that b has sufficiently rapid convergence to zero as $|x| \rightarrow \infty$.

Next we have a theorem concerning removability of isolated singularities. Such a theorem was proved by Bers [1] for solutions of the minimal surface equations and by Finn [3] for a class of divergence-form equations.

THEOREM 7. *Suppose $D_\rho(x_0) \sim \{x_0\} \subset \Omega$. Then u extends to be a $C^{1,\alpha}(D_\rho(x_0)) \cap W^{2,2}(D_\rho(x_0))$ function, where $\alpha \in (0, 1)$ depends only on γ .*

For a proof of this theorem the reader is referred to [9].

We will conclude this section with a pointwise curvature estimate of a type that was established by Heinz [4] for solutions of the minimal surface equation and by Jenkins [5] and Jenkins-Serrin [6] for a special class of equations of mean curvature type. In order to conveniently describe the restrictions on the coefficient functions a_{ij} which are needed here, it is necessary to introduce some further notation. We define a 3×3 matrix $(a_{ij}^*(X, \mu))$, where $X = (x, z) \in \Omega \times \mathbf{R}$ and $\mu = (1 + |p|^2)^{-1/2}(-p, 1)$ with $p \in \mathbf{R}^2$, by

$$(2.1) \quad \begin{cases} a_{ij}^*(X, \mu) = a_{ij}(x, z, p), & i, j = 1, 2, \\ a_{3i}^*(X, \mu) = a_{i3}^*(X, \mu) = \sum_{j=1}^2 a_{ij}(x, z, p)p_j, \\ a_{33}^*(X, \mu) = \sum_{i,j=1}^2 a_{ij}(x, z, p)p_i p_j. \end{cases}$$

(Cf. the functions $a_{ij}^*(x)$ of (1.6), (1.6)'). Notice that $a_{ij}^*(X, \mu)$ is thus defined for $\mu \in S_+^2$, where

$$(2.2) \quad S_+^2 = \{q = (q_1, q_2, q_3) \in \mathbf{R}^3: |q| = 1, q_3 > 0\}.$$

In the case when (1.1) arises as the nonparametric Euler-Lagrange equation of a parametric elliptic elliptic functional, the matrix $(a_{ij}^*(X, \mu))$ arises quite naturally (see Appendix 1).

THEOREM 8. *Suppose $D_\rho(x_0) \subset \Omega$ and*

$$(2.3) \quad \sum_{i,j=1}^3 |a_{ij}^*(X, \mu) - a_{ij}^*(\bar{X}, \bar{\mu})| \leq \delta \{|X - \bar{X}|/\rho + |\mu - \bar{\mu}|\}^\alpha$$

for all $X, \bar{X} \in S_\rho(X_0)$ and all $\mu, \bar{\mu} \in S_+^2$, where $\delta > 0$ and $\alpha \in (0, 1)$ are constants. Then

$$(\kappa_1^2 + \kappa_2^2)(X_0) \leq c(v(X_0))^{-2} \rho^{-2},$$

where c is a constant depending only on γ, α and δ .

Proof. For sufficiently small $\theta \in (0, 1)$, depending on γ , we know $S_{\theta\rho}(X_0)$ is connected by [8], Lemma (3.2)¹. Let $(\xi, \zeta) = (\xi_1, \xi_2, \zeta)$ denote new coordinates for R^3 defined by

$$(2.4) \quad (\xi, \zeta) = (X - X_0)Q',$$

where Q is an orthogonal matrix with rows $e_1, e_2, \nu(X_0)$, with $\{e_1, e_2\}$ any orthonormal basis for the tangent space of M at X_0 . By the Hölder estimate of Theorem 3 it is clear that there is a $\theta \in (0, 1)$, depending only on γ , such that $S_{\theta\rho}(X_0)$ can be represented, relative to the new coordinates (ξ, ζ) , in the form

$$(2.5) \quad \zeta = \tilde{u}(\xi), \quad \xi \in U,$$

where U is an open subset of R^2 , $\tilde{u} \in C^2(U)$ and

$$(2.6) \quad \sup |D\tilde{u}| \leq 1, \quad D_{\theta\rho}(0) \subset U, \quad \tilde{u}(0) = 0.$$

Furthermore, again using Theorem 3 we can infer that

$$(2.7) \quad |D\tilde{u}(\xi) - D\tilde{u}(\bar{\xi})| \leq c(v(X_0))^{-1} \{|\xi - \bar{\xi}|/\rho\}^\alpha, \quad \xi, \bar{\xi} \in D_{\theta\rho}(0),$$

provided $\theta \in (0, 1)$ is sufficiently small (depending on γ). By the discussion of §1 we can also infer from (1.1) and (1.2) that \tilde{u} satisfies an equation of the form

$$(2.8) \quad \sum_{i,j=1}^2 \tilde{a}_{ij}(\xi) D_{ij} \tilde{u} = 0 \quad \text{on } U,$$

where

¹ We will henceforth use this connectivity result whenever it is convenient to do so; note that for the inhomogeneous case the choice of θ depends also on $\mu\rho$.

$$|\lambda|^2 - \frac{(\lambda \cdot D\tilde{u})^2}{1 + |D\tilde{u}|^2} \leq \tilde{a}_{ij}(\xi) \lambda_i \lambda_j \leq \gamma \left(|\lambda|^2 - \frac{(\lambda \cdot D\tilde{u})^2}{1 + |D\tilde{u}|^2} \right)$$

for all $\lambda \in \mathbf{R}^2$ and $\xi \in U$. By (2.6) this clearly implies

$$(2.9) \quad \frac{1}{2} |\lambda|^2 \leq \sum_{i,j=1}^2 \tilde{a}_{ij}(\xi) \lambda_i \lambda_j \leq \gamma |\lambda|^2$$

(because $|\lambda|^2 - (\lambda \cdot D\tilde{u})^2 / (1 + |D\tilde{u}|^2) \geq |\lambda|^2 / (1 + |D\tilde{u}|^2)$ by Cauchy's inequality). In fact by virtue of the discussion of §1 together with (2.7) and the Hölder condition (2.3), it is clear that we may assume

$$(2.10) \quad |\tilde{a}_{ij}(\xi) - \tilde{a}_{ij}(\bar{\xi})| \leq c \{ |\xi - \bar{\xi}| / \rho \}^\tau, \quad \xi, \bar{\xi} \in D_{\theta\rho}(0),$$

where $c > 0$ and $\tau \in (0, 1)$ depend only on α, δ and γ .

Now by (2.9), (2.10) and the Schauder interior estimate for solutions of (2.8), we then have

$$(2.11) \quad \left\{ \sum_{i,j=1}^2 (D_{ij}\tilde{u}(0))^2 \right\}^{1/2} \leq c \rho^{-2} \sup_{D_{\theta\rho}(0)} |\tilde{u}|,$$

where c depends only on γ, α and δ . On the other hand, since $D\tilde{u}(0) = 0$ and $\tilde{u}(0) = 0$, we deduce from (2.7) that

$$(2.12) \quad \sup_{D_{\theta\rho}(0)} |\tilde{u}| \leq c \rho (v(X_0))^{-1},$$

where c depends only on γ . Also, again using the fact that $D\tilde{u}(0) = 0$, we have

$$(2.13) \quad \sum_{i,j=1}^2 (D_{ij}\tilde{u}(0))^2 = (\kappa_1^2 + \kappa_2^2)(X_0).$$

The theorem is now proved by combining (2.11), (2.12) and (2.13).

3. The inhomogeneous case. Here the notation will be the same as in §2, except that (1.3) is assumed in place of the condition $b \equiv 0$.

All the results of §2, except Theorem 4, have analogues in this more general setting, but in most cases either the hypotheses on the coefficient functions a_{ij} must be stronger or the conclusion weaker than for the corresponding results of §2.

We first have the following analogue of Theorem 3.

THEOREM 3'. *If $D_\rho(x_0) \subset \Omega$, then*

$$|\nu(X) - \nu(\bar{X})| \leq c \left\{ \frac{|X - \bar{X}|}{\rho} \right\}^\alpha, \quad X, \bar{X} \in S_{\rho/2}(X_0),$$

where $c > 0$ and $\alpha \in (0, 1)$ depend only on γ and $\mu\rho$.

In view of the remarks at the end of §1, this theorem is a special case of [8], Theorem (3.1). By considering examples it is easy to see that an estimate like that of Theorem 3, with the factor $(v(X_0))^{-1}$ on the right hand side, cannot hold in the present inhomogeneous setting. (One class of examples is obtained by considering the constant mean curvature equation

$$\sum_{i=1}^2 D_i \frac{D_i u}{\sqrt{1 + |Du|^2}} = H ,$$

where H is any nonzero constant.)

For the purposes of the present section it will be convenient to define the function $b^*(X, \mu)$ for

$$X = (x, z) \in \Omega \times \mathbf{R} \quad \text{and} \quad \mu = \frac{(-p, 1)}{\sqrt{1 + |p|^2}} , \quad p \in \mathbf{R}^2 ,$$

by

$$(3.1) \quad b^*(X, \mu) = (1 + |p|^2)^{-1/2} b(x, z, p) .$$

Note that b^* is thus well defined for $\mu \in S_+^2$, with S_+^2 as in (2.2). (The function b^* , like the functions α_{ij}^* of (2.1), arise quite naturally in case (1.1) is the non-parametric Euler-Lagrange equation of a parametric elliptic functional—see Appendix 1.)

Next we want to obtain analogues of Theorems 1, 2 for the inhomogeneous case. We will impose the following restrictions on the functions α_{ij}^* , b^* :

$$(3.2) \quad \rho \sqrt{1 + |p|^2} \left| \frac{\partial \alpha_{ij}^*}{\partial z}(x, z, \mu) \right| + \rho \sum_{k=1}^2 \left| \frac{\partial}{\partial x_k} \alpha_{ij}^*(x, z, \mu) \right| \\ + |\mu - \bar{\mu}|^{-1} |\alpha_{ij}^*(x, z, \mu) - \alpha_{ij}^*(x, z, \bar{\mu})| \leq \delta ,$$

$$(3.3) \quad \rho \sqrt{1 + |p|^2} \left| \frac{\partial b^*}{\partial z}(x, z, \mu) \right| + \rho \sum_{k=1}^2 \left| \frac{\partial}{\partial x_k} b^*(x, z, \mu) \right| \\ + |\mu - \bar{\mu}|^{-1} |b^*(x, z, \mu) - b^*(x, z, \bar{\mu})| \leq \delta \rho^{-1}$$

for all $(x, z) \in \Omega \times \mathbf{R}$ and $\mu, \bar{\mu} \in S_+^2$ with $\mu \neq \bar{\mu}$ and

$$\mu = \frac{(-p, 1)}{\sqrt{1 + |p|^2}} .$$

In these inequalities δ denotes a fixed constant.

THEOREM 1'. *If $D_\rho(x_0) \subset \Omega$ and if (3.2), (3.3) hold, then*

$$\sup_{S_{\rho/2}(X_0)} v \leq c \inf_{S_{\rho/2}(X_0)} v ,$$

where $c > 0$ depends only on $\gamma, \mu\rho, \delta$.

Before giving the proof of this theorem we point out that, by an argument like that used to prove Theorem 2, we can infer the following from Theorem 1'.

THEOREM 2'. *Under the hypotheses of Theorem 1',*

$$|Du(x_0)| \leq c_1 \exp \{c_2 u(x_0)/\rho\},$$

provided $u \geq 0$ on $D_\rho(x_0)$. Here c_1, c_2 are constants depending on $\gamma, \mu\rho$ and δ .

Proof of Theorem 1'. We consider two cases:

Case I. $|Du(x_0)| \leq 2$. In this case the Hölder estimate of Theorem 3' can be used to deduce $\sup_{D_{\rho\theta}(x_0)} |Du| \leq 3$ for suitable $\theta \in (0, 1)$ depending only on γ and $\mu\rho$. Hence the required result is established in this case.

Case II. $Du(x_0) > 2$. In this case we introduce new coordinates (ξ, ζ) as in (2.4), where now Q has the form

$$(3.4) \quad Q = \begin{pmatrix} 0 & 0 & 1 \\ \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \end{pmatrix}$$

for some constant α to be chosen. Since $|Du(x_0)| > 2$, which guarantees $\nu(X_0) \cdot (0, 0, 1) < 1/\sqrt{5}$, it is clear from the Hölder estimate of Theorem 3' that α can be chosen such that there is a representation of the form (2.5), (2.6) for suitable $\theta \in (0, 1)$. Also by (1.1)–(1.3) and by the discussion of §1, we know that

$$(3.5) \quad \sum_{i,j=1}^2 \tilde{a}_{ij}(\xi) D_{ij} \tilde{u} = \tilde{b}(\xi) \text{ on } U$$

(cf. (2.8)), where (2.9) holds and where $|\tilde{b}(\xi)| \leq \mu$ on U . Here U is as in (2.5).

Now let $\zeta = \zeta(\xi)$ be a C^1 function with compact support in U , multiply by $D_1 \zeta$ in (3.5) and integrate over U . After making use of the relations

$$\int_U D_{22} \tilde{u} D_1 \zeta d\xi = - \int_U D_2 \tilde{u} D_{12} \zeta d\xi = \int_U D_{21} \tilde{u} D_2 \zeta,$$

this gives

$$\int_{U, j=1}^2 \alpha_{ij} D_i \gamma \psi D_j \zeta d\xi = \int_U (D_1 \beta) \zeta d\xi ,$$

where $\gamma = D_1 \tilde{u}$, $\alpha_{11} = \tilde{a}_{11}/\tilde{a}_{22}$, $\alpha_{22} \equiv 1$, $\alpha_{12} \equiv 0$, $\alpha_{21} = 2\tilde{a}_{12}/\tilde{a}_{22}$, and $\beta = \tilde{b}/\tilde{a}_{22}$. Thus $\gamma = D_1 \tilde{u}$ is a weak solution of the equation

$$\sum_{i,j=1}^2 D_i (\alpha_{ij} D_j \gamma) = -D_1 \beta .$$

Since (2.4) and (3.4) imply the relations

$$\begin{aligned} \xi_1 &= u(x) - u(x_0) , \\ x - x_0 &= (\xi_1 \cos \alpha + \tilde{u}(\xi) \sin \alpha, -\xi_2 \sin \alpha + \tilde{u}(\xi) \cos \alpha) , \end{aligned}$$

one can easily check that

$$\begin{aligned} (3.6) \quad D_1 \beta &= -\tilde{a}_{22}^{-2}(\xi) b^*(x, u, \nu) \sum_{i,j=1}^3 \lambda_i \lambda_j \left\{ \frac{\partial}{\partial z} a_{ij}^*(x, u, \nu) \right. \\ &+ \frac{\partial}{\partial x_1} a_{ij}^*(x, u, \nu) \sin \alpha D_1 \tilde{u}(\xi) + \frac{\partial}{\partial x_2} a_{ij}^*(x, u, \nu) \cos \alpha D_1 \tilde{u}(\xi) \\ &+ \left. \frac{\partial}{\partial \mu} a_{ij}^*(x, u, \nu) \cdot D_1 \tilde{\nu} Q \right\} + \tilde{a}_{22}^{-1}(\xi) \left\{ \frac{\partial}{\partial z} b^*(x, u, \nu) \right. \\ &+ \frac{\partial}{\partial x_1} b^*(x, u, \nu) \sin \alpha D_1 \tilde{u}(\xi) + \frac{\partial}{\partial x_2} b^*(x, u, \nu) \cos \alpha D_1 \tilde{u}(\xi) \\ &+ \left. \frac{\partial}{\partial \mu} b^*(x, u, \nu) \cdot D_1 \tilde{\nu} Q \right\} . \end{aligned}$$

Here $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ denotes the second row of Q (so that $\tilde{a}_{22} = \sum_{i,j=1}^3 a_{ij}^* \lambda_i \lambda_j$ in accordance with the discussion of §1). Also, $\tilde{\nu} = (-D\tilde{u}, 1)/\sqrt{1 + |D\tilde{u}|^2}$, $\partial/\partial \mu$ denotes the gradient operator on S_+^2 , and we have used the relation $\tilde{\nu} Q = \nu$. Then, since

$$D_1 \tilde{\nu}_k = (1 + |D\tilde{u}|^2)^{-1/2} \sum_{j=1}^2 (\delta_{kj} - \tilde{\nu}_k \tilde{\nu}_j) D_{j_1} \tilde{u}$$

we see (by using the conditions (3.2), (3.3) together with (2.6) and the identity $D_1 \tilde{u}/\sqrt{1 + |D\tilde{u}|^2} = \nu^{-1}$) that (3.6) implies

$$(3.7) \quad \int_U \left(\sum_{i,j=1}^2 \alpha_{ij} D_j \gamma \psi D_i \zeta + \sum_{i=1}^2 \beta_i D_i \gamma \psi \zeta + \tau \gamma \psi \zeta \right) d\xi = 0 , \quad \zeta \in C_0^1(U) ,$$

where

$$\begin{aligned} (3.8) \quad \frac{1}{2} |\theta|^2 &\leq \sum_{j,i=1}^2 \alpha_{ij} \theta_i \theta_j \leq \gamma |\theta|^2 , \quad \theta = (\theta_1, \theta_2) \in \mathbf{R}^2; \text{ and} \\ \rho \sum_{i=1}^2 |\beta_i| + \rho^2 |\tau| &\leq c_1 , \end{aligned}$$

where c_1 depends only on $\mu\rho$ and δ . Clearly we can apply the De Giorgi, Nash, Moser theory and deduce that

$$(3.9) \quad \sup_{D_{\theta\rho/2}(0)} \psi \leq c_2 \inf_{D_{\theta\rho/2}(0)} \psi$$

for non-negative solutions ψ of (3.7), where c_2 depends on $\gamma, \mu\rho$ and δ . However $D_1\tilde{u} = \sqrt{1 + |D\tilde{u}|^2}/v \geq 0$, as one easily checks from the relation $\tilde{\nu}Q = \nu$. Thus we can apply (3.9) to $D_1\tilde{u}$. Because of (2.6) we then deduce the required Harnack inequality for v . This completes the proof of Theorem 1'.²

An unsatisfactory feature of Theorem 2' is that the hypotheses on b^* are such as to exclude certain important examples. For instance, the capillary surface equation

$$\sum_{i=1}^2 D_i \left(\frac{D_i u}{\sqrt{1 + |Du|^2}} \right) = \kappa u, \quad \kappa > 0$$

is excluded from the above discussion. This defect is remedied in the following theorem, in which the following condition is assumed in place of (3.3):

$$(3.10) \quad -\rho\sqrt{1 + |p|^2} \frac{\partial b^*}{\partial z}(x, z, \mu) + \rho \sum_{k=1}^2 \left| \frac{\partial}{\partial x_k} b^*(x, z, \mu) \right| \\ + |\mu - \bar{\mu}|^{-1} |b^*(x, z, \mu) - b^*(x, z, \bar{\mu})| \leq \delta\rho^{-1}$$

for all $(x, z) \in \Omega \times \mathbf{R}$ and $\mu, \bar{\mu} \in S_+^2$ with $\mu \neq \bar{\mu}$ and

$$\mu = \frac{(-p, 1)}{\sqrt{1 + |p|^2}}, \quad p \in \mathbf{R}^2.$$

THEOREM 2''. *If $D_\rho(x_0) \subset \Omega$, if $u \geq 0$ on $D_\rho(x_0)$ and if (3.2), (3.10) hold, then*

$$|Du(x_0)| \leq c_1 \exp \{c_2 u(x_0)/\rho\},$$

where c_1, c_2 are constants depending on $\gamma, \mu\rho$ and δ .

Proof. In the proof c_1, c_2, \dots will denote constants depending only on $\gamma, \mu\rho$ and δ .

We consider the same two cases as in the proof of Theorem 1. In Case I the required result is trivially satisfied. The argument for Case II begins as before, except that in place of (3.7) we now deduce that $\psi = D_1\tilde{u}$ is a *supersolution* of the equation (3.7). That

² Actually we have only proved $\sup_{S_{\theta\rho}(x_0)} v \leq c \inf_{S_{\theta\rho}(x_0)} v$ for some $\theta \in (0, 1)$ depending on $\gamma, \mu\rho$ and δ . The required result (with $\theta=1/2$) follows because we can vary X_0 .

is, we deduce

$$(3.11) \quad \int_U \left(\sum_{i,j=1}^2 \alpha_{ij} D_j \psi D_i \zeta + \sum_{i=1}^2 \beta_i D_i \psi \zeta + \tau \psi \zeta \right) d\xi \geq 0$$

for any nonnegative $\zeta \in C_0^1(U)$, where (3.8) still holds. Replacing ζ by ζ^2/ψ , we then deduce that

$$(3.12) \quad \int_U \left\{ \zeta^2 \sum_{j,i=1}^2 \alpha_{ij} D_i w D_j w + 2\zeta \sum_{i,j=1}^2 \alpha_{ij} D_i w D_j \zeta + \zeta^2 \left(\sum_{i=1}^2 \beta_i D_i w - \tau \right) \right\} d\xi \leq 0$$

for all $\zeta \in C_0^1(U)$, where $w = \log \psi^{-1}$. Using well-known arguments from the De Giorgi-Nash theory of uniformly elliptic equations, one can easily deduce from (3.12) that

$$(3.13) \quad w(0) \leq c_1 + c_2 \int_{D^+} w d\xi,$$

where $D^+ = \{\xi \in D_{\theta\rho/4}(0) : w(\xi) > 1\}$. The remainder of the proof consists in estimating the integral on the right of (3.13). We begin by noting that inequality (3.11) implies

$$\int_{D_{\theta\rho/2}(X_0)} |Dw|^2 d\xi \leq c_3.$$

If we let ω be defined by $\omega = \log v$ on M , then it is clear (by (2.6)) that this last inequality implies

$$(3.14) \quad \int_{S_{\theta\rho/2}(X_0)} |\delta\omega|^2 dA \leq c_4,$$

where δ denotes the tangential gradient operator on M . Now define

$$E = \{X = (x, u(x)) \in M \cap (D_{\theta\rho/4}(x_0) \times \mathbf{R}) : u(x) < u(x_0) + \theta\rho/4\}.$$

We can choose points $X_1, \dots, X_N \in E$ such that $E \subset \bigcup_{i=1}^N S_{\theta\rho/4}(X_i)$ and

$$(3.15) \quad N \leq c_5(1 + u(x_0)/\rho).$$

Using (3.14) with X_i in place of X_0 , summing over i , and also using (3.15), we then deduce that

$$(3.16) \quad \int_E |\delta\omega|^2 dA \leq c_6(1 + u(x_0)/\rho).$$

We now recall the fact (see e.g. [8], (4.6)) that

$$(1 - \nu_3^2) \min \{\kappa_1^2, \kappa_2^2\} \leq |\delta\nu_3|^2$$

³ Such a equality also holds, of course, if $|Du(x_0)| < 2$; in this case one simply essentially repeats the previous argument *without* introducing new coordinates ξ .

at each point of M . Hence at points where $\nu_3^2 < 1/2$ (that is, where $\omega > \log \sqrt{2}$) we have

$$(3.17) \quad \kappa_1^2 + \kappa_2^2 \leq c_7(|\delta\nu_3|^2 + \mu^2) \leq c_7(|\delta\omega|^2 + \mu^2)$$

by virtue of (1.7). Hence, since $\log \sqrt{2} < 1$,

$$(3.18) \quad \int_{E \cap \{X: \omega(X) > 1\}} H^2 dA \leq c_9 \left(\int_E |\delta\omega|^2 dA + \mu^2 |E| \right), \quad H = \kappa_1 + \kappa_2.$$

Also, we know that

$$(3.19) \quad |E| \leq c_{10}(1 + u(x_0)/\rho)\rho^2$$

by virtue of (3.15) and the area bounds $|S_{\theta\rho/4}(X_i)| \leq c_{11}\rho^2$. Next we have, by the first variation formula for the surface M ,

$$(3.20) \quad \int_M \delta_3 h dA = \int_M \nu_3 H h dA,$$

whenever h is a C^1 function with compact support in M . In particular, we can choose h of the form

$$h(X) = f(x_3)g(\omega(X))\zeta(x_1, x_2), \quad X = (x_1, x_2, x_3) \in M,$$

where f, g are $C^1(\mathbf{R})$ functions and $\zeta \in C^1(\mathbf{R}^2)$ with

$$\begin{cases} g(\omega) = \omega \text{ for } \omega > 2, g(\omega) = 0 \text{ for } \omega < 1, \text{ and } 0 \leq g'(\omega) \leq L \text{ for } \omega \in \mathbf{R}; \\ f(t) \equiv 0 \text{ for } t > u(x_0) + \sigma, f'(t) \equiv -1 \text{ for } t \in (u(x_0) - \sigma/2, u(x_0) + \sigma/2), \\ 0 \leq -f'(t) \leq 2 \text{ for } t \in \mathbf{R}, f(t) \equiv \sigma \text{ for } t < u(x_0) - \sigma \\ \zeta(x) \equiv 1 \text{ for } x \in D_{\sigma/2}(x_0), \zeta(x) \equiv 0 \text{ for } x \in \sim D_\sigma(x_0), \\ |D\zeta(x)| \leq 3/\sigma \text{ for } x \in \mathbf{R}^2. \end{cases} \text{ Here } \sigma = \theta\rho/4.$$

With such a choice of h one easily deduces

$$\int_{S_{\theta\rho/8}(X_0)} \omega dA \leq c_{12} \left\{ \rho \int_{E \cap \{X: \omega(X) > 1\}} (|\delta\omega| + \nu_3 \omega |H|) dA + |E| \right\}.$$

Since $\nu_3 \omega < 1$ we then use (3.16), (3.18) and (3.19) to deduce

$$(3.21) \quad \int_{S_{\theta\rho/8}(X_0)} \omega dA \leq c_{13}(1 + u(x_0)/\rho).$$

The required result now follows from this and (3.13). (One needs to note that $S_{\theta\rho/8}(X_0) \supset \{(\xi, \tilde{u}(\xi))Q: \xi \in D_{\theta\rho/16}(0)\}$ by virtue of (2.6).)

Next we present an analogue of Theorem 8. Note that the estimate obtained in the theorem here is weaker than that of Theorem 8 in that there is no factor of $(v(X_0))^{-2}$ on the right hand side. (Consideration of graphs with constant mean curvature shows that one cannot expect to have such a factor on the right hand side in the

nonhomogeneous case.)

THEOREM 8'. *Suppose $D_\rho(x_0) \subset \Omega$, suppose (2.3) holds, and suppose*

$$(3.22) \quad |b^*(X, \mu) - b^*(\bar{X}, \bar{\mu})| \leq \rho^{-1} \delta \{ |X - \bar{X}|/\rho + |\mu - \bar{\mu}| \}^\alpha$$

for all $X, \bar{X} \in S_\rho(X_0)$ and all $\mu, \bar{\mu} \in S_+^2$, where $\delta > 0$ and $\alpha \in (0, 1)$ are constants. Then

$$(\kappa_1^2 + \kappa_2^2)(X_0) \leq c\rho^{-2},$$

where c is a constant depending only on $\gamma, \mu\rho, \delta$ and α .

Proof. As in the proof of Theorem 8, we introduce new coordinates as in (2.4) and infer from (1.1) an equation like (2.8) for the function \tilde{u} of (2.5), (2.6). However notice that here we have to use the inequality of Theorem 3' instead of the stronger inequality of Theorem 3. Also, the equation for \tilde{u} corresponding to (2.8) now has the form

$$\sum \tilde{a}_{ij}(\xi) D_{ij} \tilde{u} = \tilde{b}(\xi) \quad \text{on } U,$$

where (2.9) still holds, where $|\tilde{b}| \leq \mu$ on U and where (by virtue of (2.3), (3.22) and the estimate of Theorem 3')

$$\sum_{i,j=1}^2 |\tilde{a}_{ij}(\xi) - \tilde{a}_{ij}(\bar{\xi})| + \rho |\tilde{b}(\xi) - \tilde{b}(\bar{\xi})| \leq c \{ |\xi - \bar{\xi}|/\rho \}^\tau$$

for $\xi, \bar{\xi} \in D_{\theta\rho}(0)$. Here $\theta \in (0, 1)$, $\tau \in (0, 1)$ and $c > 0$ depend on α, δ, γ and $\mu\rho$. Then by applying Schauder's interior estimate as in the proof of Theorem 8, we obtain the required inequality.

The next theorem generalizes Theorem 6; notice that there are no continuity hypotheses on the functions a_{ij}, b in the theorem.

THEOREM 6'. *Suppose $\Omega = \mathbf{R}^2 \sim K$, where K is compact, and suppose that*

$$(3.23) \quad |b^*(X, \mu)| \leq \mu_0 |X|^{1+\tau}, \quad X \in \mathbf{R}^3 \sim \{0\}, \mu \in S_+^2,$$

where $\mu_0 > 0$ and $\tau \in (0, 1)$ are given constants. Then there is a vector $\nu^0 = (\nu_1^0, \nu_2^0, \nu_3^0) \in \tilde{S}_+^3$ such that

$$\nu(X) = (1 + |Du(x)|^2)^{-1/2} (-Du(x), 1) \longrightarrow \nu^0$$

uniformly for $|x| \rightarrow \infty$. In case $b \equiv 0$, then $\nu_3^0 > 0$ and hence there is an $a \in \mathbf{R}^2$ such that $Du(x) \rightarrow a$ uniformly for $|x| \rightarrow \infty$.

Proof. The proof relies havily on the techniques of [8]. In the

proof, constants which depend at most on γ, ρ_0, μ_0 and τ will be denoted c_1, c_2, \dots .

To begin, let R_0 be such that $K \subset D_{R_0}(0)$, define

$$\rho_0 = \sup \{(u^2(x) + |x|^2)^{1/2} : x \in \partial D_{R_0}(0)\}$$

and for $R > \rho > \rho_0$ let $T_{\rho,R}$ be defined by

$$T_{\rho,R} = \{X \in M : \rho < |X| < R\}.$$

We will repeatedly make use of the fact that, for $\rho \geq \sigma > 0$, $T_{\rho,\sigma+\sigma}$ is covered by $S_\sigma(X_1), \dots, S_\sigma(X_N)$, where X_1, \dots, X_N are points of $T_{\rho,\sigma+\sigma}$ and $N \leq c\rho/\sigma$, where c is an absolute constant.

From the discussion of §4 of [8], have the identity

$$\kappa_1 \kappa_2 dA = d\omega^*$$

where κ_1, κ_2 are the principal curvatures of M and ω^* is the 1-form on M defined by

$$\omega^* = (1 + \nu_3)^{-1}(-\nu_2 d\nu_1 + \nu_1 d\nu_2).$$

Using Stokes' Theorem we then have, for almost all ρ, R with $\rho_0 < \rho < R$,

$$(3.24) \quad \int_{T_{\rho,R}} \zeta^2 \kappa_1 \kappa_2 dA = - \int_{T_{\rho,R}} d\zeta^2 \wedge \omega^* + \int_{\partial T_{\rho,R}} \zeta^2 \omega^*.$$

Using (1.7) (and noting that (3.23) implies that we can take μ to be the variable quantity $\mu_0/|X|^{1+\tau}$), we then deduce

$$(3.25) \quad |A_1|^{-1} \int_{T_{\rho,R}} \zeta^2 |\delta\nu|^2 dA \leq \left| \int_{T_{\rho,R}} 2\zeta d\zeta \wedge \omega^* \right| + \left| \int_{\partial T_{\rho,R}} \zeta^2 \omega^* \right| \\ + A_2 |A_1|^{-1} \int_{T_{\rho,R}} \zeta^2 |X|^{-2-2\tau} dA.$$

If ζ is chosen so that $\zeta \equiv 0$ on $\partial T_{\rho,R}$, then this gives

$$\int_{T_{\rho,R}} \zeta^2 |\delta\nu|^2 dA \leq c_1 \int_{T_{\rho,R}} (\zeta |\delta\nu| |\delta\zeta| + \zeta^2 |X|^{-2-2\tau}) dA,$$

which clearly implies (by Cauchy's inequality)

$$(3.26) \quad \int_{T_{\rho,R}} \zeta^2 |\delta\nu|^2 dA \leq c_2 \int_{T_{\rho,R}} (|\delta\zeta|^2 + \zeta^2 |X|^{-2-2\tau}) dA.$$

Next we note that because of the area bounds

$$(3.27) \quad |S_\sigma(X_0)| \leq c_3 \sigma^2,$$

which by Lemma (3.1) of [8] are valid when $X_0 \in M$ and $|X_0| > \rho_0 +$

2σ , we can deduce

$$(3.28) \quad |T_{\rho, \rho+\sigma}| \leq c_4 \rho \sigma, \quad \rho > 2\rho_0, \quad 0 < \sigma \leq \rho.$$

By using a suitable choice of ζ together with (3.28), it is now not difficult to see that (3.26) implies

$$\int_{T_{2\rho, R/2}} |\delta\nu|^2 dA \leq c_5, \quad R > 4\rho.$$

Thus (since c_5 is independent of R) we deduce

$$(3.29) \quad \int_{T_{\rho, \infty}} |\delta\nu|^2 dA \leq c_5$$

for $\rho > 2\rho_0$. By again choosing ζ appropriately, we can now deduce from (3.25) and (3.28)

$$(3.30) \quad \int_{T_{\rho, \infty}} |\delta\nu|^2 dA \leq c_6 \left\{ \left| \int_{\partial T_{\rho, \infty}} \omega^* \right| + \rho^{-2\tau} \right\}.$$

By a straightforward modification of the argument of [8], Theorem (2.1), we can infer from this that

$$(3.31) \quad \hat{\mathcal{D}}(\rho) \leq -c_7 \rho \hat{\mathcal{D}}'(\rho) + c_8 \rho^{-2\tau},$$

for almost all $\rho > 4\rho_0$, where

$$\hat{\mathcal{D}}(\rho) = \int_{T_{\rho, \infty}} |\delta\nu|^2 dA.$$

(It is necessary to note the fact that

$$(3.32) \quad -\frac{d}{d\rho} \int_{T_{\rho, \infty}} |\delta|X|^2 dA \leq c_9 \left\{ \rho^{-1} |T_{\rho, 2\rho}| + \int_{T_{\rho, 2\rho}} |H| dA \right\},$$

where $H = \kappa_1 + \kappa_2$. This follows by using the first variation formula for M in a manner similar to the argument leading to (A.2) to (A.2) of [8].) By integrating (3.32) from $4\rho_0$ to ρ , we now conclude

$$(3.33) \quad \hat{\mathcal{D}}(\rho) \leq c_{10} \rho^{-\beta}, \quad \rho > 4\rho_0,$$

where $\beta \in (0, 2\tau)$ depends only on γ, ρ_0, μ_0 and τ . In particular we note that

$$(3.33)' \quad \int_{S_\rho(X_0)} |\delta\nu|^2 dA \equiv \mathcal{D}(X_0, \rho) \leq c_{10} \rho^{-\beta}$$

for all $X_0 \in M$ with $|X_0| > 2\rho > 8\rho_0$.

Now an examination of the proof of the Hölder estimate of theorem (in particular see (2.11), (2.12) of [8]) will show that in the present setting we can assert

$$(3.34) \quad |\nu(X) - \nu(X_0)| \leq c_{11} \{ \mathcal{D}(\rho, X_0) + \rho^{-2r} \}^{1/2} \{ \sigma / \rho \}^\alpha$$

for any $X \in S_\sigma(X_0)$, $\sigma \in (0, \rho)$. Combining (3.33)' and (3.34), with $\sigma = \rho$, gives

$$|\nu(X) - \nu(X_0)| \leq c_{12} \rho^{-\beta'}, \quad X \in S_\rho(X_0), \beta' = \beta/2,$$

and this clearly implies (since we can vary X_0)

$$\sup_{X, \bar{X} \in T_{\rho, 2\rho}} |\nu(X) - \nu(\bar{X})| \leq c_{13} \rho^{-\beta'}$$

for any $\rho > 4\rho_0$. Then for any integer $k \geq 1$ we have

$$\sup_{X, \bar{X} \in T_{\rho, 2^k \rho}} |\nu(X) - \nu(\bar{X})| \leq c_{13} \rho^{-\beta'} \left(\sum_{r=0}^{k-1} 2^{-r\beta'} \right),$$

where upon we deduce that

$$\sup_{X, \bar{X} \in T_{\rho, \infty}} |\nu(X) - \nu(\bar{X})| \leq c_{14} \rho^{-\beta'}.$$

The first conclusion of the theorem clearly follows from this.

We now consider the case $b \equiv 0$. We then have (1.7) with $A_2 = 0$, so that we can use the theory developed in §4 of [8]. In particular we use the identity

$$(3.35) \quad ((1 - \nu_3)\gamma(\nu_3))' \kappa_1 \kappa_2 dA = -d(\gamma(\nu_3)(1 + \nu_3)^{-1}(-\nu_2 d\nu_1 + \nu_1 d\nu_2)),$$

where γ is an arbitrary $C^1(\mathbf{R})$ function. (This identity is the pointwise version of identity (4.5) of [8], as one easily checks by using Stokes' Theorem.) We multiply each side of (3.35) by a cut-off function ζ^2 which vanishes for $|X| > R$, and integrate (3.35) over $T_{\rho, \infty}$. After using Stokes' Theorem, this gives

$$(3.36) \quad \begin{aligned} & \int_{T_{\rho, \infty}} \zeta^2 ((1 - \nu_3)\gamma(\nu_3))' \kappa_1 \kappa_2 dA \\ &= - \int_{\partial T_{\rho, \infty}} \zeta^2 \gamma(\nu_3)(1 + \nu_3)^{-1} (-\nu_2 d\nu_1 + \nu_1 d\nu_2) \\ & \quad - \int_{T_{\rho, \infty}} 2\zeta \gamma(\nu_3)(1 + \nu_3)^{-1} d\zeta \wedge (-\nu_2 d\nu_1 + \nu_1 d\nu_2). \end{aligned}$$

Now from Theorem 1 we know that

$$(3.37) \quad \sup_{\partial T_{\rho, \infty}} \nu_3 \leq c_{15} \inf_{\partial T_{\rho, \infty}} \nu_3$$

for each $\rho > 2\rho_0$.

Also if there exists a sequence $\{X_j\} \subset M$ with $|X_j| \rightarrow \infty$ and with $\{\nu_3^{-1}(X_j)\}$ bounded, we could immediately deduce $\nu_3^0 > 0$, and the required result would be established. We may therefore assume $\nu_3(X) < 1/2$

for all $|X| > \rho_1$, where $\rho_1 > \rho_0$ is a fixed constant. Then for $\rho > \rho_1$ we can replace $\gamma(\nu_3)$ in (3.36) by $(\nu_3)^{-1}/(1 - \nu_3)$, thus giving

$$(3.38) \quad \int_{T_{\rho,\infty}} \zeta^2 \nu_3^{-2} \kappa_1 \kappa_2 dA = - \int_{T_{\rho,\infty}} \nu_3^{-1} (1 - \nu_3^2)^{-1} d\zeta^2 \wedge (-\nu_2 d\nu_1 + \nu_1 d\nu_2) \\ - \int_{\partial T_{\rho,\infty}} \zeta^2 \nu_3^{-1} (1 - \nu_3^2)^{-1} (-\nu_2 d\nu_1 + \nu_1 d\nu_2).$$

Using inequality (4.8) of [8] and also using Cauchy's inequality, one easily checks that (3.38) implies

$$\int_{T_{\rho,\infty}} \zeta^2 |\delta w|^2 dA \leq c_{16} \left\{ \int_{T_{\rho,\infty}} |\delta \zeta|^2 dA + \int_{\partial T_{\rho}} |\delta w|^2 \zeta^2 ds \right\}.$$

Here, and subsequently, $w = \log \nu_3^{-1}$. Choosing $\zeta(X) \equiv 1$ for $|X| < R/2$ and $\zeta(X) \equiv 0$ for $|X| > R$, and letting $R \rightarrow \infty$, we then deduce that

$$(3.39) \quad \int_{T_{\rho,\infty}} |\delta w|^2 < \infty$$

for $\rho > \rho_1$. Making a similar choice of ζ in (3.38) and again letting $R \rightarrow \infty$, we can conclude, for almost all $\rho > \rho_1$,

$$(3.40) \quad \int_{T_{\rho,\infty}} |\delta w|^2 dA \leq c_{17} \left| \int_{\partial T_{\rho,\infty}} -\omega_1 d\nu_1 + \omega_2 d\nu_2 \right|,$$

where

$$\omega_i = \nu_i \nu_3^{-1} (1 - \nu_3^2)^{-1}, \quad i = 1, 2.$$

Now for almost all $\rho > \rho_1$ we have $\partial T_{\rho,\infty} = \bigcup_{i=1}^{N(\rho)} \Gamma_{\rho}^{(j)}$, where $N(\rho)$ is a positive integer and $\Gamma_{\rho}^{(j)}$ is a smooth Jordan curve such that $\delta|X|$ does not vanish of $\Gamma_{\rho}^{(j)}$. Since $\int_{\Gamma_{\rho}^{(j)}} d\nu_i = 0$, we can write (3.40) in the form

$$(3.41) \quad \int_{T_{\rho,\infty}} |\delta w|^2 dA \leq c_{17} \left| \sum_{j=1}^{N(\rho)} \int_{\Gamma_{\rho}^{(j)}} -(\omega_1 - \omega_1(X^{(j)})) d\nu_1 \right. \\ \left. + (\omega_2 - \omega_2(X^{(j)})) d\nu_2 \right|,$$

where $X^{(j)}$ denotes a fixed point on $\Gamma_{\rho}^{(j)}$. Now

$$(3.42) \quad \sup_{\Gamma_{\rho}^{(j)}} |\omega_i - \omega_i(X^{(j)})| \leq \int_{\Gamma_{\rho}^{(j)}} \left| \frac{d\omega_i}{ds} \right| ds \leq c_{18} \int_{\Gamma_{\rho}^{(j)}} \nu_3^{-2} |\delta \nu| ds.$$

Combining (3.41) and (3.42), we obtain

$$\int_{T_{\rho,\infty}} |\delta w|^2 dA \leq c_{19} \left(\int_{\partial T_{\rho,\infty}} |\delta \nu| ds \right)^2 \sup_{\partial T_{\rho}} \nu_3^{-2}.$$

In view of inequality (4.8) of [8] and inequality (3.37) above, we then deduce

$$\int_{T_{\rho,\infty}} |\delta w|^2 dA \leq c_{20} \left(\int_{\partial T_{\rho,\infty}} |\delta w| ds \right)^2.$$

Using (3.32) and the Hölder inequality one can then use an argument like that of [8] Theorem (2.1) (the argument is like that needed to obtain the estimate (3.33) above); we thus obtain

$$\int_{T_{\rho,\infty}} |\delta w|^2 dA \leq c_{21} \rho^{-\beta'}$$

where $\beta' \in (0, 1)$ depends only on ρ_0 and γ . In particular we have

$$\int_{S_{\rho}(X_0)} |\delta w|^2 dA \leq c_{21} \rho^{-\beta'}$$

whenever $Y_0 \in M$ and $|X_0| > 2\rho$, $\rho > \rho_1$. Using this last inequality in combination with the inequalities (4.15), (4.16) of [8], we then deduce

$$\sup_{S_{\rho,2}(X_0)} w - \inf_{S_{\rho,2}(X_0)} w \leq c_{22} \rho^{-\beta'/2}.$$

In view of the arbitrariness of X_0 , this gives

$$\sup_{T_{\rho,2\rho}} w - \inf_{T_{\rho,2\rho}} w \leq c_{23} \rho^{-\beta'/2}.$$

Iterating, we obtain

$$\sup_{T_{\rho,2^k\rho}} w - \inf_{T_{\rho,2^k\rho}} w \leq c_{23} \rho^{-\beta'/2} \left(\sum_{r=0}^{\infty} 2^{-r\beta'/2} \right)$$

for each integer $k \geq 1$. Hence

$$\sup_{T_{\rho,\infty}} w - \inf_{T_{\rho,\infty}} w \leq c_{24} \rho^{-\beta'};$$

that is, w is bounded for $|X| > \rho_1$. The theorem now follows.

It is not clear whether or not an estimate like that obtained in Theorem 5 holds for general equation (1.1)–(1.3), (3.2), (3.3), (3.23). However, for a class of divergence-form equations (with $n \geq 2$ independent variables) such a theorem is obtained in [10]; from the discussion of §1 of [10] it is clear that the structural conditions imposed there certainly hold in case the equation arises as the non-parametric Euler-Lagrange equation (equation (A.6) in Appendix 1) of an elliptic parametric functional with integrand $F(X, q)$ in case $F(X, q)$ is independent of X . More generally, it suffices that $F(X, q) = F(x, z, q)$ is such that $F_z(x, z, q) \equiv 0$. (Actually the condition (1.3) of [10] is not quite stated in a weak enough form to include this latter case; however, one can check that all the results of [10]

remain valid if the first inequality of [10], (1.3) is replaced by the weaker condition

$$\left| \sum_{i=1}^n u_{x_i} D_{ik} \right| \leq \beta_5, \quad \sum_{i,k=1}^n |D_{ik}| \leq \beta_5,$$

where (in the notation of [10])

$$D_{ik} = u_{x_i} A_{ix_1 p_k} + u_{x_k} A_{iz} + u_{x_i} B_{p_k} + |Du|^2 A_{iz} p_k,$$

and this condition is weak enough to include the case when F in (A.6) satisfies $F_z(x, z, q) \equiv 0$.)

Finally we state the following analogue of Theorem 6; a proof will be found in [9].

THEOREM. *Suppose $D_\rho(x_0) \sim \{x_0\} \subset \Omega$. Then the closure \bar{M} (taken in $\Omega \times \mathbf{R}$) of the graph of u is a $C^{1,\alpha}$ surface such that $(\bar{D}_\sigma(x_0) \times \mathbf{R}) \cap \bar{M}$ is compact for each $\sigma < \rho$.*

One can show by example that even though the graph of u can thus be extended to a $C^{1,\alpha}$ surface in \mathbf{R}^3 , nevertheless the function $u(x)$ (as a function of $x \in \Omega$) may have no C^1 extension to $D_\rho(x_0)$. (Because it may happen that $|Du(x)|$ is unbounded for x in a neighbourhood of x_0 .)

APPENDIX: *Elliptic Parametric Functionals.*

Let Ω be a bounded domain in \mathbf{R}^2 and consider the functional I , defined for C^1 mappings $Y = (Y_1, Y_2, Y_3): \bar{\Omega} \rightarrow \mathbf{R}^3$ by

$$(A1) \quad I(Y) = \int_{\Omega} G(x, Y, D_1 Y, D_2 Y) dx,$$

where $G = G(x, X, p)$ is a given continuous function of $(x, X, p) \in \mathbf{R}^2 \times \mathbf{R}^3 \times \mathbf{R}^6$. (Here of course $D_i Y = (D_i Y_1, D_i Y_2, D_i Y_3)$ for $i = 1, 2$.) Now let us consider the possibility that I remains invariant under orientation preserving diffeomorphism of \mathbf{R}^2 ; that is, whenever ψ is a diffeomorphism of \mathbf{R}^2 onto itself with positive Jacobian, we would have

$$\int_{\Omega} G(\xi, \tilde{Y}(\xi), D_1 \tilde{Y}(\xi), D_2 \tilde{Y}(\xi)) d\xi = \int_{\Omega} G(x, Y(x), D_1 Y(x), D_2 Y(x)) dx,$$

where $\Omega' = \psi(\Omega)$ and $\tilde{Y} = Y \circ \psi^{-1}$. A simple computation (cf. [7], p. 349) shows that this would be true for all such diffeomorphisms ψ and domains Ω if and only if there is a real-valued function F on $\mathbf{R}^3 \times \mathbf{R}^3$ such that

$$(A2) \quad G(x, X, p) = F(X, P), \quad (x, X, p) \in \mathbf{R}^2 \times \mathbf{R}^3 \times \mathbf{R}^6,$$

where

$$P = (p_3p_5 - p_2p_6, p_1p_6 - p_3p_4, p_2p_4 - p_1p_5);$$

and

$$(A3) \quad F(X, \lambda q) = \lambda F(X, q), \quad (X, q) \in \mathbf{R}^3 \times \mathbf{R}^3, \quad \lambda > 0.$$

Note in particular that (A2) implies that $G(x, X, p)$ cannot depend on x ; that is, $G(x, X, p) = G(0, X, p)$ for $(x, X, p) \in \mathbf{R}^2 \times \mathbf{R}^3 \times \mathbf{R}^6$. In case $p = (D_1Y, D_2Y)$, where Y is a C^1 map from $\bar{\Omega}$ into \mathbf{R}^3 , P is given by

$$P = (D_1Y_3 \cdot D_2Y_2 - D_1Y_2 \cdot D_2Y_3, D_1Y_1 \cdot D_2Y_3 - D_1Y_3 \cdot D_2Y_1, \\ D_1Y_2 \cdot D_2Y_1 - D_1Y_1 \cdot D_2Y_2).$$

As is well known, in case Y is one-to-one and such that the Jacobian matrix $[D_jY_i(x)]$ has rank 2 for each $x \in \Omega$, this last identity can be written

$$P = \chi \nu,$$

where ν is the unit normal of the embedded surface $S = \{Y(x) | x \in \Omega\}$ and χ is the area magnification factor of the mapping Y . Thus, assuming that we orient S with unit normal ν such that $\chi > 0$, we can write

$$I(Y) = \int_S F(X, \nu(X)) dA(X);$$

that is, we can express $I(Y)$ completely in terms of the oriented surface S and independently of the particular mapping Y that is used to represent S . Through this discussion we are led to consider the functional J , defined for any smooth oriented surface S in \mathbf{R}^3 having finite area, by

$$(A4) \quad J(S) = \int_S F(X, \nu(X)) dA(X);$$

this functional has the property that $J(S) = I(Y)$ whenever Y is a one-to-one C^1 mapping from Ω into \mathbf{R}^3 such that $[D_jY_i]$ has rank 2 at each point of Ω and $S = \{Y(x) | x \in \Omega\}$.

If F satisfies (A3), we call a functional of the form (A4) a *parametric functional*. The functional J is called *elliptic* if F is C^2 on $\mathbf{R}^3 \times (\mathbf{R}^3 - \{0\})$ and if the convexity condition

$$(A5) \quad |q| D_{q_i q_j} F(X, q) \xi_i \xi_j \geq |\xi'|^2, \quad \xi' = \xi - \left(\xi \cdot \frac{q}{|q|} \right) \frac{q}{|q|},$$

holds for all $X \in \mathbf{R}^3$, $q \in \mathbf{R}^3 - (0)$ and $\xi \in \mathbf{R}^3$. Notice that, up to a scalar factor, (A5) is the strongest convexity condition possible for F in view of the homogeneity condition (A3).

If we now consider a nonparametric surface M given by

$$M = \{(x, u(x)) \in \mathbf{R}^3 \mid x \in \Omega\},$$

where $u \in C^2(\bar{\Omega})$, then, taking ν to be the downward unit normal $(Du, -1)/\sqrt{1 + |Du|^2}$, we have

$$J(S) = \int_{\Omega} F(x, u(x), Du(x), -1) dx.$$

Notice that here we have used the relation $dA = \sqrt{1 + |Du|^2} dx$. The expression on the right can be considered as a nonparametric functional, defined for any $u \in C^2(\bar{\Omega})$. The Euler-Lagrange equation for this nonparametric functional is

$$(A6) \quad \sum_{i=1}^2 D_i [F_{q_i}(x, u, Du, -1)] - D_{x_3} F(x, u, Du, -1) = 0.$$

By using the chain rule and the homogeneity condition (A3), one can easily check that this equation can be written in the form

$$a_{ij}(x, u, Du) D_{ij} u = b(x, u, Du),$$

where

$$(A7) \quad a_{ij}(x, u, Du) = |q| D_{q_i q_j} F(x, u, Du, -1), \quad i, j = 1, 2,$$

$$(A8) \quad b(x, u, Du) = -|q| \sum_{i=1}^3 D_{q_i x_i} F(x, u, Du, -1).$$

By using (A3), (A5) it is not difficult to check that (1.2), (1.3) hold with constants γ and μ depending on F . That is, the *nonparametric Euler-Lagrange equation for an elliptic parametric functional is an equation of mean curvature type*.

Jenkins [5] and Jenkins-Serrin [6] consider equations of the form (A6) in case $F(X, q)$ does not depend on X . Note (A6) has the form (1.1) with $b \equiv 0$ in this case; hence all the results of §2 apply.

Finally we wish to point out that the functions a_{ij}^* , b^* introduced in §2 have a natural interpretation in the present context. In fact one can easily check that in case a_{ij} , b are as in (A7), (A8), then a_{ij}^* , b^* are given by

$$a_{ij}^*(X, \nu) = D_{q_i q_j} F(X, \nu), \quad i, j = 1, 2, 3,$$

$$b^*(X, \nu) = -|q| \sum_{i=1}^3 D_{q_i x_i} F(X, \nu);$$

furthermore the conditions (2.3), (3.22) hold automatically with δ, α determined by F , provided that $F \in C^3(\mathbf{R}^3 \times (\mathbf{R}^3 - \{0\}))$.

REFERENCES

1. L. Bers, *Isolated singularities of minimal surfaces*, Ann. of Math., (2) **53** (1951), 364-386.
2. R. Finn, *Equations of minimal surface type*, Annals of Math., **60** (1954), 397-416.
3. ———, *Isolated singularities of solutions of non-linear partial differential equations*, Trans. Amer. Math. Soc., **75** (1953), 383-404.
4. E. Heinz, *Über die Lösungen der Minimal flächengleichung*, Nachr. Akad. Wiss. Göttingen Math. Phys. Kl., **II** (1952), 51-56.
5. H. Jenkins, *On 2-dimensional variational problems in parametric form*, Arch. Rat. Mech. Anal., **8** (1961), 181-206.
6. H. Jenkins, J. Serrin, *Variational problems of minimal surface type I*, Arch. Rat. Mech. Anal., **12** (1963), 185-212.
7. C. B. Morrey, *Multiple integrals in the calculus of variations*, Springer-Verlag.
8. L. Simon, *A Hölder estimate for quasiconformal maps between surfaces in Euclidean space*. To appear in Acta Math.
9. ———, *Isolated singularities of graphs with quasiconformal Gauss map*. In preparation.
10. ———, *Global estimates of Hölder continuity for a class of divergence-form elliptic equations*, Arch. Rat. Mech. Anal., **56** (1974), 253-272.
11. J. Spruck, *Gauss curvature estimates for surfaces of constant mean curvature*, Comm. Pure Appl. Math., **27** (1974).

Received August 16, 1976. This research was supported in part by NSF Grant MPS-75-23332 and a Sloan Foundation Fellowship.

STANFORD UNIVERSITY
STANFORD, CA 94305

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

V. V. Anh and P. D. Tuan, <i>On starlikeness and convexity of certain analytic functions</i>	1
Willard Ellis Baxter and L. A. Casciotti, <i>Rings with involution and the prime radical</i>	11
Manuel Phillip Berriozabal, Hon-Fei Lai and Dix Hayes Pettey, <i>Noncompact, minimal regular spaces</i>	19
Sun Man Chang, <i>Measures with continuous image law</i>	25
John Benjamin Friedlander, <i>Certain hypotheses concerning L-functions</i>	37
Moshe Goldberg and Ernst Gabor Straus, <i>On characterizations and integrals of generalized numerical ranges</i>	45
Pierre A. Grillet, <i>On subdirectly irreducible commutative semigroups</i>	55
Robert E. Hartwig and Jiang Luh, <i>On finite regular rings</i>	73
Roger Hugh Hunter, Fred Richman and Elbert A. Walker, <i>Finite direct sums of cyclic valued p-groups</i>	97
Atsushi Inoue, <i>On a class of unbounded operator algebras. III</i>	105
Wells Johnson and Kevin J. Mitchell, <i>Symmetries for sums of the Legendre symbol</i>	117
Jimmie Don Lawson, John Robie Liukkonen and Michael William Mislove, <i>Measure algebras of semilattices with finite breadth</i>	125
Glenn Richard Luecke, <i>A note on spectral continuity and on spectral properties of essentially G_1 operators</i>	141
Takahiko Nakazi, <i>Invariant subspaces of weak-* Dirichlet algebras</i>	151
James William Pendergrass, <i>Calculations of the Schur group</i>	169
Carl Pomerance, <i>On composite n for which $\varphi(n) \mid n - 1$. II</i>	177
Marc Aristide Rieffel and Alfons Van Daele, <i>A bounded operator approach to Tomita-Takesaki theory</i>	187
Daniel Byron Shapiro, <i>Spaces of similarities. IV. (s, t)-families</i>	223
Leon M. Simon, <i>Equations of mean curvature type in 2 independent variables</i>	245
Joseph Nicholas Simone, <i>Metric components of continuous images of ordered compacta</i>	269
William Charles Waterhouse, <i>Pairs of symmetric bilinear forms in characteristic 2</i>	275