

Pacific Journal of Mathematics

SEMIGROUPS WITH IDENTITY ON PEANO CONTINUA

WILLIAM W. WILLIAMS

SEMIGROUPS WITH IDENTITY ON PEANO CONTINUA

W. WILEY WILLIAMS

A continuum is cell-cyclic if every cyclic element is a finite dimensional cell. We show that any finite dimensional cell-cyclic Peano continuum X admits a commutative semigroup with zero and identity, and apply this to show that if X is also homogeneous it is a point.

In [12] we showed that each cell-cyclic Peano continuum (locally connected metric continuum every cyclic element of which is a finite dimensional cell) X admits a semilattice (commutative idempotent topological semigroup). We now extend this result to show that X admits a commutative semigroup with identity and zero, and then apply this to homogeneous continua. Our extension is a partial answer to a question first raised by R. J. Koch in [6].

A semilattice is also a partially ordered Hausdorff topological space in which every two elements have a greatest lower bound and the function $(x, y) \rightarrow glb\{x, y\}$ is continuous. For $A \subset S$, let $L(A) = \{z: z \leq x \text{ for some } x \in A\}$ and $M(A) = \{y: x \leq y \text{ for some } x \in A\}$. A set A is *increasing* if $M(A) = A$. An *arc chain* is a totally ordered subset of a semilattice whose underlying space is an arc. We reserve I for the unit interval under min multiplication, and T for the quotient semilattice obtained by identifying $(0, 0)$ and $(1, 0)$ in $\{0, 1\} \times I$. Note that I^n and T^n , under coordinatewise multiplication, are semilattices with identity on the n -cell, with zero in the boundary and interior respectively.

Let X be a cell-cyclic Peano continuum. We use the cyclic element notation and results of Whyburn [10] and Kuratowski and Whyburn [8], slightly modified in the following way. In X we say a set A separates a and b if each arc from a to b meets A . $C(p, q)$ denotes the *cyclic chain from p to q* and is $\{x \in X \mid \text{some arc from } p \text{ to } q \text{ contains } x\}$. A subcontinuum A of X is an *A-set* if each arc in X having end points in A is contained in A . Cyclic elements and cyclic chains are A -sets. Given a point x and an A -set A , if $x \notin A$ there is a unique element $y \in A$ such that y separates each element of A from x . Denote this y by $P(A, x)$. If $x \in A$ set $P(A, x) = x$. Then for a fixed A -set A the function $x \rightarrow P(A, x)$ is a monotone retraction of X onto A mapping $X \setminus A$ into $Fr(A) = \{x \in A \mid x \notin D^\circ \text{ for any cyclic element } D \text{ of } A\} \cup \{\text{cut points of } A\}$. A set M is *nodal* in X if $M \cap (X \setminus M)^*$ contains at most one point. A point is an *end point* of X if it has a basis of neighborhoods having one point

boundary. A *node* of X is either (i) a true cyclic element which is a nodal set or (ii) an endpoint. By $\text{Com}(x, A)$ we mean the component of x in A . The interior of A is denoted by A° .

I. Preliminary results.

THEOREM 1.1. [12]. *Any cell-cyclic Peano continuum admits the structure of a semilattice.*

We note that in the proof of 1.1 given in [12], I^n and T^n , as defined above, could have been used for the semilattice structures on the individual cyclic elements. Thus the structure may be so constructed that each cyclic element is a semilattice with identity; also the zero may be chosen to be any predetermined point.

The following is an unpublished result due to Phyrne Bacon. We include a proof for completeness.

THEOREM 1.2. *Let X be a compact semilattice and C an arc chain containing 0. If Π_c is defined by $\Pi_c(x) = \sup\{a \in C \mid x \in M(a)\}$, then*

- (i) Π_c is a homomorphism from X onto C
- (ii) Π_c is continuous iff whenever $x, y \in C$ and $x < y$ then $y \in M(x)^\circ$.

Proof. X compact implies Π_c is well-defined. For (i), first note that Π_c is order preserving. Let $x, y \in X$ and suppose $\Pi_c(x) \leq \Pi_c(y)$. Since Π_c is order preserving we have $\Pi_c(xy) \leq \Pi_c(x)$. If $\Pi_c(xy) < \Pi_c(x)$, then there exists $z \in C$ such that $\Pi_c(xy) < z < \Pi_c(x)$. Thus $x \in M(z)$ and $xy \notin M(z)$. But $\Pi_c(x) < \Pi_c(y)$ and $x \in M(z)$ implies $y \in M(z)$. We conclude $xy \in M(z)$, a contradiction. Thus

$$\Pi_c(xy) = \Pi_c(x) = \Pi_c(x)\Pi_c(y).$$

By symmetry, if $\Pi_c(y) \leq \Pi_c(x)$ then the same conclusion is reached, and Π_c is a homomorphism.

For (ii), suppose whenever $x, y \in C$ and $x < y$, then $y \in M(x)^\circ$. For each $x \in C$ define $V(x) = X \setminus M(x)$. Then each $V(x)$ is open, and we claim that $x < y$ implies $V(x)^* \subset V(y)$. First note that $M(M(x)^\circ)$ is open by the continuity of multiplication, contains $M(x)^\circ$, and is contained in $M(x)$. Thus $M(M(x)^\circ) = M(x)^\circ$, and $M(x)^\circ$ is increasing. So if $x < y$, then $y \in M(x)^\circ$, and $M(y) \subseteq M(M(x)^\circ) = M(x)^\circ$. Thus $V(y) = X \setminus M(y)$ contains $X \setminus M(x)^\circ = [X \setminus M(x)]^* = V(x)^*$. Since C is an arc chain, $\inf\{a \in C \mid x \in V(a)\} = \sup\{a \in C \mid x \in M(a)\} = \Pi_c(x)$. Thus a proof like that for Urysohn's lemma [3] shows Π_c is continuous. This completes the proof.

It is implicit in results of Lawson [9] that if X is a semilattice on a finite dimensional Peano continuum, then (i) each point of X lies on an arc chain C containing 0, and (ii) if $x < y$ in C , then $y \in M(x)^\circ$. We conclude

COROLLARY 1.3. *Each point of a finite dimensional Peano continuum X lies on an arc chain C containing 0 and there is a homomorphic retraction of X onto C .*

THEOREM 1.4. *Any finite dimensional cell-cyclic chain $C(p, q)$ admits a semilattice with identity. Moreover, if $q \in Fr(C(p, q))$ then q can be chosen to be the identity.*

Proof. Note that the true cyclic elements of $C(p, q)$ form a countable collection $\{D_i\}$. We consider two cases:

Case 1. Some true cyclic element D_0 of $C(p, q)$ contains q . Then D_0 admits a semilattice structure with zero $a = P(D, p) \neq q$ and identity e . Moreover if $q \in Fr(C(p, q))$ then $q \in Fr(D_0)$, so we may choose $e = q$. By 1.1, $C(p, a)$ admits a semilattice in which each cyclic element D_i is a semilattice with identity e_i and zero $P(D_i, p)$. In each D_i there is an arc chain T_i from e_i to $b_i = P(D_i, q)$ and also an arc chain T_0 in D_0 from e to a and a homomorphism $h: D_0 \rightarrow T_0$ which is a retraction. Let $f_i: T_0 \rightarrow T_i$ be an onto homomorphism for each i . Now define a semilattice structure $*$ on $C(p, q)$ to agree with those on $C(p, a)$ and D_0 and such that if $x \in C(p, a)$ and $y \in D$ then

$$x*y = y*x = \begin{cases} x & \text{if } x \text{ is a cut point of } C(p, a) \\ x \cdot f_i(h(y)) & \text{if } x \in D_i \end{cases}$$

This obviously idempotent and commutative. Associativity and continuity follow since h and f_i are homomorphisms and continuous. Note that e is an identity for $*$.

Case 2. q is not in any true cyclic element of $C(p, q)$. Then there is a sequence $\{c_i\}$ of distinct cut points of $C(p, q)$ such that $\{c_i\} \rightarrow q$ and c_{i+1} separates c from q . This implies

$$C(p, q) \setminus \bigcup_{i=1}^{\infty} C(c_i, c_{i+1}) = \{q\}.$$

Endow each $C(c_i, c_{i+1})$ with a semilattice structure as in 1.1 so that c_i is the zero of $C(c_i, c_{i+1})$ and each cyclic element D_j is a semilattice with zero $P(D_j, p)$ and identity e_j , and let T_j be a (possibly degenerate)

are chain in D_j from e_j to $P(D_j, q)$. Let S_i be an arc chain in $C(c_i, c_{i+1})$ from c_i to c_{i+1} and let $h_i: C(c_i, c_{i+1}) \rightarrow S_i$ be a homomorphism and retraction. For each $i, j \in \mathbb{Z}^+$, let $f_{i,j}: S_i \rightarrow T_j$ be an onto homomorphism. Now define an operation $*$ on $C(p, q)$ to agree with that on each $C(c_i, c_{i+1})$ and such that if $x \in C(c_m, c_{m+1})$ and $y \in C(c_n, c_{n+1})$ then

$$y*x = x*y = \begin{cases} x & \text{if } x \text{ is a cut point and } n = m + 1 \\ x & \text{if } n > m + 1 \\ x f_{n,j}(h_n(y)) & \text{if } x \text{ is not a cut point} \\ & \text{(i.e., } x \in D_j \text{ for some } j \in \mathbb{Z}^+) \text{ and} \\ & n = m + 1 \\ xy & \text{if } n = m \end{cases}$$

Define q to be an identity for $C(p, q)$.

This is obviously idempotent and commutative. The proof of associativity is similar to that in Case 1 except in the following case: Suppose $x \in C(c_n, c_{n+1})$, $y \in C(c_{n+1}, c_{n+2})$ and $z \in C(c_{n+2}, c_{n+3})$. If x is a cut point, then $x*y*z = x$ in any order, and if y is a cut point then $x*y*z = x*y$ in any order. If neither is a cut point then $x \in D_z$ and $y \in D_k$ for some true cyclic elements D_j and D_k . So

$$(x*y)*z = x*y = x f_{n+1,j}(h_{n+1}(y)).$$

Now $x*(y*z) = x*(y f_{n+2,k}(h_{n+2}(z))) = x f_{n+1,j}(h_{n+1}(y f_{n+2,k}(h_{n+2}(z))))$. But $h_{n+1}(y f_{n+2,k}(h_{n+2}(z))) = h_{n+1}(y) h_{n+1}(f_{n+2,k}(h_{n+2}(z)))$ since h_{n+1} is a homomorphism. Also $h_{n+1}(y) \leq P(D_k, q) = h_{n+1}(f_{n+2,k}(h_{n+2}(z)))$ since $S_{n+1} \cap D_k$ is an arc chain with maximum element $P(D_k, q)$ and T_k is an arc chain with minimum element $P(D_k, q)$. It follows that

$$x*(y*z) = x f_{n+1,j}(h_{n+1}(y)) = x*y = (x*y)*z.$$

Suppose $x_n \rightarrow x$ and $y_n \rightarrow y$. If $x \neq q \neq y$, then one can prove $x_n*y_n \rightarrow xy$ using the continuity of the functions h_i and $f_{n,j}$ and the fact that the cyclic chains $C(c_i, c_{i+1})$ meet only at cut points. If $x = q \neq y$ and $y \in C(c_i, c_{i+1})$ then eventually $c_{i-1} \leq y_n \leq c_{i+2}$ and $c_{i+4} \leq x_n$ so that $x_n*y_n = y_n \rightarrow y = xy$. If $x = q = y$ and if $W(x_n, y_n)$ denotes the smaller of i and j where $x_n \in C(c_i, c_{i+1})$ and $y_n \in C(c_j, c_{j+1})$ then $W(x_n, y_n) \rightarrow \infty$ as $n \rightarrow \infty$. Since $x_n*y_n \in C(c_{W(x_n, y_n)}, c_{W(x_n, y_n)+1})$ and since $C(p, q)$ is locally connected we conclude that $x_n*y_n \rightarrow q = xy$. This completes the proof.

We note that in Case 2, if c_{n+1} separates x from p and c_n separates y from q then $x*y = y$.

II. Ruled continua.

DEFINITION 2.1. Suppose X is a topological space and $E \subseteq X$,

$0 \in X$. Let $A = \{[0, e] : e \in E\}$ be a collection of arcs in X satisfying:

- (i) $X = \bigcup \{[0, e] : e \in E\}$.
- (ii) $[0, e] \cap [0, f]$ is a proper subarc of each when e and f are distinct elements of E .
- (iii) For each $e \in E$, there is a unique $[0, e] \in A$.
- (iv) If $x_\alpha \rightarrow x$ then $[0, x_\alpha] \rightarrow [0, x]$ in the sense of $\lim \sup\text{-}\lim \inf$ convergence.

Then A is said to be a *ruling* of X and X is said to be a *ruled space* with zero 0 . The concept of a ruled space was introduced by Eberhart in his dissertation [4]. Spaces admitting a stronger type of ruling have been studied by Koch and McAuley [7]. We note that if X is ruled then for each $x \in X$ there is a unique arc $[0, x]$ which is contained in every $[0, e]$ containing x .

DEFINITION 2.2. A metric d is *radially convex* with respect to a partial order \leq on X if $x \leq y$, $y \leq z$ and $y \neq z$ imply $d(x, y) < d(x, z)$.

LEMMA 2.3. Let X be a compact metric ruled space. Define $x \leq y$ iff $x \in [0, y]$. Then \leq is a closed partial order on X . Moreover X admits a metric radially convex with respect to this order, so that if $r \leq d(0, e)$ there is a unique $x(r) \in [0, e]$ such that $d(0, x(r)) = r$.

Proof. This is clearly a partial order; that it is closed follows from property iv) of ruled spaces. By a result of Carruth [2], X admits a metric radially convex with respect to this order. The lemma now follows.

THEOREM 2.4. Any cell-cyclic Peano continuum X admits a ruling, and 0 may be chosen to be any point of X .

Proof. By 1.1, X admits a semilattice with zero 0 chosen arbitrarily. As in the proof of 1.1 given in [12], for each true cyclic element D of X let h_D denote the homeomorphism from I^n or T^n to D used to define this semilattice. Set $E = \text{Fr}(X) \setminus (\text{cut points of } X) \cup \{0\}$. For each $e \in E$ and each true cyclic element D of $C(0, e)$ define $T(D, e)$ to be the image under h_D of the straight line segment $[h_D^{-1}(P(D, 0)), h_D^{-1}(P(D, e))]$ in I^n or T^n . Then define $[0, e] = (\bigcup \{T(D, e) : D \in C(0, e)\}) \cup \{\text{cut points of } C(0, e)\}$. Then $[0, e]$ is a metric, compact (since $C(0, e) \setminus [0, e]$ is open in $C(0, e)$) order dense chain in the semilattice X and hence an arc. We now show the four conditions are satisfied.

(i) $X = U\{0, e\}: e \in E\}$. If $x \in X \setminus E$ then x is either an interior point of some cyclic element D of X or a cut point of X . If x is an interior point of D then $x \in h_D(h_D^{-1}(P(D, 0), h_D^{-1}(e)))$ for some $e \in \text{Fr}(D)$. If $e \in E$ then $x \in T(D, e) \subseteq [0, e]$. If $e \in E$ then choosing an end element e' of a component of $X \setminus \{e\}$ other than the one containing 0 , $x \in [0, e']$

(ii) and (iii) are clear

(iv) If $e_\alpha \rightarrow e$, then $[0, e_\alpha] \rightarrow [0, e]$. This follows from the fact that $[0, e_\alpha] \subseteq L(e_\alpha)$ and from techniques like those in [12]. We omit the details.

THEOREM 2.5. *Any cell-cyclic Peano continuum with a nodal cyclic element admits a commutative semigroup with identity and zero.*

Proof. Let X be a cell cyclic Peano continuum and suppose $X = C \cup D$, where $C \cap D = \{0\}$ and D is a true cyclic element. Then C is a cell-cyclic Peano continuum and hence admits a ruling $A = \{[0, e]: e \in E\}$ with zero 0 and a radially convex metric. Let h be a homeomorphism from I^n or T^n to D , depending on whether 0 is in the boundary or interior of D , and define a semilattice with identity e on D using h . Then there is in D an arc chain S from 0 to e and a retraction $f: D \rightarrow S$ which is a homomorphism. Moreover we may assume that S is radially convex so that for $x, y \in S$, $d(0, xy) = \min\{d(0, x), d(0, y)\}$. Without loss of generality we may assume $d(0, e)$ is maximal among $\{d(0, x) | x \in X\}$. Now define a semigroup on X by

$$y * x = x * y = \begin{cases} 0 & \text{if } x, y \in C \\ xy & \text{if } x, y \in D \\ \text{The point in } [0, x] \text{ of distance } r = \\ \min\{d(0, x), d(0, f(y))\} \text{ from } 0 & \text{if } x \in C, y \in D. \end{cases}$$

Associativity is obvious in all cases except the following: Suppose $x \in C$ and $y, z \in D$. Then $(x * y) * z$ is the point in $[0, x]$ of distance $\min\{d(0, x), d(0, f(y)), d(0, f(z))\}$ from 0 , whereas $x * (y * z)$ is the point in $[0, x]$ of distance $\min\{d(0, x), d(0, f(y, z))\}$ from 0 . But $d(0, f(yz)) = d(0, f(y)f(z)) = \min\{d(0, f(y)), d(0, f(z))\}$ so $(x * y) * z = x * (y * z)$. Continuity follows from the properties of ruled spaces and the fact that f is continuous. It is clear that e is an identity and 0 a zero. This completes the proof.

We conjecture that any X as in 2.5 admits a semilattice with identity. In fact, if X can be embedded in a plane then X can be embedded in a two-cell N and ruled in such a way that $X \cap \text{Fr}(N)$ is one of the arcs ruling X . One can now apply a theorem from

Eberhart's dissertation to show that X admits a semilattice with identity.

III. Cell-cyclic Peano continua without a nodal cyclic element. The goal of this section is a result like 2.5 for finite dimensional cell-cyclic Peano continua without a nodal cyclic element.

LEMMA 3.1. *Let X be a cell-cyclic Peano continuum. Then there exist two sequences $\{p_i\}$ and $\{q_i\}$ in $Fr(X)$, with p_1 and q_1 chosen arbitrarily, such that*

(i) *If we set $H_n = \bigcup_{i=1}^n C(p_i, q_i)$, then for each $n > 1$, $\{p_n\} = C(p_n, q_n) \cap H_{n-1}$*

(ii) *If we set $H = \bigcup_{n=1}^{\infty} H_n$, then each point of $X \setminus H$ is an end point of X , and so $H^* = X$.*

(iii) *The diameter of the components of $S \setminus H_n$ tends to 0 uniformly with $1/n$.*

Proof. This was proved by Whyburn ([10], p. 73) without the condition that $\{p_i\}$ and $\{q_i\}$ are in $Fr(X)$. We show this condition can also be assumed. Whyburn's proof considers a dense sequence $\{r_i\}$ and sets $p_1 = r_1$, $q_1 = r_2$. Clearly these may be chosen arbitrarily in $Fr(X)$. In Whyburn's proof, for $j > 1$ q_j is the r_i of smallest index such that $r_i \notin H_{j-1}$ and $p_j = P(H_{j-1}, q_j)$. Thus $p_j \in Fr(X)$. If $q_j \notin Fr(X)$, then q_j is an interior point of some true cyclic element D . Let q'_j be any point in $Fr(D)$ other than $P(D, p_j)$. Then $C(p_j, q_j) = C(p_j, q'_j)$, so we may assume $q_j \in Fr(X)$. The lemma follows.

Now let X be a finite dimensional cell-cyclic Peano continuum without a nodal cyclic element. Then X has at least 2 end points ([10], p. 77); let 0 and 1 denote end points of X . Let $\{p_i\}$, $\{q_i\}$, $\{H_n\}$, and H be as described in 3.1, with $p_1 = 0$, $q_1 = 1$. Each $C(p_i, q_i)$ admits a semilattice with zero p_i and identity q_i by 1.3. We now define inductively an algorithm for defining a semilattice with identity on H .

Let $\{c_j\}$ be the sequence of cut points of $C(0, 1)$ converging to 1 such that c_{j+1} separates c_j from 1 used in 1.3 to define the semilattice on $C(0, 1)$. Let n_1 be one more than the smallest i such that c_i separates p_2 from 1 in X . Set $Q_1 = C(p_2, q_2)$, $P_1 = [\text{Com}(1, C(0, 1) \setminus \{c_{n_1}\})]^*$, and $R_1 = [\text{Com}(0, C(0, 1) \setminus \{c_{n_1}\})]^*$. Let T_1 be an arc chain from p_2 to q_2 in Q_1 and S_1 be an arc chain from c_{n_1} to 1 in $C(0, 1)$. Let $f_1: S_1 \rightarrow T_1$ be a continuous onto homomorphism such that $f_1^{-1}(q_2) = M(c_{n_1+1}) \cap S_1$, and let $h_1: P_1 \rightarrow T_1$ be the continuous onto homomorphism obtained by composing f_1 and a homomorphic retraction r_1 of $C(c_{n_1+1}, 1)$ onto S_1 . We now define a semilattice $*$ on $H_2 = C(p_1, q_1) \cup C(p_2, q_2) = H_1 \cup Q_1$ by

$$x*y = y*x = \begin{cases} xy & \text{if } x, y \in C(0, 1) = P_1 \cup R_1 \text{ or } x, y \in Q_1 \\ xp_2 & \text{if } x \in R_1, y \in Q_1 \\ h_1(x)y & \text{if } x \in P_1, y \in Q_1 \end{cases}$$

where juxtaposition means whichever of the previously defined operations on H_1 or Q_1 fits the context.

Associativity is clear in all cases except when $r \in R_1$, $p \in P_1$, $q \in Q_1$. In this case $r*(p*q) = r*(h_1(p)q) = rp_2$, whereas

$$(r*p)*q = (rp)p_2 = r(pp_2) = rp_2$$

by the note at the end of Section I. Continuity is easily checked since P_1 , Q_1 and R_1 meet only at cut points of X . Note that any point in $C(c_{n_2+1}, 1)$ acts as an identity for any point in

$$[\text{Com}(0, H_2 \setminus \{c_{n_1}\})]^*,$$

and 1 acts as an identity for all of H_2 .

Suppose that a semilattice structure with zero 0 and identity 1 has been defined on H_{k-1} so that the structure agrees with those on $C(P_i, q_i)$ for each $i \leq k$. Also suppose $c_{n_{k-1}} \in \{c_i\}$ has been chosen so that any element of $[\text{Com}(1, H_{k-1} \setminus \{c_{n_{k-1}+1}\})]^*$ acts as an identity for any element $[\text{Com}(0, H_{k-1} \setminus \{c_{n_{k-1}}\})]^*$.

Let n_k be one more than the smallest integer greater than n_{k-1} such that c_{n_k} separates p_{k+1} from 1. Set $Q_k = C(p_{k+1}, q_{k+1})$, $P_k = C(c_{n_k}, 1) = [\text{Com}(1, H_k \setminus \{c_{n_k}\})]^*$, and $R_k = [\text{Com}(0, H_k \setminus \{c_{n_k}\})]^*$. Let T_k be an arc chain from P_{k+1} to q_{k+1} in Q_k and $S_k = S_1 \cap P_k$. Let $f_k: S_k \rightarrow T_k$ be a continuous onto homomorphism such that

$$f_k^{-1}(q_{k+1}) = M(c_{n_{k+1}}) \cap S_k$$

in P_k , and let $h_k: P_k \rightarrow T_k$ be a continuous onto homomorphism obtained by composing f_k and the homomorphic retraction $r_k = r_1|_{P_k}$ of $C(c_{n_k}, 1) = P_k$ onto S_k . We now define a semilattice $*$ with identity 1 on H_k by

$$x*y = y*x = \begin{cases} xy & \text{if } x, y \in H_{k-1} \text{ or } x, y \in Q_k \\ xp_k & \text{if } x \in R_k, y \in Q_k \\ h_k(x)y & \text{if } x \in P_k, y \in Q_k \end{cases}$$

where juxtaposition means whichever of the previously defined operations on H_k or Q_k fits the context.

Again associativity is clear in all cases except when $r \in R_k$, $p \in P_k$, $q \in Q_k$. In this case $r*(p*q) = r*(h_k(p)q) = rp_k$, whereas $(r*p)*q = (rp)p_k = r(pp_k)$ since the operation on H_{k-1} is associative. But $p \in [\text{Com}(1, H_{k-1} \setminus \{c_{n_{k-1}+1}\})]^*$ and $p_k \in [\text{Com}(0, H_{k-1} \setminus \{c_{n_{k-1}}\})]^*$ so by hypothesis $pp_k = p_k$, and $r*(p*q) = (r*p)*q$. Continuity is again

easily checked. Again any point in $[\text{Com}(1, H_k \setminus \{c_{n_k+1}\})]^*$ acts as an identity for any element $[\text{Com}(0, H_k \setminus \{c_{n_k}\})]^*$. By induction we have proved the following:

LEMMA 3.2. *Each H_n admits a semilattice with zero 0 and identity 1 so that the operations agree whenever possible.*

LEMMA 3.3. *The function $P(H_n, \cdot): H \rightarrow H_n$ is a retraction and a homomorphism for each n .*

Proof. It has been previously noted that each $P(H_n, \cdot)$ is a retraction. To show that each is a homomorphism it suffices to show that the restriction of $P(H_n, \cdot)$ to H_{n+1} is a homomorphism, since $P(H_n, \cdot)$ is the composition of this restriction and $P(H_{n+1}, \cdot)$. Let $x, y \in H_{n+1} = H_n \cup Q_n$. If $x, y \in Q_n$ then

$$P(H_n, x) * P(H_n, y) = p_n * p_n = p_n = P(H_n, x * y)$$

since $x * y \in Q_n$. If $x \in Q_n, y \in H_n$ then there are two cases. If $y \in P_n$ then $P(H_n, x) * P(H_n, y) = p_n * y = p_n$ since $p_n \in R_{n-1}$ by definition and any element of P_n acts as an identity for any element of R_{n-1} . However $P(H_n, x * y) = P(H_n, x * h_n(y)) = p_n$ since $x * h_n(y) \in Q_n$. If $y \in R_n$ then $P(H_n, x) * P(H_n, y) = p_n * y = x * y = P(H_n, x * y)$. This completes the proof of the lemma.

LEMMA 3.4. *Let X be as above and let $x, y \in X$, and suppose $\{x_n\}, \{y_n\}$ are sequences in H such that $x_n \rightarrow x, y_n \rightarrow y$. Then there exists $z \in X$ such that $\{x_n * y_n\} \rightarrow z$, where $*$ denotes the operation on any H_n containing x_n and y_n , and z is independent of the choice of the sequences.*

Proof. We distinguish four cases.

Case I. $x = y = 1$. From the definition of multiplication on H , if $a, b \in P_k = [\text{Com}(1, H \setminus \{c_k\})]^*$ then $a * b \in P_k$. Now $\{P_k\}$ forms a neighborhood basis at the end point 1. Since both $\{x_n\}$ and $\{y_n\}$ are eventually in each P_k , $\{x_n * y_n\}$ is eventually in each P_k and hence $\{x_n * y_n\} \rightarrow 1$.

Case II. x, y , and 1 all distinct. Let N be an integer so large that $P(H_0, x)$ and $P(H_0, y)$ are in $\text{Com}(0, H_0 \setminus \{c_N\})$ and that the diameter of any component of $X \setminus H_N < d(x, y)/2$. This implies $\text{Com}(x, X \setminus H_N)$ and $\text{Com}(y, X \setminus H_N)$ are disjoint open sets, and we may assume $x_N \in \text{Com}(x, X \setminus H_N)$ and $y_n \in \text{Com}(y, X \setminus H_N)$ for all n . Also we may assume $d(x_n, y_n) > d(x, y)/2$ for all n . We now show

$x_n * y_n = P(H_N, y_n) * P(H_N, y_n)$ for all n . The statement is obvious if $x_n, y_n \in H_N$. Suppose it is true whenever $x_n, y_n \in H_m$ for some $m \geq N$, and let $x_n, y_n \in H_{m+1} = H_m \cup Q_m$. If $x_n \in Q_m$ and $y_n \in H_m$ then $x_n * y_n = p_m * y_n$. By hypothesis, since $p_m, y_n \in H_m$ then

$$p_m * y_n = P(H_N, p_m) * P(H_N, y_n).$$

But $P(H_N, p_m) = P(H_N, x_n)$ since $Q_m \subset [\text{Com}(x_n, X \setminus H_N)]^*$. Thus $x_n * y_n = P(H_N, x_n) * P(H_N, y_n)$. By symmetry the statement is true if $x_n \in H_m$ and $y_n \in Q_m$. The statement is obvious if both $x_n, y_n \in H_m$, whereas the case $x_n, y_n \in Q_m$ is impossible for it implies $d(x_n, y_n) < d(x, y)/2 < d(x_n, y_n)$.

We know H_N is a semilattice and hence

$$x_n * y_n = P(H_N, x_n) * P(H_N, y_n) \longrightarrow P(H_N, x) * P(H_N, y)$$

since $P(H_N, \cdot)$ is continuous.

Case III. $x = y \neq 1$

(a) $x = y \notin H$. Then $x = y$ is an end point of X and $\{U_i\} = \{[\text{Com}(x, X \setminus H_i)]^*\}$ is a neighborhood basis at $x = y$. We show that if U_i is fixed and if $x_n, y_n \in U_i \cap H_N$ then $x_n * y_n \in U_i \cap H_N$, for any N . Note the statement is true for $N \leq i$. Suppose it is true whenever $x_n, y_n \in U_i \cap H_m$ for some $m \geq i$, and let

$$x_n, y_n \in U_i \cap H_{m+1} = U_i \cap (H_m \cup Q_m).$$

If $x_n \in Q_m$ and $y_n \in H_m$ then $x_n * y_n = p_m * y_n \in U_i \cap H_m \subset U_i \cap H_{m+1}$ by the induction hypothesis. By symmetry the statement is true if $x_n \in H_m$ and $y_n \in Q_m$. If $x_n, y_n \in Q_m$ then $x_n * y_n \in Q_m \subset U_i \cap H_{m+1}$, and if $x_n, y_n \in H_m$ the statement follows from the induction hypothesis.

Since $\{x_n\}$ and $\{y_n\}$ are eventually in each U_i , and since for each n and each i we can find $N(n, i)$ such that $x_n, y_n \in U_i \cap H_{N(n, i)}$, we conclude that $\{x_n * y_n\}$ is eventually in each U_i . Thus $\{x_n * y_n\} \rightarrow x = y$.

(b) $x = y \in H_N$, some N . Let $\varepsilon > 0$. There exists $L > N$ so that the diameter of any component of $X \setminus H_L$ is less than $\varepsilon/2$, and so that $B(x, \varepsilon/2) \cap P_L = \emptyset$. We may assume $d(x_n, x) < \varepsilon/2$ and $d(y_n, y) < \varepsilon/2$ for each n . Divide $\{x_n * y_n\}$ into two (perhaps finite) sequences: If $x_n * y_n \in H_L$ then

$$\begin{aligned} x_n * y_n &= P(H_L, x_n * y_n) \\ &= P(H_L, x_n) * P(H_L, y_n) \rightarrow P(H_L, x) * P(H_L, x) = xy = x = y, \end{aligned}$$

by Lemma 3.3 and the continuity of multiplication on H_L . If $x_n * y_n \notin H_L$, then $x_n \notin H_L$ and $y_n \notin H_L$ because $B(x, \varepsilon/2) \cap P_L = \emptyset$ and using the definition of multiplication on H . Also, using the definition of

multiplication $x_n * y_n \in \text{Com}(x_n, X \setminus H_L)$ or $x_n * y_n \in \text{Com}(y_n, X \setminus H_L)$. Thus

$$d(x, x_n * y_n) \leq d(x, x_n) + d(x_n, x_n * y_n) < \varepsilon$$

or

$$d(y, x_n * y_n) \leq d(y, y_n) + d(y_n, x_n * y_n) < \varepsilon.$$

In either case $d(x, x_n * y) = d(y, x_n * y_n) < \varepsilon$. We conclude that $\{x_n * y_n\} \rightarrow x = y$.

Case IV. $y \neq x = 1$. We first establish two facts.

(A) If $a, b \in H$ so that $P(H_0, a) \in \text{Com}(1, H_0 \setminus \{c_n\})$ and $P(H_0, b) \in \text{Com}(0, H_0 \setminus \{c_n\})$ for some n , then $a * b = P(H_0, a) * b$.

The proof is by the induction on the H_i containing a . It is clear for $a \in H_0$. Suppose the statement is true for $a \in H_m$, $m \geq 0$, and let $a \in H_{m+1} = H_m \cup Q_{m+1}$. Suppose $a \in Q_{m+1}$, for the induction hypothesis assures the statement is true if $a \in H_m$. Then since a and b are separated by c_m , $b \notin Q_{m+1}$. Hence $a * b = p_{m+1} * b$. But $p_{m+1} * b = P(H_0, p_{m+1}) * b$ by the induction hypothesis, and

$$P(H_0, p_{m+1}) = P(H_0, a),$$

so

$$a * b = P(H_0, a) * b.$$

Thus (A) is established.

(B) If $a, b \in H$ so that $a \in \text{Com}(1, H_0 \setminus \{c_n\})$ and $b \in \text{Com}(0, H_0 \setminus \{c_n\})$ for some n , then either $a * b = a * P(H_n, b)$ or $a * b \in \text{Com}(b, X \setminus H_n)^*$.

The proof is by induction on the H_i containing b . If $b \in H_n$ then $P(H_n, b) = b$ and the statement is true. Suppose the statement is true when $b \in K_m$ for some $m \geq n$, and let $b \in Q_{m+1}$. If $a \in \text{Com}(1, H_0 \setminus \{c_m\})$ then $a * b \in Q_{m+1} \subset \text{Com}(b, X \setminus H_n)^*$. If $a \in [\text{Com}(0, H_0 \setminus \{c_m\})]^*$ then $a * b = a * p_m$. But $a * p_m = a * P(H_n, p_m)$ by the induction hypothesis, and $P(H_n, p_m) = P(H_n, b)$. Thus $a * b = a * P(H_n, b)$ and (B) is established.

We now distinguish two subcases of Case IV.

Subcase 1. $y \in H_M$, some M . Let $\varepsilon > 0$. Choose M so large that c_M does not separate y from 0 and the diameter of any component of $X \setminus H_M$ is less than $\varepsilon/2$. We may assume that for each n , $P(H_0, y_n) \in \text{Com}(0, H_0 \setminus \{c_M\})$ and $P(H_0, x_n) \in \text{Com}(1, H_0 \setminus \{c_M\})$. Then by (A), $x_n * y_n = P(H_0, x_n) * y_n$, and by (B), $P(H_0, x_n) * y_n = P(H_0, x_n) * P(H_M, y_n)$ or $P(H_0, x_n) * y_n \in \text{Com}(b, X \setminus H_n)^*$. If the former then

$$x_n * y_n = P(H_0, x_n) * P(H_M, y_n) \longrightarrow 1 * P(H_M, y) = y$$

by the continuity of the multiplication on H_M and Lemma 3.3. In the latter case $d(P(H_0, x_n)*y_n, y_n) < \varepsilon/2$. We may assume $d(y_n, y) < \varepsilon/2$, so $d(y, P(H_0, x_n)*y_n) < \varepsilon$. Thus we conclude that $\{x_n*y_n\} \rightarrow y$.

Subcase 2. $y \notin H$. If $V_k = [\text{Com}(y, X \setminus H_k)]^*$ then $\{V_k\}$ is a neighborhood basis, so we need only show $\{x_n*y_n\}$ is eventually in each V_k . Fix a V_k . We may assume again that for each n , $P(H_0, y_n) \in \text{Com}(0, H_0 \setminus \{c_M\})$, $P(H_0, x_n) \in \text{Com}(1, H_0 \setminus \{c_M\})$, and $y_n \in V_k$ for some $M \geq k$. By (A) and (B), $x_n*y_n = P(H_0, x_n)*P(H_M, y_n)$ or $x_n*y_n \in \text{Com}(y_n, X \setminus H_M)^* \subset V_k$. However $P(H_M, y_n) \in V_k$, and $P(H_0, x_n) \in H_0$, so $P(H_0, x_n)*P(H_M, y_n) \in V_k$. This completes the proof of the lemma.

THEOREM 3.5. *Let X be a finite dimensional cell-cyclic Peano continuum without a nodal element. Then X admits a semilattice with identity.*

Proof. By the above, the dense set H admits a semilattice with identity. For each $x, y \in X$ let $\{x_n\} \rightarrow x$, $\{y_n\} \rightarrow y$ where $\{x_n\}, \{y_n\}$ are sequences in H . Define $xy = \lim \{x_n*y_n\}$. By 3.4 this limit exists and is independent of the choice of the sequences. It follows that this operation is a semilattice with identity on X . Combining this with Theorem 2.3 we have

COROLLARY 3.6. *Let X be a finite dimensional cell-cyclic Peano continuum. Then X admits a commutative semigroup with identity and zero.*

COROLLARY 3.6. *Any retract of a two-cell admits a commutative semigroup with identity.*

Proof. Borsuk [1] has shown that a subset X of a two-cell A is a retract of A if and only if A is a locally connected continuum which does not separate the plane. Whyburn [11] has shown that for locally connected continua in the plane, not separating the plane is equivalent to every cyclic element being a simple closed curve with interior, i.e., a two-cell. Thus a retract of a two-cell is a cell-cyclic Peano continuum, and the result follows from Corollary 3.6.

DEFINITION 3.8. A space X is homogeneous if for each pair of points x and y in X there is a homeomorphism of X onto itself carrying x to y .

THEOREM 3.9. *Any finite dimensional homogeneous cell-cyclic*

Peano continuum (in particular, any homogeneous retract of a two-cell) is a point.

Proof. By a result of Hudson and Mostert [5], any homogeneous compact connected semigroup with identity is a group. Combining this with Corollaries 3.6 and 3.7, unless X is a point X admits the structure of a group with two idempotents, a contradiction.

REFERENCES

1. K. Borsuk, *Theory of Retracts*, Polska Acad. Nauk Monografie Mat., Tom **44** (1967).
2. J. H. Carruth, *A note on partially ordered compacta*, Pacific J. Math., **24** (1968), 229-231.
3. J. Dugundji, *Topology*, Allyn and Bacon, 1966.
4. C. A. Eberhart, *Some classes of continua related to clan structures*, Doctoral Dissertation, Louisiana State University, 1966.
5. A. L. Hudson and P. L. Mostert, *A finite dimensional homogeneous clan is a group*, Annals of Math., **78** (1963), 41-46.
6. R. J. Koch, *Some open questions in topological semigroups*, An. Acad. Brasil. Ci., **41** (1969), 19-20.
7. R. J. Koch and L. F. McAuley, *Semigroups on continua ruled by arcs*, Fund. Math., **56** (1964), 1-8.
8. C. Kuratowski and G. T. Whyburn, *Sur les elements cycliques et leurs applications*, Fund. Math., **16** (1930), 305-331.
9. J. D. Lawson, *Topological semilattices with small semilattices*, J. London Math. Soc., (2), **1** (1969), 719-724.
10. G. T. Whyburn, *Analytic Topology*, Amer. Math. Soc. Colloq. Pub., **28** (1942).
11. ———, *Concerning the structure of the continuous curve*, Amer. J. Math., **50** (1928), 167-194.
12. W. W. Williams, *Semilattices on Peano Continua*, Proc. Amer. Math. Soc., **49** (1975), 495-500.

Received November 3, 1975. Portions of this paper generalize unpublished parts of a dissertation presented to the Graduate School of Louisiana State University. The author wishes to thank Professor R. J. Koch for encouraging the work on these results.

UNIVERSITY OF LOUISVILLE
LOUISVILLE, KY 40208

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, CA 90024

R. A. BEAUMONT
University of Washington
Seattle, WA 98105

C. C. MOORE
University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Carol Alf and Thomas Alfonso O'Connor, <i>Unimodality of the Lévy spectral function</i>	285
S. J. Bernau and Howard E. Lacey, <i>Bicontractive projections and reordering of L_p-spaces</i>	291
Andrew J. Berner, <i>Products of compact spaces with bi-k and related spaces</i>	303
Stephen Richard Bernfeld, <i>The extendability and uniqueness of solutions of ordinary differential equations</i>	307
Marilyn Breen, <i>Decompositions for nonclosed planar m-convex sets</i>	317
Robert F. Brown, <i>Cohomology of homomorphisms of Lie algebras and Lie groups</i>	325
Jack Douglas Bryant and Thomas Francis McCabe, <i>A note on Edelstein's iterative test and spaces of continuous functions</i>	333
Victor P. Camillo, <i>Modules whose quotients have finite Goldie dimension</i>	337
David Downing and William A. Kirk, <i>A generalization of Caristi's theorem with applications to nonlinear mapping theory</i>	339
Daniel Reuven Farkas and Robert L. Snider, <i>Noetherian fixed rings</i>	347
Alessandro Figà-Talamanca, <i>Positive definite functions which vanish at infinity</i>	355
Josip Globevnik, <i>The range of analytic extensions</i>	365
André Goldman, <i>Mesures cylindriques, mesures vectorielles et questions de concentration cylindrique</i>	385
Richard Grassl, <i>Multisectioned partitions of integers</i>	415
Haruo Kitahara and Shinsuke Yorozu, <i>A formula for the normal part of the Laplace-Beltrami operator on the foliated manifold</i>	425
Marvin J. Kohn, <i>Summability R_r for double series</i>	433
Charles Philip Lanski, <i>Lie ideals and derivations in rings with involution</i>	449
Solomon Leader, <i>A topological characterization of Banach contractions</i>	461
Daniel Francis Xavier O'Reilly, <i>Cobordism classes of fiber bundles</i>	467
James William Pendergrass, <i>The Schur subgroup of the Brauer group</i>	477
Howard Lewis Penn, <i>Inner-outer factorization of functions whose Fourier series vanish off a semigroup</i>	501
William T. Reid, <i>Some results on the Floquet theory for disconjugate definite Hamiltonian systems</i>	505
Caroll Vernon Riecke, <i>Complementation in the lattice of convergence structures</i>	517
Louis Halle Rowen, <i>Classes of rings torsion-free over their centers</i>	527
Manda Butchi Suryanarayana, <i>A Sobolev space and a Darboux problem</i>	535
Charles Thomas Tucker, II, <i>Riesz homomorphisms and positive linear maps</i>	551
William W. Williams, <i>Semigroups with identity on Peano continua</i>	557
Yukinobu Yajima, <i>On spaces which have a closure-preserving cover by finite sets</i>	571