

Pacific Journal of Mathematics

ONE-SIDED HEEGAARD SPLITTINGS OF 3-MANIFOLDS

J. HYAM RUBINSTEIN

ONE-SIDED HEEGAARD SPLITTINGS OF 3-MANIFOLDS

J. H. RUBINSTEIN

For a large class of closed orientable 3-manifolds, we define a new decomposition method which uses embedded one-sided surfaces and is analogous to Heegaard splittings. The technique is most useful for studying some "small" 3-manifolds (i.e., which have finite fundamental group or are not sufficiently large). We give several general criteria for existence of these splittings and some results on nonorientable surfaces in lens spaces. Also stable equivalence (as for Heegaard splittings) and a result of Waldhausen's are shown to carry over to the one-sided case.

O. Introduction. In [7] an example is given showing that the loop theorem is not valid for one-sided surfaces in 3-manifolds. For this reason, such surfaces are difficult to handle and have not been the object of much work. We would like to present a new approach based on the following:

DEFINITION. Let M be a closed orientable 3-manifold. A pair (M, K) is called a *one-sided Heegaard splitting* if K is a closed non-orientable surface embedded in M such that $M - K$ is an open handlebody.

REMARKS. (1) If (M, K) is a one-sided Heegaard splitting and if $N(K)$ is a small closed regular neighborhood of K , then $N(K)$ is homeomorphic to a twisted line-bundle over K . Also $M - \text{int } N(K)$ is a handlebody which we denote by Y .

(2) There is a double cover $p: \tilde{M} \rightarrow M$ naturally associated with a one-sided Heegaard splitting (M, K) . It is the covering of M corresponding to the subgroup $i_*\pi_1(M - K)$ of $\pi_1(M)$, where i is the inclusion. The surface $p^{-1}(K)$, which we denote by \tilde{K} , is the orientable double cover of K and (\tilde{M}, \tilde{K}) is a Heegaard decomposition (i.e., the closures of the components of $\tilde{M} - \tilde{K}$ are handlebodies). If $g: \tilde{M} \rightarrow \tilde{M}$ is the covering transformation for p then \tilde{K} is g -invariant and g interchanges the components of $\tilde{M} - \tilde{K}$.

We work in the PL category and let M be a closed orientable 3-manifold throughout. In §1 it is proved that there are one-sided splittings associated with any nonzero class in $H_2(M, \mathbb{Z}_2)$. In §2 one-sided decompositions are discussed where the surface K is (geometrically) incompressible. This is the most useful setting for the theory (cf. [4] for some results employing this approach).

In § 3 a method based on [1] is given for studying the relation in a lens space between a genus 1 Heegaard splitting and an embedded incompressible nonorientable surface. As applications, it is proved that any two incompressible surfaces in a lens space are isotopic and a simple derivation of the algorithm of [1] for the genus of these surfaces is given.

In § 4 we show that two one-sided splittings associated with the same class in $H_2(M, Z_2)$ are stably equivalent (cf. [6]). Finally in § 5 the standard form for maps of degree 1 in [9] is adapted to one-sided decompositions.

1. Existence of one-sided splittings.

DEFINITION. Suppose K is a closed nonorientable surface embedded in M . We denote by $[K]$ the class in $H_2(M, Z_2)$ which is the image of the generator of $H_2(K, Z_2)$ by the map induced by the embedding of K in M . If (M, K) is a one-sided Heegaard splitting then we call $[K]$ the class *associated* with (M, K) .

THEOREM 1. *For any element $\alpha \neq 0$ in $H_2(M, Z_2)$, there is a one-sided Heegaard splitting (M, K) with $[K] = \alpha$.*

Proof. The existence of a closed nonorientable surface K embedded in M with $[K] = \alpha$ follows as in [1]. Explicitly let $f: M \rightarrow RP^4$ be a map such that $f_*: H_1(M, Z_2) \rightarrow H_1(RP^4, Z_2)$ gives the element of $H^1(M, Z_2)$ corresponding to α by Poincaré duality. Assume f is transverse on the submanifold RP^3 . We can then join up the components of the surface $f^{-1}(RP^3)$ using elementary surgery, by a homotopy of f . This gives K .

As in Remark 2 above, there is a double cover $p: \tilde{M} \rightarrow M$ associated with K . Suppose M is triangulated with K as a full subcomplex. Then there is a lifted g -invariant triangulation of \tilde{M} . Let $\tilde{K} = p^{-1}(K)$ and let the vertices in \tilde{K} be divided into disjoint classes V and W with $W = gV$. We denote the closures of the components of $\tilde{M} - \tilde{K}$ by Y_1 and Y_2 .

Let Γ_i be the union of the 1-simplices of the triangulation of \tilde{M} with interior in $\text{int } Y_i$, $i = 1, 2$. Let $N(\Gamma_i)$ denote the closed simplicial regular neighborhood of Γ_i in Y_i , with respect to the second derived triangulation. Then $Y_i - \text{int } N(\Gamma_i)$ is the simplicial neighborhood of the dual 1-skeleton of Y_i and so is a handlebody.

Clearly $N(\Gamma_1) \cap \tilde{K} = N(\Gamma_2) \cap \tilde{K} = N(V) \cup N(W)$, with $N(V) \cap N(W) = \emptyset$ and $gN(V) = N(W)$. Let $\bar{N}(V)$ be a small closed regular neighborhood of $N(V)$ in Y_1 , chosen so that $\bar{N}(V) \cap g\bar{N}(V) = \emptyset$. By an ambient isotopy of Y_1 with support in $\bar{N}(V)$, we can shift

Γ_1 to Γ_1^* and $N(\Gamma_1)$ to $N(\Gamma_1^*)$ so that $N(\Gamma_1^*) \cap gN(\Gamma_1^*) = \emptyset$. We denote $g\Gamma_1^*$ by Γ_2^* and $gN(\Gamma_1^*)$ by $N(\Gamma_2^*)$.

Define Y_1^* as $(Y_1 - \text{int } N(\Gamma_1^*)) \cup N(\Gamma_2^*)$. Then Y_1^* , which is obtained by adding the handles of $N(\Gamma_2^*)$ to the handlebody $Y_1 - \text{int } N(\Gamma_1^*)$, is a handlebody. Also $\tilde{K}^* = \partial Y_1^*$ is g -invariant and so (\tilde{M}, \tilde{K}^*) is a g -invariant Heegaard splitting. Therefore (M, K^*) is a one-sided Heegaard splitting, where $K^* = p(\tilde{K}^*)$. It is clear that $[K] = [K^*]$, since the same double covering \tilde{M} is associated with K and K^* .

Note. $H_2(M, Z_2) \cong H^1(M, Z_2) \cong H_1(M, Z_2)$. So M has a one-sided Heegaard splitting if and only if $H_1(M, Z_2) \neq 0$.

2. Incompressible one-sided splittings.

DEFINITION. A closed surface $K \neq S^2$ embedded in M is *incompressible* if there is no disk D embedded in M with $D \cap K = \partial D$ a noncontractible curve in K .

REMARK. If $K \neq S^2$ is two-sided in M , then as is well-known by the loop theorem, K is incompressible if and only if the map $i_*: \pi_1(K) \rightarrow \pi_1(M)$ induced by the embedding is an injection. No similar such a nice criterion has been found for one-sided surfaces.

DEFINITION. M is *irreducible* if any 2-sphere embedded in M bounds a 3-cell.

LEMMA 2. *Suppose M is irreducible and K is an incompressible nonorientable surface embedded in M . The following are equivalent:*

- (1) $M - K$ is an open handlebody.
- (2) The map $i_*: \pi_1(K) \rightarrow \pi_1(M)$ is onto.
- (3) There is no incompressible orientable surface in M which is disjoint from K .

Proof. $1 \Rightarrow 2$ is clear.

$2 \Rightarrow 3$. Suppose J is an orientable incompressible surface embedded in $M - K$ and i_* is onto. If J is nonseparating in M then there is a loop C which intersects J in a single point. But C is homotopic to a loop in K , because $\pi_1(K) \rightarrow \pi_1(M)$ is onto. Therefore since $J \cap K = \emptyset$, we get a contradiction to the invariance of intersection number under homotopy.

On the other hand if J is separating in M then $J = \partial W$ with W contained in $M - K$. As i_* is onto, it follows that $\pi_1(J) \rightarrow \pi_1(W)$ is onto. This implies that J is a 2-sphere (since as is well-known,

$2\beta_1(W) \leq \beta_1(\partial W)$), contrary to our assumption.

$3 \Rightarrow 1$. Let $N(K)$ be a small closed regular neighborhood of K with $Y = M - \text{int } N(K)$ and $L = \partial N(K)$. We compress L to get L_1 , each of whose components is incompressible or is a 2-sphere. All the compressions can be made to occur in Y since K is incompressible, as in 14.12 of [3]. Consequently we obtain that L_1 is contained in Y .

By assumption there are no incompressible orientable surfaces in Y . Therefore we conclude that L_1 is a union of 2-spheres. Then as M is irreducible, these 2-spheres bound 3-cells and it easily follows that Y is a handlebody.

LEMMA 3. *Suppose $H_1(M, Z)$ is finite. If $\alpha \neq 0$ is in $H_2(M, Z_2)$ then there are disjoint nonorientable incompressible surfaces K_i in M with $\sum_{i=1}^n [K_i] = \alpha$. In particular there is a generating set of classes of $H_2(M, Z_2)$ represented by such surfaces.*

Proof. As in Theorem 1, given α there is a closed nonorientable surface K embedded in M with $[K] = \alpha$. We compress K to K_1 satisfying each component is incompressible or a 2-sphere. If some two-sided component of K_1 is nonseparating in M , then a loop meeting this surface in a single point gives an element of infinite order in $H_1(M, Z)$, contrary to assumption. So all such components are null-homologous and the one-sided components have associated classes in $H_2(M, Z_2)$ which sum to give α . We conclude that among all such classes there is a set of generators for $H_2(M, Z_2)$.

EXAMPLE. $S^2 \times S^1$ has no embedded incompressible nonorientable surfaces, though by Theorem 1 it has one-sided Heegaard splittings.

Proof. Suppose K is incompressible and one-sided* in $S^2 \times S^1$. Assume K is transverse to $S^2 \times \{x\}$. Then by induction on the number of curves of $K \cap S^2 \times \{x\}$, it is easy to achieve $S^2 \times \{x\} \cap K = \emptyset$ by an isotopy of K . So K is contained in a 3-cell which is impossible.

DEFINITION. We call (M, K) an *incompressible one-sided Heegaard splitting* if (M, K) is a one-sided Heegaard splitting and K is incompressible in M .

THEOREM 4. *Suppose M is irreducible and $H_1(M, Z)$ is finite. The following are equivalent:*

(1) *For any orientable incompressible surface J embedded in M , the map $H_1(J, Z_2) \rightarrow H_1(M, Z_2)$ induced by the inclusion is onto.*

(2) For any nonorientable incompressible surface K embedded in M , $M - K$ is an open handlebody.

If these conditions are satisfied, then there is an incompressible one-sided splitting associated with any nonzero class in $H_2(M, Z_2)$.

Proof. $1 \Rightarrow 2$. Suppose K is nonorientable and incompressible in M . If J is orientable, incompressible and disjoint from K in M , then the image of $H_1(J, Z_2) \rightarrow H_1(M, Z_2)$ is in the kernel of the epimorphism $H_1(M, Z_2) \rightarrow Z_2$ which is the element of $H^1(M, Z_2)$ dual to $[K]$. So the map $H_1(J, Z_2) \rightarrow H_1(M, Z_2)$ is not onto, contrary to assumption. We conclude that $J \cap K = \emptyset$ does not occur and therefore by Lemma 2, $M - K$ is an open handlebody.

$2 \Rightarrow 1$. Suppose J is an orientable and incompressible surface in M , for which $H_1(J, Z_2) \rightarrow H_1(M, Z_2)$ is not onto. Then there is an epimorphism $H_1(M, Z_2) \rightarrow Z_2$ with kernel containing the image of $H_1(J, Z_2)$. Let $f: M \rightarrow RP^4$ be a map realizing the epimorphism. Since f restricted to J is null-homotopic, without loss of generality we can assume f is transverse on RP^3 and $f(J) \cap RP^3 = \emptyset$.

Because $H_1(M, Z)$ is finite, all the two-sided components of $f^{-1}(RP^3)$ are separating and there must be some one-sided components. As in Lemma 3, we compress one of these latter surfaces to obtain a one-sided incompressible surface K . Since J is incompressible and M is irreducible, all the compressions can be made to occur in $M - J$. By assumption $M - K$ is an open handlebody containing J and this gives a contradiction.

We now consider the last assertion of the theorem. Since $H_1(M, Z)$ is finite, given $\alpha \neq 0$ in $H_2(M, Z_2)$ Lemma 3 shows there are disjoint nonorientable incompressible surfaces K_i in M with $\sum_{i=1}^n [K_i] = \alpha$. But for each of these, $M - K_i$ is an open handlebody by Condition 2. For a handlebody X , since $H_2(X, Z_2) = 0$ it follows that there are no closed nonorientable surfaces embedded in X . We conclude that $n = 1$ and $[K_1] = \alpha$ as desired.

COROLLARY 5. *Suppose M is irreducible and not sufficiently large. Then there is an incompressible one-sided splitting associated with any nonzero class in $H_2(M, Z_2)$.*

REMARK. The theorem shows that M has many incompressible one-sided splittings if and only if the existence of incompressible two-sided surfaces is restricted.

PROPOSITION 6. *Suppose M has an incompressible one-sided Heegaard splitting. Then M is irreducible.*

Proof. Suppose S is a 2-sphere embedded in M . Assume S is

transverse to K , where (M, K) is an incompressible one-sided splitting. Since $M - K$ is irreducible, it follows easily by induction on the number of curves of $S \cap K$ that S bounds a 3-cell.

THEOREM 7. *Suppose M is an orientable Seifert manifold with fiber giving a nonzero class in $H_1(M, Z_2)$. Assume M is neither a union of two twisted line-bundles over a nonorientable surface nor a surface-bundle over a circle. Then M has an incompressible one-sided Heegaard splitting.*

Proof. For the properties of Seifert manifolds, see [5]. By Proposition 1 of [2], M is irreducible. Let h be a fiber of M . Since $[h] \neq 0$ in $H_1(M, Z_2)$, there is an epimorphism $H_1(M, Z_2) \rightarrow Z_2$ with $[h]$ not in the kernel. As in Theorem 1, we can find a closed non-orientable surface K embedded in M with $[K]$ dual to the corresponding element of $H^1(M, Z_2)$. Consequently K and h have odd intersection number.

Compressing K , we get a component K_1 which has odd intersection number with h and is either a 2-sphere or is incompressible in M . Suppose that the orbit surface for M is nonorientable. Then we lift K_1, h to \tilde{K}_1, \tilde{h} in a double covering \tilde{M} of M , so that \tilde{M} has an orientable orbit surface. Since \tilde{h} covers h once, it follows that \tilde{K}_1, \tilde{h} also have odd intersection number. The homotopy class $\{\tilde{h}\}$ is central in $\pi_1(\tilde{M})$. There are two possibilities.

Case 1. \tilde{K}_1 is two-sided in \tilde{M} .

After compressing \tilde{K}_1 if necessary, we obtain a two-sided incompressible surface J which has odd intersection number with \tilde{h} . Therefore $\{\tilde{h}\}$ is not in the image of $\pi_1(J)$ in $\pi_1(\tilde{M})$. Exactly as in Theorem 2 of [2] it follows that \tilde{M} is a J -bundle over S^1 and M is either a union of two twisted line-bundles over a nonorientable surface, or a surface-bundle over S^1 , contrary to assumption.

Case 2. \tilde{K}_1 is one-sided in \tilde{M} .

Then K_1 is one-sided in M . Suppose that J is an orientable incompressible surface in $M - K_1$. Since h and K_1 have an odd intersection number, we see that $\{h\}$ is not in the image of $\pi_1(J)$ in $\pi_1(M)$. Passing to \tilde{M} , we can apply the same argument as in Case 1. This gives a contradiction and so by Lemma 2, we conclude that $M - K_1$ is a handlebody.

If the orbit surface for M is orientable then a similar method applies.

REMARKS. (1) Exactly the same argument as in (4.6) of [1]

can be employed to show that if M is an orientable Seifert manifold with orientable orbit surface and if K is an embedded nonorientable surface which has odd intersection number with the fiber in M , then $\pi_1(K) \rightarrow \pi_1(M)$ is onto.

(2) To get examples where $[h] \neq 0$ in $H_1(M, Z_2)$, take M such that all the μ -invariants (multiplicities of the exceptional fibers) are odd and the b -invariant differs from the sum of the ν -invariants by an even number.

(3) The 3-manifolds in this theorem are in general sufficiently large. They contain incompressible tori formed by the ordinary fibers projecting to suitable circles in the orbit surface. Note that such surfaces to some extent satisfy the conditions of Theorem 4; the image of $H_1(T, Z_2)$ in $H_1(M, Z_2)$ certainly contains $[h]$, where T is a torus of ordinary fibers.

PROPOSITION 8. *Let M be a Seifert manifold with S^2 as orbit surface and exceptional fibers of multiplicity $(2, 2, n)$ excluding $S^2 \times S^1$, or $(2, 4, n)$, for $n \geq 1$. Then there is an incompressible one-sided splitting (M, K) such that K has genus 2 or 3 respectively.*

Proof. For the $(2, 2, n)$ case, see [4]. In the other case let N_1 and N_2 be small fibered neighborhoods of the exceptional fibers of multiplicity 2 and 4 respectively and let Φ be the projection from M to the orbit surface. Choose a nonsingular arc λ in the orbit surface from $\Phi(\partial N_1)$ to $\Phi(\partial N_2)$ missing the image of the third exceptional fiber and $\Phi(\text{int } N_1 \cup \text{int } N_2)$. Then $\Phi^{-1}(\lambda)$ is an annulus in M with $\partial\Phi^{-1}(\lambda) = \Phi^{-1}(\partial\lambda)$ consisting of a fiber in ∂N_1 and one in ∂N_2 . By [1] or § 3 it follows that the component of $\partial\Phi^{-1}(\lambda)$ in ∂N_1 bounds a Mobius band in N_1 , while the other curve of $\partial\Phi^{-1}(\lambda)$ bounds a Klein bottle with a hole in N_2 . Therefore we obtain a nonorientable surface K of genus 3 embedded in M .

If K is compressible in M then M contains an embedded one-sided RP^2 . By Proposition 1 of [2], $M \approx RP^3$ or $M \approx RP^3 \# RP^3$ follows. The presentation of $\pi_1(M)$ given by the fibering (see [5]) shows that $\pi_1(M) = Z_2$ or $Z_2 * Z_2$ is impossible. So K is incompressible.

Finally we prove that $M - K$ is an open handlebody. Now it is easy to see as in § 3 or [1] that the map $\pi_1(K) \rightarrow \pi_1(N_1 \cup N_2 \cup \Phi^{-1}(\lambda))$ induced by the inclusion is onto. Since the complement of $N_1 \cup N_2 \cup \Phi^{-1}(\lambda)$ in M is an open solid torus, it follows that $\pi_1(K) \rightarrow \pi_1(M)$ is onto and so by Lemma 2 the result is established.

REMARKS. (1) Note that a Seifert space with exceptional fibers of multiplicity $(2, 2, 4)$ and S^2 as orbit surface has incompressible

splittings of genus 2 and genus 3. These are associated with different classes in $H_2(M, Z_2)$ however.

(2) The Seifert manifolds with S^2 as orbit surface and exceptional fibers of multiplicity $(2, 2, n)$ excluding $S^1 \times S^2$, or $(2, 4, 3)$, all have finite fundamental groups.

3. One-sided surfaces in lens spaces. Let M be the lens space $L(2k, q)$, where $(2k, q) = 1$. Since $H_2(M, Z_2) = Z_2$, M contains embedded nonorientable surfaces. In [1] an algorithm is given for the minimal genus, denoted $N(2k, q)$, of such surfaces. Note that an embedded nonorientable surface of minimal genus is incompressible, since $H_2(M, Z) = 0$.

Let K denote an incompressible surface in M throughout this section. Since M is not sufficiently large, K must be nonorientable and $M - K$ is an open handlebody by Corollary 5. Let (M, L) be a Heegaard splitting of genus 1, with $M = T \cup T'$ where $T \cup T' = \partial T = \partial T' = L$ and T, T' are solid tori.

LEMMA 9. *After an isotopy, $K \cap T$ can be transformed into a disk.*

Proof. (cf. [1, p. 99]). By shrinking T , it can be assumed that $K \cap T$ is a collection of n meridian disks for T . Let D' be a meridian disk for T' . By an isotopy of K , we can suppose that D' is transverse to K and $K \cap D'$ contains no simple closed curves, since K is incompressible. Also without loss of generality D' can be chosen so that no arc in $\partial D'$ together with an arc in $K \cap L$ bounds a disk in L .

Let λ be a component of $K \cap D'$. Then λ is an arc with ends on curves of $K \cap L$. If μ is an arc of $\partial D'$ with $\partial \lambda = \partial \mu$ then we choose λ so that $(\text{int } \mu) \cap K = \emptyset$.

Case 1. Both ends of λ are on a curve C of $K \cap L$.

Then either $n = 1$ or there is an arc γ of C with $\partial \mu = \partial \gamma$ and $\mu \cup \gamma$ bounds a disk on L . The latter contradicts our choice of D' .

Case 2. λ joins curves C and C' of $K \cap L$.

There is an isotopy of K with support in a neighborhood of D' , which carries λ to μ along D' . After this isotopy K meets T in $n - 2$ meridian disks and in a disk parallel to a disk on L . So a further isotopy gives $n - 2$ meridian disks in $K \cap T$ and the result follows by induction on n . Note $K \cap T = \emptyset$ implies K is contained in T' , which is impossible.

LEMMA 10. *By an isotopy, $K \cap T$ can be changed into a Mobius*

band.

Proof. By Lemma 9 we can assume $K \cap T$ is a single meridian disk. Denote $K \cap L$ by C and let D' be a meridian disk for T' . We can again suppose that K is transverse to D' , there are no closed curves in $K \cap D'$ and no arc in $\partial D'$ together with an arc in C bounds a disk in L .

Let λ be an arc of $K \cap D'$, chosen so that an arc μ of $\partial D'$ with $\partial\mu = \partial\lambda$ satisfies $(\text{int } \mu) \cap K = \emptyset$. If μ has both ends on the same side of C , then we get a contradiction to our choice of D' . Hence μ goes from one side of C to the other. The isotopy of K taking λ to μ along D' transforms $K \cap T$ into a Mobius band as required.


PROPOSITION 11. *There are disjoint tori L_j , $1 \leq j \leq g - 1$, in M ($g = \text{genus } K$) which are all parallel to each other and give Heegaard splittings. The tori satisfy $L_j \cap K = C_j$ is a single curve and $K - \bigcup_j C_j$ consists of 2 open Mobius bands and $g - 2$ open Mobius bands with single holes.*

Proof. By Lemma 10 there is a torus L_1 with $L_1 \cap K = C_1$ and $T_1 \cap K$ equal to a Mobius band, where $L_1 = \partial T_1 = \partial T'_1$ and T_1, T'_1 are solid tori. We can assume K is transverse to D'_1 , a meridian disk for T'_1 , $K \cap D'_1$ contains no loops and no arc in $\partial D'_1$ together with an arc in C_1 bounds a disk in L_1 . Then arcs μ and λ can be found with properties as in Lemma 10. Let L_2 be the image of L_1 by the isotopy taking μ to λ along D'_1 and pushing L_1 a small distance into $\text{int } T'_1$. Then L_2 has the required properties and all the tori L_j are constructed similarly. Note that after g steps the meridian disk D'_g can be chosen to miss K and the process stops.

THEOREM 12. *Any two incompressible surfaces embedded in M are isotopic.*

Proof. Suppose K_1, K_2 are both incompressible surfaces in M and let (M, L) be a Heegaard splitting of genus 1 with $L = \partial T = \partial T'$. By Lemma 9 we can assume $K_1 \cap T$ and $K_2 \cap T$ are disjoint meridian disks. Let C_i denote the curve $K_i \cap L$, $i = 1, 2$.

Let D' be a meridian disk for T' and apply the argument for Lemma 10. Then arcs μ_i, λ_i with the properties in that proof for K_i can be found, $i = 1, 2$. Without loss of generality we can suppose that no arc of $\partial D'$ with interior disjoint from C_i has both ends on the same side of C_i , for $i = 1$ or 2 . It can be assumed also that $\mu_1 \cap C_2$ and $\mu_2 \cap C_1$ are both single points. There are three possibilities as shown in Figure 1.



In (A) μ_1 and μ_2 “spiral” oppositely, while in (B) and (C) they “spiral” in the same direction. Note that in (C), $\mu_1 \cap \mu_2 \neq \emptyset$. It is easy to see that in cases (B) and (C), after the isotopies of K_i taking λ_i to μ_i along D' , the new curves $K_i \cap L$ are isotopic on L . Consequently by a further isotopy it can be achieved that $K_1 \cap T$ and $K_2 \cap T$ are equal to the same Mobius band.

Suppose on the other hand that (A) is true. The four components of $L - C_1 - C_2 - \mu_1 - \mu_2$ are open disks. Let B_1 and B_2 be the closures of two of these, chosen as shown in (A) of Figure 1. Assume that $\partial D'$ is oriented so that it enters B_1 at the end-point of μ_1 . Then $\partial D'$ must leave B_1 at a point of $C_2 \cap B_2$, since otherwise there is an arc of $\partial D'$ with both ends on the same side of C_1 , contrary to assumption. So $\partial D'$ enters B_2 and similarly leaves it at a point of $C_1 \cap B_1$. This implies that $\partial D'$ remains in $B_1 \cup B_2$, which is clearly false. We conclude that (A) does not occur.

To continue the procedure, it is only necessary to slightly disjoin K_1 and K_2 near L by an isotopy. After g steps it follows that K_1 and K_2 are isotopic.

DEFINITION. One-sided Heegaard splittings (M, K_1) and (M, K_2) are strongly equivalent if K_1 and K_2 are isotopic.

REMARKS. (1) The theorem implies that any incompressible surface in a lens space has minimal genus.

(2) An equivalent statement to the theorem is that any two incompressible one-sided splittings of a lens space are strongly equivalent. We conjecture that this is true for arbitrary one-sided splittings for which the surfaces have the same genus in a lens space [cf. [10)].

(3) Let M be a lens space and (M, K) be an incompressible one-sided splitting. Let $\mathcal{H}(M)$ denote the quotient of the homeomorphism group of M by the normal subgroup of homeomorphisms isotopic to the identity. The theorem implies that any class in

$\mathcal{H}(M)$ is represented by a homeomorphism which preserves the splitting (M, K) . This can be useful for studying $\mathcal{H}(M)$ (cf. [4] for the case $M = L(4n, \pm(2n - 1))$).

THEOREM 13 ((8.3) of [1]). $N(2k, q) = 1 + N(2(k - Q), q - 2m)$, where m, Q satisfy $2km - qQ = \pm 1$, $0 < Q < k$.

Proof. Let K be an incompressible surface in M . By Lemma 10 we can assume $K \cap T$ is a Mobius band denoted E , where as usual $M = T \cup T'$, $T \cap T' = L$ and T, T' are solid tori. Let T_1 be a solid torus with meridian disk D_1 . We form a new lens space M_1 as the union of T_1 and T' identified along their boundaries by a homeomorphism which takes ∂D_1 to ∂E . Then M_1 contains a non-orientable surface $K_1 = (K - \text{int } E) \cup D_1$, which has genus 1 less than K .

Suppose K_1 is compressible in M_1 . Then there is a nonorientable surface K_2 embedded in M_1 with smaller genus than K_1 . By Lemma 9, after an isotopy $K_2 \cap T_1$ can be transformed into a meridian disk. But then $K_2 - \text{int } T_1$ has a boundary curve which bounds a Mobius band in T . Consequently there is a nonorientable surface in M with genus 1 greater than K_2 , which contradicts our assumption that K has minimal genus. Therefore if $M_1 = L(2k_1, q_1)$ we conclude that $N(2k, q) = 1 + N(2k_1, q_1)$.

In $H_1(L, Z)$ let a, a' be elements satisfying $a \sim 0$ in T and $a' \sim 0$ in T' . Also let classes b, b' be chosen so that $\{a, b\}, \{a', b'\}$ are bases for $H_1(L, Z)$. Without loss of generality we can suppose that $a = qa' + 2kb'$ and $b = ma' + Qb'$, with m, Q chosen as above, since $M = L(2k, q)$.

Now a, a' have intersection number $\pm 2k$. In the argument for Lemma 10, clearly ∂E has two less intersection points with the curve $\partial D'$ than ∂D does, where D, D' are the meridian disks for T, T' respectively. So $[\partial E]$ and a' have intersection number with absolute value strictly less than $2k$. Also by a suitable orientation of ∂E it can be assumed that $[\partial E] = -2b + Na$ where N is odd, since ∂E and ∂D have intersection number ± 2 . Hence $[\partial E] = (Nq - 2m)a' + (2kN - 2Q)b'$ and $[\partial E], a'$ have intersection number $\pm(2kN - 2Q)$. We conclude that $N = 1$. Since ∂E is a meridian curve for T_1 , it follows that $M_1 = L(2k - 2Q, q - 2m)$ as desired.

4. Stable equivalence. Let (M, K) be a one-sided Heegaard splitting and (M', K') a one- or two-sided Heegaard splitting. Let B, B' be 3-cells in M, M' such that $B \cap K = D$ and $B' \cap K' = D'$ are disks. Suppose $M - \text{int } B$ is attached to $M' - \text{int } B'$ by a homeomorphism of the boundary 2-spheres, which takes ∂D to $\partial D'$. The

resulting 3-manifold is the connected sum $M \# M'$ and contains the closed nonorientable surface $K \# K'$. Clearly $(M \# M', K \# K')$ is a one-sided Heegaard splitting which we will denote by $(M, K) \# (M', K')$ and call the *connected sum* of the splittings (M, K) , (M', K') .

DEFINITION. One-sided Heegaard splittings (M, K) , (M', K') are *equivalent* if there is a homeomorphism from M to M' , which maps K to K' .

Note that $(M, K) \# (M', K')$ is independent of the choice of the 3-cells B, B' up to equivalence. Let (S^3, L) denote the standard Heegaard splitting of genus 1 of S^3 and let $(M, K) \# n(S^3, L)$ be the connected sum with n copies of (S^3, L) .

DEFINITION. One-sided Heegaard splittings (M, K) , (M', K') are *stably equivalent* if $(M, K) \# n(S^3, L)$ is equivalent to $(M', K') \# m(S^3, L)$ for some m, n .

THEOREM 14. *Suppose (M, K) , (M, K') are one-sided Heegaard splittings with $[K] = [K']$. Then they are stably equivalent.*

Proof. There is a double cover \tilde{M} of M associated with both splittings, as in § 0. Let \tilde{K}, \tilde{K}' denote the lifts of K, K' to \tilde{M} . Since $M - K'$ is an open handlebody, there is a wedge of circles, denoted A' , which is embedded in $M - K'$ as a deformation retract. Assume that A' is transverse to K and let A'_1, A'_2 be the lifts of A' to \tilde{M} .

We do the procedure of Theorem 1, using a triangulation of M which contains A' as a subcomplex. Let $\Gamma_i, N(\Gamma_i), Y_i, V, W, \bar{N}(V)$ be defined as in Theorem 1, with respect to K . It can be supposed that the partition $V \cup W$ of the vertices of \tilde{K} is chosen so that V contains $A'_2 \cap \tilde{K}$. Instead of moving Γ_1 just in $\bar{N}(V)$, we need to use an isotopy with support in $\bar{N}(V) \cup \bar{N}(A'_2 \cap Y_1)$, where $\bar{N}(A'_2 \cap Y_1)$ is a small regular neighborhood of the second derived simplicial neighborhood of $A'_2 \cap Y_1$ in Y_1 . The isotopy is chosen to shift Γ_1 to Γ_1^* so that $N(\Gamma_1^*) \cap gN(\Gamma_1^*) = \emptyset$ and $N(\Gamma_1^*) \cap A'_2 = \emptyset$.

Defining \tilde{K}^* as in Theorem 1, we see that $\tilde{K}^* \cap (A'_1 \cup A'_2) = \emptyset$ has been achieved. Also by [6] it follows that (M, K^*) and (M, K) are stably equivalent, where $K^* = p(\tilde{K}^*)$. So there is no loss of generality in replacing (M, K) by (M, K^*) , which we now do. Then $K \cap A' = \emptyset$ and hence K is contained in a closed regular neighborhood $N(K')$ of K' .

LEMMA 15. *Suppose K is closed nonorientable surface embedded in N which is a twisted line-bundle over a closed nonorientable surface K' . Then $\text{genus } K \geq \text{genus } K'$. Equality holds if and only if K is incompressible in N . In this case K and K' are isotopic.*

Proof. We compress K so that every component is either incompressible or a 2-sphere. Since every two-sided surface in N separates, some component K_1 is nonorientable. It suffices to prove that K_1 is isotopic to K' .

Suppose without loss of generality that K_1 is transverse to K' and that $K_1 \cap K'$ contains only loops which are noncontractible in both K_1 and K' (as K_1, K' are incompressible). Let $N(K')$ be a small closed regular neighborhood with $K_1 \cap N(K')$ consisting of Mobius bands and annuli. Since $H_2(N, \mathbb{Z}_2) = \mathbb{Z}_2$, it is easy to see that $K' - K_1$ and $K_1 - K'$ are orientable.

$N - \text{int } N(K')$ is homeomorphic to $\partial N \times I$ and will be denoted by X . Every component R_1 of $K_1 \cap X$ is orientable and incompressible in X . By [8] it follows that there is a region R' in $\partial N(K')$ with $\partial R_1 = \partial R'$ and R_1 parallel to R' . Clearly then R_1 can be chosen so that $(\text{int } R') \cap K_1 = \emptyset$. There are unique components \bar{R}_1, \bar{R}' , of $K_1 - K'$ and $K' - K_1$ containing R_1, R' respectively, and there is an isotopy of K_1 taking \bar{R}_1 to \bar{R}' .

If $K_1 \cap K'$ consists entirely of orientation-reversing curves then the closure of \bar{R}_1 or \bar{R}' is K_1 or K' . So K_1 and K' are isotopic. On the other hand if $K_1 \cap K'$ contains some orientation-preserving curves then there must be at least one in $\partial \bar{R}_1 = \partial \bar{R}'$. In this case, after the isotopy moving \bar{R}_1 to \bar{R}' we can slightly disjoint K_1 and K' so that the surfaces are again transverse and have fewer curves of intersection. By induction on the number of these curves, the lemma is proved.

Returning to the proof of the theorem, we have K included in $N(K')$. So by the lemma there is a unique one-sided component K_1 obtained from compressing K in $N(K')$, and K_1 is isotopic to K' . Any two-sided incompressible surface in $N(K')$ must be parallel to $\partial N(K')$ by [8] and so is completely compressible in M . We conclude that K' can be assumed to be the result of a sequence of compressions of K .

Consequently without loss of generality there is a collection of disks D_1, \dots, D_r embedded in M with the following properties

— The curves ∂D_i are all contained in K and form a nonseparating set of loops in K (i.e., $K - \cup \partial D_i$ is connected).

— $D_i \cap D_j$ is a union of disks D_k in the collection, for all i, j .

— D_i intersects K transversely for all i , and each component of $K \cap D_i$ is one of the curves ∂D_k .

— K' is obtained by successively compressing K using the disks D_i in order of size, starting with the smallest.

Next the *complexity* n_i of a disk D_i is defined to be the number of disks D_j which are contained in D_i . Our aim is to decrease all the complexities to zero. Suppose there is a complexity which is

nonzero. $N(K)$ can be chosen small enough so that every component of $D_i \cap N(K)$ is a small annulus containing exactly one curve of $D_i \cap K$, for all i .

At least one of the components of $D_i \cap Y$ is not a disk, for some i . Let us consider the intersections of the meridian disks for Y with the disks D_i . Note that in a handlebody, a loop which is disjoint from a complete set of meridian disks must be contractible. Therefore it easily follows that for some i , either one of the components of $D_i \cap Y$ is compressible in Y or there is a disk D embedded in Y with $\partial D = \lambda \cup \gamma$, where λ, γ are arcs in $\partial Y, D_i$ respectively and both join two different curves of $D_i \cap \partial Y$. In the former case it is easy to see that we can alter the choice of the compressing disks D_i so as to decrease some of the numbers n_i .

It now suffices to suppose that we have a disk D as above. Let μ_1 and μ_2 be arcs in $D_i \cap N(K)$ which join the end-points of γ to K , with $(\text{int } \mu_1) \cap K = (\text{int } \mu_2) \cap K = \emptyset$. Clearly $\mu_1 \cap K$ and $\mu_2 \cap K$ lie in two different components of $D_i \cap K$. An embedded disk D' in $N(K)$ can be found with $\partial D' = \mu_1 \cup \lambda \cup \mu_2 \cup \lambda'$, where λ' is an arc in K and $K \cap \text{int } D' = \emptyset$. To see this we identify $N(K)$ with the mapping cylinder of the double covering of K . Then the mapping cylinder restricted to λ intersects K in a singular arc, in general. However the singularities may be pushed off the ends of this arc to give D' .

Let $D_0 = D \cup D'$ and let γ' denote the arc $\mu_1 \cup \gamma \cup \mu_2$. Then $\partial D_0 = \gamma' \cup \lambda'$. We perform an elementary surgery on K along the path γ' . Consequently the effect of this is to add a trivial handle to K . Also γ' is contained in D_i and without loss of generality, D_i can be chosen so that γ' joins ∂D_i to some curve ∂D_j for $j \neq i$. We now observe that a set of compressing disks for K with the trivial handle along γ' added, can be obtained by replacing D_i by two disks D'_i, D''_i which have complexities both strictly less than n_i . For the choices of D'_i, D''_i , see Figures 2 and 3.





FIGURE 2



Note that $D_2 \supset D_3$ still and $D'_2 \supset D_3$.

FIGURE 3

By induction, this completes the proof that all the complexities can be reduced to zero. In this case it can be assumed that K is obtained from K' by elementary surgeries along a collection Γ of arcs, each of which intersects K' precisely at its endpoints.

Now M can be triangulated so that K is a full subcomplex and the 1-skeleton contains Γ . Applying the procedure of Theorem 1, we add a suitably shifted boundary of a small neighborhood of the 1-skeleton of $M - K$ to K , and similarly for K' using the 1-skeleton of $M - K - \Gamma$. The same new splitting (M, K'') is obtained in both cases, and (M, K'') is stably equivalent to (M, K) and (M, K') by [6]. Hence the theorem is proved.

5. Mappings of degree 1.

LEMMA 17. *Let $f: M' \rightarrow M$ be an odd degree map between closed orientable 3-manifolds. If (M, K) is a one-sided Heegaard splitting then there is a one-sided Heegaard splitting (M', K') and a map f' homotopic to f such that $f'^{-1}(K) = K'$.*

Proof. Using elementary surgery, we can obtain $f^{-1}(K)$ is connected by a homotopy of f , since $\pi_1(M, K) = \{1\}$. Suppose $f^{-1}(K)$ is orientable. Then f factors through the double cover of M associated with the splitting (M, K) . This contradicts the assumption that f is of odd degree. Therefore $f^{-1}(K)$ is nonorientable.

We now do the procedure of Theorem 1 using M' and the surface $f^{-1}(K)$. As $\pi_1(M, K) = \{1\}$, all the 1-handles can be added to $f^{-1}(K)$ by a homotopy of f . If $f': M' \rightarrow M$ is the new map then $f'^{-1}(K) = K'$ gives a one-sided Heegaard splitting of M' as required.

THEOREM 18. *With the hypotheses of Lemma 17, suppose that f is of degree 1. Then f' and K' can be chosen so that*

$$f': M' - \text{int } N(K') \longrightarrow M - \text{int } N(K)$$

is standard (in the sense of [9]), where $f'^{-1}(N(K)) = N(K')$.

Proof. The map f' found in Lemma 17 lifts to a degree 1 map $\tilde{f}': \tilde{M}' \rightarrow \tilde{M}$, where \tilde{M}' , \tilde{M} are the double covers associated with the splittings (M', K') , (M, K) . If \tilde{K} , \tilde{K}' are the lifts of K , K' to M , M' respectively then $\tilde{f}'^{-1}(\tilde{K}) = \tilde{K}'$ and (\tilde{M}, \tilde{K}) , (\tilde{M}', \tilde{K}') are Heegaard splittings.

Let Y'_1 , Y'_2 be the closures of the components of $\tilde{M}' - \tilde{K}'$. In Theorem 2.1 of [9] it is proved that \tilde{f}' can be made standard on the handlebodies Y'_1 , Y'_2 by the operation of drilling holes. It is only necessary to do this in a manner which is invariant under the covering transformation to obtain the result.

REFERENCES

1. G. Bredon and J. Wood, *Nonorientable surfaces in orientable 3-manifolds*, Invent. Math., **7** (1969), 83-110.
2. W. Heil, *Almost sufficiently large Seifert fiber spaces*, Michigan Math. J., **20** (1973), 217-223.
3. J. Hempel, *3-Manifolds*, Annals of Math. Studies, No. 86, Princeton, 1976.
4. J. H. Rubinstein, *On 3-manifolds that have finite fundamental group and contain Klein bottles*, to appear in Trans. Amer. Math. Soc.
5. H. Seifert, *Topologie dreidimensionaler gefaserner Räume*, Acta Math., **60** (1933), 147-238.
6. J. Singer, *Three-dimensional manifolds and their Heegaard diagrams*, Trans. Amer. Math. Soc., **35** (1933), 88-111.
7. J. Stallings, *On the loop theorem*, Annals of Math., **72** (1960), 12-19.
8. F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, Annals of Math., **87** (1968), 56-88.
9. ———, *On Mappings of Handlebodies and of Heegaard Splittings*, Topology of Manifolds, Markham, Chicago 1970, 205-211.
10. ———, *Heegaard-Zerlegungen der 3-sphäre*, Topology **7** (1968), 195-203.

Received January 15, 1977 and in revised form June 23, 1977. The author was supported by a Rothman's Fellowship during this research.

UNIVERSITY OF MELBOURNE
PARKVILLE, VICTORIA, 3052, AUSTRALIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

C. W. CURTIS

University of Oregon
Eugene, OR 97403

C. C. MOORE

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Ata Nuri Al-Hussaini, <i>Potential operators and equimeasurability</i>	1
Tim Anderson and Erwin Kleinfeld, <i>Semisimple nil algebras of type δ</i>	9
Stephen LaVern Campbell, <i>Linear operators for which T^*T and $T + T^*$ commute. III</i>	17
Robert Jay Daverman, <i>Special approximations to embeddings of codimension one spheres</i>	21
Donald M. Davis, <i>Connective coverings of BO and immersions of projective spaces</i>	33
V. L. (Vagn Lundsgaard) Hansen, <i>The homotopy type of the space of maps of a homology 3-sphere into the 2-sphere</i>	43
James Victor Herod, <i>A product integral representation for the generalized inverse of closed operators</i>	51
A. A. Iskander, <i>Definability in the lattice of ring varieties</i>	61
Russell Allan Johnson, <i>Existence of a strong lifting commuting with a compact group of transformations</i>	69
Heikki J. K. Junnila, <i>Neighborhoods</i>	83
Klaus Kalb, <i>On the expansion in joint generalized eigenvectors</i>	109
F. J. Martinelli, <i>Construction of generalized normal numbers</i>	117
Edward O'Neill, <i>On Massey products</i>	123
Vern Ival Paulsen, <i>Continuous canonical forms for matrices under unitary equivalence</i>	129
Justin Peters and Terje Sund, <i>Automorphisms of locally compact groups</i>	143
Duane Randall, <i>Tangent frame fields on spin manifolds</i>	157
Jeffrey Brian Remmel, <i>Realizing partial orderings by classes of co-simple sets</i>	169
J. Hyam Rubinstein, <i>One-sided Heegaard splittings of 3-manifolds</i>	185
Donald Charles Rung, <i>Meier type theorems for general boundary approach and σ-porous exceptional sets</i>	201
Ryōtarō Satō, <i>Positive operators and the ergodic theorem</i>	215
Ira H. Shavel, <i>A class of algebraic surfaces of general type constructed from quaternion algebras</i>	221
Patrick F. Smith, <i>Decomposing modules into projectives and injectives</i>	247
Sergio Eduardo Zarantonello, <i>The sheaf of outer functions in the polydisc</i>	267