Pacific Journal of Mathematics

A NOTE ON EXPONENTIALS OF DISTRIBUTIONS

ALAN SLOAN

Vol. 79, No. 1 May 1978

A NOTE ON EXPONENTIALS OF DISTRIBUTIONS

ALAN SLOAN

Nonstandard analysis is used to discuss nonlinear functions of distributions. An application is given to obtain a generalized Trotter product formula. The strong resolvent topology is discussed from a nonstandard point of view.

Enlarging the real number system to include infinite and infinitesimal quantities enabled Laugwitz [5] to view the delta function distribution as a point function. Independently Robinson [7] demonstrated that distributions could be viewed as generalized polynomials. Luxemburg [6] presented an alternate picture of distributions as generalized functions within the context of Robinson's theory of nonstandard analysis and it is a special case of this point of view that we take here. Once one accepts distributions as generalized functions, the composition map provides a natural method of defining nonlinear functions of distributions. Unfortunately, the difficulties in the standard attempt to define a function, f, of a distribution, τ , (by first writing τ as a limit, in some appropriate sense, of smooth standard functions τ_n , next composing these approximations with f, and finally taking the limit of $f \circ \tau_n$) still remain in the nonstandard theory. In particular, this procedure may not lead to a distribution and even when it does the distribution obtained may depend on the representation of the original distribution as a generalized function. Thus, at present, there is no comprehensive theory of nonlinear functions of distributions.

The development of such a theory may well proceed along alternate tracks depending on the applications intended. One use for distributions occurs in the study of perturbations of selfadjoint operators in Hilbert space. In some perturbation problems the pathology of obtaining compositions which do not represent distributions may be avoided by considering only bounded functions of distributions. For example, in the Trotter product formula one works with bounded exponentials. Even though, in this case, the exponentials of distributions may be identified with standard distributions it is the conclusion of this paper that such an identification should not be made. It is the author's view that functions of distributions should be regarded as generalized functions but not distributions. The utility of this view is illustrated by a non-standard version of the Trotter product formula.

For an introduction to nonstandard analysis and its relation to distributions, see [10]. In [11] a square root for the delta function

was defined nonstandardly.

Let *X be a δ -incomplete ultrapower of a structure X containing the real numbers, R. We shall use the convention that $A \subset {}^*A$, for all sets A considered. If T is a topological space with u in T and v in *T then we shall write $u \approx v$ if and only if v is in $\bigcap \{{}^*\mathcal{U} : \mathcal{U} \text{ is an open set containing } u\}$. In this case we say u is the standard part of v, relative to the given topology and that v is near standard. We write u = st(v). We always assume that R^p has the Euclidean topology and all Hilbert spaces have the norm topology.

If f and g are Lebesgue measurable on *S for some standard set S which is measurable, we shall write $\langle f,g\rangle$ for $\int_{*S}fgdx$, providing the integral exists. If, in fact, f and g are standard functions then $\langle f,g\rangle=\int_Sfgdx$, by the transfer principle. Let W be an open subset of R^p . Let B_n be an increasing

Let W be an open subset of R^p . Let B_n be an increasing sequence of relatively compact subsets of W whose union is W and whose closures are in W. If $\mathscr{D} = \bigcup_{n=1}^{\infty} C_c^{\circ}(B_n)$ is given the inductive limit topology then the elements of \mathscr{D}' , the dual of \mathscr{D} , are the distributions on W. In order to define a function of a distribution we first regularize the distribution.

Let ρ be a $C_c^{\infty}(\mathbf{R}^p)$ function satisfying $0 \le \rho \le 1$, $\rho(x) = 1$ if $||x|| \le 1$ and $\rho(x) = 0$ if $||x|| \ge 2$. Set

$$\gamma_{n}(x) =
ho(n_{\scriptscriptstyle 0}x) \Bigl(\int
ho(n_{\scriptscriptstyle 0}t) dt \Bigr)^{-1}$$

where n_0 is chosen so that

distance
$$(\bar{B}_n, \overline{B_{n+1}^c}) > 2/n_0$$
.

Choose β_n to be a $C_c^{\infty}(\mathbb{R}^p)$ function satisfying $\beta_n=1$ on B_{n-1} , $0 \le \beta_n \le 1$ and support $(\beta_n) \subset B_n$.

If x is in B_n and $\gamma_n(x-y)\neq 0$ then $||x-y||\leq 2/n_0$ so that y is in B_{n+1} . If $\gamma_n(x-\cdot)$ denotes the function $y\to\gamma_n(x-y)$ then $\gamma_n(x-\cdot)$ is in $C_c^\infty(B_{n+1})$ if $x\in B_n$. For any distribution T on W define a function T_n by

$$T_n(x) = \beta_n(x) T(\gamma_n(x-\cdot))$$

for n in N. Then T_n is in $C_c^{\infty}(\mathbb{R}^p)$ and support $(T_n) \subset B_n$. By the transfer principle T_n is in ${}^*C_c^{\infty}(\mathbb{R}^p)$ and support $(T_n) \subset {}^*B_n$ for n in *N . Further, since $T_n \to T$ in \mathscr{D}' as $n \to \infty$ in N it follows that

$$(1) T_n(f) \approx T(f)$$

for all n in N-N and f in $\mathcal D$ where $T_n(f)=\langle T_n,f\rangle$.

The T_n are called the regularizations of T.

DEFINITION 1. Fix n infinite in *N. Let g be a complex valued function of a complex variable defined on the range of T_n . Then we define g(T) to be $g \circ T_n$.

Thus, g(T) is a generalized function in the sense that it is an internal *C-valued function defined on *W. Since $T_n(x) = 0$ if x is not in B_n we find that g(T)(x) = g(0) if x is not in *B_n.

Certain generalized functions $h: {}^*W \to {}^*C$ define distributions τ according to the rule

$$\tau(f) = st(\langle h, f \rangle)$$

for f in \mathscr{D} . Here, st (λ) is the standard part of λ , if λ is a finite hypercomplex number, and undefined otherwise. Thus not every function of a distribution will determine a distribution according to (2). If g is a bounded measurable function then g(T) always defines a distribution by (2).

If T is given by a locally L^1 function F on W, (i.e., if $T(f) = \langle F, f \rangle$ for all f in \mathscr{D}) then g of T is naturally defined within standard distribution theory as the distribution $f \to \langle g \circ F, f \rangle$ providing the composition is defined. A simple example suffices to show that the distribution determined by g(T) need not be $g \circ F$.

EXAMPLE 2. Let E be the complement of the Cantor set, in [0,1], of measure 1/2 obtained by repeated deletions of 2^{n-1} open intervals of length 2^{-2n} , $n=1,2,\cdots$ from [0,1]. Let F be the characteristic function of E and define $T(f)=\langle F,f\rangle$. Let W=(-2,2) in \mathbb{R}^1 . Choosing B_1 so that $[0,1]\subset B_1$ gives

$$egin{aligned} T_{\mathbf{n}}(x) &= T(\gamma_{\mathbf{n}}(x-m{\cdot})) = \langle F, \gamma_{\mathbf{n}}(x-m{\cdot})
angle \ &= \int & F(y) \gamma_{\mathbf{n}}(x-y) dy > 0 \quad ext{for all n in N} \end{aligned}$$

and all x in [0,1]. By the transfer principle $T_n(x) > 0$ for all n in *N and x in ${}^*[0,1]$. Let g(x) = 1 if $x \neq 0$ and g(0) = 0. Then g(T)(x) = 1 if x is in ${}^*[0,1]$ while $g \circ F(x) = 1$ if x is in E and $g \circ F(x) = 0$ if x is not in E. Choosing f in $C_c^{\infty}(-2,2)$ with f = 1 on [0,1] shows $g(T)(f) = \langle g(T), f \rangle = 1$ while

$$egin{aligned} \langle g\circ F,\,f
angle &=\int_{\mathbb{R}}g(F(x))f(x)dx \ &=\int_{\mathbb{R}}g(F(x))f(x)dx \ &=\int_{\mathbb{R}}f(x)dx= ext{measure}\,(E)=1/2 \;. \end{aligned}$$

Thus

$$g \circ F(f) = \langle g \circ F, f \rangle \neq \langle g(T), f \rangle = g(T)(f)$$
.

The necessity of considering only bounded "smooth" g's is now demonstrated, if one is to obtain a generalization of composition of functions. In Lemma 5 we obtain such a generalization.

LEMMA 3. Let f be in $^*L^1(K, dx)$ where K is a subset of \mathbb{R}^p . Let $||f||_1 \approx 0$. Then $M = \{x \ in \ ^*K: f(x) \neq 0\}$ is contained in an internal measurable subset P of *K and measure $(P) \approx 0$.

Proof. Let $A_m=\{x \text{ in } ^*K; |f(x)|>1/m\}$. Then $M=igcup_{m \text{ in } ^{V}}A_m$. Since

$$||f||_{\scriptscriptstyle 1} > \int_{A_m} |f| \, dx > ext{meas} \, (A_m)/m$$
 ,

we find that meas $(A_m) \approx 0$ for each m in N. By definition $A_1 \subset A_2 \subset A_3 \subset \cdots$.

Extend $\{A_m\}$ to an internal sequence. Let $I=\{m \text{ in } *N \text{: } 2 \leq k \leq m \text{ implies meas } (A_k) < 1/k \text{ and } A_{k-1} \subset A_k\}$. I is internal and contains N. Since N is external there is an ω in $(*N-N) \cap I$. Choose $P=A_\omega$. Then meas $(P) \approx 0$ and $\bigcup_{m \text{ in } N} A_m \subset P$.

LEMMA 4. Let F be a locally L^1 function on W. Let T be the distribution $T(f) = \langle F, f \rangle$. For every compact K in W and every infinite n in *N

$$||T_n - F||_1 \approx 0$$

where $|| \cdot ||_1$ is the L^1 norm on *K.

Proof. Choose an open set Q with compact closure such that

$$K \subset Q \subset ar{Q} \subset W$$
 .

For n in N sufficiently large, $\beta_n=1$ on \overline{Q} , and x in K implies support $(\gamma_n(x-\cdot))\subset \overline{Q}$. Consequently, for n in N sufficiently large $T_n(x)=(\gamma_n*F)(x)$. But $\gamma_n*F\to F$ in $L^1(K)$ so for n infinite $||F-\gamma_n*F||_1\approx 0$ while by definition $T_n=\gamma_n*F$ so $||F-T_n||_1\approx 0$.

LEMMA 5. Let g be a bounded uniformly continuous function defined on a closed interval containing 0 and the range of a locally L^1 function F. Let T be the distribution defined by F: $T(f) = \langle F, f \rangle$ for f in \mathscr{D} . Then $g \circ F$ is the distribution determined by g(T).

Proof. We first must show g(T) is defined. There is a closed interval on which g is defined and which contains the range of F. Since $0 \le \beta_n \le 1$ and $\int \gamma_n dx = 1$ it follows that the range of $\beta_n(\gamma_n^*F) = T_n$ is in the closed interval and so the range of T_n is in the domain of g for all n in N. By transfer $g(T_n)$ is defined for all n in N.

Fix n infinite and define $g(T) = g(T_n)$.

We are to prove that for every f in \mathscr{D}

$$\langle g(T), f \rangle \approx \langle g \circ F, f \rangle$$
.

Let K be a compact set containing the support of f. Let A be an internal subset of *K containing $\{x \text{ in } {}^*K \colon T_n(x) \not\approx F(x)\}$ and having measure $\lambda \approx 0$. Since g is uniformly continuous, x in ${}^*K - A$ implies $g(T_n(x)) \approx g(F(x))$. Let M in N be a bound for g. Let ε be a positive real number. Let m be the Lebesgue measure of K. Then

$$\begin{split} |\langle g(T),F\rangle - \langle g\circ F,f\rangle| \\ & \leqq \int_{A} |g(T_{n}) - g(F)| \; |f| \, dx + \int_{*_{K-A}} |g(T_{n}) - g(F)| \, |f| \, dx \\ & \leqq 2M \, ||f||_{\scriptscriptstyle \infty} \lambda + \varepsilon \, ||f||_{\scriptscriptstyle \infty} m < 2\varepsilon \, ||f||_{\scriptscriptstyle \infty} m \; . \end{split}$$

Since ε is arbitrary, this completes the proof.

EXAMPLE 6. Let δ be the delta function $\delta(f)=f(0)$, f in $\mathscr{D}(\mathbf{R}^1)$. Choose $n_0=n$ and $B_n=(-n,n)$. Then, $\delta_n(x)=\gamma_n(x)$. For all n in N, support $(\delta_n)\subset (-2/n,2/n)$ and $0\leq \delta_n<\infty$. Thus for t>0 in R, $0< e^{-t\delta_n}\leq 1$ and if |x|>2/n then $e^{-t\delta_n}(x)=1$. By the transfer principle choosing n in N-N and setting $g(T)=g(T_n)$ we find 0< g(T)< 1 and g(T)(x)=1 if |x|>2/n. Thus

$$\langle g(T), f \rangle \approx \langle 1, f \rangle$$
, f in \mathscr{D} so

1 is the distribution determined by $e^{-t\varrho}$.

Next we wish to discuss exponentials of distributions in connection with the Trotter product formula. This requires a discussion of the topology of strong resolvent convergence, which we now give. More details on this topology are to be found in the appendix.

DEFINITION. Let SA(K) be the set of all selfadjoint operators on a Hilbert space K. A neighborhood basis of a selfadjoint operator S in the topology of strong resolvent convergence is given by the collection

$$\{G(S, E, \varepsilon): E \text{ is a finite subset of } K, \ 0 < \varepsilon < \infty\}$$

where $G(S,E,\varepsilon)=\{T \text{ in } SA(K)\colon ||(T+i)^{-i}h-(S+i)^{-i}h\,||<\varepsilon,h \text{ in } E\}.$ We refer to the elements of *SA(K) as the internal selfadjoint operators on *K. Information about *SA(K) may be obtained by transfer. For example, "For all T in *SA(K), λ in *C, $\lambda\neq 0$, $||(T+\lambda)^{-1}||\leq |\operatorname{Im}(\lambda)|^{-i}$ " is obtained by transfer from "For all S in SA(K), λ in C, $\lambda\neq 0$, $||(S+\lambda)^{-1}||\leq |\operatorname{Im}(\lambda)|^{-i}$.

LEMMA 7. An internal selfadjoint operator T is near standard in the topology of generalized strong convergence if and only if there is an S in SA(K) such that

$$||(T+i)^{-1}h - (S+i)^{-1}h|| \approx 0 \text{ for all } h \text{ in } K.$$

Proof. S = st(T) iff T is in

$$\{ \cap {}^*G(S, E, r) \colon E \text{ is finite, } 0 < r < \infty \}$$
 .

But T is in *G(S, E, r) iff

$$||(T+i)^{-1}h - (S+i)^{-1}h|| < r$$
 for all h in E, since E is finite.

LEMMA 8. Let S belong to SA(K) and T to *SA(K). Then $T \approx S$ if and only if

$$(3) e^{itT}h \approx e^{itS}h$$

for all finite t in *R and h in K.

If T and S are bounded from below by some finite number. Then (3) may be replaced with

$$(4) e^{-tT}h \approx e^{-tS}$$

for all finite nonnegative t in *R and h in K.

Proof. We give the proof that $T \simeq S$ implies (3) for t positive and finite. For any P in *SA(K) let $R(P) = (-iT+1)^{-1}$ and $E(P) = e^{iP}$. Let h be an arbitrary element of K. Then, by definition of $T \approx S$, one has $R(T)h \approx R(S)h$. Since R(T), R(S) are contractions it follows that $R(T)a \approx R(S)b$ for all near standard a, b in *K with $a \approx b$. In particular, since $s \to E(sS)$ is strongly continuous we conclude that E(sS)h is near standard for all finite s in *R, so that $||(R(T) - R(S))E(sS)h|| \approx 0$. Consequently, for all positive real λ , $||E((t-s)T)(R(T) - R(S))E(sS)h|| < \lambda$. Transferring equation (2.27) of Reference 3, page 501, gives

$$R(T)(E(tT)-E(tS))R(S)=i\int_0^t E((t-s)T)(R(T)-R(S))E(sS)ds$$

and provides the estimate $||R(T)(E(tT) - E(tS))R(S)h|| \leq \lambda t$ for all positive real λ so that $R(T)E(tT)h \approx E(tS)R(S)h$. For all t in *R, E(tS)R(S) = R(S)E(tS), follows by transferring the corresponding statement with t in R. Since E(tS)h is near standard we conclude

(5)
$$R(T)E(tT)h \approx R(T)E(tS)h.$$

Next we argue

$$E(tT)R(S)h \approx E(tT)R(T)h = R(T)E(tT)h \approx R(T)E(tS)h$$

 $\approx R(S)E(tS)h = E(tS)R(S)h$.

Since the range of R(S) is dense in K and since E(tS), E(tT) have finite norms we conclude that $E(tT)h \approx E(tS)h$.

If t is finite and negative the proof follows as above upon replacing T and S by -T and -S respectively.

For the proof of the converse we assume $e^{itT}he^{itS}h$ for all finite t in *R and h in K. From Lemma 12, appendix, it suffices to prove

$$(-iT+1)^{-1}h \approx (-iS+1)^{-1}h$$

for all h in K. This is done via the transferred Laplace transform formula $(-iT+1)^{-1}h=\int_0^\infty e^{-t}e^{itT}hdt$. Let $\lambda, \in >0$ be arbitrary real numbers. First choose a positive real number c so that $\int_c^\infty e^{-t}dt < \varepsilon$.

Thus, $(-iT+1)^{-1}h \approx (-iS+1)^{-1}h$ for all h in K.

The proof for the case of T and S finitely bounded below follows similarly.

REMARK. The proof above closely parallels Kato's proof [3, p. 502] of an analogous standard result.

REMARK. The restriction of (3) to finite t in *R is not superfluous. Let n be a positive infinite integer, T=1/n, S=0, and t=n. Then $T\approx S$ but

$$e^{itT}h = e^ih \approx h = e^{itS}h$$
.

EXAMPLE 8. Let $h_0: \mathbf{R}^p \to [0, \infty)$ be continuous and suppose e^{-th_0} is in L^1 and is positive definite. Let $H_0 = FM(h_0)F^{-1}$ where F is the unitary Fourier transform on $L^2(\mathbf{R}^p)$ and $M(h_0)$ is the selfadjoint

operator given as multiplication by the function h_0 . Examples of such operators are given by $h_0(k) = |k|^2$ when $H_0 = -\Delta$ and $h_0(k) = \sqrt{|k|^2 + m^2}$, m > 0. For a complete discussion of such operators, see [2]. Let τ be a real distribution on R^p which satisfies for some 0 < a < 1 and b > 0

$$\pm \tau(|f|^2) \leq a(H_0 f, f) + b||f||^2$$

for all f in $C_c^{\infty}(\mathbf{R}^p)$. Such a τ is called a small form perturbation of H_0 . As shown in [2] the form sum of H_0 and the continuous extension of τ to the form domain of H_0 is selfadjoint and bounded below. We denote this standard operator by $H_0 + \tau$. As concrete examples choose $H_0 = -\Delta$ and τ the delta function concentrated on the surface of a sphere in p > 1 dimensions and $\tau = \delta$, the delta function in p = 1 dimensions. Also when p = 1, $\tau = e^x \cos(e^x)$ is a small form perturbation of $-\Delta$. See [2] for more examples.

Let τ_n denote the regularizations of τ . Then, defining $H_0 + \tau_n$ as the operator sum, it follows from the strong convergence of the resolvents of $H_0 + \tau_n$ to those of $H_0 + \tau$, [2], that

$$H_0 + \tau_n \approx H_0 + \tau$$

for all positive infinite integers, n.

EXAMPLE 9. Let H_0 , V be nonnegative selfadjoint operators on a Hilbert space K. Suppose $D((H_0)^{1/2}) \cap D((V)^{1/2})$ is dense. Define the truncations of V by

$$V_n = egin{cases} V & ext{if} & V \leq n \ o & ext{if} & V > n \end{cases}$$

via the spectral theorem. Let H_0+V denote the standard self-adjoint operator defined as the form sum of H_0 and V in [1]. Then, since $(H_0+V_n+i)^{-1}$ converges strongly to $(H_0+V+i)^{-1}$, [1], we conclude

$$H_0 + V \approx H_0 + V_n$$

for all positive infinite integers, n. See [1] and [4] for concrete examples.

We now proceed to discuss the Trotter product formula for given form sums. A discussion of form sums may be found in [1]. Here H_0 is a selfadjoint operator on a Hilbert space K and H is the selfadjoint operator on K given as the form sum of H_0 and V in the following three cases:

Case 1. V is a selfadjoint operator and the operator sum H_0+

V is essentially selfadjoint.

Case 2. H_0 and V are bounded below selfadjoint operators with dense form domain intersection; and

Case 3. V is a small Hermitian form perturbation of H_0 .

In Case 1, H is just the closure of $H_0 + V$ and the Trotter product formula holds [12]:

(6)
$$e^{\lambda t H} = \text{strong limit } (e^{\lambda t H_0/n} e^{\lambda t V/n})^n$$

uniformly for t in compact subsets of $[0, \infty)$, where $\lambda = \pm i$. If H_0 , V are bounded below then (6) also holds with $\lambda = -1$.

In Case 2 Kato has shown [4] that (6) holds for $\lambda = -1$ but in general (6) is not known for $\lambda = \pm i$ and in fact there has been essentially no progress in extending (6) for $\lambda = \pm i$ beyond Case 1.

In Case 3, not only is Trotter's product formula unknown for $\lambda=\pm i$ and $\lambda=-1$ but in general it makes no sense. For example, if $H_0=-d^2/dx^2$ and $V=\delta$, the Dirac delta function on $L^2(\mathbf{R}^1)$, then $e^{\lambda V/n}$ is undefined in any standard sense. In some examples of Case 3 such as $H_0=-d^2/dx^2$ and $V(x)=e^x\cos{(e^x)}$ the formula (6) makes sense but its validity is unknown for $\lambda=-1$.

Elementary nonstandard analysis provides a convenient framework for obtaining a type of product formula in the cases discussed above.

THEOREM 10. A Product Formula:

Let H_0 , H be in SA(K) and V in *SA(K) where K is a separable Hilbert space. Suppose the closure $[H_0 + V]$ of the operator sum of H_0 and V is in *SA(K). Also assume $H \approx [H_0 + V]$.

Then there is an N in *N such that for all n > N, h in K and nonnegative finite t in *R:

$$(7) e^{\lambda t H} h \approx (e^{\lambda t H_0/n} e^{\lambda t V/n})^n h$$

where $\lambda = \pm i$.

If also, H, H_0 , V > c for some c in R then (7) holds with $\lambda = -1$.

Proof. By Lemma 8

$$(8) e^{\lambda t H} h \approx e^{\lambda t [H_0 + V]} h$$

for all h in K and all finite t in R.

Fix α to be a positive infinitesimal. Fix h in K and M in N.

It follows from the transferred Trotter product formula of Case 1 that there is a Q(h, M) in *N, (we suppress the α dependence), such that q > Q(h, M) implies

$$||e^{\lambda t[M_0+V]}h-(e^{\lambda tH_0/q}e^{\lambda tV/q})^qh||$$

for all t in *[0, M]. Let Q(h) be an upper bound for $\{Q(h, M): M \text{ is in } N\}$, which exists by [8, page 59]. Then q > Q(h) implies

$$(10) e^{\lambda t II} h \approx (e^{\lambda t H_0/q} e^{\lambda t V/q})^q h$$

for all finite t in *[0, ∞].

Let C be a countable basis for K. Let Q be an upper bound for $\{Q(h): h \text{ is in } C\}$, which again exists by [8]. Let q > Q, t in $*[0, \infty)$ be finite and h in K be arbitrary. Given β in $(0, \infty)$ choose k in C so that $||h - k|| < \beta$. Then

$$\begin{aligned} ||e^{\lambda t H}h - (e^{\lambda t H_0/q}e^{\lambda t V/q})^q h|| \\ & \leq ||(e^{\lambda t H}(h-k))|| \\ & + ||(e^{\lambda t H} - (e^{\lambda t H_0/q}e^{\lambda t V/q})^q)k|| \\ & + ||(e^{\lambda t H_0/q}e^{\lambda t V/q})^q (k-h)|| < 3\beta \end{aligned},$$

since all operators appearing are contractions and the 2nd term being infinitesimal is infact less than β . As β is arbitrary in $(0, \infty)$ the difference in (11) is infinitesimal and the proof is complete.

- REMARKS. (a) The theorem applies to Example (8) with $\lambda=\pm i$ and in case the distribution is nonnegative, such as with the delta functions, λ may also be taken as -1. The theorem also applies to Example (9) with $\lambda=\pm i,-1$. H is the form sum and in these examples, the V is either a truncation or a regularization with positive infinite integral index and so is a bounded element of *SA(K). Consequently $[H_0 + V] = H_0 + V$.
- (b) In case $\lambda = -1$ it suffices to assume H_0 , V > C as we will see in the appendix, Example 6.
- (c) If V is in SA(K) rather than *SA(K) then the formula (7) is equivalent to the standard formula (6).
- (d) In [9] we used a preliminary form of this product formula to express the dynamics of a singularly perturbed quantum system in terms of a nonstandard Feynman path integral.
- (e) If K is not separable, one still obtains a product formula but then the N in N depends on the N in K.

EXAMPLE 11. In Example 6 we observed that $e^{-t\delta}$ could be identified with 1. However, this identification may not be made in

the present context, for we have just seen that

$$e^{-t(-d^2/dx^2+\delta)}h pprox (e^{t(d^2/dx^2/n)}e^{-t\delta/n})^n h$$

or all n sufficiently large. Had we also identified $e^{-t\delta/n}$ with 1 we would have obtained

$$e^{-t(-d^2/dx^2+\delta)}h\approx e^{t(d^2/dx^2)}h$$

which implies $-d^2/dx^2 + \delta = d^2/dx^2$ and this is false.

APPENDIX. The Strong Resolvent Topology.

In this appendix T will denote an element of *SA(K) and S an element of SA(K), for K a complex Hilbert space. Conditions for and consequences of $T \approx S$ will be investigated. There are closely related standard results which are obtained by considering a sequence, $\{T_n\}$, in SA(K) and its (possible) limit S, in the strong resolvent topology. One could obtain corresponding nonstandard results by transferring to $T = T_n$, n infinite and such a procedure would be sufficient for the examples considered in this paper: $T_n = H_0 + V_n$. However, in this appendix we are not assuming that T in *SA(K) has this special form and so, in general simple transfer arguments are not necessarily sufficient to establish the desired results.

DEFINITION. The finite resolvent set of T, denoted $\rho(T)$, is the set of all finite λ in *C such that $T-\lambda$ is 1-1, onto and $||(T-\lambda)^{-1}||$ is finite.

EXAMPLES. By transfer of the resolvent equation

- (a) {finite $\lambda \varepsilon^* C$: Im $(\lambda) \approx 0$ } $\subset \rho(T)$;
- (b) If $T \ge 0$ then

{finite
$$\lambda \varepsilon^* R: \lambda < 0$$
, $\lambda \not\approx 0$ } $\subset \rho(T)$.

Lemma 12. Let ρ denote the intersection of the finite resolvent sets of S and T. Then

(i) $T \approx S$ if there is a λ in ρ so that

(12)
$$||(T-\lambda)^{-1}h - (S-\lambda)^{-1}h|| \approx 0$$

for all h in K.

(ii) If $T \approx S$ then (12) holds for all λ in ρ .

Proof. The proof follows from a resolvent equation as in [3, page 429] together with the fact that operators with finite norm take infinitesimal vectors into infinitesimal vectors.

DEFINITION. Let D be a subspace of the domain of S, D(S). D is core for S if S restricted to D has exactly one selfadjoint extension.

LEMMA 13. Let D be a core for S. If $D \subset D(T)$ and $||Sf - Tf|| \approx 0$ for all f in D then $T \approx S$.

Proof. E=(S+i)(D) is dense in K. Let g in K be arbitrary and let δ be a positive real number. Choose f in E so that $||f-g||<\delta/3$. Then

$$\begin{split} \|(T+i)^{-1}g - (S+i)^{-1}g\| \\ & \leq \|(T+i)^{-1}(g-f)\| + \|[(T+i)^{-1} - (S+i)^{-1}]f\| \\ & + \|(S+i)^{-1}(f-g)\| \\ & \leq \|g-f\| + \|(T+i)^{-1}(T-S)(S+i)^{-1}f\| + \|f-g\| \\ & \leq 2\delta/3 + \|(T-S)(S+i)^{-1}f\| < \delta \ . \end{split}$$

REMARK. See [3, page 424] for analogous standard result.

EXAMPLE 14. Let H_0 , V be in SA(K). Let V_n be the truncations of V as in Example 9. If the operator sum $H_0 + V$ is essentially selfadjoint then $H_0 + V_n \approx [H_0 + V]$, the closure of $H_0 + V$, for all positive infinite n. This follows directly from Lemma 13 because if $f \in D(V)$ then $Vf \approx V_n f$, $n \in {}^*N - N$, by dominated convergence. Thus,

$$(H_0 + V_n)f \approx [H_0 + V]f$$

for all f in the core $D(H_0 + V)$ for $[H_0 + V]$.

REMARK. It may happen that $T \approx S$ but that there is no core D for S on which T and S nearly agree. For example let V be a nonnegative locally L^1 function on R^1 which is not L^2 on any interval. Let V_n be the truncation of $V: V_n(x) = V(x)$ if $V(x) \leq n$ and $V_n(x) = 0$ otherwise. Let $S = -d^2/dx^2$ and T = 1 the form sum of S and V_n , n infinite. If f is in D(T) = D(S) and in K then f is continuous and unless f is identically zero, |f| is bounded away from zero on some interval. Consequently $||V_n f||$ is infinite. Since ||Sf|| is finite it follows that

$$f \in K \cap D(T) \longrightarrow ||Tf||$$
 infinite.

Thus, there is no converse to the previous lemma. On the other hand we do have

Lemma 15. If $T \approx S$ then to each h in D(S) there is an h' in

D(T) with $h' \approx h$ and $T(h') \approx S(h)$.

Proof. Define $h' = (T+i)^{-1}(S+i)h$. Then $h' \in D(T)$ and $h' = h + ((T+i)^{-1} - (S+i)^{-1})(S+i)h \approx h$. Further, (T+i)h' = (S+i)h so that $Th' \approx Sh$.

EXAMPLE 16. Let c belong to R. Let $H_0 \in SA(K)$, $V \in {}^*SA(K)$ be * bounded and H_0 , $V \ge c$. Suppose $H \in SA(K)$ and $H \approx H_0 + V$. By Lemma 15, for each h in D(H) there is an h' in $D(H_0 + V)$ such that $Hh \approx (H_0 + V)h'$ and $h \approx h'$. Thus

$$(Hh, h) pprox (Hh, h') pprox ((H_0 + V)h', h') \ \geq c ||h'||^2 \ pprox c ||h||^2 .$$

Since (Hh, h) and $c ||h||^2$ are in R we conclude $H \ge c$.

For bounded operators there are no domain problems so we expect

Lemma 17. Let S be bounded and let T be everywhere defined with finite norm.

- (a) If $Th \approx Sh$ for all h in a dense subspace D of K, then $T \approx S$.
 - (b) If $T \approx S$ then $Th \approx Sh$ for all h in K.

Proof. Since D is dense and S is bounded, D is a core for S so part (a) follows from Lemma 13.

For part (b), assume $T \approx S$ so that $(T+i)^{-1}h \approx (S+i)^{-1}h$ for all h in K. Since T has finite norm $h \approx (T+i)(S+i)^{-1}h$ for all h in K. Let $g = (S+i)^{-1}h$. Then $(S+i)g \approx (T+i)g$ for all g in (the range of $(S+i)^{-1}) = K$.

One of the reasons for the utility of the topology of strong resolvent convergence in perturbation theory is

LEMMA 18. Let ϕ be a bounded Borel function on the real line which is continuous except on a closed set of S spectral measure zero. If $T \approx S$ then $\phi(T)h \approx \phi(S)h$, for all h in K.

Proof. With only minor modifications, (replacing certain "="s with " \approx "s), the proof of Faris, [1, pages 40-42] for an analogous standard result works here.

LEMMA 19. Let T be near standard in the strong resolvent

topology. Then for all h in K and all $\lambda \approx 0$

$$e^{i\lambda T}h \approx h$$
.

If also $T \ge c$, c in R, then

$$e^{-\lambda T}h \approx h$$
.

Proof. By Lemma 8

$$e^{i\lambda T}h \approx e^{i\lambda S}h \approx h$$

by strong continuity of $t \to e^{itS}$.

EXAMPLE 20. Let T=n, n a positive, infinite integer. Let $\lambda=1/n\approx 0$. Then $e^{i\lambda T}h=e^ih\not\approx h$ so T is not near standard even though $e^{itT}h$ is near standard for all t in *R.

Similarly $(1+i\lambda T)^{-\imath}h\not\approx h$ while from Lemma 19 and the Laplace transform formula

$$(1+i\lambda G)^{\scriptscriptstyle -1}=\int_{\scriptscriptstyle 0}^{\scriptscriptstyle \infty}\!e^{-t}e^{-\imath\lambda tG}hdt$$

it follows that $(1+i\lambda G)^{-1}h\approx h$ for all near standard G. Thus, $(T+c)^{-1}h$ near standard for all h in K does not imply that T is near standard in ${}^*SA(K)$. We next discuss when $(T+c)^{-1}h$ near standard does imply T is near standard. We shall use the following hybrid notation.

For a_n in *R we let " $a_n \to 0$, n in N" indicate that for every real $0 < \varepsilon < \infty$ there is an N in N such that n > N implies $|a_n| < \varepsilon$.

LEMMA 21. If h_n is near standard in *K, if h is in *K and if

$$||h_n-h||\longrightarrow 0$$
, n in N

then h is near standard.

Proof. Though this lemma is a special case of [10, 8.4.29]. We include the following elementary proof:

Let $h_n=k_n+\varepsilon_n$, k_n in K, $\varepsilon_n\approx 0$. Then $||k_n-h||\leq ||k_n-h_n||+||h_n-h||\to 0$, n in N, so that $||k_n-k_m||<||k_n-h||+||h-k_m\to 0||$ as $n,m\to\infty$ in N. Thus $\{k_n\}$ is a Cauchy sequence in K. Let $k=\lim_{n\to\infty}k_n$. Then k is in K and for every δ in $(0,\infty)||h-k||\leq ||h-h_n||+||h_n-k_n||+||k_n-k||<\delta$, by choosing n sufficiently large. Thus, $h\approx k$.

LEMMA 22. Suppose that for some finite λ in *C with Im $(\lambda) \approx$ 0, $(T - \lambda)^{-1}h$ is near standard for all h in K. Then, for all finite

ho in *C with $\operatorname{Im}(
ho) \approx 0$ and $\operatorname{sign}(\operatorname{Im}(\lambda)) = \operatorname{sign}(\operatorname{Im}(
ho))$ and for all h in K, $(T -
ho)^{-1}h$ is near standard.

Proof. It suffices to consider those ρ 's satisfying $|\rho-\lambda|<|\text{Im}(\lambda)|$ and $|\rho-\lambda|\not\approx|\text{Im}(\lambda)|$. For such ρ 's there is the formula $(T-\rho)^{-1}=\sum_{j=0}^\infty(\rho-\lambda)^k(T-\lambda)^{-(k+1)}$. The lemma then follows from the observation that since $(T-\lambda)^{-1}$ maps standard vectors into near standard vectors and since $||(T-\lambda)^{-1}||$ is finite, $(T-\lambda)^{-1}$ in fact maps near standard vectors into near standard vectors so that $(T-\lambda)^{-(k+1)}$ maps standard vectors into near standard ones.

REMARK. See Kato [3, page 427] for an analogous standard result.

THEOREM 23. In order that T be near standard it is necessary and sufficient that both

- (a) for some finite, positive and noninfinitesimal c, $(T\pm ic)^{-1}h$ be near standard for all h in K; and
- (b) $(1+i\in T)^{-i}h\approx h$ for all positive infinitesimal ε and all h in K, be true.

Proof. The necessity follows from Lemma 12 and the discussion in Example 20. For sufficiency, define, by Lemma 22,

$$R(\pm ia)h = \operatorname{st}((T \pm ia)^{-1}h)$$

for all positive, finite noninfinitesimal a and all h in K.

Then

$$\begin{split} (R(x) &- R(y))h \ &pprox ((T+x)^{-1} - (T+y)^{-1})h \ &= (y-x)(T+x)^{-1}(T+y)^{-1}h \;. \end{split}$$

But

$$\begin{split} ||(T+x)^{-1}(T+y)^{-1}h - R(x)R(y)h|| \\ &\leq ||(T+x)^{-1}((T+y)^{-1} - R(y))h|| \\ &+ ||((T+x)^{-1} - R(x))R(y)h|| \approx 0 \end{split}$$

because $||(T+x)^{-1}||$ has finite norm. Thus $(R(x)-R(y))h\approx (y-x)R(x)R(y)h$ and since both sides are standard the pseudo-resolvent equation results:

$$R(x) - R(y) = (y - x)R(x)R(y) .$$

The proof may now be completed as in Kato's proof [3, page 503] of an analogous standard result.

THEOREM 24. In order that T be near standard it is necessary and sufficient that both

$$(13) e^{\lambda sT} h \approx h$$

for all positive infinitesimal s; and

$$(14) e^{\lambda tT}h$$

is near standard for all finite $t \ge 0$ in *R be true for all h in K with both $\lambda = +i$ and $\lambda = -i$.

If $T \ge c$ for some c in R and if (13) and (14) both hold with $\lambda = -1$ then T is near standard.

Proof. We give the proof in case $\lambda = \pm i$. The case of $\lambda = -1$ follows similarly. The necessity follows from Lemmas 8 and 19.

For the proof of sufficiency, fix h in K, finite c>0 in *R , $c \approx 0$. The Riemann integrals on ${}^*[0, \infty)$

$$\int_{0}^{\infty} e^{-ct} e^{\pm itT} h dt$$

are defined by the strong continuity of the integrand.

Fix $\varepsilon > 0$ in R. Choose b > 0 in R so that

$$\int_{b}^{\infty}\!e^{-arepsilon t}dt .$$

Let I be the collection of z's in $*[0, \infty)$ which satisfy

$$t$$
, s in $[0, b]$, $|t - s| < z$ imply

$$||\,(e^{\imath t T}\,-\,e^{\imath s T})h\,|| < \varepsilon c/(2(1\,-\,e^{-\imath b}))$$
 .

I is internal. The semigroup property and (13) show that I contains all positive infinitesimals. Consequently, there is a δ in $I \cap (0, \infty)$.

Let $a_k = k\delta$ for $k = 0, 1, \dots, n$, where n in N is chosen so that $n\delta < b$ while $(n+1)\delta \ge b$. Set $a_{n+1} = b$. Let

$$h_{arepsilon} = \sum\limits_{k=0}^{n-1} e^{ik\delta T} h \!\! \int_{a_k}^{a_{k+1}} \!\! e^{-ct} dt$$
 .

 h_{ε} is near standard being a finite sum of near standard vectors.

$$igg|\int_0^\infty e^{-ct}e^{itT}hdt-h_arepsilonigg|\ \le \sum_{k=0}^{n-1}\int_{a_k}^{a_{k+1}}e^{-ct}||(e^{itT}-e^{ik\delta T})h||dt$$

$$egin{align} &+\int_{b}^{\infty} e^{-ct} \,||\, e^{itT} h\,||\, dt \ &\leq \int_{0}^{b} (e^{-ct} arepsilon c/2 (1\,-\,e^{-cb})) dt \,+\, rac{arepsilon}{2} \ &= arepsilon \;. \end{gathered}$$

By Lemma 21,

$$\int_{0}^{\infty} e^{-\varepsilon t} e^{\pm itT} h dt$$

is near standard. From the Laplace transform formulas

$$(\mathit{T}\pm\mathit{ic})^{\scriptscriptstyle{-1}}=\mp\mathit{i}\!\int_{\scriptscriptstyle{0}}^{\scriptscriptstyle{\infty}}\!e^{\scriptscriptstyle{-\mathit{ct}}}e^{\scriptscriptstyle{\pm\mathit{itT}}}dt$$

we conclude that $(T \pm ic)^{-i}h$ is near standard. We have thus verified (a) of Theorem 23. We next verify (b).

Let $\varepsilon > 0$, $\varepsilon \approx 0$. Choose $\beta > 0$ to be infinite in such a way that $\varepsilon \beta \approx 0$. Then

$$egin{aligned} &||(1+iarepsilon T)^{-1}h-h||\ &=\left\|\int_0^\infty e^{-t}e^{+iarepsilon tT}hdt-h
ight\|\ &\leq \int_0^\beta e^{-t}||\,e^{iarepsilon tT}h-h\,||\,dt\ &+2||\,h\,||\int_{eta}^\infty e^{-t}dt\ &pprox 0 \end{aligned}$$

since $0 < t < \beta \rightarrow \varepsilon t \approx 0 \rightarrow e^{i\varepsilon tT} h \approx h$. Thus $(1 + i\varepsilon T)^{-1} h \approx h$ and this theorem follows from Theorem 23.

See Example 20 for a remote T. Notice that (13) fails, in other words, the semigroup is not strongly S-continuous.

REFERENCES

- 1. W. Faris, Self-Adjoint Operators, Springer-Verlag, (1975).
- 2. I. Herbst and A. Sloan, Perturbation of translation invariant positivity preserving semigroups on $L^2(\mathbb{R}^N)$, Trans. Amer. Math. Soc., 236 (1978), 325-360.
- 3. T. Kato, Perturbation Theory for Linear Operators, Springer-Verlag, (1966).
- 4. ——, Trotter's product formula for an arbitrary pair of selfadjoint contraction semigroups, Berkeley preprint.
- 5. D. Laugwitz, Eine einfuhrung der δ -funktionen, S.-B. Bayerische Akad. Wiss., 4 (1958), 41-59.
- 6. W. Luxemburg, Nonstandard Analysis. Lectures on A. Robinson's Theory of Infinitesimal and Infinitely Large Numbers, Mimeographed Notes, California Institute of Technology, (1962).

- 7. A. Robinson, Nonstandard Analysis, North-Holland, (1974).
- 8. A. Robinson and A. Lightstone, Non-Archimedian Fields and Asymptotic Expansions, North-Holland, (1975).
- 9. A. Sloan, An application of the nonstandard Trotter product formula, J. Math. Phys., 18 (1977), 2495-2496.
- 10. K. Stroyan and W. Luxemburg, Introduction to the Theory of Infinitesimals, Academic Press, (1976).
- 11. J. Thurber and J. Katz, Applications of Fractional Powers of Delta Functions, in "Victoria Symposium on Nonstandard Analysis", A. Hurd and P. Loeb, eds., Springer-Verlag, (1974).
- 12. H. Trotter, Approximation of semigroups of operators, Pacific J. Math., 8 (1958), 887-919.

Received August 4, 1977 and in revised form January 11, 1978. Supported in part by NSF MCS 76-07543.

GEORGIA INSTITUTE OF TECHNOLOGY ATLANTA, GA 30332

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California Los Angeles, California 90024

C. W. CURTIS

University of Oregon Eugene, OR 97403

C. C. MOORE

University of California Berkeley, CA 94720 J. Dugundji

Department of Mathematics University of Southern California Los Angeles, California 90007

R. FINN AND J. MILGRAM Stanford University

Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. Wolf

K. Yoshida

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF HAWAII UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 79, No. 1 May, 1978

Teófilo Abuabara, <i>A remark on infinitely nuclearly differentiable</i> functions	1
David Fenimore Anderson, <i>Projective modules over subrings of $k[X, Y]$</i>	-
generated by monomials	5
Joseph Barback and Thomas Graham McLaughlin, On the intersection of regressive sets	19
Murray Bell, John Norman Ginsburg and R. Grant Woods, <i>Cardinal</i>	
inequalities for topological spaces involving the weak Lindelof number	37
Laurence Richard Boxer, <i>The space of ANRs of a closed surface</i>	47
Zvonko Cerin, Homotopy properties of locally compact spaces at	
infinity-calmness and smoothness	69
Isidor Fleischer and Ivo G. Rosenberg, <i>The Galois connection between partial</i>	93
functions and relations.	93
John R. Giles, David Allan Gregory and Brailey Sims, <i>Geometrical</i>	
implications of upper semi-continuity of the duality mapping on a Banach space	99
Troy Lee Hicks, Fixed-point theorems in locally convex spaces	111
Hugo Junghenn, Almost periodic functions on semidirect products of	111
transformation semigroups	117
Victor Kaftal, On the theory of compact operators in von Neumann algebras.	11,
II	129
Haynes Miller, A spectral sequence for the homology of an infinite	
delooping	139
Sanford S. Miller, Petru T. Mocanu and Maxwell O. Reade, <i>Starlike integral</i>	
operators	157
Stanley Stephen Page, Regular FPF rings	169
Ghan Shyam Pandey, Multipliers for C, 1 summability of Fourier series	177
Shigeo Segawa, Bounded analytic functions on unbounded covering	
surfaces	183
Steven Eugene Shreve, Probability measures and the C-sets of	
Selivanovskij	189
Tor Skjelbred, Combinatorial geometry and actions of compact Lie	
groups	197
Alan Sloan, A note on exponentials of distributions	207
Colin Eric Sutherland, Type analysis of the regular representation of a	
nonunimodular group	225
Mark Phillip Thomas, Algebra homomorphisms and the functional	
calculus	251
Sergio Eduardo Zarantonello, A representation of H ^p -functions with	
$0 $	271