

Pacific Journal of Mathematics

**INTERSECTIONS OF TERMS OF POLYCENTRAL SERIES OF
FREE GROUPS AND FREE LIE ALGEBRAS**

TED HURLEY

INTERSECTIONS OF TERMS OF POLYCENTRAL SERIES OF FREE GROUPS AND FREE LIE ALGEBRAS

T. C. HURLEY

This paper is concerned with deriving bases for the lower central factors of a free group modulo the intersection of certain terms of the polycentral series with the lower central series, and with the identification of the intersections themselves. The results from the group case are used to derive analogous results for the free Lie Algebra.

Let G be an arbitrary group. The *commutator group* $[A, B]$ of the subsets A and B of G is the subgroup generated by all commutators $[a, b] = a^{-1}b^{-1}ab$, $a \in A$, $b \in B$. The *lower central series* of G is the chain of normal subgroups

$$G = G_1 \geq G_2 \geq \dots \geq G_i \geq G_{i+1} \geq \dots$$

where $G_{i+1} = [G_i, G]$.

Let m_1, \dots, m_n, \dots be a sequence of positive integers. For $n=1$ we define G_{m_1} , as above, to be the m_1 th term of the lower central series of G and then define the *polycentral series*,

$$G \geq G_{m_1} \geq \dots \geq G_{m_1, \dots, m_i} \geq G_{m_1, \dots, m_i, m_{i+1}} \geq \dots$$

of G relative to this sequence by the rule that $G_{m_1, \dots, m_{i+1}}$ is the m_{i+1} th term of the lower central series of G_{m_1, \dots, m_i} .

Let L be a Lie Algebra. By analogy to the group case we define the *polycentral series of L relative to the sequence m_1, \dots, m_i, \dots*

$$L \geq L_{m_1} \geq \dots \geq L_{m_1, \dots, m_i} \geq L_{m_1, \dots, m_i, m_{i+1}} \geq \dots$$

in the following way: $L_1 = L$ and $L_{i+1} = L_i$. L (where in L , $A.B =$ the ideal generated by all $a.b$, $a \in A$, $b \in B$ and $.$ denotes the multiplication in L). Then, inductively, define

$$L_{m_1, \dots, m_{i+1}} = (L_{m_1, \dots, m_i})_{m_{i+1}}.$$

A group G is *nilpotent* iff $G_n = 1_G$ for some n and *polynilpotent* iff $G_{m_1, \dots, m_n} = 1_G$ for some (finite) sequence m_1, \dots, m_n . Similarly define a nilpotent Lie Algebra and a polynilpotent Lie Algebra. The n th lower central factor of G is G_n/G_{n+1} . A group G is said to be *residually- P* , where P is some group property, iff for each $g \in G$,

$g \neq 1$, there exists a normal subgroup N_g of G such that G/N_g has the property P and $g \notin N_g$.

If F is a free group, then $F/F_{m_1, \dots, m_n}$ is the *free polynilpotent group relative to the sequence* m_1, \dots, m_n . Similarly define $\mathcal{L}/\mathcal{L}_{m_1, \dots, m_n}$, the *free polynilpotent Lie Algebra*, where \mathcal{L} is a free Lie Algebra.

Free polynilpotent groups have been studied extensively by Gruenberg [1], who shows for example that they are residually finite p -groups for every prime p , and by Smelkin [5] who determines their lower central factors and also obtains bases for the additive groups of free polynilpotent Lie Algebras. Results along these lines have also been obtained by Ward [6].

It is a consequence of Gruenberg's result that $G = F/(F_{m_1, m_2, \dots, m_n} \cap F_{m_1 m_2 \dots m_n + r})$, ($r \geq 0$), is residually nilpotent and it may also be verified that for $s \geq m_1 m_2 \dots m_n + r$ the s th lower central factor of G and of $F/F_{m_1, m_2, \dots, m_n}$ are isomorphic—see Lemma 7 below. In this paper, all the lower central factors of $F/(F_{m, n} \cap F_{m n + r})$ ($r \geq 0$), are computed (Theorem E below). This includes Smelkin's result on $F_{m, n}$ (the case $r = 0$) and the proof here is independent of his relying solely on computations with basic group commutators. Also a classification of $F_{m, n} \cap F_{m n + r}$ as a product of certain commutator subgroups of F is obtained (Theorem A)—see Ward [6, Theorem 17.2] for the case $n = 2$. The lower central factors of $F/(F_{m, n} \cap F_{m n + r})$ turn out to be free abelian and consequently this group is residually torsion-free-and-nilpotent and therefore by Gruenberg [1] is residually a finite p -group for all primes p .

Using the results from the group case, $\mathcal{L}_{m, n} \cap \mathcal{L}_{m n + r}$ is identified (Theorem A') and a basis for the additive group of $\mathcal{L}/(\mathcal{L}_{m, n} \cap \mathcal{L}_{m n + r})$ is obtained (Theorem F).

2. Further notation and preliminary lemmas. Define, for any group G , with $n \geq 2$,

$$G(m, n; r) = \prod [G_{m+r_1}, G_{m+r_2}, \dots, G_{m+r_n}]$$

where the product is over all nonnegative integers r_1, \dots, r_n with $r_1 + r_2 + \dots + r_n = r$.

If $n = 1$, define $G(m, l; r) = G_{m+r}$, (for $m \geq 1, r \geq 0$).

Then it is clear that

$$G(m, n; 0) = G_{m, n}.$$

THEOREM A. $F_{m, n} \cap F_{m n + r} = F(m, n; r)$.

The special case

$$F_{m,2} \cap F_{2m+r} = \prod_{r_1+r_2=r} [F_{m+r_1}, F_{m+r_2}]$$

has been shown by Ward [6, Theorem 17.2].

To prove this and to describe the lower central factors of $F/F(m, n; r)$ I need some properties of *free groups*, *free Lie Algebras* and *basic commutators*, and the reader is referred to either Magnus, Karrass and Solitar [4, Chapter 5] or to M. Hall [2, Chapter 11]. An excellent account may also be found in P. Hall's notes [3, Chapter 5].

For a precise formulation of *bracket arrangement* β^s see [5, § 5.2]; imprecisely $\beta^s(A_1, \dots, A_s)$ means the commutator subgroup obtained by bracketing A_1, \dots, A_s in a certain fixed defined way. Thus, for example, if $\beta^4 = ((), ())$ then $\beta^4(A_1, A_2, A_3, A_4) = [[A_1, A_2], [A_3, A_4]]$.

The bracket arrangement is used only in Lemma 1 below, which is probably well-known and is in any case a simple application of P. Hall's 3-subgroup lemma [5, Theorem 5.2].

LEMMA 1. *Let A_1, \dots, A_s be normal subgroups of a group G , β^s a bracket arrangement of weight s and ρ a (fixed) permutation of $1, 2, \dots, s$. Then*

$$\beta^s(A_{\rho(1)}, A_{\rho(2)}, \dots, A_{\rho(s)}) \subseteq \prod_{\sigma} [A_{\sigma(1)}, A_{\sigma(2)}, \dots, A_{\sigma(s)}]$$

where the product is over all permutations σ of $1, 2, \dots, s$.

Proof. It is clear that we may assume ρ is the identity permutation.

The cases $s = 2$ and $s = 3$ are obvious. Now use induction on s . Let $s > 3$ and suppose $\beta^s = \beta^t \beta^q$ so that

$$\beta^s(A_1, \dots, A_s) = [\beta^t(A_1, \dots, A_t), \beta^q(A_{t+1}, \dots, A_s)].$$

By induction $\beta^t(A_1, \dots, A_t) \subseteq \prod_{\sigma} [A_{\sigma(1)}, \dots, A_{\sigma(t)}]$ and

$$\beta^q(A_{t+1}, \dots, A_s) \subseteq \prod_{\rho} [A_{\rho(t+1)}, \dots, A_{\rho(s)}]$$

and so it is only necessary to show that

$$(1) \quad [[A_{\sigma(1)}, \dots, A_{\sigma(t)}], [A_{\rho(t+1)}, \dots, A_{\rho(s)}]] \subseteq \prod_{\sigma} [A_{\sigma(1)}, \dots, A_{\sigma(s)}]$$

since for normal subgroups $[AB, C] \subseteq [A, C][B, C]$.

We can assume, since $[A, B] = [B, A]$, that $t \geq q$, and (1) can be proved by induction on q . If $q = 1$, then $t = s - 1$ and (1) is obvious. If $q > 1$ then by the 3-subgroup lemma

$$\begin{aligned} & [[A_{\sigma(1)}, \dots, A_{\sigma(t)}], [A_{\rho(t+1)}, \dots, A_{\rho(s)}]] \\ & \subseteq [[A_{\sigma(1)}, \dots, A_{\sigma(t)}, A_{\rho(s)}], [A_{\rho(t+1)}, \dots, A_{\rho(s-1)}]] \\ & \times [[A_{\sigma(1)}, \dots, A_{\sigma(t)}], [A_{\rho(t+1)}, \dots, A_{\rho(s-1)}], A_{\rho(s)}]. \end{aligned}$$

Applying the inductive hypothesis to the groups on the right hand side of the above gives the required result.

$G(m, n; r)$ as defined presents some notational difficulties and so in what follows I shall use the convention that \prod_r (where the expression after the product is expressed as a function of r_1, \dots, r_n) to mean the product over all nonnegative integers r_1, \dots, r_n with $r_1 + \dots + r_n = r$.

- LEMMA 2. (i) $G(m, n; r) \subseteq G(m, n; r - 1)$, ($r \geq 1$).
(ii) $[G(m, n; r), G(m, p; s)] \subseteq G(m, n + p; r + s)$.
(iii) $[G(m, n; r), G(m, n; s)] \subseteq G(m, n; r + s + 1)$.
(iv) $G(m, n + 1; r) \subseteq G(m, n; r + 1)$.

Proof. (i) is obvious.

(ii):- $[G(m, n; r), G(m, p; s)] = [\prod_r [G_{m+r_1}, \dots, G_{m+r_n}], \prod_s [G_{m+s_1}, \dots, G_{m+s_p}]]$. (If $n = 1$ interpret $\prod_r = G_{m+r}$ and similarly if $p = 1$). Hence $[G(m, n; r), G(m, p; s)] \subseteq \prod_{r,s} [[G_{m+r_1}, \dots, G_{m+r_n}], [G_{m+s_1}, \dots, G_{m+s_p}]]$ since for normal subgroups, $[AB, C] \subseteq [A, C][B, C]$ and $[A, BC] \subseteq [A, B][A, C]$. Now use Lemma 1 to complete the proof.

(iii):- $[G(m, n; r), G(m, n; s)] \subseteq \prod_{r,s} [[G_{m+r_1}, \dots, G_{m+r_n}], [G_{m+s_1}, \dots, G_{m+s_n}]]$.

$$\begin{aligned} \text{If } n = 1, \text{ then } \prod_{r,s} &= [G_{m+r}, G_{m+s}] \\ &\subseteq G_{m+r+m+s} \\ &\subseteq G_{m+r+s+1} = G(m, n; r + s + 1). \end{aligned}$$

If $n > 1$

$$\begin{aligned} [G_{m+r_1}, \dots, G_{m+r_n}] &\subseteq [G_{m+r_1+m+r_2}, G_{m+r_3}, \dots, G_{m+r_n}] \\ &\subseteq [G_{m+r_1+r_2+1}, G_{m+r_3}, \dots, G_{m+r_n}], \end{aligned}$$

and so

$$\begin{aligned} & [G(m, n; r), G(m, n; s)] \\ & \subseteq \prod_{r,s} [[G_{m+r_1+r_2+1}, G_{m+r_3}, \dots, G_{m+r_n}], [G_{m+s_1}, \dots, G_{m+s_n}]] \\ & \subseteq \prod_t [G_{m+t_1}, \dots, G_{m+t_{2n-1}}], \quad t = r + s + 1. \end{aligned}$$

The last step comes, as before, from Lemma 1.

(iv):- The proof is similar to (iii) and is omitted.

LEMMA 3. *Let A, B, C be normal subgroups of the group G . Suppose $a \in A, b \in B, c \in C$. Then*

$$[a, b, c][b, c, a][c, a, b] \equiv 1 \text{ modulo } [[A, B], [A, C]] \cdot [[A, B], [B, C]] \cdot [[A, C], [B, C]] .$$

Proof. By well-known commutator identities (see e.g., [4, Theorem 5.1]),

$$\begin{aligned} [a, b, c^a][c, a, b^c][b, c, a^b] &= 1 \\ \implies [a, b, c][c, a][c, a, b][b, c][b, c, a][a, b] &= 1 \\ \implies [a, b, [c, a]][a, b, c][a, b, c, [c, a]][c, a, [b, c]] \\ &\times [c, a, b][c, a, b, [b, c]][b, c, [a, b]][b, c, a][b, c, a, [a, b]] = 1 . \end{aligned}$$

Now since

$$\begin{aligned} [a, b, [c, a]] \text{ and } [a, b, c, [c, a]] &\in [[A, B], [A, C]] , \\ [c, a, [b, c]] \text{ and } [c, a, b, [b, c]] &\in [[A, C], [B, C]] , \\ [b, c, [a, b]] \text{ and } [b, c, a, [a, b]] &\in [[A, B], [B, C]] , \end{aligned}$$

and the groups are all normal we have that

$$\begin{aligned} [a, b, c][b, c, a][c, a, b] &\in [[A, B], [A, C]] \cdot \\ &[[A, B], [B, C]] \cdot [[A, C], [B, C]] . \end{aligned}$$

This lemma can be thought of as a generalization of the Jacobi identity for groups.

COROLLARY. *Suppose $a \in G(m, n; r), b \in G(m, p; s), c \in G(m, q; t)$. Then*

$$[a, b, c][b, c, a][c, a, b] \equiv 1 \text{ modulo } G(m, n + p + q; r + s + t + 1) .$$

Proof. Let $A = G(m, n; r), B = G(m, p; s), C = G(m, q; t)$. Then by Lemma 2,

$$\begin{aligned} [[A, B], [A, C]] &\subseteq [G(m, n + p; r + s), G(m, n + q; r + t)] \\ &\subseteq G(m, n + p + q; r + s + r + t) . \\ &\subseteq G(m, n + p + q; r + s + t + 1) . \end{aligned}$$

Similarly for $[[A, B], [B, C]]$ and $[[A, C], [B, C]]$.

3. Bases. Let c be a basic commutator with $c = [c_1, c_2]$ for

basic commutators c_1, c_2 . Say c is *structurally contained* in $F(m, 1; r) = F_{m+r}$ iff $c \in F_{m+r}$. Suppose it is defined what is meant by saying that c is structurally contained in $F(m, t; r)$ for all $1 \leq t < n$ and for all r . Then say c is *structurally contained* in $F(m, n; r)$ iff c_1 is structurally contained in $F(m, s; r_1)$, and c_2 is structurally contained in $F(m, t; r_2)$ for some $s, t \geq 1$ and some $r_1, r_2 \geq 0$ with $s + t = n$ and $r_1 + r_2 = r$. (Note: If $s \geq 2$, then automatically $c_2 \in F_m$ since $[c_1, c_2]$ is basic.)

For a basic commutator c , write $c \bar{\in} F(m, n; r)$ iff c is structurally contained in $F(m, n; r)$.

The following lemmas are easily verified.

LEMMA 4. If $c \bar{\in} F(m, n; r)$ then $c \in F(m, n; r)$.

LEMMA 5. (i) If $c \bar{\in} F(m, n; r)$ then (ii) $c \bar{\in} F(m, n - 1; r + m)$, ($n \geq 2$). $c \bar{\in} F(m, n; r - 1)$. ($r \geq 1$).

Proofs. These come directly from the definition and Lemma 2.

LEMMA 6. Suppose b, c are basic commutators such that $b \bar{\in} F(m, s; i)$, and $c \bar{\in} F(m, t; k)$, with $s + t = n$ and $i + k = r$. Then $[b, c]$ is congruent, modulo $F(m, n; r + 1)$, to a product of basic commutators, each of which is structurally contained in $F(m, n; r)$.

Proof. We may assume that F is finitely generated since b and c only involve a finite number of generators of F .

We may also assume that $b > c$ since $[b, c] = [c, b]^{-1}$. If b is a generator of F then clearly $[b, c]$ is a basic commutator. If $b = [b_1, b_2]$ where $b_2 \leq c$ then again $[b, c]$ is a basic commutator and there is nothing to be shown.

Define the *difference number* (d.n.) of a commutator $[b, c]$ where b and c are basic commutators with $b > c$, in the following way: Let b have weight w and suppose there exists v basic commutators of weight $\leq w$. Let c be the l th basic commutator. Then $l < v$ and define

$$\text{d.n. } [b, c] = v - l.$$

Now use induction on d.n. $[b, c]$. (This is a type of "backward induction").

Suppose $b = [b_1, b_2]$ where b_1, b_2 are basics and $b_2 > c$. Then weight $b_2 \geq$ weight c , giving that $b_2 \in F_m$ and consequently $b_1 \in F_m$.

Thus, we may assume $s \geq 2$ and therefore suppose

$$b_1 \bar{\in} F(m, s_1; i_1), \quad b_2 \bar{\in} F(m, s_2; i_2),$$

with $s_1 + s_2 = s$, $i_1 + i_2 = i$, $(s_1, s_2 \geq 1, i_1, i_2 \geq 0)$. Then by corollary to Lemma 3,

$$(3) \quad [b, c] = [b_1, b_2, c] \equiv [b_1, c, b_2][b_2, c, b_1]^{-1} \text{ modulo } F(m, n; r + 1).$$

Now d.n. $[b_1, c] < \text{d.n. } [b, c]$ and so by the inductive hypothesis $[b_1, c]$ is congruent, modulo $F(m, s_1 + t; i_1 + k + 1)$, to a product, $e_1^{\alpha_1} \cdots e_q^{\alpha_q}$ say ($\alpha_i \in \mathbf{Z}$), of basic commutators e_i with each $e_i \bar{\in} F(m, s_1 + t; i_1 + k)$.

Then using the commutator identity $[xy, z] = [x, z][x, z, y][y, z]$ and Lemmas 2 and 3, we see that

$$(4) \quad [b_1, c, b_2] = \prod_{j=1}^q [e_j, b_2]^{\alpha_j} \text{ modulo } F(m, n; r + 1).$$

Now for each j , $1 \leq j \leq q$, weight $e_j \leq \text{weight } b$. Also $b_2 > c$ and hence d.n. $[e_j, b_2] < \text{d.n. } [b, c]$. Therefore by induction each $[e_j, b_2]$, and consequently $[b_1, c, b_2]$, can be written, modulo $F(m, n; r + 1)$, as a product of basic commutators structurally contained in $F(m, n; r)$.

Also d.n. $[b_2, c] < \text{d.n. } [b, c]$ and therefore $[b_2, c]$ is congruent, modulo $F(m, s_2 + t; i_2 + k + 1)$, to a product, $f_1^{\beta_1} \cdots f_h^{\beta_h}$ say ($\beta_i \in \mathbf{Z}$), of basic commutators f_i such that $f_i \bar{\in} F(m, s_2 + t; i_2 + k)$.

For a particular f_i , note that there are two possibilities, either $f_i \geq b_1$ or $f_i < b_1$. In any case, as above, get

$$(5) \quad [b_2, c, b_1] \equiv \prod_{j=1}^h [f_j, b_1]^{\beta_j} \text{ modulo } F(m, n; r + 1).$$

Now consider the two cases.

Case 1. $f_j \geq b_1$. Now weight $f_j \leq \text{weight } b$ and also $b_1 > b_2 > c$ giving that d.n. $[f_j, b_1] < \text{d.n. } [b, c]$.

Case 2. $b_1 > f_j$. Now weight $b_1 < \text{weight } b$ and also, since f_j has bigger weight than c , $f_j > c$ giving that d.n. $[b_1, f_j] < \text{d.n. } [b, c]$.

Hence, by induction, each $[f_j, b_1]$ is congruent, modulo $F(m, n; r + 1)$, to a product of basic commutators structurally contained in $F(m, n; r)$ and therefore from (5), so is $[b_2, c, b_1]$. Thus $[b_1, c, b_2]$ and $[b_2, c, b_1]^{-1}$ are products, modulo $F(m, n; r + 1)$, of basic commutators of the required form and using (3) gives the required result.

THEOREM B. $F(m, n; r)/F(m, n; r + 1)$ is free abelian, freely

generated by those basic commutators c of weight $mn + r$ such that $c \in F(m, n; r)$.

Proof. It follows from Lemma 2 that the group is abelian. Also since $F(m, n; r) \subseteq F_{mn+r}$, and $F(m, n; r+1) \subseteq F_{mn+r+1}$ the freeness will follow from the Basis Theorem for Basic Commutators [4, Theorem 5.13A or 2, Theorem 11.2.4].

Therefore it is only necessary to show that the stated basic commutators generate the group.

If $n = 1$, this is equivalent to showing that F_{m+r}/F_{m+r+1} is generated by basic commutators of weight $m + r$ and this is part of the Basis Theorem. Let $n > 1$ and proceed by induction on n . If $a \in F(m, n; r)$ then a is a product of elements of the form

$$b = [b_1, \dots, b_n], \quad b_i \in F_{m+r_i} \quad \text{with} \quad r_1 + \dots + r_n = r.$$

By induction $[b_1, \dots, b_{n-1}]$ is congruent, modulo $F(m, n-1; r-r_n+1)$ to a product of basic commutators structurally contained in $F(m, n-1; r-r_n)$ and b_n is also congruent modulo F_{m+r_n+1} to a product of basic commutators contained in F_{m+r_n} . Now from Lemma 2,

$$[F(m, n-1; r-r_n+1), F_{m+r_n}] \subseteq F(m, n; r+1)$$

and

$$[F(m, n-1; r-r_n), F_{m+r_n+1}] \subseteq F(m, n; r+1)$$

and so the problem reduces to showing that $[a, b]$, with $a \in F(m, n-1; r-r_n)$ and $b \in F(m, 1; r_n)$, is congruent, modulo $F(m, n; r+1)$, to a product of basic commutators of weight $mn + r$ each of which is structurally contained in $F(m, n; r)$. Lemma 6 does this for us.

COROLLARY. *Let b be a basic commutator. Then*

$$b \in F(m, n; r) \iff b \in F(m, n; r).$$

Proof. We have already seen (Lemma 4) that if $b \in F(m, n; r)$ then $b \in F(m, n; r)$.

Suppose now $b \in F(m, n; r)$. Then b , being a basic commutator, is not 1 and so $b \notin \bigcap_{i=1}^{\infty} F_i$, since free groups are residually nilpotent. Thus $b \notin \bigcap_{i=1}^{\infty} F(m, n; i)$ and suppose $b \in F(m, n; s)$, $b \notin F(m, n; s+1)$. So $s \geq r$ and therefore by Theorem B, b is a unique product, modulo $F(m, n; s+1)$ of basic commutators structurally contained in $F(m, n; s)$. Therefore since b is also a basic commutator, this unique product must be precisely b , giving that $b \in F(m, n; s)$ and therefore $b \in (m, n; r)$.

THEOREM C. *Every element of $F_{m,n}$ can be written uniquely as*

$$c^{\alpha_1} \cdots c^{\alpha_s} \text{ mod } F(m, n; r),$$

where c_1, \dots, c_s are the basic commutators of weight $< mn + r$ structurally contained in $F_{m,n}$, $\alpha_i \in \mathbf{Z}$, and $c_1 < \dots < c_s$ in the ordering of the basic commutators.

REMARK. This theorem reduces to P. Hall's Basis Theorem for the case $n = 1$ and $m = 1$. This then can be considered as a "basis theorem" for certain terms of the polycentral series of F . Whereas in the Basis Theorem uniqueness modulo F_{r+1} is obtained, here F_{r+1} must be replaced by the complicated expression $F(m, n; r)$.

Proof of Theorem C. The proof is immediate from Theorem B. I omit spelling out the details.

THEOREM A.

$$F_{m,n} \cap F_{mn+r} = F(m, n; r).$$

Proof. The proof is now easy. Use induction on r . The case $r = 0$ is trivial.

It is clear that $F(m, n; r) \subseteq F_{m,n} \cap F_{mn+r}$. Suppose $r > 0$ and $a \in F_{m,n} \cap F_{mn+r}$. Then $a \in F(m, n; r - 1)$ by the inductive hypothesis, and by Theorem B, a is a product, modulo $F(m, n; r)$, of certain basic commutators of weight $mn + r - 1$. Since also $a \in F_{mn+r}$, this product must be 1 by the Basis Theorem, and so $a \in F(m, n; r)$. This proves Theorem A.

It is clear that these arguments work for the free Lie Algebra and so I state without proof the following two theorems.

Let T_i denote the set of basic commutators of weight i .

THEOREM B'. $\mathcal{L}_{m,n} / \mathcal{L}(m, n; r)$ has as additive basis those basic commutators of weight $< mn + r$ which are structurally contained in $\mathcal{L}_{m,n}$.

THEOREM A'.

$$\mathcal{L}_{m,n} \cap \mathcal{L}_{mn+r} = \mathcal{L}(m, n; r).$$

Theorems A' and B' may also be deduced easily from Theorem F and its corollary below.

THEOREM D. (Smelkin [5]). *Let $G = F/F_{m,n}$. Then G_i/G_{i+1} is free abelian, freely generated by the set $S_i = T_i \setminus R_i$, where R_i denotes the basic commutators of weight i structurally contained in $F_{m,n}$.*

Proof. Clearly G_i/G_{i+1} is generated by S_i since the basics structurally contained in $F_{m,n}$ vanish in G .

We need only consider $i \geq mn$ and hence suppose $i = mn + r$, $r \geq 0$.

Suppose a product Π of elements from S_i is in $F_{i+1}F_{m,n}$. Then

$$\Pi = ab, \quad a \in F_{m,n}, \quad b \in F_{i+1}.$$

Hence $a \in F_{m,n} \cap F_i = F(m, n; r)$ by Theorem A. Therefore by Theorem B, a is a product, modulo $F(m, n; r+1) (\subseteq F_{i+1})$, of basic commutators of weight i which are structurally contained in $F(m, n; r)$. The set of basic commutators structurally contained in $F(m, n; r)$ of weight i has empty intersection with S_i and so $a \equiv 1$ modulo F_{i+1} and hence $\Pi \equiv 1$ modulo F_{i+1} , giving that Π is identically 1.

THEOREM D'. (Smelkin [5]). *The set consisting of those basic commutators which are not structurally contained in $\mathcal{L}_{m,n}$ is an additive basis for $\mathcal{L}/\mathcal{L}_{m,n}$.*

This may be proved from Theorem A' by analogy to the group case. It is also a trivial consequence of Theorem F below.

The proof of the following theorem includes that of Theorem D.

THEOREM E. *Let $G = F/F(m, n; r)$. Then G_i/G_{i+1} is free abelian, freely generated by the set $S_i = T_i \setminus R_i$, where R_i denotes the basic commutators of weight i structurally contained in $F(m, n; r)$.*

Proof. Clearly S_i generates G_i/G_{i+1} . It is only necessary to consider $i \geq mn + r$ and hence let $i = mn + r + s$, $s \geq 0$. Suppose a product Π of elements from S_i is contained in $F(m, n; r)F_{i+1}$.

Then

$$\Pi = ab, \quad a \in F(m, n; r), \quad b \in F_{i+1}.$$

Therefore $a \in F_i \cap F(m, n; r) = F(m, n; r+s)$, by Theorem A. Thus a , by Theorem B, is a product, modulo $F(m, n; r+s+1)$, of basic commutators of weight i which are structurally contained in

$F(m, n; r + s)$. An element of S_i cannot be structurally contained in $F(m, n; r + s)$ for if so it would be constructurally contained in $F(m, n; r)$ —see Lemma 5. Hence Π is identically 1.

If G is any group then as e.g., in Magnus, Karrass and Solitar [4, Chapter 5] $\bigoplus_{i=1}^{\infty} G_i/G_{i+1}$ can be made into a Lie Algebra L by defining addition, $+$, in L by

$$\sum a_i G_{i+1} + \sum a'_i G_{i+1} = \sum a_i a'_i G_{i+1}, (a_i, a'_i \in G_i)$$

(i.e., using the group multiplication componentwise) and multiplication, \cdot , in L by first defining

$$a_i G_{i+1} \cdot a_j G_{j+1} = [a_i, a_j] G_{i+j+1}, (a_i \in G_i, a_j \in G_j)$$

and then using the distributive laws to extend this definition to all of $\bigoplus G_i/G_{i+1}$.

Let $G = F/F(m, m; r)$. Then there is the obvious surjective Lie Algebra homomorphism $\sigma: \mathcal{L}/\mathcal{L}(m, n; r) \rightarrow \bigoplus G_i/G_{i+1}$. The basic commutators structurally contained in $\mathcal{L}(m, n; r)$ vanish in $\mathcal{L}/\mathcal{L}(m, n; r)$ and the others are linearly independent in $\bigoplus G_i/G_{i+1}$ by Theorem E. This gives the following theorem.

THEOREM F. *The additive group of $\mathcal{L}/\mathcal{L}(m, n; r)$ has as basis the set of those basic commutators which are not structurally contained in $\mathcal{L}(m, n; r)$.*

COROLLARY. *$\mathcal{L}/\mathcal{L}(m, n; r)$ and $\bigoplus G_i/G_{i+1}$ are isomorphic Lie Algebras under the obvious map.*

LEMMA 7. (i) *If F/R is residually nilpotent so is $F/(R \cap F_n)$.*

(ii) *The m th lower central factor of F/R is isomorphic to that of $F/(R \cap F_n)$ for any $m \geq n$.*

Proof. (i) If F/R is residually nilpotent, then $\bigcap_i F_i R = R$. Now $\bigcap_i F_i(R \cap F_n) \subseteq \bigcap_i F_i R = R$. Also for $i \geq n$, $F_i(R \cap F_n) \subseteq F_n$ giving that $\bigcap_i F_i(R \cap F_n) = R \cap F_n$. Thus $F/(R \cap F_n)$ is residually nilpotent when F/R is.

(ii) Since $F_{m+1}(R \cap F_n) \subseteq F_{m+1}R$ we have a surjective map

$$F_m(R \cap F_n)/F_{m+1}(R \cap F_n) \longrightarrow F_m R/F_{m+1}R.$$

The kernel of this map is $F_m(R \cap F_n) \cap F_{m+1}R = F_{m+1}(R \cap F_n)$, for $m \geq n$, giving that the two groups are isomorphic.

COROLLARY 1. *$F/(F_{m_1, \dots, m_n} \cap F_{m_1 \dots m_n + r})$ is residually nilpotent.*

COROLLARY 2. $F/(F_{m_1, n} \cap F_{m_1 + r})$ is residually torsion-free and nilpotent.

COROLLARY 3. $F/(F_{m_1, n} \cap F_{m_1 + r})$ is residually a finite p -group for all primes p .

Proof of corollaries. By Gruenberg [1], $F/F_{m_1, \dots, m_n}$ is residually nilpotent, giving that $F/(F_{m_1, \dots, m_n} \cap F_{m_1 \dots m_n + r})$ is residually nilpotent. By Theorem E, $F/(F_{m_1, n} \cap F_{m_1 + r})$ has torsion-free lower central factors and is therefore residually torsion-free-and-nilpotent and hence again by Gruenberg [1] it is residually a finite p -group for all primes p .

It is also a corollary that the s th lower central factor of $F/F_{m_1, \dots, m_n}$ is isomorphic to that of $F/(F_{m_1, \dots, m_n} \cap F_{m_1 \dots m_n + r})$, for $s \geq m_1 \dots m_n + r$.

REFERENCES

1. K. W. Gruenberg, *Residual properties of infinite soluble groups*, Proc. London Math. Soc., (3) **7** (1957), 29–62.
2. M. Hall, *The Theory of Groups*, Macmillan, New York, 1959.
3. ———, *The Edmonton Notes on Nilpotent Groups*, Queen Mary College Lecture Notes, London, 1969.
4. W. Magnus, A. Karrass and D. Solitar, *Combinatorial Group Theory*, Interscience, New York-London-Sydney, 1966.
5. A. L. Smelkin, *Free Polynilpotent Groups*, Izv. Akad. Nauk SSSR Ser. Mat., **28** (1964), 91–122; English transl., Amer. Math. Soc. Transl., (2) **55** (1966), 270–304.
6. M. A. Ward, *Basic Commutators*, Phil. Trans. Royal Soc. of London, Ser. A, **264** (1969), 343–412.

Received September 13, 1977 and in revised form June 9, 1978.

UNIVERSITY COLLEGE
BELFIELD, DUBLIN 4, IRELAND

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Werner Bäni, <i>Subspaces of positive definite inner product spaces of countable dimension</i>	1
Marilyn Breen, <i>The dimension of the kernel of a planar set</i>	15
Kenneth Alfred Byrd, <i>Right self-injective rings whose essential right ideals are two-sided</i>	23
Patrick Cousot and Radhia Cousot, <i>Constructive versions of Tarski's fixed point theorems</i>	43
Ralph S. Freese, William A. Lampe and Walter Fuller Taylor, <i>Congruence lattices of algebras of fixed similarity type. I</i>	59
Cameron Gordon and Richard A. Litherland, <i>On a theorem of Murasugi</i>	69
Mauricio A. Gutiérrez, <i>Concordance and homotopy. I. Fundamental group</i>	75
Richard I. Hartley, <i>Metabelian representations of knot groups</i>	93
Ted Hurley, <i>Intersections of terms of polycentral series of free groups and free Lie algebras</i>	105
Roy Andrew Johnson, <i>Some relationships between measures</i>	117
Oldřich Kowalski, <i>On unitary automorphisms of solvable Lie algebras</i>	133
Kee Yuen Lam, <i>K O-equivalences and existence of nonsingular bilinear maps</i>	145
Ernest Paul Lane, <i>PM-normality and the insertion of a continuous function</i>	155
Robert A. Messer and Alden H. Wright, <i>Embedding open 3-manifolds in compact 3-manifolds</i>	163
Gerald Ira Myerson, <i>A combinatorial problem in finite fields. I</i>	179
James Nelson, Jr. and Mohan S. Putcha, <i>Word equations in a band of paths</i>	189
Baburao Govindrao Pachpatte and S. M. Singare, <i>Discrete generalized Gronwall inequalities in three independent variables</i>	197
William Lindall Paschke and Norberto Salinas, <i>C*-algebras associated with free products of groups</i>	211
Bruce Reznick, <i>Banach spaces with polynomial norms</i>	223
David Rusin, <i>What is the probability that two elements of a finite group commute?</i>	237
M. Shafii-Mousavi and Zbigniew Zielezny, <i>On hypoelliptic differential operators of constant strength</i>	249
Joseph Gail Stampfli, <i>On selfadjoint derivation ranges</i>	257
Robert Charles Thompson, <i>The case of equality in the matrix-valued triangle inequality</i>	279
Marie Angela Vitulli, <i>The obstruction of the formal moduli space in the negatively graded case</i>	281