

# Pacific Journal of Mathematics

## **ON THE ASYMPTOTIC BEHAVIOR OF LARGE PRIME FACTORS OF INTEGERS**

KRISHNASWAMI ALLADI AND PAUL ERDŐS

# ON THE ASYMPTOTIC BEHAVIOR OF LARGE PRIME FACTORS OF INTEGERS

K. ALLADI AND P. ERDÖS

**We prove results on the asymptotic behavior of large prime factors of the integers. The basic idea of the paper is that if  $k$  is any fixed integer, then the  $k$ th largest prime factor of  $n$ , denoted by  $P_k(n)$  is generally much bigger than  $\sum_{j>k} P_j(n)$ . We give precise estimates of this phenomenon. This paper is a sequel to an earlier paper by the authors.**

**1. Notations and definitions.** Throughout this paper the letters  $p$  and  $q$ , with or without subscript will denote primes.

Let  $n = \prod_{i=1}^r p_i^{\alpha_i}$ ,  $p_1 > p_2 > \dots > p_r$  be the canonical decomposition of an integer  $n > 1$  into primes. We set

$$(1.1) \quad A(n) = \sum_{i=1}^r \alpha_i p_i, \quad A^*(n) = \sum_{i=1}^r p_i$$

and

$$(1.2) \quad \Omega(n) = \sum_{i=1}^r \alpha_i, \quad \omega(n) = r.$$

Let  $A(1) = A^*(1) = \Omega(1) = \omega(1) = 0$ .

We may define the  $k$ th largest prime factor in two ways depending on whether we want to count prime factors according to multiplicity or not. To be more precise set

$$(1.3) \quad \begin{aligned} P_k^*(n) &= p_k \quad \text{for } k \leq \omega(n) \\ &= 0 \quad \text{for } k > \omega(n). \end{aligned}$$

We may also define

$$(1.4) \quad \begin{aligned} P_1(n) &= p_1 \\ P_k(n) &= P_1 \left( \frac{n}{P_1(n) \cdot P_2(n) \cdots P_{k-1}(n)} \right), \quad 1 < k \leq \Omega(n) \\ P_k(n) &= 0 \quad \text{for } k > \Omega(n). \end{aligned}$$

Observe that  $P_1(n) = P_1^*(n)$ .

The terms “average order” and “normal order” will mean the following: Let  $f$  be an arithmetic function and set

$$(1.5) \quad F(x) = \sum_{1 \leq n \leq x} f(n).$$

Suppose  $g$  is a monotonic function such that

$$(1.6) \quad \lim_{x \rightarrow \infty} \frac{G(x)}{F(x)} = 1$$

where

$$(1.7) \quad G(x) = \sum_{1 \leq n \leq x} g(n) ,$$

then  $f$  has average order  $g$ . Next, we say that two functions  $f$  and  $g$  are “nearly the same almost always” if for each  $\varepsilon > 0$

$$(1.8) \quad \lim_{x \rightarrow \infty} \frac{\psi_\varepsilon(x)}{x} = 1 ,$$

where

$$(1.9) \quad \psi_\varepsilon(x) = \sum_{\substack{1 \leq n \leq x \\ 1 - \varepsilon < (f(n)/g(n)) < 1 + \varepsilon}} 1 .$$

If in (1.9) and (1.8), the function  $g$  is monotonic, we say that  $f$  has normal order  $g$ .

Consider the sum

$$(1.10) \quad \psi(x, y) = \sum_{\substack{1 \leq n \leq x \\ P_1(n) \leq y}} 1 .$$

If  $\alpha \geq 1$  is a real number and  $y = x^{1/\alpha}$ , it is well known (see [10]) that

$$(1.11) \quad \rho(\alpha) = \lim_{x \rightarrow \infty} \frac{\psi(x, x^{1/\alpha})}{x}$$

exists. The limit in (1.11) is also defined if  $-\infty < \alpha < 1$  and

$$(1.12) \quad \rho(\alpha) = \begin{cases} 1 & 0 \leq \alpha < 1 \\ 0 & -\infty < \alpha < 0 . \end{cases}$$

The function  $\rho(\alpha)$  is a monotonic decreasing continuous function of  $\alpha$  for  $\alpha \geq 1$ .

Finally we define the sums

$$(1.13) \quad \begin{aligned} S_1(x, k) &= \sum_{2 \leq n \leq x} \frac{A(n) - P_1(n) - \cdots - P_{k-1}(n)}{P_1(n)} , & k \geq 1 \\ S_2(x, k) &= \sum_{2 \leq n \leq x} \frac{A^*(n) - P_1^*(n) - \cdots - P_{k-1}^*(n)}{P_1(n)} , & k \geq 1 \\ S_3(x, k) &= \sum_{2 \leq n \leq x} \frac{P_k(n)}{P_1(n)} , & k \geq 1 \\ S_4(x, k) &= \sum_{2 \leq n \leq x} \frac{P_k^*(n)}{P_1(n)} , & k \geq 1 . \end{aligned}$$

The aim of this paper is to obtain estimates for these sums  $S_i(x, k)$ ,  $i = 1, 2, 3, 4$ .

**2. General background and main theorem.** The results in this paper are in continuation of those in § 2 of [2].

It is a well known theorem of Hardy and Ramanujan [6], [7] that the functions  $\Omega(n)$  and  $\omega(n)$  both have average and normal order  $\log \log n = g(n)$ . This means that a number  $n$  usually has  $\log \log n$  prime factors and most of them occur square free. Thus it is natural to expect the large prime factors to occur with multiplicity one, most of the time. So one should be able to show that the functions  $A$  and  $A^*$  have the same average order. In an earlier paper [2] we showed this to be true and much more.

Not only do  $A$  and  $A^*$  have the same average order, but the function  $P_1(n)$  dominates the sums in (1.1) to such an extent that  $A, A^*$  and  $P_1$  have the same average order. More generally  $A(n) - P_1(n) - \dots - P_{k-1}(n)$  and  $P_k(n)$  have the same average order. It was observed in [1] that the functions  $P_k^*(n)$  and  $A^*(n) - P_1^*(n) - \dots - P_{k-1}^*(n)$  also have the same average order as  $P_k(n)$ , since the asymptotic analysis in [2] remains unaffected if the weak inequalities are replaced by strict ones. Thus we restate (without proof) the main theorem in [2] in a more complete form:

**THEOREM A.** *If  $k$  is a fixed positive integer then*

$$(2.1) \quad \begin{aligned} \sum_{1 \leq n \leq x} \{A(n) - P_1(n) - \dots - P_{k-1}(n)\} &\sim \sum_{1 \leq n \leq x} P_k(n) \sim \sum_{1 \leq n \leq x} P_k^*(n) \\ &\sim \sum_{1 \leq n \leq x} \{A^*(n) - P_1^*(n) - \dots - P_{k-1}^*(n)\} \sim a_k \frac{x^{1+1/k}}{(\log x)^{\alpha}} \end{aligned}$$

where  $a_k$  is a constant depending only on  $k$ , and is a rational multiple of  $\zeta(1 + 1/k)$  where  $\zeta$  is the Riemann zeta function. In addition for each  $k \geq 1$

$$(2.2) \quad \sum_{1 \leq n \leq x} \{A(n) - A^*(n)\} = x \log \log x + O(x) = o\left(\sum_{1 \leq n \leq x} P_k^*(n)\right).$$

Theorem A says that the average order in (2.1) is  $g(n) = a_k^* \cdot n^{1/k} / (\log n)^k$  where  $a_k^* = a_k \cdot (1 + 1/k)$ . An average is essentially influenced by two things—(i) the abnormally large values of a function, which certainly contribute to (2.1) and (ii) the values a function takes most often.

The question now arises whether  $A, A^*$ , and  $P_1$  are nearly the same almost always. The main theorem stated below answers this question in the affirmative.

**THEOREM B.** *If  $k$  is a fixed positive integer then*

$$(2.3) \quad S_1(x, k) \sim S_2(x, k) \sim S_3(x, k) \sim S_4(x, k) \sim a'_k \frac{x}{(\log x)^{k-1}}$$

where  $a'_1 = 1$  and  $a'_k$  for  $k > 1$  is a constant depending only on  $k$ , and is a rational multiple of  $e^\gamma$  where  $\gamma$  is Euler's constant. In addition for each  $k \geq 1$

$$(2.4) \quad \sum_{2 \leq n \leq x} \frac{A(n) - A^*(n)}{P_1(n)} = O\left(\frac{x}{e^c \sqrt{\log x \log \log x}}\right) = o\left(\sum_{2 \leq n \leq x} \frac{P_k^*(n)}{P_1(n)}\right)$$

where  $c$  is an absolute constant  $> 0$ .

**3. Consequences and motivation.** Statements (2.3) and (2.4) may be looked upon as analogues to (2.1) and (2.2). Theorem A said that  $A, A^*$  and  $P_1$  have the same average order,  $\pi^2 n / 6 \log n$ , ( $a_1 = \pi^2 / 12$ , see [2]). We can deduce from Theorem B the following.

**COROLLARY.** *The functions  $A, A^*$  and  $P_1$  are all nearly the same almost always. Also all three functions fail to possess a normal order.*

*Proof.* Consider two arithmetic functions  $f, g$  satisfying  $f(n) \geq g(n) > 0$ . Suppose that

$$(3.1) \quad \sum_{1 \leq n \leq x} \frac{f(n)}{g(n)} \sim x.$$

We rewrite (3.1) as

$$(3.2) \quad \sum_{1 \leq n \leq x} \left\{ \frac{f(n)}{g(n)} - 1 \right\} = o(x).$$

Since  $f/g \geq 1$  we infer from (3.2) that

$$(3.3) \quad \left| \frac{\psi_\varepsilon(x)}{x} - 1 \right| < \frac{o(x)}{\varepsilon \cdot x} \longrightarrow 0 \quad \text{as } x \longrightarrow \infty$$

for each  $\varepsilon > 0$ , where  $\psi_\varepsilon(x)$  is as in (1.8). So  $f$  and  $g$  are nearly the same almost always. (We can deduce (3.3) also if  $f(n) \leq g(n)$  for all  $n$ ).

Setting  $k = 1$  in (2.3) we see that (3.1) is true with  $f = A(n)$  and  $g(n) = P_1(n)$ . Therefore  $A$  and  $P_1$  are nearly the same almost always. Since  $A \geq A^* \geq P_1$ , the same is true for all three functions.

Now to show that these three functions do not have normal orders it suffices to show that one of them does not. It follows easily from a theorem of Elliott [5] on additive functions

$$(3.4) \quad f(n) = \sum_{p|n} f(p),$$

that  $A^*$  does not have a normal order. That proves the corollary.

REMARK. Since  $A(n) \geq \log n$ , it follows from (2.2) that

$$(3.5) \quad \left| \sum_{2 \leq n \leq x} \left\{ \frac{A^*(n)}{A(n)} - 1 \right\} \right| \leq \sum_{2 \leq n \leq x} \frac{A(n) - A^*(n)}{\log n} \\ = O\left(\frac{x \log \log x}{\log x}\right).$$

From (3.2), (3.3) and (3.5) we can deduce that  $A$  and  $A^*$  are nearly the same almost always.

Let us look a little more closely at (2.3) which for  $f = A$  or  $A^*$  and  $g = P_1$  is a more accurate form of (3.1). We may rewrite (2.3) as

$$(3.6) \quad \sum_{2 \leq n \leq x} \frac{A^*(n)}{P_1(n)} = \sum_{2 \leq n \leq x} 1 + \sum_{2 \leq n \leq x} \frac{P_2^*(n)}{P_1(n)} + \dots \\ + \sum_{1 \leq n \leq x} \frac{P_k^*(n)}{P_1(n)} + \dots$$

where

$$(3.7) \quad \sum_{2 \leq n \leq x} \frac{P_k^*(n)}{P_1(n)} \sim a'_k x / (\log x)^k.$$

We show in §5 that

$$(3.8) \quad a'_k = \int_1^\infty \rho(s - k) s^{k-2} ds$$

where  $\rho$  is defined in (1.11). We deduce from (3.8) in §6 that  $a'_k$  is a rational multiple of  $e^\gamma$  for  $k > 1$ . The integral representation is investigated in §6 and this leads to pretty connections with some related problems.

The next section is devoted to obtaining upper and lower bounds for  $S_i(x, k)$ ,  $i = 1, 2, 3, 4$ . This enables us to deduce the first four asymptotic relations in (2.3). It is only §5 that we prove (3.7) and (3.8). But the upper bound method in §4 is used in §5 to take care of the error terms arising out of (3.6) and (2.3). For the reader who does not want to go through the detailed proof, see [1], where some of the ideas of this paper and an earlier paper by the authors [2] are summarized.

We now move on to the proofs of our results.

4. Upper and lower bounds. In what follows,  $c_1, c_2, c_3, \dots$

denote absolute positive constants whose precise values will not be our concern. Also  $\exp \{x\} = e^x$ . We begin by proving

**THEOREM 1.** *There exists for each positive integer  $k$  a constant  $b_k$  and a real number  $x_0 = x_0(k)$  such that if  $x \geq x_0$  then  $S_i(x, k) > b_k \cdot x / (\log x)^{k-1}$  for  $i = 1, 2, 3, 4$ .*

To prove this we need

**LEMMA 1.** *Let  $s$  be a positive real number. Then*

$$\sum_{p > x} \frac{1}{p(\log p)^s} = \frac{1}{s(\log x)^s} + O(\exp\{-c_1 \sqrt{\log x}\}).$$

*Proof.* We use the Prime Number Theorem [4], [9] in the form

$$(4.1) \quad |\pi(x) - li(x)| = O(x \exp\{-c_2 \sqrt{\log x}\}).$$

Now write

$$\begin{aligned} \sum_{p > x} \frac{1}{p(\log p)^s} &= \int_{x+}^{\infty} \frac{d\pi(y)}{y(\log y)^s} = \int_x^{\infty} \frac{dy}{y(\log y)^{s+1}} + \int_{x+}^{\infty} \frac{d\{\pi(y) - li(y)\}}{y(\log y)^s} \\ (4.2) \quad &= \frac{1}{s(\log x)^s} + \frac{\pi(y) - li(y)}{y(\log y)^s} \Big|_{x+}^{\infty} \\ &\quad + O\left(\int_{x+}^{\infty} \frac{\{\pi(y) - li(y)\}}{y^2(\log y)^{s+1}} dy\right). \end{aligned}$$

Lemma 1 follows from (4.1) and (4.2).

*Proof of Theorem 1.* It suffices to prove Theorem 1 for the smallest of the four sums  $S_i(x, k)$ .

Assume first that  $k > 1$ . For  $x$  sufficiently large choose a prime  $p_1$  in the interval

$$(4.3) \quad k! x^{1/k+1} \leq p_1 \leq x^{1/k}.$$

Now choose primes  $p_2, p_3, \dots, p_k$  satisfying

$$(4.4) \quad \frac{p_1}{k} < p_k < \frac{p_1}{k-1} < p_{k-1} < \frac{p_1}{k-2} < \dots < \frac{p_1}{3} < p_3 < \frac{p_1}{2} < p_2 < p_1.$$

Consider any multiple  $m \leq x$  of  $p_1 p_2 \dots p_k$

$$(4.5) \quad m = n' p_1 p_2 \dots p_k.$$

Because of (4.3) and (4.4) we have

$$(4.6) \quad p_1 p_2 \dots p_k \leq x$$

and

$$(4.7) \quad n' \leq \frac{m}{p_1 \cdots p_k} \leq \frac{x}{(k!)^{n-1} x^{k/k+1}} < x^{1/k+1}.$$

By (4.7) and (4.4)

$$(4.8) \quad P_1(n') \leq n' < x^{1/k+1} \leq (k-1)! x^{1/k+1} < p_k.$$

Thus by (4.5) and (4.8) we see that  $P_k^*(m) = p_k$ . So any multiple  $\leq x$  of  $p_1 \cdots p_k$  has  $p_k$  as its  $k$ th largest prime factor ( $P_k^*$ ). So

$$(4.9) \quad S_4(x, k) = \sum_{2 \leq n \leq x} \frac{P_k^*(n)}{P_1(n)} \geq \sum_{\substack{2 \leq n \leq x \\ n = n' p_1 \cdots p_k \\ (p_i \text{ satisfying (4.4)}}} \frac{P_k^*(n)}{P_1(n)}.$$

We can estimate the second sum in (4.9) by using the well known result [11]

$$(4.10) \quad \sum_{p \leq k} \frac{1}{p} = \log \log x + c_3 + O(\exp \{-c_4 \sqrt{\log x}\}).$$

Observe that the second sum in (4.9) is

$$(4.11) \quad \begin{aligned} & \sum_{k! x^{1/(k+1)} \leq p_1 \leq x^{1/k}} \sum_{p_1/2 < p_2 < p_1} \sum_{p_1/3 < p_3 < p_1/2} \cdots \sum_{p_1/k < p_k < p_1/(k-1)} \frac{x}{p_1 \cdots p_k} \cdot \frac{p_k}{p_1} \\ &= x \sum_{k^n x^{1/(k+1)} \leq p_1 \leq x^{1/k}} \frac{1}{p_1^2} \left\{ \left( \prod_{j=2}^{k-1} \left( \sum_{p_1/j < p_j < p_1/(j-1)} \frac{1}{p_j} \right) \right) \cdot \left( \sum_{p_1/k < p_k < p_1/(k-1)} 1 \right) \right\} \\ &\geq b'_k x \sum_{k^n x^{1/(k+1)} \leq p_1 \leq x^{1/k}} \frac{1}{p_1 (\log p_1)^{k-1}} > b_k x / (\log x)^{k-1} \end{aligned}$$

by virtue of Lemma 1 and (4.10). Theorem 1 follows from (4.9) and (4.11), for  $k > 1$ . For  $k = 1$ , Theorem 1 is trivially true.

Now for an upper bound.

**THEOREM 2.** *All four sums  $S_i(x, k)$ ,  $i = 1, 2, 3, 4$  are  $O(x/(\log x)^{k-1})$  where  $k$  is an integer  $\geq 1$ , and the  $O$ -constant depends only on  $k$ .*

We need a few preliminary results before proving Theorem 2.

**LEMMA 2.** *Let  $k$  be a nonnegative integer and*

$$S_k^*(x) = \sum_{1 \leq p \leq x} (\log \log x - \log \log p)^k.$$

*Then*

$$S_k^*(x) = \frac{k! x}{(\log x)^{k+1}} + O_k \left( \frac{x \cdot \log \log x}{(\log x)^{k+2}} \right).$$



*Proof.* If we write  $S_k^*(x)$  as a Stieltjes integral, use the fact that

$$d\pi(y) = \frac{dy}{\log y} + d\{\pi(y) - li(y)\},$$

integrate the second integral by parts and then use (4.1) we get

$$(4.12) \quad S_k^*(x) = \int_4^x \frac{(\log \log x - \log \log y)^k}{\log y} dy + O\left(\frac{x}{(\log x)^{k+2}}\right).$$

Next

$$\begin{aligned} T_k(x) &= \int_4^x \frac{(\log \log x - \log \log y)^k}{\log y} dy \\ &= (\log \log x - \log \log y)^k li(y) \Big|_4^x \\ (4.13) \quad &+ k \int_4^x \frac{li(y)(\log \log x - \log \log y)^{k-1}}{y \log y} dy \\ &= O((\log \log x)^k) + k \int_4^x \frac{li(y)(\log \log x - \log \log y)^{k-1}}{y \log y} dy. \end{aligned}$$

But

$$(4.14) \quad li(y) = \frac{y}{\log y} + O\left(\frac{y}{\log^2 y}\right).$$

So the integral in (4.13) becomes

$$\begin{aligned} (4.15) \quad &k \int_4^x \frac{(\log \log x - \log \log y)^{k-1}}{\log^2 y} dy + O\left(\int_4^x \frac{(\log \log x - \log \log y)^{k-1}}{\log^3 y} dy\right) \\ &= I_1 + I_2. \end{aligned}$$

We split  $I_1$  into

$$(4.16) \quad I_1 = k \int_4^{x/(\log x)^{k+3}} + k \int_{x/(\log x)^{k+3}}^x.$$

Clearly in (4.16)

$$(4.17) \quad \int_4^{x/(\log x)^{k+3}} = O\left(\frac{x(\log \log x)^{k-1}}{(\log x)^{k+3}}\right) = O\left(\frac{x}{(\log x)^{k+2}}\right).$$

Regarding the second integral in (4.16) we observe that

$$\begin{aligned} (4.18) \quad &k \int_{x/(\log x)^{k+3}}^x = k \left\{ \frac{1}{\log x} + O\left(\frac{\log \log x}{\log^2 x}\right) \right\} \\ &\times \int_{x/(\log x)^{k+3}}^x \frac{(\log \log x - \log \log y)^{k-1}}{\log y} dy. \end{aligned}$$

Now the last integral in (4.18) is

$$(4.19) \quad T_{k-1}(x) + O\left(\frac{x(\log \log x)^{k-1}}{(\log x)^{k+3}}\right) = T_{k-1}(x) + O\left(\frac{x}{(\log x)^{k+2}}\right).$$

From the definition of  $T_k$  we have

$$(4.20) \quad T_0(x) = \frac{x}{\log x} + O\left(\frac{x}{\log^2 x}\right).$$

Now make the induction hypothesis that for  $k \geq 1$

$$(4.21) \quad T_{k-1}(x) = \frac{(k-1)!x}{(\log x)^k} + O\left(\frac{x(\log \log x)}{(\log x)^{k+1}}\right).$$

Then from equations (4.16) through (4.21) we deduce that

$$(4.22) \quad I_1 = \frac{k!x}{(\log x)^{k+1}} + O\left(\frac{x \log \log x}{(\log x)^{k+2}}\right).$$

By analysis very similar to the above one can show that

$$(4.23) \quad I_2 = O\left(\frac{x}{(\log x)^{k+2}}\right).$$

So from (4.22), (4.23), (4.15) and (4.13) we see that (4.21) is true for  $T_k(x)$  and so by induction for all  $k \geq 1$ . Lemma 2 follows from (4.12) and (4.21).

LEMMA 3. *Let  $x, y \geq 4$  be real numbers and  $k \geq 0$  an integer. Then*

$$\sum_{y \leq p \leq x} \frac{(\log \log x - \log \log p)^k}{p} = \frac{(\log \log x - \log \log y)^{k+1}}{k+1} + O_k((\log \log x - \log \log y)^k \exp\{-c_k \sqrt{\log y}\}).$$

*Proof.* As in the beginning of the proof of Lemma 2 we convert the above sum into a Stieltjes integral and replace  $d\pi(y)$  by  $dy/\log y$ . Lemma 3 can be easily proved by making the substitution  $\log \log x - \log \log y = t$ . We do not go through the details.

*Proof of Theorem 2.* It suffices to prove Theorem 2 for the largest of the four sums  $S_1(x, k)$ . That is we will show

$$(4.24) \quad S_1(x, k) = \sum_{2 \leq n \leq x} \frac{A(n) - P_1(n) - \cdots - P_{k-1}(n)}{P_1(n)} = O\left(\frac{x}{(\log x)^{k-1}}\right)$$

for  $k \geq 1$  an integer. We claim that it suffices to prove (4.24), for  $k > 1$  because for  $k = 1$  we have

$$\begin{aligned}
 (4.25) \quad S_1(x, 1) &= \sum_{2 \leq n \leq x} \frac{A(n)}{P_1(n)} = \sum_{2 \leq n \leq x} 1 + \sum_{2 \leq n \leq x} \frac{A(n) - P_1(n)}{P_1(n)} \\
 &= x - 1 + S_1(x, 2) \\
 &= O(x)
 \end{aligned}$$

assuming that (4.24) is true for  $k = 2$ . So from now on we assume that  $k > 1$ .

We write

$$\begin{aligned}
 (4.27) \quad \frac{A(n) - P_1(n) - P_2(n) - \dots - P_{k-1}(n)}{P_1(n)} &= \frac{P_k(n)}{P_1(n)} \\
 &\quad + \frac{P_{k+1}(n)}{P_1(n)} + \dots
 \end{aligned}$$

Let us denote a general nonzero term of (4.27) by  $p_k/p_1$ . We would like to know how often this term occurs in  $S_1(x, k)$ . The term  $p_k/p_1$  occurs as often as we can find integers  $n = p_1 p_2 \dots p_{k-1} \cdot p_k m \leq x$  where the  $p_i$  satisfy  $p_k \leq p_{k-1} \leq \dots \leq p_1$  and  $P_1(m) \leq p_{k-1}$ . If we fix the primes  $p_i$  to satisfy these conditions then the number of such  $n$  is given by

$$(4.28) \quad \psi\left(\frac{x}{p_1 p_2 \dots p_k}, p_{k-1}\right)$$

where  $\psi$  is defined in (1.10).

Thus we may rewrite (4.27) as

$$\begin{aligned}
 (4.29) \quad S_1(x, k) &= \sum_{2 \leq p_1 \leq x} \sum_{p_k \leq p_1} \sum_{p_k \leq p_{k-1} \leq p_1} \sum_{p_{k-1} \leq p_{k-2} \leq p_1} \dots \\
 &\quad \dots \sum_{p_3 \leq p_2 \leq p_1} \psi\left(\frac{x}{p_1 \dots p_k}, p_{k-1}\right) \frac{p_k}{p_1}.
 \end{aligned}$$

We first consider a subsum of (4.29) with a restriction on  $p_1$ . That is we choose  $\beta$  with  $0 < \beta < 1$ , whose value will be specified later, and consider  $p_1$  in (4.29) satisfying  $x^\beta \leq p_1 \leq x$ . We shall get an upper bound for this sum.

Observe that the sum in (4.29) with this extra condition on  $p_1$  is

$$\begin{aligned}
 (4.30) \quad &\leq \sum_{x^\beta \leq p_1 \leq x} \sum_{p_k \leq p_1} \sum_{p_k \leq p_{k-1} \leq p_1} \dots \sum_{p_3 \leq p_2 \leq p_1} \frac{x}{p_1 \dots p_k} \cdot \frac{p_k}{p_1} \\
 &= x \sum_{x^\beta \leq p_1 \leq x} \frac{1}{p_1^2} \sum_{p_k \leq p_1} 1 \sum_{p_k \leq p_{k-1} \leq p_1} \frac{1}{p_{k-1}} \dots \sum_{p_3 \leq p_2 \leq p_1} \frac{1}{p_2}.
 \end{aligned}$$

(Note: If  $k = 2$  in (4.30) we have only

$$(4.31) \quad x \sum_{x^\beta \leq p_1 \leq x} \frac{1}{p_1^2} \sum_{p_2 \leq p_1} 1$$

and no other terms. For  $k > 2$ , there is no confusion in (4.30). Because of this difference assume for the moment that  $k > 2$ . Then if we use Lemma 3 we infer

$$(4.32) \quad \sum_{p_3 \leq p_2 \leq p_1} \frac{1}{p_2} = O(\log \log p_1 - \log \log p_3).$$

Again by Lemma 3 and (4.32)

$$(4.33) \quad \sum_{p_4 \leq p_3 \leq p_1} \frac{1}{p_3} \sum_{p_3 \leq p_2 \leq p_1} \frac{1}{p_2} = O((\log \log p_1 - \log \log p_4)^2).$$

Iterating this process we get in (4.30) for  $k > 2$

$$(4.34) \quad O\left(x \sum_{x^\beta \leq p_1 \leq x} \frac{1}{p_1^2} \sum_{p_k \leq p_1} (\log \log x - \log \log p_k)^{k-2}\right)$$

by repeated use of Lemma 3. Now observe that because of (4.31) we see that (4.34) is true even for  $k = 2$ . Thus for  $k \geq 2$ , we may replace (4.30) by (4.34). Thus from now on we drop the assumption  $k > 2$ , but of course still assume  $k > 1$ .

To estimate (4.34) we use Lemma 2 which gives

$$(4.35) \quad O\left(x \sum_{x^\beta \leq p_1 \leq x} \frac{1}{p_1 (\log p_1)^{k-1}}\right).$$

Finally Lemma 1 and (4.35) imply that the sum in (4.35) and hence in (4.30) is

$$(4.36) \quad O\left(\frac{x}{\beta^{k-1} (\log x)^{k-1}}\right)$$

where the constant on the  $O$ -term in (4.36) depends only on  $k$  and not on  $\beta$ .

So (4.36) gives a bound for the sum in (4.29) with the condition  $x^\beta \leq p_1 \leq x$ . For the sum corresponding to  $p_1 \leq x^\beta$  we write

$$(4.37) \quad \sum_{2 \leq p_1 \leq x^\beta} = \sum_{m=0}^{\infty} \sum_{x^{\beta/2^m+1} \leq p_1 \leq x^{\beta/2^m}}.$$

To estimate (4.37) we use the following result of de Bruijn [3]; If  $y = x^{1/\alpha}$  then

$$(4.38) \quad \psi(x, y) = O(x \exp\{-c_6 \alpha\}).$$

In (4.37) consider the case

$$(4.39) \quad x^{\beta/2^m+1} \leq p_1 \leq x^{\beta/2^m}.$$

Then in (4.29) with the restriction (4.39) on  $p_1$  we have from (4.38)

the following:

$$(4.40) \quad \alpha = \frac{\log(x/p_1 \cdots p_k)}{\log p_{k-1}} \geq \left(1 - \frac{k\beta}{2^m}\right) \bigg/ \left(\frac{\beta}{2^m}\right) = \frac{2^m - k\beta}{\beta}.$$

We choose  $\beta = \beta(k)$ , depending on  $k$ , so small that

$$(4.41) \quad \alpha \geq \frac{2^m - k\beta}{\beta} > \frac{2^{m-1}}{\beta}.$$

Then by (4.38), (4.39), and (4.41) we will have in (4.29) for the subsum corresponding to (4.39)

$$(4.42) \quad \psi\left(\frac{x}{p_1 \cdots p_k}, p_{k-1}\right) = O\left(\frac{x}{p_1 \cdots p_k} \exp\{-c_6 2^{m-1}/\beta\}\right).$$

If we substitute (4.42) in (4.29) and analyze this sum just the way we derived (4.36) we get

$$(4.42) \quad \begin{aligned} & O\left(\frac{x}{\{\log(x^{\beta/2^{m-1}})\}^{k-1}} \exp\{-c_6 2^{m-1}/\beta\}\right) \\ &= O\left(\frac{x(2^{m+1}/\beta)^{k-1}}{(\log x)^{k-1} \exp\{c_6 2^{m-1}/\beta\}}\right). \end{aligned}$$

But then

$$(4.43) \quad \sum_{m=0}^{\infty} \frac{(2^{m+1}/\beta)^{k-1}}{\exp\{c_6 2^{m-1}/\beta\}} < \infty.$$

This means that (4.43), (4.42), and (4.36) imply that in (4.29)

$$S_1(x, k) = O(x/(\log x)^{k-1})$$

for  $k > 1$ . That completes the proof of Theorem 2.

It is interesting to note that Theorems 1 and 2 actually imply the first four asymptotic relations in Theorem B, as will be shown below. Before establishing this we prove the last part of Theorem B namely

**THEOREM 3.** *For each positive integer  $k$  we have*

$$\begin{aligned} \sum_{2 \leq n \leq x} \frac{A(n) - A^*(n)}{P_1(n)} &= O(x \exp\{-c_7 \sqrt{\log x \log \log x}\}) \\ &= o\left(\sum_{2 \leq n \leq x} \frac{P_k^*(n)}{P_1(n)}\right). \end{aligned}$$

*Proof.* First let  $1 \leq y \leq x$  and  $y = x^{1/\alpha}$ . N. G. de Bruijn [3] showed that if  $3 < \alpha < 4y^{1/2}/\log y$  then

$$(4.44) \quad \psi(x, y) = O(x \log^2 y \exp \{-\alpha \log \alpha - \alpha \log \log \alpha + c_8 \alpha\}) .$$

Take  $y = \exp \{\sqrt{\log x \log \log x}\}$ . Then from (4.44) we have

$$(4.45) \quad \psi(x, y) = O(x \exp \{-c_9 \sqrt{\log x \log \log x}\}) .$$

Next observe that

$$(4.46) \quad \frac{A(n) - A^*(n)}{P_1(n)} \leq \Omega(n) = O(\log n) .$$

We now split

$$(4.47) \quad \sum_{2 \leq n \leq x} \frac{A(n) - A^*(n)}{P_1(n)} = \sum_{\substack{2 \leq n \leq x \\ P_1(n) \leq y}} + \sum_{\substack{2 \leq n \leq x \\ P_1(n) > y}} = \sum_1 + \sum_2 .$$

Clearly from (4.46) and (4.45)

$$(4.48) \quad \sum_1 = O(\log x \cdot \psi(x, y)) = O(x \exp \{-c_{10} \sqrt{\log x \log \log x}\}) .$$

But then by Theorem A, (2.2), we have

$$(4.49) \quad \begin{aligned} \sum_2 &\leq \exp \{-\sqrt{\log x \log \log x}\} \sum_{2 \leq n \leq x} (A(n) - A^*(n)) \\ &= O(x \exp \{-c_{11} \sqrt{\log x \log \log x}\}) . \end{aligned}$$

The first equation in Theorem 3 follows from (4.47), (4.48) and (4.49). The second equation is a consequence of Theorem 1. That proves Theorem 3.

**THEOREM 4.** *For every integer  $k \geq 1$  we have*

$$S_1(x, k) \sim S_2(x, k) \sim S_3(x, k) \sim S_4(x, k) .$$

*Proof.* The smallest of the four sums is  $S_4(x, k)$ . By Theorem 1

$$(4.50) \quad S_1(x, k) \geq S_4(x, k) > b_k x / (\log x)^{k-1} .$$

The largest of the four sums is  $S_1(x, k)$ . Consider the difference

$$(4.51) \quad \begin{aligned} S_1(x, k) - S_4(x, k) &= \sum_{2 \leq n \leq x} \frac{A(n) - P_1(n) - \cdots - P_{k-1}(n) - P_k^*(n)}{P_1(n)} \\ &= \sum_{2 \leq n \leq x} \frac{A(n) - P_1(n) - \cdots - P_k(n)}{P_1(n)} \\ &\quad + \sum_{2 \leq n \leq x} \frac{P_k(n) - P_k^*(n)}{P_1(n)} \\ &= S_1(x, k+1) + \sum_{2 \leq n \leq x} \frac{P_k(n) - P_k^*(n)}{P_1(n)} . \end{aligned}$$

By Theorem 2

$$(4.52) \quad S_1(x, k+1) = O(x/(\log x)^k) .$$

But then

$$(4.53) \quad A(n) - A^*(n) = \sum_{j=1}^{\infty} P_j(n) - \sum_{j=1}^{\infty} P_j^*(n) = \sum_{j=1}^{\infty} \{P_j(n) - P_j^*(n)\} \\ \geq P_k(n) - P_k^*(n) .$$

So by (4.53) and Theorem 3 we have

$$(4.54) \quad \sum_{2 \leq n \leq x} \frac{P_k(n) - P_k^*(n)}{P_1(n)} \leq \sum_{2 \leq n \leq x} \frac{A(n) - A^*(n)}{P_1(n)} \\ = O(x \exp \{-c_7 \sqrt{\log x \log \log x}\}) .$$

Clearly from (4.51), (4.52) and (4.54)

$$(4.55) \quad S_1(x, k) - S_4(x, k) = O(x/(\log x)^k) .$$

Thus from (4.55) and (4.50) we deduce

$$(4.56) \quad S_1(x, k) \sim S_4(x, k) .$$

But since these are the smallest and largest sums, Theorem 4 follows from (4.56).

While proving Theorem 2 we did not use Lemmas 1, 2, and 3 in the forms in which they were stated, but used only the upper bounds they implied. These lemmas will play a role in obtaining asymptotic estimates, which we take up in the next section. We refer to the method of proof of Theorem 2 (namely the choice of  $\beta$  and the convergence of the series (4.43)), as the “upper bound method” and use this method to take care of the error terms arising out of the asymptotic estimates in what follows.

5. Asymptotic estimates. Our goal in this section is to prove

**THEOREM 5.** *Let  $k$  be a positive integer. Then all the four sums  $S_i(x, k)$ ,  $i = 1, 2, 3, 4$  are asymptotically equal to*

$$a'_k x / (\log x)^{k-1}$$

where

$$a'_k = \int_1^{\infty} \rho(s-k) s^{k-2} ds .$$

We need some lemmas before we go to the proof.

**LEMMA 4.** *If  $\alpha \geq 1$  and  $\varepsilon > 0$  then*

$$|\rho(\alpha + \varepsilon) - \rho(\alpha)| = O\left(\frac{\varepsilon}{\Gamma(\alpha + 1)}\right).$$

*Proof.* It is well known (see [10]) that  $\rho$  satisfies

$$(5.1) \quad \rho(\alpha) = 1 - \int_1^\alpha \frac{\rho(t-1)dt}{t}.$$

Furthermore (see [10], [3])

$$(5.2) \quad \rho(\alpha) < \frac{c_{12}}{\Gamma(\alpha + 1)}.$$

Combining (5.1) and (5.2) we get

$$0 \leq \rho(\alpha) - \rho(\alpha + \varepsilon) = \int_\alpha^{\alpha+\varepsilon} \frac{\rho(t-1)}{t} dt \leq \frac{\rho(\alpha-1)\varepsilon}{\alpha} = O\left(\frac{\varepsilon}{\Gamma(\alpha + 1)}\right)$$

because  $\rho$  is monotonic decreasing.

**LEMMA 5.** *There exists constants  $c_{13}$ ,  $c_{14}$  and  $c_{15}$  such that if  $\alpha \geq 1$  and  $y = x^{1/\alpha}$ ,  $x \geq 1$ , then*

$$|\psi(x, x^{1/\alpha}) - x\rho(\alpha)| \leq 2 \max\left(\frac{c_{13}x\alpha^2}{\exp\{c_{14}\sqrt{\log y}\}}, \frac{c_{15}x\alpha}{e^{\alpha/4} \cdot \log x}\right).$$

*Proof.* Lemma 5 is obtained by combining certain results of de Bruijn [3]. For the function  $\Lambda(x, x^{1/\alpha})$  defined by de Bruijn, it is known

$$(5.3) \quad |\psi(x, x^{1/\alpha}) - \Lambda(x, x^{1/\alpha})| < c_{13}x\alpha^2 \exp\{-c_{14}\sqrt{\log y}\}$$

and

$$(5.4) \quad |\Lambda(x, x^{1/\alpha}) - x\rho(\alpha)| < c_{15}x\alpha/(e^{\alpha/4} \cdot \log x).$$

Lemma 5 follows from (5.3) and (5.4).

*Proof of Theorem 5.* Because of Theorem 4 it suffices to prove Theorem 5 for one of sums  $S_i(x, k)$ . We consider  $S_1(x, k)$ . So we start with (4.29). (We assume  $k > 1$  since Theorem 5 is trivially true for  $k = 1$ . (See (5.1), (5.2) and Theorem 4.)

In (4.29) we first look at the contribution due to numbers for which

$$(5.5) \quad \frac{p_k}{p_1} < \frac{1}{(\log p_1)^{k+1}}.$$

We will get an upper bound for the contribution due to such numbers.



Let  $0 < \beta < 1$  be a real number whose value will be specified later. Then write

$$(5.6) \quad \sum_{2 \leq p_1 \leq x} = \sum_{x^\beta \leq p_1 \leq x} + \sum_{m=0}^{\infty} \sum_{x^{\beta/2^m+1} \leq p_1 \leq x^{\beta/2^m}}.$$

In the interval  $x^{\beta/2^m+1} \leq p_1 < x^{\beta/2^m}$  one has an upper bound for  $\psi$  given in (4.42), while for  $x^\beta \leq p_1 \leq x$  we use the trivial upper bound

$$(5.7) \quad \psi\left(\frac{x}{p_1 \cdots p_k}, p_{k-1}\right) \leq \frac{x}{p_1 \cdots p_k}.$$

Then for numbers satisfying (5.5) together with  $x^\beta \leq p_1 \leq x$ , we have the following bound in (4.29)

$$(5.8) \quad O\left(x \sum_{x^\beta < p_1 \leq x} \frac{1}{p_1(\log p_1)^{k+1}} \sum_{p_k \leq p_1} \frac{1}{p_k} \sum_{p_k \leq p_{k-1} \leq p_1} \frac{1}{p_{k-1}} \cdots \sum_{p_3 \leq p_2 \leq p_1} \frac{1}{p_2}\right).$$

Analysis similar to (4.32), (4.33) and (4.34) yields

$$(5.9) \quad \begin{aligned} & O\left(x \sum_{x^\beta \leq p_1 \leq x} \frac{1}{p_1(\log p_1)^{k+1}} \sum_{p_k \leq p_1} \frac{1}{p_k} (\log \log p_1 - \log \log p_k)^{k-2}\right) \\ &= O\left(x \sum_{x^\beta \leq p_1 \leq x} \frac{(\log \log p_1)^{k-1}}{p_1(\log p_1)^{k+1}}\right) = O\left(\frac{x(\log \log x)^{k-1}}{\beta^{k+1}(\log x)^{k+1}}\right) \end{aligned}$$

using Lemma 1. To estimate the contribution due to integers satisfying (5.5) for the case  $p_1 \leq x^\beta$ , we use the decomposition of the last sum of (5.6). Then the upper bound method yields

$$(5.10) \quad O\left(\frac{x(\log \log x)^{k-1}}{(\log x)^{k+1}}\right) = O\left(\frac{x}{(\log x)^k}\right)$$

provided  $\beta$  is suitably chosen. Thus from (5.9) and (5.10) we conclude that the contribution due to terms satisfying (5.5) is given by (5.10), and is smaller than the asymptotic term we are seeking.

Next we observe that the contribution due to terms for which  $p_1 = p_1(n)$  is small is negligible. For that purpose set

$$(5.11) \quad y = (\exp\{(\log x)^{2/3}\}).$$

With  $y$  as in (5.11) we have by (4.38)

$$(5.12) \quad \psi(x, y) = O(x \exp\{-c_{16}(\log x)^{1/3}\}).$$

So, if  $p_1 = P_1(n) \leq y$ , then

$$(5.13) \quad \begin{aligned} & \sum_{2 \leq n \leq x} \frac{A(n) - P_1(n) - \cdots - P_{k-1}(n)}{P_1(n)} \leq \sum_{\substack{2 \leq n \leq x \\ P_1(n) \leq y}} \Omega(n) \leq O\left(\sum_{\substack{2 \leq n \leq x \\ P_1(n) \leq y}} \log x\right) \\ &= O(\log x \psi(x, y)) = O(x \exp\{-c_{17}(\log x)^{1/3}\}). \end{aligned}$$

Because of (5.13) and (5.10), we assume from now on that

$$(5.14) \quad \frac{p_1}{(\log p_1)^{k+1}} \leq p_k \leq p_1; P_1(n) = p_1 > \exp \{(\log x)^{2/3}\}.$$

Once we assume (5.14) we can rewrite Lemma 5 as

$$(5.15) \quad \begin{aligned} \gamma\left(\frac{x}{p_1 p_2 \cdots p_k}, p_{k-1}\right) &= x \rho\left(\frac{\log(x/p_1 \cdots p_k)}{\log p_{k-1}}\right) \\ &+ O\left(\frac{x e^{-\alpha/4} \cdot \alpha}{p_1 \cdots p_k \log p_{k-1}}\right) \end{aligned}$$

where  $\alpha = \log(x/p_1 \cdots p_k)/\log p_{k-1}$ .

The idea is to substitute (5.15) in (4.29). It is then easy to take care of the contribution due to the error term in (5.15) in (4.29) by observing that (5.14)

$$(5.16) \quad \log p_{k-1} \geq \log p_k \sim \log p_1 > \frac{1}{2} \log p_1, \quad x \geq x_0.$$

This means if we substitute the  $O$ -term of (5.15) in (4.29), and use the upper bound method we get

$$(5.17) \quad O(x/(\log x)^k).$$

The convergence of a series like (4.43) is ensured this time by the  $e^{-\alpha/4}$  term in (5.15). Since (5.17) is smaller than the asymptotic term we are seeking, we may forget the contribution of the  $O$ -term in (5.15), in the sum (4.29).

As to the leading term of (5.15) we observe that

$$(5.18) \quad \rho\left(\frac{\log(x/p_1 \cdots p_k)}{\log p_{k-1}}\right) = \rho\left(\frac{\log x - \sum_{i=1}^k \log p_i}{\log p_{k-1}}\right).$$

By (5.14) we have

$$(5.19) \quad \log p_i = \log p_1 + O(\log \log p_1), \quad 1 \leq i \leq k.$$

Substituting (5.19) in (5.18) we get

$$(5.20) \quad \rho\left(\frac{\log(x/p_1 \cdots p_k)}{\log p_{k-1}}\right) = \rho\left\{\frac{\log x}{\log p_1} - k + O\left(\frac{\log x \cdot \log \log p_1}{\log^2 p_1}\right)\right\}.$$

Using Lemma 4 to estimate (5.20) we get

$$(5.21) \quad \rho\left(\frac{\log x}{\log p_1} - k\right) + O\left(\frac{\log x \cdot \log \log p_1}{\log^2 p_1 \cdot \Gamma(\alpha)}\right)$$

where  $\alpha$  is as in (5.15).

Thus the factor  $\rho$  in the leading term of (5.15) is equal to the quantity in (5.21). Recall that our idea is to substitute (5.15) in (4.29) and estimate the sum. The contribution of the  $O$ -terms in (5.21) can be obtained by the upper bound method. There is a  $\log x$  in the numerator, but a  $\log^2 p_1$  in the denominator. This time the presence of  $\Gamma(\alpha)$  in the denominator ensures convergence in a series like (4.43). Thus the upper bound method yields

$$(5.22) \quad O(x \log \log x / (\log x)^k)$$

as the contribution due to the  $O$ -term of (5.21). Thus we deduce that the main contribution from (4.29) comes by assuming (5.14) and replacing  $\psi(x/p_1 \cdots p_k, p_{k-1})$  by

$$(5.23) \quad \frac{x}{p_1 \cdots p_k} \rho \left( \frac{\log x}{\log p_1} - k \right).$$

So we replace (4.29) by

$$(5.24) \quad x \sum_{\exp\{(\log x)^{2/3}\} \leq p_1 \leq x} \frac{\rho((\log x / \log p_1) - k)}{p_1^2} \\ \times \sum_{p_1 / (\log p_1)^{k+1} \leq p_k \leq p_1} \sum_{p_k \leq p_{k-1} \leq p_1} \frac{1}{p_{k-1}} \cdots \sum_{p_3 \leq p_2 \leq p_1} \frac{1}{p_2}.$$

To estimate (5.24) we use Lemma 3. First we get

$$(5.25) \quad \sum_{p_3 \leq p_2 \leq p_1} \frac{1}{p_2} = (\log \log p_1 - \log \log p_3) + O(\exp\{-c_3 \sqrt{\log p_3}\}).$$

The contribution due to the  $O$ -term in (5.25) in (5.24) is taken care of by the upper bound method. This time the presence of  $\rho$  in (5.24) ensures convergence of a series like (4.43), because of (5.2). Actually every error term that arises in (5.24) by repeated use of Lemma 3 can be estimated by the upper bound method, yielding

$$(5.26) \quad O(x \exp\{-c_{18} \sqrt{\log x}\}).$$

So we need only look at the leading terms arising out of Lemma 3 in (5.24). After  $k-2$  applications of the lemma we are left with

$$(5.27) \quad x \sum_{\exp\{(\log x)^{2/3}\} \leq p_1 \leq x} \frac{\rho((\log x)/(\log p_1) - k)}{p_1^2} \\ \times \sum_{p_1 / (\log p_1)^{k+1} \leq p_k \leq p_1} \frac{(\log \log p_1 - \log \log p_k)^{k-2}}{(k-2)!}.$$

In (5.27) we use Lemma 2 to get

$$(5.28) \quad x \sum_{\exp\{(\log x)^{2/3}\} \leq p_1 \leq x} \frac{\rho((\log x)/(\log p_1)) - k}{p_1^2} \\ \times \left[ \frac{p_1}{(\log p_1)^{k-1}} + O\left(\frac{p_1(\log \log p_1)}{(\log p_1)^k}\right) \right].$$

As before, the  $O$ -term in (5.28) contributes

$$(5.29) \quad O(x \log \log x / (\log x)^k)$$

by use of the upper bound method. Finally the leading term in (5.28) is estimated by writing it as a Stieltjes integral. That is

$$(5.30) \quad x \sum_{\exp\{(\log x)^{2/3}\} \leq p_1 \leq x} \frac{\rho((\log x)/(\log p_1)) - k}{p_1(\log p_1)^{k-1}} \\ = x \int_{\exp\{(\log x)^{2/3}\}^-}^{x^+} \frac{\rho((\log x)/(\log y)) - k}{y(\log y)^{k-1}} d\pi(y) \\ = x \int_{\exp\{(\log x)^{2/3}\}}^x \frac{\rho((\log x)/(\log y)) - k}{y(\log y)^k} dy \\ + x \int_{\exp\{(\log x)^{2/3}\}^-}^{x^+} \frac{\rho((\log x)/(\log y)) - k}{y(\log y)^{k-1}} \\ \times d\{\pi(y) - li(y)\} = I_3 + I_4.$$

We can bound  $I_4$  rather easily. First observe that  $|\rho| \leq 1$ . Ignoring  $o$ , we integrate by parts, and use (4.1) to deduce

$$(5.31) \quad I_4 = O((x \exp\{-c_{19}(\log x)^{1/3}\})).$$

To estimate  $I_3$  write  $y = x^{1/s}$ . Then

$$(5.32) \quad I_3 = \frac{x}{(\log x)^{k-1}} \int_1^{(\log x)^{1/3}} \rho(s - k) s^{k-2} ds \\ = \frac{x}{(\log x)^{k-1}} \left[ \int_1^\infty - \int_{(\log x)^{1/3}}^\infty \right] \\ = a'_k \frac{x}{(\log x)^{k-1}} + O\left(\frac{x}{(\log x)^k}\right)$$

because of (5.2). So Theorem 5 follows from (5.32) and the preceding estimates.

REMARKS. Note that we have actually shown that

$$(5.33) \quad S_1(x, k) = a'_k \frac{x}{(\log x)^{k-1}} + O\left(\frac{x \log \log x}{(\log x)^k}\right).$$

Observe that  $S_1(x, k)$  is the largest of the four sums and  $S_4(x, k)$  is the smallest. Therefore, because of (4.55), we deduce a stronger form of Theorem 5, namely

$$(5.34) \quad S_i(x, k) = a'_k \frac{x}{(\log x)^{k-1}} + O\left(\frac{x \log \log x}{(\log x)^k}\right)$$

for  $i = 1, 2, 3, 4$ .

Thus we have proved all the statements of Theorem B, except the relation between  $a_k$  and  $e^\gamma$ . We do this in the next section.

**6. The constants  $a'_k$ .** It is obvious from Theorem 4 or (5.1) and (5.2) that  $a'_1 = 1$ . So we suppose  $k \geq 2$ . For  $k \geq 2$  write

$$(6.2) \quad \begin{aligned} a'_k &= \int_1^\infty \rho(s-k) s^{k-2} ds = \int_k^\infty \rho(s-k) s^{k-2} ds = \int_0^\infty \rho(t)(t+k)^{k-2} dt \\ &= \sum_{j=0}^{k-2} \binom{k-2}{j} k^{k-2-j} \int_0^\infty \rho(t) t^j dt = \sum_{j=0}^{k-2} \binom{k-2}{j} k^{k-2-j} f_j \end{aligned}$$

where

$$(6.3) \quad f_j = \int_0^\infty \rho(t) t^j dt.$$

In a recent paper, Knuth and Pardo [8], have studied the behavior of

$$(6.4) \quad \psi_k(x, y) = \sum_{\substack{1 \leq n \leq x \\ P_k^*(n) \leq y}} 1.$$

In the course of their investigations they show

$$(6.5) \quad f_j = e^\gamma g_j$$

where  $\gamma$  is Euler's constant and the  $g_j$  are recursively defined by

$$(6.6) \quad g_0 = g_1 = 1, \quad g_j = \frac{1}{j} \sum_{1 \leq i \leq j} \binom{j}{i} g_{j-i}, \quad j \geq 0.$$

Combining (6.2), (6.5), and (6.6) we infer that  $a'_k$  is a rational multiple of  $e^\gamma$  for  $k \geq 2$ . For instance

$$a'_2 = f_0 = e^\gamma g_0 = e^\gamma.$$

That completes the proof of Theorem B.

**Acknowledgments.** K. Alladi is grateful to Professors R. Miech and E. G. Straus for their advice and encouragement. They went through an early draft of this paper and made helpful suggestions. We also thank the referee for his critical comments and suggestions.

## REFERENCES

1. K. Alladi, *Analogues to the Hardy-Ramanujan theorems*, in Proceedings of the

- Conference of Number Theory and Computational Methods, Publications of "Matscience", Madras, India (1977).
2. K. Alladi and P. Erdős, *On an additive arithmetic function*, Pacific J. Math., **71** (2), (1977), 275-294.
  3. N. G. de Bruijn, *On the number of positive integers  $\leq x$  and free of prime factors  $> y$* , Indag. Math., **13** (1951), 50-60.
  4. K. Chandrasekharan, *Arithmetical Functions*, Springer-Verlag, New York, 1970.
  5. P. D. T. A. Elliott, *On a problem of Hardy and Ramanujan*, Mathematika **23**, (1976), 10-17.
  6. G. H. Hardy and S. Ramanujan, *The normal number of prime factors of a number  $n$* , Quarterly J. Math., **58** (1917), 76-92.
  7. G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, Oxford, Clarendon, 1960, 4-Ed.
  8. D. E. Knuth and L. T. Pardo, *Analysis of a simple factorisation algorithm*, (to appear).
  9. W. J. LeVeque, *Topics in Number Theory*, Vol. 2, Addison-Wesley, Reading, Mass., 1956.
  10. K. Norton, *Numbers with small prime factors and the least  $k$ th-power non-residue*, Memoirs of the A. M. S., **106**, Providence, Rhode Island, 1971.
  11. J. B. Rosser and L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, Illinois J. Math., **6** (1962), 64-94.

Received April 3, 1978 and in revised form October 16, 1978. This work was supported in part under National Science Foundation Grants No. MCS 77-01780 and MCS 78-02685.

UNIVERSITY OF CALIFORNIA  
 DEPARTMENT OF MATHEMATICS  
 LOS ANGELES, CALIFORNIA 90024  
 AND  
 MATHEMATICS INSTITUTE  
 THE HUNGARIAN ACADEMY OF SCIENCES  
 BUDAPEST, HUNGARY

*Current Address of first author:* University of Michigan  
 Department of Mathematics  
 Ann Arbor, MI 48109



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Older back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).

8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1979 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan



Krishnaswami Alladi and Paul Erdős, <i>On the asymptotic behavior of large prime factors of integers</i> .....	295
Alfred David Andrew, <i>A remark on generalized Haar systems in <math>L_p</math>, <math>1 &lt; p &lt; \infty</math></i> .....	317
John M. Baker, <i>A note on compact operators which attain their norm</i> .....	319
Jonathan Borwein, <i>Weak local supportability and applications to approximation</i> .....	323
Tae Ho Choe and Young Soo Park, <i>Wallman's type order compactification</i> .....	339
Susanne Dierolf and Ulrich Schwanengel, <i>Examples of locally compact noncompact minimal topological groups</i> .....	349
Michael Freedman, <i>A converse to (Milnor-Kervaire theorem) <math>\times R</math> etc.</i> ..	357
George Golightly, <i>Graph-dense linear transformations</i> .....	371
H. Groemer, <i>Space coverings by translates of convex sets</i> .....	379
Rolf Wim Henrichs, <i>Weak Frobenius reciprocity and compactness conditions in topological groups</i> .....	387
Horst Herrlich and George Edison Strecker, <i>Semi-universal maps and universal initial completions</i> .....	407
Sigmund Nyrop Hudson, <i>On the topology and geometry of arcwise connected, finite-dimensional groups</i> .....	429
K. John and Václav E. Zizler, <i>On extension of rotund norms. II</i> .....	451
Russell Allan Johnson, <i>Existence of a strong lifting commuting with a compact group of transformations. II</i> .....	457
Bjarni Jónsson and Ivan Rival, <i>Lattice varieties covering the smallest nonmodular variety</i> .....	463
Grigori Abramovich Kolesnik, <i>On the order of Dirichlet L-functions</i> .....	479
Robert Allen Liebler and Jay Edward Yellen, <i>In search of nonsolvable groups of central type</i> .....	485
Wilfrido Martínez T. and Adalberto Garcia-Maynez Cervantes, <i>Unicoherent plane Peano sets are <math>\sigma</math>-unicoherent</i> .....	493
M. A. McKiernan, <i>General Pexider equations. I. Existence of injective solutions</i> .....	499
M. A. McKiernan, <i>General Pexider equations. II. An application of the theory of webs</i> .....	503
Jan K. Pachl, <i>Measures as functionals on uniformly continuous functions</i> .....	515
Lee Albert Rubel, <i>Convolution cut-down in some radical convolution algebras</i> ....	523
Peter John Slater and William Yslas Vélez, <i>Permutations of the positive integers with restrictions on the sequence of differences. II</i> .....	527
Raymond Earl Smithson, <i>A common fixed point theorem for nested spaces</i> .....	533
Indulata Sukla, <i>Generalization of a theorem of McFadden</i> .....	539
Jun-ichi Tanaka, <i>A certain class of total variation measures of analytic measures</i> .....	547
Kalathoor Varadarajan, <i>Modules with supplements</i> .....	559
Robert Francis Wheeler, <i>Topological measure theory for completely regular spaces and their projective covers</i> .....	565