

Pacific Journal of Mathematics

**INTEGRAL REPRESENTATION FOR ELEMENTS OF THE
DUAL OF $ba(S, \Sigma)$**

MICHAEL KEISLER

INTEGRAL REPRESENTATION FOR ELEMENTS OF THE DUAL OF $ba(S, \Sigma)$

MICHAEL KEISLER

D. Mauldin presented conditions under which the dual of $ca(S, \Sigma)$ could be given an integral representation. W.D.L. Appling gave a "pseudo-representation" for the dual of $ba(S, \Sigma)$, but the latter was not the type obtained by Mauldin. This paper gives conditions that are necessary and sufficient for the existence of the classical type integral representation for the dual of $ba(S, \Sigma)$.

1. Introduction. Mauldin [11] found that an integral representation is possible for the elements of the dual of $ca(S, \Sigma)$ provided the cardinality of $ca(S, \Sigma)$ is $\leq 2^{\aleph_0}$ and the continuum hypothesis holds. Edwards and Wayment noted in [7] that for the function space of real-valued, absolutely continuous functions on $[0, 1]$, AC , the dual cannot be represented using Riemann or Lebesgue-type integrals. Using a different type of integral, a " v -integral", the latter authors then presented an integral representation for elements of AC^* . W.D.L. Appling [1] observed that an analogous result held for the subspace of $ba(S, \Sigma)$ made up of functions absolutely continuous with respect to a given member μ of $ba(S, \Sigma)$. The latter paper went on to give a "pseudo-representation" of elements of the dual of $ba(S, \Sigma)$. The representation was not the same type as obtained, conditionally, by Mauldin for $ca(S, \Sigma)$. Dunford and Schwarz [6], p. 374, noted that no "satisfactory representation" for members of $ba(S, \Sigma)^*$ was known.

In this paper it is proven that a representation such as that obtained by Mauldin for $ca(S, \Sigma)$ is possible for $ba(S, \Sigma)$ if and only if $ba(S, \Sigma)$ does not contain a "continuous" element, a condition which is shown by application of a theorem of Horn and Tarski [8], to be equivalent to the existence of a subfield of Σ that is isomorphic to the smallest field of subsets of $(0, 1]$ containing the intervals of the form $(k/2^n, (k+1)/2^n]$, for nonnegative integers k and n , hereafter denoted Σ_0 .

The integral here will be the same as that used by both Mauldin and Appling. The reader is referred to [2] for its definition and properties.

2. Filters. Properties of filters make them very convenient for several of the arguments. Filters are discussed in [12], and the first two lemmas below, stated without proof, can be found there.

The terminology used here differs slightly, so definitions are given first.

DEFINITION. Let Σ be a field of subsets of S . $F \subseteq \Sigma$ is a *filter* iff $v, w \in F$ implies $v \cap w \neq \emptyset$, $v \cap w \in F$ and for $z \in \Sigma$, $v \cup z \in F$. A filter F is an *ultrafilter* if it is contained properly in no other filter. B is a *base* for a filter F if $B \subseteq F$ and for $v \in F$ there is $w \in B$ such that $w \subseteq v$.

LEMMA 1. *Every filter is contained in an ultrafilter.*

LEMMA 2. *F is an ultrafilter iff $v \in \Sigma$ implies $v \in F$ or $v' \in F$ (where v' is the complement of v).*

For an ultrafilter F we can define $\mu(v) = 1$, if $v \in F$, and $\mu(v) = 0$, if $v \notin F$. It easily seen that $\mu \in ba(S, \Sigma)$. Therefore the lemma below can be used to construct some special members of $ba(S, \Sigma)$. Note, however, that there is no guarantee that the μ constructed above is in $ca(S, \Sigma)$, should Σ happen to be a σ -algebra.

LEMMA 3. *Let Q be a property that if possessed by $v \in \Sigma$, then for a subdivision D of v , there is $w \in D$ having property Q . A filter F , with a base of elements having property Q , is contained in an ultrafilter with a base of elements having property Q .*

Proof. Let \mathcal{L} be the family of all filters having a base of elements having property Q . \mathcal{L} is partially ordered by inclusion, and by Kuratowski's lemma [9] there is a maximal chain \mathcal{H} in \mathcal{L} containing F . Clearly $F^* = \bigcup \mathcal{H}$ is a filter, and since $v \in F^*$ implies $G \in \mathcal{H}$ such that $v \in G$, and hence a w with property Q in G such that $w \subseteq v$, then it follows that $F^* \in \mathcal{L}$. Suppose $z \in \Sigma$ and neither z nor z' is in F^* . Then $z \cap v \neq \emptyset$, for $v \in F^*$, and it follows that $M = \{y \in \Sigma \mid y \supseteq z \cap v, \text{ for some } v \in F^*\}$ is a filter containing F^* . The maximality of \mathcal{H} implies $M \notin \mathcal{L}$, and hence there is y_M in M such that $k \in M$ and $k \subseteq y_M$ implies k does not have property Q . Similarly for $N = \{y \in \Sigma \mid y \supseteq z' \cap v, \text{ for some } v \in F^*\}$. If $w \in F^*$ such that $z \cap w \subseteq y_M$ and $z' \cap w \subseteq y_N$, then $\{z \cap w, z' \cap w\}$ subdivides w , from which it follows that w does not have property Q . Since F^* is in \mathcal{L} , the latter must be false, and we have $z \in F^*$ or $z' \in F^*$; i.e., by Lemma 2, F^* is an ultrafilter.

3. Continuous and discrete elements. There are various uses of the words "atom" and "continuous" in the literature. Their use here is generally consistent with the literature, but we provide definitions for clarity.

DEFINITION. Let $\mu \in ba(S, \Sigma)$. μ is said to be an *atom* if $\{v \mid \mu(v) \neq 0\}$ is an ultrafilter.

DEFINITION. Let $\mu \in ba(S, \Sigma)$. μ is said to be *continuous* if for $\varepsilon > 0$ there is a subdivision D of S such that if E refines D , then for $v \in E$, $|\mu(v)| < \varepsilon$.

DEFINITION. Let $\mu \in ba(S, \Sigma)$. μ is said to be *discrete* if for λ such that $0 < \lambda$ and $\lambda \leq \int |\mu|$, then λ is not continuous.

The set of nonnegative members of $ba(S, \Sigma)$ will be denoted $ba(S, \Sigma)^+$. The next lemma is a restatement of results by Appling, [4], and is a consequence of a decomposition theorem in [5] since the continuous elements form a normal subspace.

LEMMA 4. Let $\mu \in ba(S, \Sigma)^+$. There is μ_0 continuous and μ' discrete such that $\mu = \mu_0 + \mu'$, and $\mu'(v) = \int_v \mu^*$, for $v \in \Sigma$, where $\mu^*(v) = \inf \{\sup \{\mu(w) \mid w \in D\} \mid D \text{ is a subdivision of } v\}$, for $v \in \Sigma$.

Lemma 5 reveals the relationship between atoms and discrete elements.

LEMMA 5. Let $\mu \in ba(S, \Sigma)^+$. μ is discrete iff there is a sequence of atoms, $\{\mu_n\}$, such that $\mu = \sum \mu_n$.

Proof. Lemma 4 implies that if μ is discrete, then $\mu = \int \mu^*$. For $v \in \Sigma$, if D is a subdivision of v , then there is w in D such that $\mu^*(w) = \mu^*(v)$. Thus for $v \in \Sigma$, by Lemma 3, there is an ultrafilter F_v with a base of elements on which μ^* equals $\mu^*(v)$; let μ_v be the atom on F_v such that $\mu_v(S) = \mu^*(v)$. It is immediate that $\mu^*(w) = \mu^*(v)$, for w in F_v and $w \subseteq v$. Therefore, $\mu_v \neq \mu_w$ implies $F_v \neq F_w$, and hence there is z such that $z \in F_v$ and $z' \in F_w$. Thus $\mu_v \vee \mu_w = \int \max \{\mu_v, \mu_w\} = \mu_v + \mu_w$. Since $ba(S, \Sigma)$ is a complete vector lattice, [5], and $\mu \geq \mu_v$, for every v , it follows that $\lambda = \sup M$, where $M = \{\mu_v \mid \mu_v > 0\}$, exists, equals $\sum M$, and $\lambda \leq \mu$. M summable implies that M is countable. And since $\lambda(v) \geq \mu_v(v) = \mu^*(v)$, for $v \in \Sigma$, it follows that $\lambda \geq \mu$.

For the converse, suppose $\mu = \sum \mu_n$, for some sequence $\{\mu_n\}$ of atoms. For $v \in \Sigma$, D a subdivision of v , and n a positive integer, $\mu_n(v) = \sup \{\mu_n(w) \mid w \in D\}$; thus $\sup \{\mu(w) \mid w \in D\} \geq \mu_n(v)$. It follows that $\int \mu^* \geq \sum^N \mu_n$, for every N , since there is a subdivision D such that for $w \in D$, $\sum^N \mu_n(w) = \sup \{\mu_n(w)\}^N$. Therefore, $\int \mu^* \geq \sum \mu_n = \mu$. From Lemma 4 we conclude that $\int \mu^* = \mu$, and hence μ is discrete.

4. **On integral representation.** Discrete and continuous elements play important roles insofar as integral representations of the dual are concerned. One of the most familiar examples of a field of sets, that generated by the half open intervals in $(0, 1]$, gives rise to a continuous element, the interval length measure. On the other hand, it is not difficult to construct a field for which no continuous elements exist; given a set S , let Σ be the field consisting of all finite subsets of S and their complements, then $ba(S, \Sigma)$ contains no nonzero continuous elements. Theorem 2 demonstrates that the first mentioned example has no classical integral representation for members of the dual, and Theorem 3 shows that the second example does have such a representation for members of the dual.

THEOREM 1. Σ contains a subfield Σ' isomorphic to Σ_0 iff there is a positive, continuous $\mu \in ba(S, \Sigma)$.

Proof. The "interval length" function can be extended to a member μ' of $ba(S, \Sigma')$ and, by Theorem 1.22 of [8], μ' can be extended to a $\mu \in ba(S, \Sigma)$. Clearly μ is a positive and continuous element of $ba(S, \Sigma)$. Conversely, a positive and continuous $\mu \in ba(S, \Sigma)$ can be used via an induction argument to construct a subfield Σ' of Σ isomorphic to Σ_0 .

DEFINITION. A member T of $ba(S, \Sigma)^*$ will be said to be *representable* provided there is $f: \Sigma \rightarrow R$ such that $T(\mu) = \int_S f \mu$, for $\mu \in ba(S, \Sigma)$. For nonnegative $\mu \in ba(S, \Sigma)$, P_μ represents the Lebesgue decomposition projection operator (see [2] for definition and properties).

THEOREM 2. If there is a positive $\mu \in ba(S, \Sigma)$ for which $I \in \Sigma$ such that $\mu(I) > 0$ implies there are disjoint $J, K \in \Sigma$ such that $I = J \cup K$ and $\mu(I) \notin \{\mu(J), \mu(K)\}$, then $T(\lambda) = P_\mu(\lambda)(S)$, for $\lambda \in ba(S, \Sigma)$, defines a nonrepresentable member of $ba(S, \Sigma)^*$.

Proof. Suppose T is representable, with $f: \Sigma \rightarrow R$ such that $\int_S f \lambda = T(\lambda)$, for each $\lambda \in ba(S, \Sigma)$. If for $I \in \Sigma$, $\mu'(V) = \mu(I \cap V)$, for $V \in \Sigma$, then $T(\mu') = \int_S f \mu' = \int_I f \mu$ and since $P_\mu(\mu')(S) = \mu(I)$, we have $\mu(I) = \int_I f \mu$. Applying a theorem of Kolmogoroff, [10],

$$0 = \int_S \left| f \mu - \int_S f \mu \right| = \int_S \left| f \mu - \mu \right| = \int_S |f - 1| \mu.$$

Let $Q = \{I \in \Sigma \mid \text{each subdivision of } I \text{ contains } J \text{ such that } \mu(J) >$

0 and $f(J) \geq 1/2$, and note that if $V \in Q$ and D subdivides V , there is $J \in D \cap Q$. Suppose $V \in \Sigma$ such that $\mu(V) > 0$. Since $\int_V |f - 1| \mu = 0$, there is a subdivision D of V such that if E refines D , then $\sum_E |f - 1| \mu < \mu(V)/2$, from which it follows that there is $I \in D \cap Q$. Since to each $V \in \Sigma$ on which $\mu(V) > 0$ there are disjoint $J, K \in \Sigma$ such that $V = J \cup K$, $\mu(J) > 0$ and $\mu(K) > 0$, then we may construct, inductively, a sequence $\{D_n\}$ of disjoint subsets of Q such that $I \in D_n$ implies there are $J, K \in D_{n+1}$ such that $I \supseteq J \cup K$ and $I \cap V = \emptyset$ for $V \in D_{n+1}$ and $V \notin \{J, K\}$. Then the collection of infinite maximal chains in $\cup D_n$ has cardinality $\geq 2^{\aleph_0}$ and any two contain disjoint sets. Each gives rise, by Lemma 3, to an ultrafilter with a base in Q and since $\mu \not\rightarrow 0$ on at most \aleph_0 ultrafilters, we conclude there is ultrafilter F with a base in Q (implying $f \not\rightarrow 0$) and on which $\mu \rightarrow 0$. Thus $\lambda(I) = 1$, for $I \in F$, and $= 0$, otherwise, defines $\lambda \in ba(S, \Sigma)$ for which $P_\mu(\lambda) = 0$, although $\int_S f \lambda \geq 1/2$. Therefore T is not representable.

Let $\{r_n\}_{n=1}^\infty$ be an enumeration of the rationals in $(0, 1)$. For $I \in \Sigma_0$, let $\mu_n(I) = 1/2^n$, if $r_n \in I$, and $= 0$, if $r_n \notin I$. $\{\sum_{n=1}^N \mu_n\}_{N=1}^\infty$ forms a Cauchy sequence in the Banach space $ba(S, \Sigma_0)$ (with variation norm). Let $\mu = \lim_{N \rightarrow \infty} \sum_{n=1}^N \mu_n$, and note that μ is a discrete function satisfying the hypothesis of Theorem 2. Thus the latter can produce non-representable members of $ba(S, \Sigma)^*$ through both continuous and discrete elements. However, applying it to continuous elements alone yields the following result.

COROLLARY. *If each member of $ba(S, \Sigma)^*$ is representable, then there is no positive continuous member of $ba(S, \Sigma)$.*

Lemma 4 assures us that if a $ba(S, \Sigma)$ contains no positive continuous elements, then all members of $ba(S, \Sigma)$ are discrete and can be written as in Lemma 5. This representation will be useful in Theorem 3.

THEOREM 3. *If $ba(S, \Sigma)$ contains no positive continuous element, then each T in $ba(S, \Sigma)^*$ is representable.*

Proof. For $B \subseteq U = \{\text{all ultrafilters in } \Sigma\}$, let $B^0 = \{F \in B \mid B(V) = \{F' \in B \mid V \in F'\} \text{ is infinite for } V \in F\}$. Let $g(1) = U$ and if g is defined for all ordinals less than an ordinal x , then let

$$g(x) = (\cap \{g(y) \mid y < x\})^0.$$

By transfinite induction (see [13]) g can be defined for all ordinals $\leq 2^{2^U}$. Since $x < y$ implies $g(x) \supseteq g(y)$, the domain of g has cardinality

greater than that of its range, and since the ordinals are well ordered, it follows that there are ordinals x and x' such that $x < x'$ and $g(x) = g(x')$. Thus $g(x) = g(x + 1)$ and if $B = g(x)$, then $B = B^0$.

If $F \in B$ and $J \in F$, then $B(J)$ must be infinite (since $F \in B^0$!) and hence we may select disjoint I_0 and I_1 such that each belongs to a member of B and $J = I_0 \cup I_1$. Thus if $B \neq \emptyset$, then we may, by induction, construct a sequence $\{D_n\}$ of subdivisions S such that for each n , every $I \in D_n$ is contained in a member of B , and there are distinct $J, K \in D_{n+1}$ such that $I = J \cup K$. The smallest field Σ' containing $\cup D_n$ is isomorphic to Σ_0 . Theorem 1 implies there is a positive continuous member of $ba(S, \Sigma)$. Thus $B = \emptyset$.

Let $A = \{\mu \mid \mu \text{ is an atom and } \mu(S) = 1\}$ and $F_\mu = \{V \mid \mu(V) = 1\}$, for $\mu \in A$. For $V \in \Sigma$, let $V^* = \{F \in \cap \{g(y) \mid y < z(V)\} \mid V \in F\}$, where $z(V) = \inf \{y \mid V \notin \cup g(y)\}$ (which exists since $g(x) = \emptyset$). Finally, define $f(V) = \sup \{T(\mu) \mid \mu \in A \text{ and } F_\mu \in V^*\}$, $V^* \neq \emptyset$, and $f(V) = 0$, if $V^* = \emptyset$. For $\mu \in A$, let $h(\mu) = \inf \{y \mid F_\mu \notin g(y)\}$ and note that $F_\mu \notin g(h(\mu))$ and $F_\mu \in \cap \{g(y) \mid y < h(\mu)\}$. Thus there is $V \in F_\mu$ such that $M = \{F \in \cap \{g(y) \mid y < h(\mu)\} \mid V \in F\}$ is finite. In fact, we can find V such that $M = \{F_\mu\}$, and since $V \notin \cup g(h(\mu))$, it follows that $h(\mu) = z(V)$ and hence $V^* = M$. Thus $f(V) = T(\mu)$, and for $W \in F_\mu$ and $W \subseteq V$, $f(W) = T(\mu)$, from which it follows that $\int_S f \mu = T(\mu)$. Since each $\mu \in ba(S, \Sigma)$ is discrete, there is a sequence $\{x_n\} \subseteq R$ and a sequence $\{\mu_n\} \subseteq A$ such that $\mu = \sum x_n \mu_n$. Since T is continuous, $T(\mu) = \sum x_n T(\mu_n)$. And since $\|\lambda\| = 1$, for $\lambda \in A$, it follows that f is bounded by $\|T\|$. A result in [3] implies that $\int_S f \mu$ exists and equals $\lim \int_S f \sum_N x_n \mu_n = \lim \sum_N x_n \int_S f \mu_n = \lim \sum_N x_n T(\mu_n) = T(\mu)$.

Theorems 1, 2 and 3 may be combined to produce the following characterization of those $ba(S, \Sigma)$ with representable duals.

THEOREM 4. *The following are equivalent.*

- (i) *Each $T \in ba(S, \Sigma)^*$ is representable.*
- (ii) *Each element of $ba(S, \Sigma)$ is discrete.*
- (iii) *There is no positive continuous member of $ba(S, \Sigma)$.*
- (iv) *There is no subfield of Σ isomorphic to Σ_0 .*

Thus for spaces with a dual representable in the sense of this paper one must turn to subspaces of $ba(S, \Sigma)$ (as Mauldin [11] has done with $ca(S, \Sigma)$, where Σ is a σ -algebra, and as Applying [1] has done with the space of functions absolutely continuous with respect to a given member of $ba(S, \Sigma)$) or be content to deal with a $ba(S, \Sigma)$ having no positive continuous element.

The author wishes to express his appreciation to Wayne C. Bell for his counsel in revising this paper from earlier versions.

REFERENCES

1. W.D.L. Appling, *An isomorphism and isometry theorem for a class of linear functionals*, Trans. Amer. Math. Soc., **199** (1974), 131-140.
2. ———, *Concerning a class of linear transformations*, J. London Math. Soc., **44** (1969), 385-396.
3. ———, *Summability of real-valued set functions*, Riv. Math. Univ. Parma, (2) **8** (1967), 77-100.
4. ———, *Continuity and the subdivision norm infimum function*, Notices Amer. Math. Soc., **17** (1970), 438.
5. S. Bochner and R.S. Phillips, *Additive set functions and vector lattices*, Ann. of Math., (2) **42** (1941), 316-324.
6. N. Dunford and J.T. Schwarz, *Linear Operators, Part I*, Interscience, New York, 1958.
7. J.R. Edwards and S.G. Wayment, *Representations of transformations continuous in BV norm*, Trans. Amer. Math. Soc., **154** (1971), 251-265.
8. A. Horn and A. Tarski, *Measures in Boolean algebras*, Trans. Amer. Math. Soc., **64** (1948), 467-497.
9. J.L. Kelley, *General Topology*, D. van Nostrand, New York, 1955.
10. A. Kolmogoroff, *Untersuchen uber den Integralbegriff*, Math. Ann., **103** (1930), 654-696.
11. R.D. Mauldin, *A representation theorem of the second dual of $C[0, 1]$* , Studia Math., **46** (1973), 197-200.
12. R.R. Stoll, *Sets, logic and axiomatic theories*, 2nd edition, W.H. Freeman, San Francisco, 1974.
13. P. Suppes, *Axiomatic Set Theory*, D. van Nostrand, Princeton, N. J., 1960.

Received January 11, 1977 and in revised form March 3, 1979.

ARKANSAS TECH UNIVERSITY
 RUSSELLVILLE, AR 72801

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

| | |
|--|-----|
| Richard Neal Ball, <i>Topological lattice-ordered groups</i> | 1 |
| Stephen Berman, <i>On the low-dimensional cohomology of some infinite-dimensional simple Lie algebras</i> | 27 |
| R. P. Boas and Gerald Thomas Cargo, <i>Level sets of derivatives</i> | 37 |
| James K. Deveney and John Nelson Mordeson, <i>Splitting and modularly perfect fields</i> | 45 |
| Robert Hugh Gilman and Ronald Mark Solomon, <i>Finite groups with small unbalancing 2-components</i> | 55 |
| George Grätzer, Andras Hajnal and David C. Kelly, <i>Chain conditions in free products of lattices with infinitary operations</i> | 107 |
| Benjamin Rigler Halpern, <i>Periodic points on tori</i> | 117 |
| Dean G. Hoffman and David Anthony Klarner, <i>Sets of integers closed under affine operators—the finite basis theorem</i> | 135 |
| Rudolf-Eberhard Hoffmann, <i>On the sobrification remainder $^s X - X$</i> | 145 |
| Gerald William Johnson and David Lee Skoug, <i>Scale-invariant measurability in Wiener space</i> | 157 |
| Michael Keisler, <i>Integral representation for elements of the dual of $ba(S, \Sigma)$</i> | 177 |
| Wayne C. Bell and Michael Keisler, <i>A characterization of the representable Lebesgue decomposition projections</i> | 185 |
| Wadi Mahfoud, <i>Comparison theorems for delay differential equations</i> | 187 |
| R. Daniel Mauldin, <i>The set of continuous nowhere differentiable functions</i> | 199 |
| Robert Wilmer Miller and Mark Lawrence Teply, <i>The descending chain condition relative to a torsion theory</i> | 207 |
| Yoshiomi Nakagami and Colin Eric Sutherland, <i>Takesaki's duality for regular extensions of von Neumann algebras</i> | 221 |
| William Otis Nowell, <i>Tubular neighborhoods of Hilbert cube manifolds</i> | 231 |
| Mohan S. Putcha, <i>Generalization of Lentin's theory of principal solutions of word equations in free semigroups to free product of copies of positive reals under addition</i> | 253 |
| Amitai Regev, <i>A primeness property for central polynomials</i> | 269 |
| Saburou Saitoh, <i>The Rudin kernels on an arbitrary domain</i> | 273 |
| Heinrich Steinlein, <i>Some abstract generalizations of the Ljusternik-Schnirelmann-Borsuk covering theorem</i> | 285 |