

Pacific Journal of Mathematics

EMBEDDING LATTICES INTO LATTICES OF IDEALS

GEORGE GRÄTZER, CRAIG ROBERT PLATT AND GEORGE WILLIAM SANDS

EMBEDDING LATTICES INTO LATTICES OF IDEALS

G. GRÄTZER, C. R. PLATT, AND B. SANDS

A lattice L is transferable iff, whenever L can be embedded in the ideal lattice of a lattice M , then L can be embedded in M . This concept was introduced by the first author in 1965 who also proved in 1966 that in a transferable lattice there are no doubly reducible elements. In fact, he proved that every lattice can be embedded in the ideal lattice of a lattice containing no doubly reducible elements. In a recent paper of the first two authors, the idea emerged that one should study transferability via classes K of lattices with the property that every lattice is embeddable in the ideal lattice of a lattice in K . This approach was used to establish that transferable lattices are semi-distributive. This investigation is carried further in this paper. Our main result shows that every lattice can be embedded in the ideal lattice of a lattice satisfying the two semi-distributive properties and two variants of Whitman's condition.

1. Introduction. It was shown by G. Grätzer ([6], [7]) that every transferable lattice L satisfies the condition

(X) L has no doubly reducible element.

In fact, he proved a stronger result, namely, that every lattice can be embedded in the ideal lattice of a lattice satisfying (X).

In general, if (P) is a lattice-theoretic property which is preserved by sublattices and which satisfies the assertion

$\mathcal{E}(P)$: every lattice can be embedded in the ideal lattice of a lattice satisfying (P) ,

then (P) is a property of all transferable lattices. In addition to (X), properties of a lattice L for which this assertion is known to hold include

(SF) L is sectionally finite (that is, all principal ideals are finite);

(SD $_{\wedge}$) for $a, b, c \in L$, $a \wedge b = a \wedge c$ implies that $a \wedge b = a \wedge (b \vee c)$;

(SD $_{\vee}$) for $a, b, c \in L$, $a \vee b = a \vee c$ implies that $a \vee b = a \vee (b \wedge c)$.

That $\mathcal{E}(SF)$ holds is a consequence of P. M. Whitman's embedding

theorem [10] and the observation that the partition lattice on a set S is isomorphic to the ideal lattice of the lattice of all finite partitions of S ; that $\mathcal{E}(SD_{\wedge})$ and $\mathcal{E}(SD_{\vee})$ hold is the content of a recent paper of G. Grätzer and C. R. Platt [8].

Consider the properties

- (W) for $a, b, c, d \in L$, $a \wedge b \leq c \vee d$ implies that $[a \wedge b, c \vee d] \cap \{a, b, c, d\} \neq \emptyset$;
- (W_l) for $a, b, c, d \in L$, $c \leq a \wedge b \leq c \vee d$ implies that $[a \wedge b, c \vee d] \cap \{a, b, c, d\} \neq \emptyset$;
- (W_u) for $a, b, c, d \in L$, $a \wedge b \leq c \vee d \leq a$ implies that $[a \wedge b, c \vee d] \cap \{a, b, c, d\} \neq \emptyset$.

K. Baker and A. W. Hales [2] proved that if a lattice satisfies (W), then so does its ideal lattice. Hence $\mathcal{E}(W)$ fails; however, in this paper, we will show that $\mathcal{E}(W_l)$ and $\mathcal{E}(W_u)$ hold. In fact, we will prove that every lattice can be embedded in the ideal lattice of a lattice satisfying the four properties (SD_{\vee}), (SD_{\wedge}), (W_l), and (W_u) simultaneously. More succinctly, our main result is

THEOREM. $\mathcal{E}((SD_{\vee}) \& (SD_{\wedge}) \& (W_l) \& (W_u))$ holds.

It follows from the theorem and the preceding remarks that every transferable lattice is sectionally finite and satisfies (SD_{\vee}), (SD_{\wedge}), (W_l), and (W_u). By a result of R. Antonius and I. Rival [1], we conclude:

COROLLARY. *Every transferable lattice satisfies (W).*

The proof of the theorem is contained in §2. In §3 we shall settle the truth or falsity of $\mathcal{E}(P)$ for most remaining combinations (P) of the above properties. In particular, it will be shown that $\mathcal{E}((SD_{\vee}) \& (SF) \& (X))$ holds and that $\mathcal{E}((SF) \& (SD_{\wedge}))$ and $\mathcal{E}((SF) \& (W_u))$ fail. With these results, we can determine the status of $\mathcal{E}(P)$ for all but two combinations (P) of the properties (X), (SF), (SD_{\vee}), (SD_{\wedge}), (W_l), and (W_u). These two will be given at the end of the paper.

2. Proof of the theorem.

DEFINITION 1. Let L be a lattice and let $\langle a, b, c, d \rangle$ be an ordered quadruple of elements of L . Then we will say that

- (i) $\langle a, b, c, d \rangle$ is a (W_l)-failure if $c \leq a \wedge b \leq c \vee d$ and $[a \wedge b, c \vee d] \cap \{a, b, c, d\} = \emptyset$;

(ii) $\langle a, b, c, d \rangle$ is a (W_u) -failure if $a \wedge b \leq c \vee d \leq a$ and $[a \wedge b, c \vee d] \cap \{a, b, c, d\} = \emptyset$;

(iii) $\langle a, b, c, d \rangle$ is an (SD_\wedge) -failure if $a \wedge b = a \wedge c = d$ and $a \wedge (b \vee c) \neq d$;

(iv) $\langle a, b, c, d \rangle$ is an (SD_\vee) -failure if $a \vee b = a \vee c = d$ and $a \vee (b \wedge c) \neq d$;

(v) $\langle a, b, c, d \rangle$ is a failure if it is any of the above four types of failures.

DEFINITION 2. Let L be a lattice, let $\langle a, b, c, d \rangle$ be a failure in L , and let φ be a homomorphism from a lattice M onto L . Then φ repairs $\langle a, b, c, d \rangle$, or $\langle a, b, c, d \rangle$ is repaired in M by φ , iff $\langle a', b', c', d' \rangle$ is never a failure in M of the same type as $\langle a, b, c, d \rangle$, for any $a' \in \varphi^{-1}(a)$, $b' \in \varphi^{-1}(b)$, $c' \in \varphi^{-1}(c)$, and $d' \in \varphi^{-1}(d)$.

LEMMA 3. Let K , L , and M be lattices, let $\varphi_1: M \rightarrow L$ and $\varphi_2: L \rightarrow K$ be onto homomorphisms, and let $\langle a, b, c, d \rangle$ be a failure in K . If $\langle a, b, c, d \rangle$ is repaired in L by φ_2 , then it is repaired in M by $\varphi_2 \circ \varphi_1$.

Proof. Each of the four conditions (SD_\vee) , (SD_\wedge) , (W_l) , and (W_u) can be expressed in the form $P(x, y, z, w) \Rightarrow Q(x, y, z, w)$, where P and Q are disjunctions of polynomial equations and hence are preserved under homomorphisms. Since $\langle a, b, c, d \rangle$ is a failure in K , there exist appropriate P and Q such that $P(a, b, c, d)$ holds but $Q(a, b, c, d)$ fails. Suppose that $\langle a, b, c, d \rangle$ is not repaired in M by $\varphi_2 \circ \varphi_1$; then there are elements $a', b', c', d' \in M$ such that $(\varphi_2 \circ \varphi_1)(x') = x$ for $x \in \{a, b, c, d\}$, $P(a', b', c', d')$ holds, and $Q(a', b', c', d')$ fails. Consequently, $P(\varphi_1(a'), \varphi_1(b'), \varphi_1(c'), \varphi_1(d'))$ holds in L . Since $\langle a, b, c, d \rangle$ is repaired in L by φ_2 , this implies that $Q(\varphi_1(a'), \varphi_1(b'), \varphi_1(c'), \varphi_1(d'))$ holds in L . But now $Q(a, b, c, d) = Q(\varphi_2(\varphi_1(a')), \varphi_2(\varphi_1(b')), \varphi_2(\varphi_1(c')), \varphi_2(\varphi_1(d')))$ holds in K , a contradiction.

Part of the proof of our theorem involves showing how to repair all failures in a lattice. Before describing the constructions by which this is accomplished, we make some observations.

Denote the lattice of ideals of a lattice L by $\mathcal{I}(L)$. Let L and K be lattices and let φ be a homomorphism of L onto K . For $I \in \mathcal{I}(K)$, consider the set

$$\varphi^{-1}(I) = \{x \in L \mid \varphi(x) \in I\}.$$

$\varphi^{-1}(I)$ is an ideal of L , and hence φ^{-1} is a map of $\mathcal{I}(K)$ into $\mathcal{I}(L)$

which is easily seen to be order preserving and one-to-one. Moreover, since meets of ideals are defined by set intersection, φ^{-1} is also meet preserving.

LEMMA 4. *The map $\varphi^{-1}: \mathcal{J}(K) \rightarrow \mathcal{J}(L)$ is an embedding if and only if φ satisfies the condition*

(*) *if $y \in L$, $x_1, x_2 \in K$, and $\varphi(y) \leq x_1 \vee x_2$, then $y \leq y_1 \vee y_2$ for some $y_1, y_2 \in L$ satisfying $\varphi(y_1) = x_1$, $\varphi(y_2) = x_2$.*

Proof. To prove the “if” direction, by the above remarks we need only show that for $I, J \in \mathcal{J}(K)$, $\varphi^{-1}(I \vee J) \subseteq \varphi^{-1}(I) \vee \varphi^{-1}(J)$. Let $x \in \varphi^{-1}(I \vee J)$; then $\varphi(x) \in I \vee J$, so there exist $x_1 \in I$, $x_2 \in J$ such that $\varphi(x) \leq x_1 \vee x_2$. By (*) there are $y_1, y_2 \in L$ such that $\varphi(y_1) = x_1$, $\varphi(y_2) = x_2$, and $x \leq y_1 \vee y_2$. But then $y_1 \in \varphi^{-1}(I)$, $y_2 \in \varphi^{-1}(J)$, so $x \in \varphi^{-1}(I) \vee \varphi^{-1}(J)$, as desired.

Conversely, suppose that φ^{-1} is an embedding, and let $y \in L$ and $x_1, x_2 \in K$ be such that $\varphi(y) \leq x_1 \vee x_2$. Then $(\varphi(y)) \leq (x_1] \vee (x_2]$ in $\mathcal{J}(K)$, and since φ^{-1} is join preserving we have that $y \in \varphi^{-1}((\varphi(y))) \subseteq \varphi^{-1}((x_1]) \vee \varphi^{-1}((x_2])$. Thus there exist $y_1 \in \varphi^{-1}((x_1])$, $y_2 \in \varphi^{-1}((x_2])$ such that $y \leq y_1 \vee y_2$. Clearly we may assume that $\varphi(y_1) = x_1$ and $\varphi(y_2) = x_2$.

The next three propositions allow us to repair all failures in a lattice. The constructions used in these results are slight modifications of constructions that have appeared elsewhere; that of Proposition 5 is taken from Theorem 4.4 of H. S. Gaskill, G. Grätzer, and C. R. Platt [5], and that of Propositions 6 and 7 is taken from Theorem 3.1 of T. G. Kucera and B. Sands [9]. We have included Figures 1 and 2 to illustrate the constructions in Propositions 5 and 6 respectively.

PROPOSITION 5. *Let $\mathbf{x} = \langle a, b, c, d \rangle$ be a failure of (W_l) or (W_u) in the lattice L . There exists a lattice L_x and a homomorphism φ_x of L_x onto L satisfying (*) such that \mathbf{x} is repaired in L_x by φ_x .*

REMARK. One method of repairing failures of (W_l) or (W_u) is already in the literature; namely, the “interval construction” of A. Day [3]. However, it will be crucial for the proof of our main theorem that the homomorphisms we use to repair failures satisfy (*), and it is easy to verify that the homomorphism associated with the interval construction does not enjoy this necessary property.

Proof of Proposition 5. Let \mathbf{Z} be the integers with their natural order and let E and O denote the sets of even and odd integers,

respectively. Extend Z to $Z_b = Z \cup \{-\infty, \infty\}$ where $-\infty$ is the least and ∞ the greatest element of Z_b . Setting $u = a \wedge b$ and $v = c \vee d$, we define a subset L_x of $L \times Z_b$ by

$$L_x = ([b] \times \{\infty\}) \cup ([d] \times \{-\infty\}) \cup (((v) - (d]) \times E) \\ \cup (((L - (v)) \cup [u]) - [b]) \times O).$$

(Figure 1(c) shows L_x for the case when L is the lattice of Figure 1(a) and $x = \langle a, b, c, d \rangle$. Figure 1(b) shows L_x as a subset of $L \times Z_b$.) It is not hard to verify that each element of $L \times Z_b$ that




FIGURE 1

is not of the form $\langle y, -\infty \rangle$ for $y \not\leq d$ has a least upper bound in L_x . This and a dual observation shows that L_x , with the partial order inherited from $L \times \mathbf{Z}_b$, is a lattice. Also, the projection $\pi_1: L \times \mathbf{Z}_b \rightarrow L$ restricts to a homomorphism φ_x of L_x onto L .

We first claim that φ_x satisfies (*). Let $\langle y, t \rangle \in L_x$ and $x_1, x_2 \in L$ be such that $y = \varphi_x(\langle y, t \rangle) \leq x_1 \vee x_2$. There exist $i, j \in \mathbf{Z}_b$ such that $y_1 = \langle x_1, i \rangle$ and $y_2 = \langle x_2, j \rangle$ are in L_x . If $y \leq d$ then $t = -\infty$, so $\langle y, t \rangle \leq y_1 \vee y_2$, as desired. Also, if $x_1 \vee x_2 \geq b$ then $y_1 \vee y_2 = \langle x_1 \vee x_2, \infty \rangle$, whence again $\langle y, t \rangle \leq y_1 \vee y_2$. Thus we may assume $y \not\leq d$ and $x_1 \vee x_2 \not\geq b$, and without loss of generality we have both y and x_1 in $L - ((d] \cup [b))$, implying that $t, i \in \mathbf{Z}$. Since, for any $x \in L$ and $n \in \mathbf{Z}$, $\langle x, n \rangle \in L_x$ implies $\langle x, n \pm 2 \rangle \in L_x$, we may choose $i \geq t$, and so $\langle y, t \rangle \in y_1 \vee y_2$ holds in any case, showing that φ_x satisfies (*).

Secondly, we show that x is repaired in L_x by φ_x . Let $a', b', c', d' \in L_x$ be such that $\varphi_x(a') = a$, $\varphi_x(b') = b$, $\varphi_x(c') = c$, and $\varphi_x(d') = d$. It follows that $a' = \langle a, i \rangle$ where $i \in O$, $c' = \langle c, j \rangle$ where $j \in E$, $b' = \langle b, \infty \rangle$, and $d' = \langle d, -\infty \rangle$. Thus $a' \wedge b' = \langle a \wedge b, i \rangle$ and $c' \vee d' = \langle c \vee d, j \rangle$, whence if $a' \wedge b' \leq c' \vee d'$ we have $i \leq j$. If x is failure of (W_l) , assume that $\langle a', b', c', d' \rangle$ is a failure of (W_l) in L_x ; then $c' \leq a' \wedge b'$, yielding $j \leq i$ and thus $i = j$, which is impossible since i is odd and j is even. Thus $\langle a', b', c', d' \rangle$ cannot be a failure of (W_l) . Similarly, if x is a failure of (W_u) , $\langle a', b', c', d' \rangle$ is not a failure of (W_u) in L_x . Hence x is repaired in L_x by φ_x .


PROPOSITION 6. *Let $x = \langle a, b, c, d \rangle$ be a failure of (SD_\vee) in the lattice L . There exists a lattice L_x and a homomorphism φ_x of L_x onto L satisfying (*) such that x is repaired in L_x by φ_x .*

Proof. Let \mathbf{Z}_b, E , and O be as in Proposition 5, and set $p = a \vee (b \wedge c)$. Define a subset L_x of $L \times \mathbf{Z}_b$ by

$$L_x = ((p] \times \{-\infty\}) \cup ((L - ((p] \cup (b))) \times E) \\ \cup ((L - ((p] \cup (c))) \times O).$$

Figure 2(c) shows L_x when L in the lattice of 2(a) and $x = \langle a, b, c, d \rangle$, and Figure 2(b) shows L_x as a subset of $L \times \mathbf{Z}_b$. It is easy to see that L_x is a join-semilattice, and that each element of $L \times \mathbf{Z}_b$ that is not of the form $\langle y, \infty \rangle$ has a greatest lower bound in L_x ; whence L_x , with the partial order inherited from $L \times \mathbf{Z}_b$, is a lattice. Furthermore, the projection $\pi_1: L \times \mathbf{Z}_b \rightarrow L$ again restricts to a homomorphism φ_x of L_x onto L .

To show that φ_x satisfies (*), let $\langle y, t \rangle \in L_x$ and $x_1, x_2 \in L$ be such that $y = \varphi_x(\langle y, t \rangle) \leq x_1 \vee x_2$. There exist $i, j \in \mathbf{Z}_b$ such that $y_1 = \langle x_1, i \rangle$ and $y_2 = \langle x_2, j \rangle$ are in L_x . If $y \leq p$, then $t = -\infty$, so $\langle y, t \rangle \leq$



$y_1 \vee y_2$. On the other hand, if $y \not\leq p$, then without loss of generality we may let $x_1 \not\leq p$, and both t and i are in \mathbf{Z} . We may now choose i such that $\langle x_1, i \rangle \in L_c$ and $t \leq i$, and thus $\langle y, t \rangle \leq y_1 \vee y_2$ follows in either case, proving that φ_x satisfies (*).

To show that x is repaired in L_x by φ_x , let $a', b', c', d' \in L_x$ be such that $\varphi_x(a') = a$, $\varphi_x(b') = b$, $\varphi_x(c') = c$, and $\varphi_x(d') = d$. Then $b' = \langle b, i \rangle$ where $i \in \mathbf{O}$, $c' = \langle c, j \rangle$ where $j \in \mathbf{E}$, and since $a \leq p$, $a' = \langle a, -\infty \rangle$. Thus $a' \vee b' = \langle a \vee b, i \rangle$ and $a' \vee c' = \langle a \vee c, j \rangle$. Since i is odd and j is even, $a' \vee b' \neq a' \vee c'$, so $\langle a', b', c', d' \rangle$ is not a failure of (SD_\vee) in L_x .

PROPOSITION 7. *Let $x = \langle a, b, c, d \rangle$ be a failure of (SD_\wedge) in the lattice L . There exists a lattice L_x and a homomorphism φ_x of L_x onto L satisfying (*) such that x is repaired in L_x by φ_x .*

Proof. Let $p = a \wedge (b \vee c)$, and define a subset L_x of $L \times \mathbf{Z}_b$ by

$$L_x = ([p] \times \{\infty\}) \cup ((L - ([p] \cup [b])) \times \mathbf{E}) \\ \cup ((L - ([p] \cup [c])) \times \mathbf{O}).$$

This construction is just the dual of the one in the previous proposition, so L_x is a lattice and we have the natural homomorphism φ_x of L_x onto L . An argument dual to that in Proposition 6 shows that x is repaired in L_x by φ_x , so we need only show that φ_x satisfies (*). Let $\langle y, t \rangle \in L_x$ and $x_1, x_2 \in L$ be such that $\varphi_x(\langle y, t \rangle) \leq x_1 \vee x_2$. There exist $i, j \in \mathbf{Z}_i$ such that $y_1 = \langle x_1, i \rangle$ and $y_2 = \langle x_2, j \rangle$ are in L_x . If $x_1 \vee x_2 \geq p$ then $y_1 \vee y_2 = \langle x_1, i \rangle \vee \langle x_2, j \rangle = \langle x_1 \vee x_2, \infty \rangle \geq \langle y, t \rangle$; therefore, we assume $x_1 \vee x_2 \not\geq p$, which implies that $t, i, j \in \mathbf{Z}$. Now we can choose $i \in \mathbf{Z}$ such that $\langle x_1, i \rangle \in L_x$ and $i \geq t$, whence $\langle y, t \rangle \leq y_1 \vee y_2$ follows.

Before continuing with the proof of the theorem, we recall the following construction.

DEFINITION 8. Let $(L_i | i \in I)$ be a family of lattices, let L be a lattice, and let $\varphi_i: L_i \rightarrow L$ be a lattice homomorphism for each $i \in I$. Form the direct product $\prod(L_i | i \in I)$, and consider the subset

$$K = \{x \in \prod L_i \mid \varphi_i(x(i)) = \varphi_j(x(j)) \text{ for all } i, j \in I\}.$$

Then K is a sublattice of $\prod L_i$, and is called the *pullback* of the family $(\varphi_i | i \in I)$. Letting $\pi_i: K \rightarrow L_i$ be the restriction of the projection of $\prod L_i$ onto L_i , we have $\varphi_i \circ \pi_i = \varphi_j \circ \pi_j$ for all $i, j \in I$; hence there is a natural homomorphism $\varphi = \varphi_i \circ \pi_i$ of K into L . If φ_i is onto for all $i \in I$, then φ is onto.

PROPOSITION 9. *For any lattice L , there exists a lattice L^* and a homomorphism φ^* of L^* onto L satisfying (*) that repairs all failures in L .*

Proof. Let $\mathcal{F}(L)$ be the set of all failures in L . From Propositions 5, 6, and 7, we obtain a family $(L_x \mid x \in \mathcal{F}(L))$ of lattices and a family $(\varphi_x: L_x \rightarrow L \mid x \in \mathcal{F}(L))$ of onto homomorphisms satisfying (*) such that for each $x \in \mathcal{F}(L)$, x is repaired in L_x by φ_x . Let L^* be the pullback of $\{\varphi_x \mid x \in \mathcal{F}(L)\}$ and let φ^* be the natural homomorphism of L^* onto L . Then by Lemma 3, φ^* repairs all failures in L . To show that φ^* satisfies (*), let $p \in L^*$ and $u, v \in L$ be such that $\varphi^*(p) \leq u \vee v$. Letting p_x denote the x th component of p , for each $x \in \mathcal{F}(L)$, we have that $\varphi_x(p_x) = \varphi^*(p) \leq u \vee v$. For each $x \in \mathcal{F}(L)$, since φ_x satisfies (*), there exist $u_x, v_x \in L_x$ such that $\varphi_x(u_x) = u$, $\varphi_x(v_x) = v$, and $p_x \leq u_x \vee v_x$ in L_x . Then the elements $u = (u_x \mid x \in \mathcal{F}(L))$ and $v = (v_x \mid x \in \mathcal{F}(L))$ are clearly in L^* ; moreover $\varphi^*(u) = u$, $\varphi^*(v) = v$, and $p \leq u \vee v$.

Finally, we are in a position to prove our main result.

THEOREM 10. *Every lattice can be embedded in the ideal lattice of a lattice satisfying (SD_\vee) , (SD_\wedge) , (W_l) , and (W_u) .*

Proof. Let L be a lattice. Set $L_0 = L$, and inductively let $L_{n+1} = L_n^*$ and $\varphi_n^*: L_n^* \rightarrow L_n$, $n \geq 0$, be the lattice and homomorphism of Proposition 9. Let L_∞ be the inverse limit of the system of lattices $(L_n \mid n < \omega)$ and homomorphisms $(\varphi_n^* \mid n < \omega)$, and let $\varphi_n: L_\infty \rightarrow L_n$ be the natural projection for each n .

We first claim that L_∞ satisfies (SD_\vee) , (SD_\wedge) , (W_l) , and (W_u) . Again note that each of these four conditions is expressible in the form $P(x, y, z, w) \Rightarrow Q(x, y, z, w)$, where P and Q are disjunctions of polynomial equations. Let $\langle a, b, c, d \rangle$ be a failure in L_∞ ; then there exist appropriate P and Q such that $P(a, b, c, d)$ holds but $Q(a, b, c, d)$ fails. It follows that $P(\varphi_n(a), \varphi_n(b), \varphi_n(c), \varphi_n(d))$ holds for all $n \in \omega$, and, since L_∞ is a sublattice of $\prod(L_n \mid n \in \omega)$, $Q(\varphi_m(a), \varphi_m(b), \varphi_m(c), \varphi_m(d))$ fails for some $m \in \omega$. Therefore $\langle \varphi_m(a), \varphi_m(b), \varphi_m(c), \varphi_m(d) \rangle$ is a failure in L_m . But by construction $\langle \varphi_{m+1}(a), \varphi_{m+1}(b), \varphi_{m+1}(c), \varphi_{m+1}(d) \rangle$ is not a failure in L_{m+1} , which contradicts Lemma 3. Thus there can be no failures in L_∞ ; that is, L_∞ satisfies (SD_\vee) , (SD_\wedge) , (W_l) , and (W_u) .

Next we prove that the homomorphism φ_0 of L_∞ onto L satisfies (*). Let $x \in L_\infty$ and $u_0, v_0 \in L = L_0$ be such that $\varphi_0(x) \leq u_0 \vee v_0$. Then $\varphi_0(x) = \varphi_0^*(\varphi_1(x)) \leq u_0 \vee v_0$, and since φ_0^* satisfies (*) there exist $u_1, v_1 \in L_1$ such that $\varphi_0^*(u_1) = u_0$, $\varphi_0^*(v_1) = v_0$, and $\varphi_1(x) \leq u_1 \vee v_1$. Proceeding by induction, assume that we have $u_n, v_n \in L_n$ such that $\varphi_n(x) \leq u_n \vee v_n$. Then $\varphi_n(x) = \varphi_n^*(\varphi_{n+1}(x)) \leq u_n \vee v_n$, and since φ_n^* satisfies (*) there exist $u_{n+1}, v_{n+1} \in L_{n+1}$ such that $\varphi_n^*(u_{n+1}) = u_n$, $\varphi_n^*(v_{n+1}) = v_n$, and $\varphi_{n+1}(x) \leq u_{n+1} \vee v_{n+1}$. Now let $u = \langle u_n \mid n < \omega \rangle$ and $v = \langle v_n \mid n < \omega \rangle$. It follows that $u, v \in L_\infty$, $\varphi_0(u) = u$, $\varphi_0(v) = v$, and $x \leq u \vee v$, whence φ_0 satisfies (*).

From Lemma 4, $\mathcal{J}(L)$ is embedded in $\mathcal{J}(L_\infty)$. Since L is embedded in $\mathcal{J}(L)$, the theorem is proved.

As mentioned earlier, we have the following corollary.

COROLLARY 11. *Every transferable lattice satisfies (W) .*

REMARK. The use of homomorphisms, pullbacks, and inverse limits to repair failures stems from a proof in a recent paper of A. Day, namely, the proof (see Theorem 3.2 in [4]) that every lattice is a bounded homomorphic image of a lattice satisfying (W) .

3. Additional results. In this section we investigate the status of $\mathcal{E}(P)$ for most other combinations (P) of the properties defined

in the introduction. First, we shall indicate how certain techniques in a paper of G. Grätzer and C. R. Platt [8] can be modified so as to prove that $\mathcal{E}((SD_{\vee}) \& (SF) \& (X))$ holds.

Let L be a lattice. It has already been observed that there is a lattice K satisfying (SF) such that L is embeddable in $\mathcal{J}(K)$. Hence we need only show that for every lattice K satisfying (SF) there is a lattice M satisfying (SD_{\vee}) , (SF) , and (X) such that $\mathcal{J}(K)$ is embeddable in $\mathcal{J}(M)$.

Let K be a lattice satisfying (SF) . In [8], Grätzer and Platt construct a lattice $L(K_I)$ satisfying (SD_{\vee}) such that K can be embedded in $\mathcal{J}(L(K_I))$. From Lemma 3 and their proof it is clear that they in fact embed $\mathcal{J}(K)$ in $\mathcal{J}(L(K_I))$. The lattice $L(K_I)$ consists of certain subsets (called *closed* subsets) of $K \times \mathbf{Z}$, ordered by inclusion.

Now we replace \mathbf{Z} by ω , and consider the set $L_f(K_I)$ of all *finitely generated* closed subsets of $K \times \omega$, that is, all closed subsets which are closures of finite subsets of $K \times \omega$. Since K satisfies (SF) , each element of $L_f(K_I)$ is finite. Hence $L_f(K_I)$, ordered by inclusion, is a lattice; in fact $L_f(K_I)$ is embeddable in $L(K_I)$ and therefore satisfies (SD_{\vee}) . Furthermore, $L_f(K_I)$ is sectionally finite. Next, it can be proved as in [8] that $\mathcal{J}(K)$ is embeddable in $\mathcal{J}(L_f(K_I))$, and moreover the image under this embedding of each ideal in K is a nonprincipal ideal of $L_f(K_I)$. Therefore (G. Grätzer [7]) the elements of $L_f(K_I)$ may all be “split” to yield a lattice M satisfying (X) such that $\mathcal{J}(K)$ is embeddable in $\mathcal{J}(M)$. It is easy to see that M will still satisfy (SD_{\vee}) and (SF) . Thus we have:

THEOREM 12. $\mathcal{E}((SD_{\vee}) \& (SF) \& (X))$ holds.

In contrast to the above, we now establish two negative results.

LEMMA 13. *If a lattice L satisfies (SF) and (SD_{\wedge}) , then $\mathcal{J}(L)$ satisfies (SD_{\wedge}) .*

Proof. Let L satisfy (SF) and (SD_{\wedge}) , and let $A, B, C \in \mathcal{J}(L)$ satisfy $A \cap B = A \cap C$. Let $p \in A \cap (B \vee C)$. There exist $b \in B$ and $c \in C$ such that $p \leq b \vee c$. By (SF) , there exist largest elements $b_0 \in B$ and $c_0 \in C$ such that $b_0, c_0 \leq b \vee c$. Since $p \in A$, $p \wedge b_0 \in A \cap B$ and $p \wedge c_0 \in A \cap C = A \cap B$. Thus the element $q = (p \wedge b_0) \vee (p \wedge c_0) \in A \cap B$, and by the choice of b_0 , $p \wedge c_0 \leq q \leq b_0$. Hence $p \wedge c_0 \leq p \wedge b_0$; by symmetry we have that $p \wedge b_0 = p \wedge c_0$. Since L satisfies (SD_{\wedge}) , $p \wedge b_0 = p \wedge (b_0 \vee c_0) = p \wedge (b \vee c) = p$. We conclude that $p \in A \cap B$, and so $A \cap (B \vee C) = A \cap B$, showing that $\mathcal{J}(L)$ satisfies (SD_{\wedge}) .

COROLLARY 14. $\mathcal{E}((SF) \ \& \ (SD_{\wedge}))$ fails.

LEMMA 15. If a lattice L satisfies (SF) and (W_u) , then $\mathcal{J}(L)$ satisfies (W_u) .

Proof. Let L satisfy (SF) and (W_u) , and suppose that $\langle A, B, C, D \rangle$ is a (W_u) -failure in $\mathcal{J}(L)$. Then there exists an element $x \in A \cap B$ such that $x \notin C$ and $x \notin D$, and an element $b \in B$ such that $b \geq x$ and $b \notin C \vee D$. Since L satisfies (SF) , there exists a largest element $x_0 \in A \cap B$ such that $x_0 \leq b$; note that $x_0 \geq x$ and so $x_0 \notin C$, $x_0 \notin D$. Since $x_0 \in A \cap B \subseteq C \vee D$, there exist $c \in C$, $d \in D$ such that $x_0 \leq c \vee d$. However, $b \notin C \vee D$, so $b \not\leq c \vee d$. Finally, since $A \supset C \vee D$, we may choose $a \in A$ such that $a > c \vee d$. But now $a \wedge b \geq (c \vee d) \wedge b \geq x_0$, and by the maximality of x_0 we obtain that $a \wedge b = x_0$. Hence the quadruple $\langle a, b, c, d \rangle$ is a failure of (W_u) in L , contradicting the hypothesis. We conclude that $\mathcal{J}(L)$ satisfies (W_u) .

COROLLARY 16. $\mathcal{E}((SF) \ \& \ (W_u))$ fails.

To end this paper we ask two questions that are still open:

- (i) Does $\mathcal{E}((SF) \ \& \ (W_l))$ hold?
- (ii) Does $\mathcal{E}((SF) \ \& \ (W_l) \ \& \ (SD_{\vee}))$ hold?

REFERENCES

1. R. Antonius and I. Rival, *A note on Whitman's property for free lattices*, Algebra Universalis, **4** (1974), 271-272.
2. K. A. Baker and A. W. Hales, *From a lattice to its ideal lattice*, Algebra Universalis, **4** (1974), 250-258.
3. A. Day, *A simple solution to the word problem for lattices*, Canad. Math. Bull., **13** (1970), 253-254.
4. ———, *Splitting lattices generate all lattices*, Algebra Universalis, **7** (1977), 163-169.
5. H. S. Gaskill, G. Grätzer, and C. R. Platt, *Sharply transferable lattices*, Canad. J. Math., **27** (1975), 1247-1262.
6. G. Grätzer, *Universal Algebra in Trends in Lattice Theory*, 173-215. New York, Van Nostrand Reinhold, 1970.
7. ———, *A property of transferable lattices*, Proc. Amer. Math. Soc., **43** (1974), 269-271.
8. G. Grätzer and C. R. Platt, *Two embedding theorems for lattices*, Proc. Amer. Math. Soc., **69** (1978), 21-24.
9. T. G. Kucera and B. Sands, *Lattices of lattice homomorphisms*, Algebra Universalis, **8** (1978), 180-190.
10. P. M. Whitman, *Lattices, equivalence relations, and subgroups*. Bull. Amer. Math. Soc., **52** (1946), 507-522.

Received May 3, 1978 and in revised form December 29, 1978. The work of all three authors was supported by the National Research Council of Canada.

UNIVERSITY OF MANITOBA
WINNIPEG, MANITOBA, CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Ralph Alexander, <i>Metric averaging in Euclidean and Hilbert spaces</i>	1
B. Aupetit, <i>Une généralisation du théorème de Gleason-Kahane-Żelazko pour les algèbres de Banach</i>	11
Lung O. Chung, Jiang Luh and Anthony N. Richoux, <i>Derivations and commutativity of rings. II</i>	19
Lynn Harry Erbe, <i>Integral comparison theorems for third order linear differential equations</i>	35
Robert William Gilmer, Jr. and Raymond Heitmann, <i>The group of units of a commutative semigroup ring</i>	49
George Grätzer, Craig Robert Platt and George William Sands, <i>Embedding lattices into lattices of ideals</i>	65
Raymond D. Holmes and Anthony Charles Thompson, <i>n-dimensional area and content in Minkowski spaces</i>	77
Harvey Bayard Keynes and M. Sears, <i>Modelling expansion in real flows</i>	111
Taw Pin Lim, <i>Some classes of rings with involution satisfying the standard polynomial of degree 4</i>	125
Garr S. Lystad and Albert Robert Stralka, <i>Semilattices having bialgebraic congruence lattices</i>	131
Theodore Mitchell, <i>Invariant means and analytic actions</i>	145
Daniel M. Oberlin, <i>Translation-invariant operators of weak type</i>	155
Raymond Moos Redheffer and Wolfgang V. Walter, <i>Inequalities involving derivatives</i>	165
Eric Schechter, <i>Stability conditions for nonlinear products and semigroups</i>	179
Jan Sørensen, <i>Symmetric shift registers</i>	201
Toshiji Terada, <i>On spaces whose Stone-Čech compactification is O_Z</i>	231
Richard Vrem, <i>Harmonic analysis on compact hypergroups</i>	239