

# Pacific Journal of Mathematics

**FIBER HOMOLOGY AND ORIENTABILITY OF MAPS**

J. WOLFGANG SMITH

# FIBER HOMOLOGY AND ORIENTABILITY OF MAPS

J. WOLFGANG SMITH

In this paper we introduce a concept of *fiber homology* for an arbitrary map  $f: X \rightarrow Y$  and coefficient module  $G$ . This is a graded module denoted by  $H_*(f_*; G)$  which reduces to  $H_*(F; G)$  when  $f$  represents an orientable fiber bundle with standard fiber  $F$ . The concept of fiber homology permits us also to define a generalized notion of orientability, and these ideas turn out to be useful in the study of submersions. Our main theorem (obtained by means of a spectral sequence) asserts that if the fibers of a submersion  $f: X \rightarrow Y$  are acyclic in dimensions smaller than  $q$ , then the rank  $r_q$  of the fiber homology  $H_q(f_*; G)$  is bounded above by the sum of the  $q$  and  $(q+1)$ -dimensional Betti numbers of  $X$  and  $Y$ , respectively. In the orientable case, the  $q$ -dimensional Betti number of an arbitrary fiber  $f^{-1}(y)$  is bounded above by  $r_q$ , and therefore also by the aforementioned sum. This leads to a number of more specialized results. For example, it is shown that the fibers of an orientable submersion  $f: R^{2m-1} \rightarrow S^m$  must be either acyclic or homology spheres, and moreover, the subspace of points in  $S^m$  corresponding to the spherical fibers must have the homology of a point.

1. Basic concepts. Let  $f: X \rightarrow Y$  denote a continuous map between topological spaces. By a *tubular neighborhood* belonging to  $f$  we will understand a homeomorphism

$$\Phi: B \times F \approx V$$

where  $B$  is an open connected subspace of  $Y$ ,  $F$  a compact space and  $V$  a subset of  $X$ , such that  $f \circ \Phi$  is the projection  $B \times F \rightarrow B$ . Given a point  $y \in B$ , we will write

$$F_y = V \cap f_y$$

where  $f_y$  denotes the preimage of  $y$  under  $f$ , and given two points  $y, y' \in B$ , we let

$$\Phi_y^{y'}: F_y \approx F_{y'},$$

denote the homeomorphism induced by  $\Phi$ . The diagram

$$\begin{array}{ccc} H_*(f_y; G) & & H_*(f_{y'}; G) \\ \uparrow i & & \uparrow i' \\ H_*(F_y; G) & \xrightarrow{\Phi_y^{y'}} & H_*(F_{y'}; G) \end{array}$$

(where  $i, i'$  are inclusion induced,  $G$  is a coefficient module and  $H_*$  designates singular homology) gives rise to a linear relation

$$i' \circ \Phi_{y_*}^{y'} \circ i^{-1}: H_*(f_y; G) \longrightarrow H_*(f_{y'}; G)$$

henceforth referred to as a *horizontal relation* induced by  $\Phi$ . We will employ the notation

$$(1.1) \quad w \not\sim w' \bmod \Phi$$

to indicate that the homology classes  $w, w'$  are horizontally related via  $\Phi$ . Now let  $W$  denote the linear subspace of the direct sum

$$(1.2) \quad \sum_{i \in Y} H_*(f_i; G)$$

generated by elements of the form  $[w - w']$  as  $(w, w')$  ranges over all pairs satisfying a relation (1.1). This makes sense, inasmuch as every element  $w \in H_*(f_y; G)$  can be identified with a corresponding element in (1.2), an identification which we shall always assume. The *fiber homology* of  $f$  (with coefficients in  $G$ ) is defined to be the quotient module

$$H_*(f_*; G) = \sum_{i \in Y} H_*(f_i; G) / W.$$

It is to be noted that  $H_*(f_*; G)$  reduces to  $H_*(F; G)$  when  $f$  represents an orientable fiber bundle (see [4]) with standard fiber  $F$ . The following simple and illustrative examples will exhibit the fiber homology in a more general case.

**EXAMPLE 1.** Let  $X \subset R^3$  denote the open subspace consisting of points  $(x_1, x_2, x_3)$  with  $|x_3| < 1$ , excluding the origin, and let  $f: X \rightarrow R$  be given by

$$f(x) = \log \|x\|.$$

This defines a differentiable submersion from a punctured 3-space to the real line whose fibers  $f_y$  are cylinders for  $y \geq 0$  and spheres for  $y < 0$ . It is easy enough to verify that the fiber homology is given by

$$H_p(f_*; G) \approx \begin{cases} G & p = 0, 2 \\ 0 & \text{otherwise.} \end{cases}$$

In particular, to see that the fiber homology vanishes in dimension 1, consider two points  $y, y' \in R$  with  $y' < 0 < y$ . The cylindrical fiber  $f_y$  intersects the plane  $x_3 = 0$  in a circle  $C_y$ , which (when oriented) represents a generator  $w \in H_1(f_y; G)$ . If we now subject

$C_y$  to a contraction (via concentric circles) in the given plane, we obtain a corresponding family of homology classes  $w_i \in H_1(f_i; G)$  as  $t$  varies from  $y$  to  $y'$ . At least for values  $t, t'$  which are not too far apart, the corresponding homology classes will clearly satisfy a horizontal relation (1.1). But by virtue of the fact that  $f_y$  is simply connected, this implies that  $w$  is annihilated by projection into  $H_1(f_*; G)$ .

EXAMPLE 2. Modifying the preceding example by deleting the positive  $x_3$ -axis, one obtains a new submersion  $f: R^3 \rightarrow R$  (after a suitable change of coordinates) whose fibers are cylinders for  $y \geq 0$  and planes for  $y < 0$ . This map is "fiber acyclic".

Returning to the case of an arbitrary map  $f: X \rightarrow Y$ , it is to be noted that for every point  $y \in Y$  one has a natural homomorphism

$$(1.3) \quad H_*(f_y; G) \longrightarrow H_*(f_*; G)$$

(canonical injection into (1.2), followed by natural projection), and we will say that  $f$  is *orientable at  $y$*  (over  $G$ ) if this homomorphism is injective. A map is said to be *orientable* if it is orientable at all points  $y \in Y$ , and it will also make sense to say that  $f$  is orientable in a particular dimension. Moreover, one observes that these definitions reduce to the classical conception of orientability in the case of fiber bundles.

A novel feature of this generalized notion of orientability lies in the fact that a map  $f: X \rightarrow Y$  may fail to be orientable over arbitrarily small neighborhoods of a given point  $y \in Y$ , a circumstance which is exemplified by both of the preceding examples. Not only do these submersions fail to be orientable, but this failure is concentrated (so to speak) in the immediate vicinity of a particular fiber (the fiber  $f_0$ , which constitutes the smallest or innermost cylinder in the given foliation of  $X$ ). Fibers of this kind will be referred to as "separatrices", and one may define this concept as follows. Let us say that a homology class  $w \in H_*(f_y; G)$  is *horizontally annihilated* over some neighborhood  $U$  of  $y$ , provided there exists a tubular neighborhood  $\Phi$  with  $B \subset U$  such that  $w \neq 0' \bmod \Phi$ , where  $0'$  denotes the zero element in  $H_*(f_{y'}; G)$  for some  $y' \in B$ . A fiber  $f_y$  will be called a *separatrix* (over  $G$ ) if it admits a nonzero homology class  $w \in H_*(f_y; G)$  which is horizontally annihilated over every neighborhood  $U$  of  $y$ .

In the present paper we shall restrict our attention to the class of submersions. Little will be lost if we presuppose a  $C^\infty$  setting and take "submersion" to mean a differentiable surjection  $f: X \rightarrow Y$  between paracompact differentiable manifolds with  $\dim X \geq \dim Y$ ,

such that the differential  $df$  has everywhere maximal rank. On the other hand, it should be pointed out that the proof of our Main Theorem does not require as much. The crucial property which is required for many steps can be formulated directly in terms of tubular neighborhoods. Given a map  $f: X \rightarrow Y$ , a subspace  $C \subset X$  and a tubular neighborhood  $\Phi: B \times F \rightarrow V$  belonging to  $f$ , we will say that  $\Phi$  *cuts*  $C$  provided that

$$C \cap f_y \subset F_y$$

for all  $y \in B$ . The property in question is this:

(1.4) *Given  $y \in Y$  and a compact subset  $C \subset X$ , there shall exist a tubular neighborhood  $\Phi$  such that  $y \in B$  and  $\Phi$  cuts  $C$ .*

The proof of our Main Theorem also utilizes a spectral sequence theorem obtained in [1], the proof of which actually requires, in addition to (1.4), that  $X$  is second countable and locally compact, and that  $Y$  is triangulable. Although these conditions on the map  $f: X \rightarrow Y$  suffice for the Main Theorem, some of the more specialized results which we shall present as corollaries presuppose also that the fibers  $f_y$  shall be manifolds. All these conditions are satisfied in the case of a differentiable submersion.

**2. Main theorem and corollaries.** We will now state our main result and develop a few of its implications. The notation  $R_p$  shall indicate Betti numbers, and in particular,  $R_p(f_*; G)$  shall denote the rank of the  $p$ -dimensional fiber homology.

**MAIN THEOREM.** *If  $f: X \rightarrow Y$  is a submersion and  $G$  a coefficient module, then*

$$R_0(f_*; G) = R_0(X; G).$$

*Moreover, if every fiber  $f_y$  is connected and  $t$ -acyclic over  $G$  for  $q - m < t < q$ , where  $m = \dim Y$ , then*

$$R_q(f_*; G) \leq R_q(X; G) + R_{q+1}(Y; G).$$

*If, in addition,  $H_q(X; G)$  and  $H_{q+1}(Y; G)$  vanish, then  $H_q(f_*; G)$  vanishes.*

In the orientable case one can say a good deal more since one has also an inequality

$$R_*(f_y; G) \leq R_*(f_*; G)$$

which holds for every point  $y \in Y$ . By a successive application of our theorem one can therefore obtain

**COROLLARY 1.** *Let  $f: X \rightarrow Y$  be a submersion which is orientable over  $G$  in dimensions  $t \leq p$ . Let  $X$  and  $Y$  be  $t$ -acyclic over  $G$  for  $t \leq p$  and  $t \leq p + 1$ , respectively. Then  $f_y$  is  $t$ -acyclic over  $G$  for all  $t \leq p$  and every point  $y \in Y$ .*

It will be shown in § 6 that a real-valued submersion  $f: X \rightarrow R$  without separatrices is orientable. This fact, together with Corollary 1, implies

**COROLLARY 2.** *Let  $f: X \rightarrow R$  be a submersion, and let  $X$  be  $t$ -acyclic over  $G$  for  $t \leq p$ . If for some  $t \leq p$  and some point  $y \in Y$  the fiber  $f_y$  is not  $t$ -acyclic over  $G$ , then  $f$  admits a separatrix over  $G$  in some dimension  $s \leq t$ .*

This result shows, in particular, that the separatrices which have been exhibited in our Examples 1 and 2 were necessitated by the existence of cylindrical fibers and the topology of  $X$ .

It is easy to see that a compact fiber cannot be a separatrix, since such a fiber can be imbedded in a tubular neighborhood (in consequence of (1.4)). This observation leads immediately to

**COROLLARY 3.** *Let  $f: X \rightarrow R$  be a submersion, and let  $X$  be  $t$ -acyclic over  $G$  for  $t < p$ . If every noncompact fiber is  $t$ -acyclic over  $G$  for  $t \leq p$ , then*

$$R_p(f_y; G) \leq R_p(X; G)$$

for all  $y \in Y$ .

Our assumptions imply that there are no separatrices in dimensions  $t \leq p$ , and this implies orientability in the same range. One may therefore conclude by Corollary 1 that the fibers are acyclic in dimensions  $t < p$ . The Main Theorem now gives  $R_p(f_*; G) \leq R_p(X; G)$ , and since  $f$  is also orientable in dimension  $p$ , one obtains the inequality in question.

Apart from real-valued submersions, another particularly simple class is given by submersions of codimension 1, which is to say, submersions having 1-dimensional fibers. Here we can mention the following result:

**COROLLARY 4.** *Let  $f: X \rightarrow Y$  be a submersion of codimension 1 having connected fibers. If  $X$  and  $Y$  are acyclic over  $Z_2$  in dimensions 1 and 2, respectively, then  $f$  admits no compact fiber.*

It should be noted that if the fibers of a submersion of codi-

mension 1 are assumed to be connected, then these fibers can only be circles or lines. Under the given acyclicity conditions on  $X$  and  $Y$ , our corollary concludes that there can be no circular fibers, and this actually implies that  $f$  is equivalent to the natural projection  $Y \times R \rightarrow Y$ . For a detailed study of the codimension 1 case, we may refer to [2].

To establish Corollary 4, we first apply our Main Theorem with  $q = 1$  to conclude that  $R_1(f_*; Z_2) = 0$ . It therefore suffices to show that  $f$  is orientable in dimension 1. The easiest way to see this, perhaps, is to consider the subset  $U \subset Y$  corresponding to circular fibers and observe that the restriction  $g$  of  $f$  to the preimage of  $U$  represents a circle bundle, and this is certainly orientable over  $Z_2$ . Now  $f$  will be orientable if for any  $y \in U$ , the homomorphism (1.3) turns out to be injective. But this homomorphism clearly factors as follows:

$$H_*(f_y; Z_2) \xrightarrow{\phi} H_*(g_*; Z_2) \xrightarrow{\psi} H_*(f_*; Z_2)$$

where  $\phi$  and  $\psi$  are the obvious homomorphisms. Since  $g$  is orientable over  $Z_2$ ,  $\phi$  will be injective, and it is also evident that  $\psi$  will be an isomorphism. The composition is therefore injective.

It may be of interest to point out in connection with Corollary 4 that a submersion  $f: R^3 \rightarrow S^2$  with connected fibers can be obtained from the Hopf fibration  $S^3 \rightarrow S^2$  by deleting a point in  $S^3$ . Moreover, we have shown in [2] that every submersion  $f: X \rightarrow S^2$  of codimension 1 with connected fibers arises from this Hopf fibration by deleting a subspace  $W \subset S^3$ .

In the remaining corollaries we will be concerned with submersions  $f: R^n \rightarrow S^m$  which are orientable over some coefficient module  $G$ . It should also be noted that these results depend on the fact that the fibers are manifolds.

**COROLLARY 5.** *There does not exist an orientable submersion  $f: R^n \rightarrow S^m$  for  $n < 2m - 1$ .*

For such a submersion one could conclude by Corollary 1 that all fibers are  $t$ -acyclic over  $G$  for  $t < m - 1$ . But since the fibers are  $(n - m)$ -manifolds and  $n - m < m - 1$ , it follows that the fibers are  $t$ -acyclic over  $G$  for all  $t$ . By the Vietoris-Begle theorem for submersions [3], this would imply that  $S^m$  is acyclic.

**COROLLARY 6.** *For an orientable submersion  $f: R^{2m-1} \rightarrow S^m$  every compact fiber must be a homology sphere and every noncompact fiber must be acyclic. Moreover, the subspace  $U \subset S^m$  corresponding to compact fibers must have the homology of a point.*

One sees as before that every fiber is  $t$ -acyclic for  $t < m - 1$ . For the noncompact fibers this implies total acyclicity. The compact fibers will be homology  $(m - 1)$ -spheres, provided they are orientable (in the sense of manifolds). This is easily established in the differentiable case if we consider the tangent bundle  $\tau$  and normal bundle  $\nu$  to  $f_y$ . Clearly  $\nu$  is a trivial bundle (since it maps to the tangent vector space  $S_y^m$  under  $df$ ) and is therefore orientable. On the other hand,

$$\tau \oplus \nu \approx \bar{\tau} | f_y$$

where  $\bar{\tau}$  denotes the tangent bundle to  $R^{2m-1}$ . Since  $\nu$  and  $\bar{\tau}$  are both orientable,  $\tau$  must be orientable, and this implies that  $f_y$  is also orientable. The conclusion about  $U$  follows now from a generalized Thom-Gysin sequence (Theorem 4.9 of [1]).

We have previously observed that submersions satisfying the hypothesis of Corollary 6 can be obtained in the case  $m = 2$  from the Hopf fibration  $S^3 \rightarrow S^2$ , and it is clear that one can do likewise for  $m = 4$  and  $8$ . It is an open question whether orientable submersions  $f: R^{2m-1} \rightarrow S^m$  exist for  $m \neq 1, 2, 4$ , and  $8$ .

**COROLLARY 7.** *Let  $f: R^n \rightarrow S^m$  be an orientable submersion with  $2m - 1 \leq n \leq 3(m - 1)$ . Then every compact fiber must be a homology sphere.*

As before, every fiber must be  $t$ -acyclic for  $t < m - 1$ , and every compact fiber  $f_y$  must be orientable. By Poincaré duality applied to  $f_y$ ,  $t$ -acyclicity for  $0 < t < m - 1$  implies  $t$ -acyclicity for  $n - 2m + 1 < t < n - m$ . But the stipulated relation between  $n$  and  $m$  implies that the union of the two intervals is precisely the interval  $0 < t < n - m$ . Using once more the fact that  $f_y$  is orientable (together with  $0$ -acyclicity), one also has  $H_{n-m}(f_y; G) \approx G$ , as was to be shown.

**3. Homology of submersions.** With every submersion  $f: X \rightarrow Y$  and coefficient module  $G$  one can associate certain homology groups  $H_{s,t}(f; G)$  which arise quite naturally and can be studied on an axiomatic basis. The importance of these groups in the context of the present paper stems from the following theorem, which is the main result established in [1].

**THEOREM.** *Let  $f: X \rightarrow Y$  be a submersion and  $G$  a coefficient module. There exists a convergent  $E^2$  spectral sequence with*

$$E_{s,t}^2 \approx H_{s,t}(f; G)$$



and  $E^\infty$  isomorphic to the bigraded group associated with a filtration of  $H_*(X, G)$ .

It is to be noted that the given spectral sequence reduces to the Serre sequence in the case of a fibration.

All that we shall need to know about the homology groups  $H_{s,t}(f; G)$  is that they can be computed by means of a certain algorithm, which is fully described in [1]. For the convenience of the reader we will now give a brief summary of this algorithm.

Firstly, one requires the notion of a *simplicial bundle* over a simplicial complex  $K$ , which may be defined as a map  $p: E \rightarrow |K|$ , together with a function which to every simplex  $\sigma \in K$  assigns a homeomorphism  ${}_s\Phi: |\sigma| \times F_\sigma \approx E_\sigma$ , where  $F_\sigma$  is compact and  ${}_s\Phi$  is the restriction of a tubular neighborhood belonging to  $p$ , such that

$$(B1) \quad E = \bigcup_{\sigma \in K} E_\sigma$$

$$(B2) \quad {}_\tau\Phi \text{ cuts } E_\sigma \text{ for every face } \tau < \sigma.$$

With every simplicial bundle  $p: E \rightarrow |K|$  and coefficient module  $G$  we will associate a chain complex  $C_*(K; H_t(F; G))$  in which the homology groups  $\{H_t(F_\sigma; G): \sigma \in K\}$  function as a local coefficient system. The  $s$ -dimensional chain group ( $s \geq 0$ ) is given by

$$(3.1) \quad C_s(K; H_t(F; G)) = \sum_{\sigma \in K^{(s)}} H_s(|\sigma|, |\dot{\sigma}|) \otimes H_t(F_\sigma; G)$$

where the right side represents a direct sum,  $K^{(s)}$  denotes the set of  $s$ -simplexes in  $K$  and  $|\dot{\sigma}| \subset |\sigma|$  the boundary. The chain boundary  $\partial$  is defined on generators  $a \otimes c$  through a formula

$$(3.2) \quad \partial(a \otimes c) = \sum_{\tau \in K^{(s-1)}} \varepsilon_\sigma^-(a) \otimes i_\tau^-(c)$$

where

$$\varepsilon_\sigma^-: H_s(|\sigma|, |\dot{\sigma}|) \longrightarrow H_{s-1}(|\tau|, |\dot{\tau}|)$$

are incidence homomorphisms and

$$(3.3) \quad i_\sigma^-: H_t(F_\sigma; G) \longrightarrow H_t(F_\tau; G)$$

are so-called *fiber projections*, obtained as follows. For each  $\sigma \in K$  we choose a point  $\hat{\sigma} \in |\sigma|$  and identify  $F_\sigma$  with  $F_{\hat{\sigma}}$  via  ${}_s\Phi$ . For  $\tau < \sigma$ , one has  $\hat{\tau} \in |\sigma|$ , and by virtue of (B2) the homeomorphism  ${}_s\Phi_{\hat{\sigma}}$  defines an injection  $F_\sigma \rightarrow F_{\hat{\tau}}$  (whose homotopy class does not depend upon  $\hat{\tau}$  and  $\hat{\sigma}$ ) which induces the fiber projection (3.3). This completes our definition of the chain complex associated with a simplicial bundle.

Now let  $f: X \rightarrow Y$  be a map. A simplicial bundle  $p: E \rightarrow |K|$  is said to *approximate*  $f$  if  $E \subset X$  and  $p$  is the restriction of  $f$  to  $E$ . A sequence  $\{\alpha p: \alpha E \rightarrow |\alpha K|\}$  of simplicial bundles is called an *approximating sequence* for  $f$  if

- (A1) each  $\alpha p$  approximates  $f$  and every compact subset of  $X$  is contained in some  $\alpha E$ ;
- (A2) for  $\alpha < \beta$ ,  $\alpha E \subset \beta E$  and some subdivision of  $\alpha K$  is a subcomplex of  $\beta K$ .

It has been shown in [1] that every submersion admits an approximating sequence.

Given an approximating sequence, one obtains chain projections

$$(3.4) \quad \phi_{\alpha}^{\beta}: C_s(\alpha K; H_t(\alpha F; G)) \longrightarrow C_s(\beta K; H_t(\beta F; G))$$

for  $\alpha < \beta$ . These are defined on generators  $a \otimes c$  by a formula of the form

$$(3.5) \quad \phi_{\alpha}^{\beta}(a \otimes c) = \sum_{\omega \in \beta K(s)} B_{\sigma, \omega}(a) \otimes j_{\sigma}^{\omega}(c),$$

where

$$B_{\sigma, \omega}: H_s(|\sigma|, |\dot{\sigma}|) \longrightarrow H_s(|\omega|, |\dot{\omega}|)$$

represents a subdivision operator for  $|\omega| \subset |\sigma|$  and is zero otherwise, and  $j_{\sigma}^{\omega}: \alpha F_{\sigma} \rightarrow \beta F_{\omega}$  denotes the composition

$$\alpha F_{\sigma} \xrightarrow{\Phi_{\sigma}^{\omega}} \alpha F_{\omega} \subset \beta F_{\omega}$$

which makes sense for  $|\omega| \subset |\sigma|$  by virtue of (A2).

The chain projections give us a direct system of chain complexes, and therefore also a direct system of homology groups. We obtain the desired homology of  $f$  as a direct limit by setting

$$H_{s,t}(f; G) \approx \lim_{\longrightarrow} H_s(\alpha K; H_t(\alpha F; G)).$$

4. Identification of the fiber homology. In this section we shall prove the following

**THEOREM.** *For every submersion  $f: X \rightarrow Y$  and coefficient module  $G$  there is a canonical isomorphism*

$$\psi_*: H_{0,*}(f; G) \approx H_*(f_*; G).$$

The proof of our Main Theorem, which will be given in the following section, depends upon this result, or more precisely, upon

the surjectivity of  $\psi_*$ .

To prove the theorem at hand, we require the algorithm of § 3 for the computation of the homology groups  $H_{s,t}(f; G)$ . We will take an approximating sequence  $\{\alpha p: {}_\alpha E \rightarrow |{}_\alpha K|\}$  for  $f$ , and for each  $\alpha$  we let  ${}_\alpha C_{s,t}$  denote the corresponding chain group (3.1). One now observes that when  $\sigma$  is a vertex of  ${}_\alpha K$ ,  $H_0(|\sigma|, |\dot{\sigma}|)$  can be identified with the additive group of integers, and this permits us to identify  ${}_\alpha C_{0,t}$  with the direct sum

$$(4.1) \quad \sum_{\sigma \in {}_\alpha K^{(0)}} H_t({}_\alpha F_\sigma; G) .$$

Considering formula (3.2) for the chain boundary  $\partial: {}_\alpha C_{1,t} \rightarrow {}_\alpha C_{0,t}$ , and bearing in mind the definition of the fiber projections  $i_\sigma^*$ , one sees that  $\partial({}_\alpha C_{1,t})$  is precisely the submodule  $W_{\alpha,t}$  of the direct sum (4.1) generated by the subset

$$\{[w - w']: w \wedge w' \bmod {}_\sigma \Phi \text{ for } \sigma \in {}_\alpha K^{(1)}\} .$$

Moreover, the inclusion induced homomorphism

$$H_t({}_\alpha F_\sigma; G) \longrightarrow H_t(f_\sigma; G) ,$$

defined for every vertex  $\sigma$  in  ${}_\alpha K$ , induces a canonical homomorphism

$$\psi_\alpha: \sum_{\sigma \in {}_\alpha K^{(0)}} H_t({}_\alpha F_\sigma; G) \longrightarrow \sum_{y \in Y} H_t(f_y; G)$$

which obviously maps  $W_{\alpha,t}$  into the submodule  $W$  (defined in § 1). Consequently one obtains a canonical homomorphism

$$(4.2) \quad \psi_{\alpha*}: {}_\alpha H_{0,t} \longrightarrow H_t(f_*; G) ,$$

where  ${}_\alpha H_{0,t}$  denotes the 0-dimensional homology of the chain complex  ${}_\alpha C_{*,t}$ .

We recall that the homology module  $H_{0,t}(f; G)$  is given as the direct limit of a direct system

$$(4.3) \quad \{{}_\alpha H_{0,t}, \phi_{\alpha*}^\beta\}$$

the projections  $\phi_{\alpha*}^\beta$  being induced by the chain projections (3.4). It is easy to verify that the homomorphisms (4.2) commute with the projections  $\phi_{\alpha*}^\beta$ . For under the representation (4.1) of the 0-dimensional chain groups, formula (3.5) assumes the form

$$(4.4) \quad \phi_\alpha^\beta(c) = \sum_{\sigma \in {}_\alpha K^{(0)}} j_{\sigma*}^\beta(c) ,$$

where  $j_{\sigma*}^\beta: {}_\alpha F_\sigma \rightarrow {}_\beta F_\sigma$  is induced by the inclusion  ${}_\alpha F_\sigma \subset {}_\beta F_\sigma$ , so that the desired commutativity holds already on the chain level (i.e.,  $\psi_\beta \circ \phi_\alpha^\beta = \psi_\alpha$  for  $\alpha < \beta$ ). The homomorphisms (4.2) constitute there-

fore a homomorphism of the direct system (4.3), and this induces a homomorphism

$$\psi_*: H_{0,t}(f; G) \longrightarrow H_t(f_*; G)$$

in the direct limit.

To establish the surjectivity of  $\psi_*$ , it will suffice to show that for any point  $y \in Y$  and homology class  $w \in H_t(f_y; G)$ , there exists an index  $\alpha$  and an element  $c \in {}_\alpha C_{0,t}$  such that

$$(4.5) \quad [\psi_\alpha(c) - w] \in W.$$

But this is very easy to prove. Let  $w$  be represented by a singular cycle  $Z$  having its support in some compact subset  $C \subset f_y$ . Property (A1) of the approximating sequence permits us to choose an index  $\alpha$  such that  $C \subset {}_\alpha E$ , and we let  $\sigma$  denote the carrier simplex of  $y$ . This means that  $y$  is an interior point of  $|\sigma|$ , and for an arbitrary vertex  $\tau$  of  $\sigma$  we can define a singular cycle  $Z'$  in  ${}_\alpha F_\tau$ , through the formula

$$Z' = {}_\sigma \Phi_y^\tau Z.$$

This determines an element  $w' \in H_t({}_\alpha F_\tau; G)$ , and recalling the representation (4.1), we obtain thus a chain  $c \in {}_\alpha C_{0,t}$ , which obviously satisfies (4.5).

A considerably more complicated argument will be required to establish the injectivity of  $\psi_*$ . To simplify the proof, it will be convenient to assume that our approximating sequence satisfies the additional condition

$$(A3) \quad \lim_{\rightarrow} \text{mesh } {}_\alpha K = 0,$$

which implies that if  $U$  is any open set in  $Y$ , there exists an index  $\alpha$  such that  $U$  contains a vertex of  ${}_\alpha K$ . It is quite easy to see that if one has an approximating sequence satisfying (A1) and (A2), the additional condition (A3) can be achieved by taking appropriate subdivisions. It will also be convenient to precede the proof of injectivity by three lemmas.

To begin with, it will be recalled that an element  $u \in W$  admits a representation of the form

$$(4.6) \quad u = \sum_{i=1}^r [w_i - w'_i]$$

where  $w_i \in H_*(f_{y_i}; G)$ ,  $w'_i \in H_*(f_{y'_i}; G)$  and

$$w_i \frown w'_i \bmod {}_i \Phi$$

for some tubular neighborhoods  ${}_i \Phi: B_i \times {}_i F \rightarrow V_i$  belonging to  $f$ . If

$K$  is a complex with  $|K| \subset Y$ , we will say that the representation (4.6) belongs to  $K$  provided the points  $y_i$  and  $y'_i$  are vertices of  $K$  for all  $i$ .

LEMMA 1. *Given  $c \in {}_\alpha C_{0,i}$  such that  $\psi_\alpha(c) \in W$ , there exists an index  $\beta > \alpha$  such that  $\psi_\alpha(c)$  admits a representation (4.6) belonging to  ${}_\beta K$ .*

To prove this, let us consider an arbitrary representation (4.6) for the element  $\psi_\alpha(c)$ . By virtue of (A3) we can choose  $\beta > \alpha$  so that each subset  $B_i$  contains a vertex of  ${}_\beta K$ . For each index  $i$  we may now choose points  $\bar{y}_i, \bar{y}'_i \in B_i \cap {}_\beta K^{(0)}$ , with the proviso that  $\bar{y}_i = y_i$  when  $y_i \in {}_\beta K^{(0)}$ , and  $\bar{y}'_i = y'_i$  when  $y'_i \in {}_\beta K^{(0)}$ . Moreover, (4.6) implies that we can represent  $w_i$  and  $w'_i$  by singular cycles  $Z_i$  and  $Z'_i$ , in  ${}_i F_{y_i}$  and  ${}_i F_{y'_i}$ , respectively such that

$$Z'_i = {}_i \Phi_{y'_{i\sharp}}^{y'_i} Z_i.$$

Setting

$$\bar{Z}_i = {}_i \Phi_{\bar{y}'_{i\sharp}}^{\bar{y}_i} Z_i$$

$$\bar{Z}'_i = {}_i \Phi_{\bar{y}'_{i\sharp}}^{\bar{y}'_i} Z'_i$$

we obtain corresponding homology classes  $\bar{w}_i \in H_*(f_{\bar{y}_i}; G)$  and  $\bar{w}'_i \in H_*(f_{\bar{y}'_i}; G)$ , and we can now rewrite the representation (4.6) for  $\psi_\alpha(c)$  in the form

$$\psi_\alpha(c) = \sum_{i=1}^r \{[\bar{w}_i - \bar{w}'_i] + [w_i - \bar{w}_i] + [\bar{w}'_i - w'_i]\}.$$

It will be shown that

$$(4.7) \quad \sum_{i=1}^r \{[w_i - \bar{w}'_i] + [\bar{w}'_i - w'_i]\} = 0.$$

To establish this, one observes that  $\psi_\alpha(c)$  must be a linear combination of elements corresponding to vertices of  ${}_\alpha K$ . Our construction insures that  $[w_i - \bar{w}_i]$  vanishes when  $y_i \in {}_\beta K^{(0)}$ , and similarly  $[\bar{w}'_i - w'_i]$  vanishes when  $y'_i \in {}_\beta K^{(0)}$ . This means that the summation in (4.7) represents a linear combination of elements corresponding to vertices which do not belong to  ${}_\beta K$ . On the other hand, since the terms  $[\bar{w}_i - \bar{w}'_i]$  involve only vertices belonging to  ${}_\beta K$ , one may now conclude (4.7). Consequently one has

$$\psi_\alpha(c) = \sum_{i=1}^r [\bar{w}_i - \bar{w}'_i],$$

which is a representation belonging to  ${}_{{}_\beta}K$ .

LEMMA 2. *Given  $c \in {}_{{}_\alpha}C_{0,t}$  such that  $\psi_\alpha(c) = 0$ , there exists an index  $\beta > \alpha$  such that  $\phi_\alpha^\beta(c) = 0$ .*

Since  ${}_{{}_\alpha}C_{0,t}$  is generated by elements  $w \in H_t({}_{{}_\alpha}F_\tau; G)$ , where  $\tau$  is a vertex of  ${}_{{}_\alpha}K$ , it will suffice to establish the lemma in the case  $c = w$ . Let  $w$  be represented by a singular cycle  $Z$  in  ${}_{{}_\alpha}F_\tau$ . Since  $\psi_\alpha$  annihilates  $w$ , there exists a compact subspace  $C \subset f_\tau$  such that  $Z$  bounds in  $C$ . If we now choose  $\beta > \alpha$  such that  $C \subset {}_\beta F_\tau$  (we are using properties (A1) and (A2)), then the inclusion induced homomorphism

$$j_{\tau*}: H_t({}_{{}_\alpha}F_\tau; G) \longrightarrow H_t({}_\beta F_\tau; G)$$

annihilates  $w$ , which implies by (4.4) that  $\phi_\alpha^\beta(w) = 0$ .

LEMMA 3. *Let  $\Phi: B \times F \rightarrow V$  and  $\bar{\Phi}: \bar{B} \times \bar{F} \rightarrow \bar{V}$  be tubular neighborhoods belonging to  $f$ , such that  $\bar{\Phi}$  cuts  $V$  and  $B \cap \bar{B}$  is connected. Let  $y, y' \in B \cap \bar{B}$ , and let  $Z$  denote a singular cycle in  $F_y$ . Then*

$$\Phi_{y*}^{y'} Z \sim \bar{\Phi}_{y*}^{y'} Z \text{ in } \bar{F}_{y'}.$$

We may assume without loss of generality that  $B = \bar{B}$  and  $F \subset \bar{F}$ , which implies, moreover, that  $V \subset \bar{V}$  and  $F_x \subset \bar{F}_x$  for every  $x \in B$ . Considering  $\Phi_{y'}^{y'}$  and  $\bar{\Phi}_{y'}^{y'}$  as maps from  $F_y$  into  $\bar{F}_{y'}$ , it will suffice to construct a homotopy  $H: I \times F_y \rightarrow \bar{F}_{y'}$  such that

$$(4.8) \quad H: \bar{\Phi}_{y'}^{y'} \simeq \Phi_{y'}^{y'}.$$

Let  $\phi: I \rightarrow B$  denote a path from  $y$  to  $y'$ . The requisite homotopy  $H$  may now be given by the formula

$$H(t, x) = \bar{\Phi}_{y'}^{y'} \circ \bar{\Phi}_{\phi(t)}^{y'} \circ \Phi_{y'}^{\phi(t)}(x)$$

which obviously gives (4.8).

At last we are ready to establish the injectivity of  $\psi_*$ . It will suffice to prove that if  $\psi_\alpha(c) \in W$  for some  $c \in {}_{{}_\alpha}C_{0,t}$ , then there exists an index  $\gamma > \alpha$  such that  $\phi_\alpha^\gamma(c) \in W_\gamma$ . We first apply Lemma 1 to conclude that there exists a  $\beta > \alpha$  such that  $\psi_\alpha(c)$  admits a representation (4.6) belonging to  ${}_{{}_\beta}K$ . By virtue of condition (A1) one may further assume that  $V_i \subset {}_\beta E$  for each index  $i$ , which means that each homology class  $w_i$  and  $w'_i$  in the representation (4.6) can be represented by a singular cycle  $Z_i$  and  $Z'_i$  in  ${}_{{}_\beta}F_{y_i}$  and  ${}_{{}_\beta}F_{y'_i}$ , respectively, such that

$$Z'_i = {}_i\Phi_{y_i^*}^{y_i'} Z_i .$$

The cycles  $Z_i$  and  $Z'_i$  induce corresponding homology classes  ${}_{{}_\beta}w_i \in H_t({}_{{}_\beta}F_{y_i}; G)$  and  ${}_{{}_\beta}w'_i \in H_t({}_{{}_\beta}F_{y_i}; G)$ , and this gives an element

$$c' = \sum_{i=1}^r [{}_{{}_\beta}w_i - {}_{{}_\beta}w'_i]$$

in  ${}_{{}_\beta}C_{0,t}$  such that  $\psi_{{}_\beta}(c') = \psi_\alpha(c)$ . Since  $\psi_\alpha = \psi_{{}_\beta} \circ \phi_\alpha^{{}_\beta}$ , it follows that  $\psi_{{}_\beta}$  annihilates  $[\phi_\alpha^{{}_\beta}(c) - c']$ . By Lemma 2, this element is also annihilated by  $\phi_{{}_\beta}^\gamma$  for some  $\gamma > \beta$ , and this leads to a formula

$$\phi_\alpha^\gamma(c) = \sum_{i=1}^r [{}_{{}_\gamma}w_i - {}_{{}_\gamma}w'_i] ,$$

in which the homology classes  ${}_{{}_\gamma}w_i$  and  ${}_{{}_\gamma}w'_i$  can once again be represented by the singular cycles  $Z_i$  and  $Z'_i$ , respectively. It remains to be shown that

$$[{}_{{}_\gamma}w_i - {}_{{}_\gamma}w'_i] \in \partial_{{}_\gamma}C_{1,t}$$

for all  $i$ . Since each subset  $B_i$  is open and connected, we may assume without loss of generality that the vertices  $y_i$  and  $y'_i$  can be joined by an edge path in  ${}_{{}_\gamma}K$  (condition (A3) guarantees that this will hold for sufficiently large values of  $\gamma$ ). Let  $\tau_0, \dots, \tau_n$  denote successive vertices of this edge path, and let

$$(4.9) \quad Z_j = {}_i\Phi_{y_j^*}^{\tau_j} Z; \quad j = 0, \dots, n .$$

The cycles  $Z_j$  determine homology classes  $\bar{w}_j \in H_t({}_{{}_\gamma}F_{\tau_j}; G)$ , and one sees that  $\bar{w}_0 = {}_{{}_\gamma}w_i$  and  $\bar{w}_n = {}_{{}_\gamma}w'_i$ . It suffices therefore to show that

$$(4.10) \quad [\bar{w}_j - \bar{w}_{j-1}] \in \partial_{{}_\gamma}C_{1,t}; \quad j = 1, \dots, n .$$

We note that

$$Z_j = {}_i\Phi_{\tau_{j-1}^*}^{\tau_j} Z_{j-1}$$

by virtue of (4.9). Moreover, if  $\sigma_j$  denotes the 1-simplex in  ${}_{{}_\gamma}K$  with vertices  $\tau_j$  and  $\tau_{j-1}$ , then  $\sigma_j\Phi$  cuts  $V_i$ . One may conclude by Lemma 3 that

$$\bar{w}_j \not\sim \bar{w}_{j-1} \bmod \sigma_j\Phi ,$$

which implies (4.10).

**5. Proof of the main theorem.** Let  $f: X \rightarrow Y$  be a submersion and  $G$  a coefficient module. According to the main result in [1], there exists a convergent  $E^2$  spectral sequence with

$$(5.1) \quad E_{s,t}^2 \approx H_{s,t}(f; G)$$

and  $E^\infty$  isomorphic to the bigraded module associated with a filtration of  $H_*(X; G)$ . By the result of § 4, (5.1) implies

$$(5.2) \quad E_{0,t}^2 \approx H_t(f_*; G) .$$

Since evidently  $E_{0,0}^2 \approx E_{0,0}^\infty \approx H_0(X; G)$ , one obtains immediately the desired equality  $R_0(f_*; G) = R_0(X; G)$ .

Under the assumption that the fibers  $f_y$  are connected, the homology groups  $H_{s,0}(f; G)$  evidently reduce to  $H_s(Y; G)$ , so that

$$(5.3) \quad E_{s,0}^2 \approx H_s(Y; G) .$$

If one assumes further that the fibers  $f_y$  are  $t$ -acyclic over  $G$  for  $q - m < t < q$ , one may conclude (by Theorem 4.1 of [1] and (5.1)) that

$$(5.4) \quad E_{s,t}^2 = 0 \text{ for all } s \text{ and } q - m < t < q .$$

Moreover, since the chain groups entering into the definition of  $H_{s,t}(f; G)$  are trivial for  $s > m = \dim Y$ , one also obtains

$$(5.5) \quad E_{s,t}^2 = 0 \text{ for } s > m \text{ and all } t .$$

The desired inequality

$$(5.6) \quad R_q(f_*; G) \leq R_q(X; G) + R_{q+1}(Y; G)$$

follows from conditions (5.2) through (5.5) by a simple spectral sequence argument, which runs as follows.

We may assume  $q \geq 1$ . The exact sequence

$$(5.7) \quad E_{q+1,0}^{q+1} \xrightarrow{d^{q+1}} E_{0,q}^{q+1} \longrightarrow \text{coker } d^{q+1} \longrightarrow 0$$

gives the rank inequality

$$(5.8) \quad \text{rank } E_{0,q}^{q+1} \leq \text{rank coker } d^{q+1} + \text{rank } E_{q+1,0}^{q+1} .$$

Since all differentials into the  $(q+1, 0)$ -position are trivial, one has  $E_{q+1,0}^{q+1} \subset E_{q+1,0}^2$ , and this implies by (5.3) that

$$(5.9) \quad \text{rank } E_{q+1,0}^{q+1} \leq R_{q+1}(Y; G) .$$

Moreover, one observes by (5.4) and (5.5) that  $d^{q+1}$  is the only non-trivial differential touching the  $(0, q)$ -position. This gives  $E_{0,q}^2 \approx E_{0,q}^{q+1}$ , which together with (5.2) implies

$$(5.10) \quad \text{rank } E_{0,q}^{q+1} = R_q(f_*; G) .$$

The preceding observation also gives



$$\text{coker } d^{q+1} \approx E_{0,q}^{q+2} \approx E_{0,q}^{\infty}.$$

But since  $E^{\infty}$  is isomorphic to the bigraded module associated to a filtration of  $H_*(X; G)$ , this implies

$$(5.11) \quad \text{rank coker } d^{q+1} \leq R_q(X; G).$$

Finally, (5.8) through (5.11) yields (5.6).

If, in addition,  $H_q(X; G)$  and  $H_{q+1}(Y; G)$  vanish, then one sees by the same considerations that the exact sequence (5.7) reduces to

$$0 \longrightarrow H_q(f_*; G) \longrightarrow 0 \longrightarrow 0,$$

and this completes our proof.

## 6. Orientability of real-valued submersions.

**THEOREM.** *A submersion  $f: X \rightarrow R$  is orientable if and only if it does not admit separatrices.*

In this section we shall prove the nontrivial half of this theorem (the “if” part). This will involve looking carefully at representations (4.6) for elements  $u \in W$  in the special case  $Y = R$ . Such a representation will be called *simple* if

$$y_1 < y'_1 \leq y_2 < y'_2 \leq y_3 < \cdots < y'_r.$$

**LEMMA 1.** *Every element of  $W$  admits a simple representation.*

To begin with, one may assume without loss of generality that one has a representation with  $y_i < y'_i$  for  $i \leq r$  and  $y_i \leq y_{i+1}$  for  $i < r$  (this can be achieved by proper “labeling”). We also note that if  $y \in (y_i, y'_i)$ , one may evidently “subdivide” the given representation by setting

$$[w_i - w'_i] = [w_i - w] + [w - w'_i],$$

where  $w$  denotes the (uniquely determined) element in  $H_*(f_y; G)$  such that

$$w_i \frown w \bmod \Phi \text{ and } w \frown w' \bmod \Phi.$$

By the use of such subdivisions one can eliminate “partial intersections” between open intervals  $(y_i, y'_i)$ , which is to say that one can achieve the following condition: two intervals  $(y_i, y'_i)$  and  $(y_j, y'_j)$  are either disjoint or identical. The resulting representation need not yet be simple, precisely on account of the second alternative. Let us therefore suppose that for all values of  $i$  in some subset  $J$  one

has  $y_i = y$  and  $y'_i = y'$ . It remains to be shown that one can construct a simple representation for the element

$$(6.1) \quad \sum_{i \in J} [w_i - w'_i] .$$

To this end let  $C \subset X$  denote a compact subset containing all images  $V_i$  corresponding to tubular neighborhoods  ${}_i\Phi$  with  $i \in J$ . By virtue of (1.4) there exists a partition

$$y = y^0 < y^1 < \cdots < y^s = y'$$

and tubular neighborhoods  $\Phi^j: B^j \times F^j \rightarrow V^j$  such that  $[y^{j-1}, y^j] \subset B^j$  and  $\Phi^j$  cuts  $C$  for  $j = 1, \dots, s$ . For each  $i \in J$  the homology class  $w_i$  can be represented by a singular cycle  $Z_i$  in  ${}_iF_y$ , and this gives rise to cycles

$$Z_i^j = {}_i\Phi_{y^j}^j Z_i ; \quad j = 0, \dots, s .$$

We let  $w_i^j \in H_*(f_{y^j}; G)$  denote the corresponding homology classes. It follows now by Lemma 3 of § 4 that

$$w_i^{j-1} \sim w_i^j \bmod \Phi^j \quad \text{for } j = 1, \dots, s .$$

Setting

$$w^i = \sum_{i \in J} w_i^j$$

one obtains a simple representation

$$\sum_{j=1}^s [w^{j-1} - w^j]$$

for the element (6.1).

LEMMA 2. Let  $w \in H_t(f_y; G)$  be nonzero, and let

$$w \sim 0_{y'} \bmod \Phi$$

where  $0_{y'}$  denotes the zero in  $H_t(f_{y'}; G)$ . Then  $f$  admits a separatrix  $f_{y^*}$  for some point  $y^*$  between  $y$  and  $y'$ .

The proof of this lemma is very simple. We may assume  $y < y'$ , and we let  $S$  denote the set of all points  $\bar{y} \in [y, y']$  such that

$$w \sim 0_{\bar{y}} \bmod \Phi .$$

It is an easy consequence of (1.4) that  $S$  is open. Since  $y' \in S$  and  $y \notin S$ , there must be a point  $y^* \in [y, y')$  belonging to the boundary of  $S$ , and this means that  $f_{y^*}$  will be a separatrix.

The proof of our theorem can now proceed as follows. Let

$f: X \rightarrow R$  be a submersion without separatrices. To show that  $f$  is orientable, one must prove that a nonzero element  $w \in H_t(f_y; G)$  cannot belong to  $W$ . But if  $w \in W$ , one concludes by Lemma 1 that it admits a simple representation

$$\sum_{i=1}^r [w_i - w'_i]$$

and it may be supposed on the strength of Lemma 2 that the elements  $w_i$  and  $w'_i$  are all nonzero (for if one element is zero, the other member of the pair must be zero as well). Since  $w'_r$  is the only element in the given representation which corresponds to the point  $y'_r$ ,  $w'_r$  can be nonzero only if  $y = y'_r$  and  $w = w'_r$ . But this would imply by the same token that  $w_1 = 0$ , and this furnishes the desired contradiction.

#### REFERENCES

1. P. C. Endicott and J. W. Smith, *A homology spectral sequence for submersions*, Pacific J. Math., to appear.
2. J. W. Smith, *Submersions of codimension 1*, J. Math. Mech., **18** (1968), 437-444.
3. J. W. Smith, *On the homology structure of submersions*, Math. Ann., **193** (1971), 217-224.
4. E. H. Spanier, *Algebraic Topology*, McGraw-Hill, 1966.

Received December 11, 1978 and in revised form May 31, 1979.

ROUTE 2, Box 16  
PHILOMATH, OR 97370

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1980 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Frank Hayne Beatrous, Jr. and R. Michael Range, <i>On holomorphic approximation in weakly pseudoconvex domains</i> .....	249
Lawrence Victor Berman, <i>Quadratic forms and power series fields</i> .....	257
John Bligh Conway and Waclaw Szymański, <i>Singly generated antisymmetric operator algebras</i> .....	269
Patrick C. Endicott and J. Wolfgang Smith, <i>A homology spectral sequence for submersions</i> .....	279
Sushil Jajodia, <i>Homotopy classification of lens spaces for one-relator groups with torsion</i> .....	301
Herbert Meyer Kamowitz, <i>Compact endomorphisms of Banach algebras</i> .....	313
Keith Milo Kendig, <i>Moiré phenomena in algebraic geometry: polynomial alternations in <math>\mathbf{R}^n</math></i> .....	327
Cecelia Laurie, <i>Invariant subspace lattices and compact operators</i> .....	351
Ronald Leslie Lipsman, <i>Restrictions of principal series to a real form</i> .....	367
Douglas C. McMahon and Louis Jack Nachman, <i>An intrinsic characterization for PI flows</i> .....	391
Norman R. Reilly, <i>Modular sublattices of the lattice of varieties of inverse semigroups</i> .....	405
Jeffrey Arthur Rosoff, <i>Effective divisor classes and blowings-up of <math>\mathbf{P}^2</math></i> .....	419
Zalman Rubinstein, <i>Solution of the middle coefficient problem for certain classes of C-polynomials</i> .....	431
Alladi Sitaram, <i>An analogue of the Wiener-Tauberian theorem for spherical transforms on semisimple Lie groups</i> .....	439
Hal Leslie Smith, <i>A note on disconjugacy for second order systems</i> .....	447
J. Wolfgang Smith, <i>Fiber homology and orientability of maps</i> .....	453
Audrey Anne Terras, <i>Integral formulas and integral tests for series of positive matrices</i> .....	471