

Pacific Journal of Mathematics

IRREDUCIBLE OPERATORS WHOSE SPECTRA ARE SPECTRAL SETS

ROBERT OLIN AND JAMES E. THOMSON

IRREDUCIBLE OPERATORS WHOSE SPECTRA ARE SPECTRAL SETS

ROBERT F. OLIN AND JAMES E. THOMSON

In this note those compact subsets of the plane that are the spectra of irreducible subnormal operators are characterized.

In December 1977 John B. Conway presented a colloquium talk at Virginia Tech and asked, "Which compact subsets of the plane are the spectra of irreducible subnormal operators?" If the adjective irreducible is replaced by pure in this question, then Clancey and Putnam [2] have the following answer: A compact set K is the spectrum of a pure subnormal operator if and only if for every open disc Δ that has a nonempty intersection with K , we have $R(K \cap \Delta^-) \neq C(K \cap \Delta^-)$. Similarly, our answer to Conway's question will be a function algebraic characterization. For the basic facts concerning this area we refer to [3] or [10].

If \mathcal{H} is a separable Hilbert space the algebra of continuous operators on \mathcal{H} will be denoted $\mathcal{B}(\mathcal{H})$. An operator $T \in \mathcal{B}(\mathcal{H})$ is irreducible if \mathcal{H} has no nonzero subspace that is invariant under T and its adjoint T^* . For the basic facts concerning subnormal operators we refer to [4]. If $T \in \mathcal{B}(\mathcal{H})$ then $\sigma(T)$ will denote the spectrum of T .

If K is a compact subset of the plane, $\mathcal{R}(K)$ will denote the collection of rational functions with poles off K ; the uniform closure of $\mathcal{R}(K)$ in $C(K)$, the algebra of continuous functions on K , is denoted $R(K)$. If φ belongs to the maximal ideal space of $R(K)$, then there exists a point $z \in K$ such that $\varphi(f) = f(z)$ for all $f \in R(K)$. Hence the Gleason parts of $R(K)$ form a partition of K .

If $T \in \mathcal{B}(\mathcal{H})$ and K is a compact set containing $\sigma(T)$, then K is called a spectral set for T if

$$(i) \quad \|f(T)\| \leq \|f\|_K$$

for all $f \in \mathcal{R}(K)$. (Here $\|f\|_K$ denotes the sup norm of f on K). If K is a spectral set for T , it is easy to define $f(T) (\in \mathcal{B}(\mathcal{H}))$ for all $f \in R(K)$. A basic fact about a subnormal operator S is that $\sigma(S)$ is a spectral set and equality holds in (i) with $K = \sigma(S)$ for all $f \in R(\sigma(S))$.

THEOREM 1. *A compact set K is the spectrum of an irreducible operator T whose spectrum is a spectral set if and only if $R(K)$ has one nontrivial Gleason part G and $G^- = K$.*

REMARK. In the sufficiency part of the proof the operator we construct is subnormal. Hence, the theorem answers the question of Conway. For the necessity part of the proof we need two (non-trivial) results and an elementary fact.

Fact. Suppose K_i is a spectral set for $T_i \in \mathcal{B}(\mathcal{H}_i)$ and $K = (\cup K_i)^-$ is compact. Then K is a spectral set for $T = \bigoplus T_i$.

We leave the proof to the reader.

THEOREM 2. (*Mlak, Lautzenheiser, Seever*). Let K be a spectral set for $T \in \mathcal{B}(\mathcal{H})$ and let G_1, G_2, \dots be the nontrivial Gleason parts of $R(K)$. Then $T = N \bigoplus (\bigoplus_i T_i)$, where N is a normal operator with spectrum contained in the boundary of K and G_i^- is a spectral set for T_i . The proof of this and related results can be found in [5, 7, 8, 9].

The following theorem (combined with the fact and Theorem 2) guarantees that if K is a spectral set for T with at least two nontrivial Gleason parts G_1, G_2 for $R(K)$ such that $\sigma(T) \cap G_i \neq \emptyset$ $i = 1, 2$, then T has a nontrivial reducing subspace.

THEOREM 3. (*Melnikov*). Let G_1 be a nontrivial Gleason part for $R(K)$. If $a \in G_1^- \setminus G_1$ then a is a peak point. In particular, if G_2 is another Gleason part then $G_2 \cap G_1^- = \emptyset$.

The proof of this result can be found in [6].

Proof of Theorem 1. The only thing left to establish is if K is a compact set with one nontrivial Gleason part G and $G^- = K$, then K is the spectrum of an irreducible subnormal operator. Choose a sequence $\{z_n\}$ of points belonging to G such that $\{z_n\}^- = K$. Let δ_n denote point mass measure at z_n . Fix $z_0 \in G$.

For each z_n choose a representing measure λ_n for $R(K)$ at z_0 such that δ_n is absolutely continuous with respect to λ_n (such a measure can be found by [10, p. 165]). Then

$$\lambda = \sum_n \frac{\lambda_n}{2^n}$$

is a representing measure for $R(K)$ at z_0 such that the support of λ is K . Let $\mathcal{H} = R^2(\lambda)$, the $L^2(\lambda)$ closure of $R(K)$, and let S be multiplication by z . Clearly $\sigma(S) = K$ so we need to show S is irreducible.

Observe first that if $f \in R(K)$ then

$$\begin{aligned}
 |f(z_0)| &= \left| \int f d\lambda \right| \\
 &= |\langle f, 1 \rangle|.
 \end{aligned}$$

(Here \langle , \rangle denotes the inner product in $R^2(\lambda)$.) It is now routine to show that

$$(ii) \quad \langle h_1 h_2, 1 \rangle = \langle h_1, 1 \rangle \langle h_2, 1 \rangle$$

for all $h_i \in R^2(\lambda) \cap L^\infty(\lambda)$ for $i = 1, 2$.

Using the techniques of [12], one can show that the commutant of S is the set $\{M_\psi: \psi \in R^2(\lambda) \cap L^\infty(\lambda)\}$, where M_ψ denotes multiplication by ψ . Hence, if P is a projection commuting with S , then $P = M_f$, where f is the characteristic function of a Borel set F . By (ii) we see that $\langle f, 1 \rangle$ equals zero or one. We assume that $\langle f, 1 \rangle = 1$ (otherwise we work with the projection $1 - P = M_{1-f}$). Hence

$$\begin{aligned}
 1 &= \langle f, 1 \rangle \\
 &= \int_F d\lambda \\
 &= \lambda(F).
 \end{aligned}$$

Since λ is a probability measure, $f = 1$ almost everywhere; hence $P = 1$.

Unfortunately there are no known topological criteria that characterize the sets which are nontrivial Gleason parts of $R(K)$. (Many necessary topological and measure theoretic facts are known: (1) \bar{G} is connected. (2) G is σ -compact. (3) G has (area) density one at each of its points. (4) G is area connected (consult [6]), etc.) However, the literature contains many interesting examples discussing various properties of these parts. Combining these examples with Theorem 1, one comes up with interesting operator theoretic consequences.

Fix an operator T whose spectrum is a spectral set. If $\sigma(T)$ has two components of its interior, say U_1 and U_2 such that $\sigma(T) = (U_1 \cup U_2)^-$ one may ask if $T = T_1 \oplus T_2$ with $\sigma(T_i) \subset U_i^-$ for $i = 1, 2$. However, since there exists a disconnected nontrivial part [10, Examples 26.24 and 26.25] the answer, in general, is no. (The first person to observe this phenomenon was Lautzenheiser in his thesis [5]. In this interesting work he also discusses how Theorem 2 subsumes many other known results.) One may also wonder if it is possible to write $T = T_1 \oplus T_2$ with $\sigma(T_1) \subset (\text{int } \sigma(T))^-$ and $\sigma(T_2) \subset (\sigma(T) \setminus \text{int } \sigma(T))$. Again the answer is no. Let $D = \{z: |z| < 1\}$, $D_+ = \{z \in D: \text{Re } z > 0\}$ and $D_- = \{z \in D: \text{Re } z < 0\}$. Construct a compact set F from the closed unit disc D^- by removing pairwise disjoint

open discs D_i from D_+ such that $\text{int}(D_+ \setminus \cup D_i) = \emptyset$ and $\sum r_i < \infty$ (r_i is the radius of D_i). Using Theorem 2 in [11] and Theorem 3 in [1], we can construct F such that zero is not a peak point of $R(F)$. Let $K = G^-$ where G is the nontrivial part of $R(F)$ containing zero. (Note: $G \cap D_+ \neq \emptyset$ and $G \supset D_-$.)

REFERENCES

1. A. Browder, *Point derivations on function algebras*, J. Functional Anal., **1** (1967), 22-27.
2. K. Clancey and C. R. Putnam, *The local spectral behavior of completely subnormal operators*, Trans. Amer. Math. Soc., **163** (1972), 239-244.
3. T. Gamelin, *Uniform Algebras*, Printice-Hall, Englewood Cliffs, NJ., 1969.
4. P. Halmos, *A Hilbert space problem book*, Van Nostrand, Princeton, NJ., 1967.
5. R. Lautzenheiser, *Spectral sets, reducing subspaces and function algebras*, Thesis, Indiana University, 1973.
6. M. Melnikov, *The Gleason parts of the algebra $R(X)$* , Math. Sb., **101** (143) (1976), 293-300.
7. W. Mlak, *Decompositions and extensions of operator valued representations of junction algebras*, Acta Sci. Math. (Szeged), **30** (1969), 181-193.
8. ———, *Partitions of spectral sets*, Ann. Pol. Math., **25** (1972), 273-280.
9. G. L. Seever, *Operator representations of uniform algebras. I.*, preprint.
10. E. L. Stout, *The theory of uniform algebras*, Bogden and Quigley, Tarrytown-on-Hudson NJ., 1971,
11. J. Wermer, *Bounded point derivations on certain Banach algebras*, J. Functional Anal., **1** (1967), 28-36.
12. T. Yoshino, *Subnormal operators with a cyclic vector*, Tohoku Math. J., **21** (1969), 47-55.

Received October 10, 1979. This research was supported in part by the National Science Foundation under Grant No. MCS77-00966.

Current address: Virginia Tech. Blacksburg, Virginia 24061

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$84.00 a year (6 Vols., 12 issues). Special rate: \$42.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1980 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Victor P. Camillo and Julius Martin Zelmanowitz, <i>Dimension modules</i>	249
Yonina S. Cooper, <i>Stable sequences in pre-abelian categories</i>	263
Chandrakant Mahadeorao Deo and H. Ship-Fah Wong, <i>On Berry-Esseen approximation and a functional LIL for a class of dependent random fields</i>	269
H. P. Dikshit and S. N. Dubey, <i>$C, 1$ summability of series associated with Fourier series</i>	277
M. Edelstein, <i>On the homomorphic and isomorphic embeddings of a semiflow into a radial flow</i>	281
Gilles Godefroy, <i>Compacts de Rosenthal</i>	293
James Guyker, <i>Commuting hyponormal operators</i>	307
Thomas Eric Hall and Peter R. Jones, <i>On the lattice of varieties of bands of groups</i>	327
Taqdir Husain and Saleem H. Watson, <i>Topological algebras with orthogonal Schauder bases</i>	339
V. K. Jain, <i>Some expansions involving basic hypergeometric functions of two variables</i>	349
Joe W. Jenkins, <i>On group actions with nonzero fixed points</i>	363
Michael Ellsworth Mays, <i>Groups of square-free order are scarce</i>	373
Michael John McAsey, <i>Canonical models for invariant subspaces</i>	377
Peter A. McCoy, <i>Singularities of solutions to linear second order elliptic partial differential equations with analytic coefficients by approximation methods</i>	397
Terrence Millar, <i>Homogeneous models and decidability</i>	407
Stephen Carl Milne, <i>A multiple series transformation of the very well poised ${}_{2k+4}\Psi_{2k+4}$</i>	419
Robert Olin and James E. Thomson, <i>Irreducible operators whose spectra are spectral sets</i>	431
Robert John Piacenza, <i>Cohomology of diagrams and equivariant singular theory</i>	435
Louis Jackson Ratliff, Jr., <i>Integrally closed ideals and asymptotic prime divisors</i>	445
Robert Breckenridge Warfield, Jr., <i>Cancellation of modules and groups and stable range of endomorphism rings</i>	457
B. J. Day, <i>Correction to: "Locale geometry"</i>	487
Stanley Stephen Page, <i>Correction to: "Regular FPF rings"</i>	487
Augusto Nobile, <i>Correction to: "On equisingular families of isolated singularities"</i>	489