

# Pacific Journal of Mathematics

## **$\mathcal{H}$ -BORELIAN EMBEDDINGS AND IMAGES OF HAUSDORFF SPACES**

BRENDA MACGIBBON

# $\mathcal{K}$ -BORELIAN EMBEDDINGS AND IMAGES OF HAUSDORFF SPACES

BRENDA MACGIBBON

The following two questions are discussed in this article:  
(1) under what conditions is a Hausdorff space embeddable as a  $\mathcal{K}$ -Borelian subset of some Hausdorff space; and (2) under what conditions is the Hausdorff 1 — 1 continuous image of a  $\mathcal{K}$ -Borelian subset of a Hausdorff space a  $\mathcal{K}$ -Borelian subset of some Hausdorff space containing it. We obtain necessary and sufficient conditions in answer to question (1) in the case of a  $\mathcal{K}_\sigma$  and necessary conditions for  $\mathcal{K}$ -Borelian subsets of each class  $\alpha$  if the containing space is a perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$ . Question 2 does not always have a positive answer as is shown by an example of a Hausdorff 1 — 1 continuous image of  $\mathcal{K}$ -Borelian subset of a compact Hausdorff space which is not  $\mathcal{K}$ -Borelian in any Hausdorff space containing it. In partial answer to question (2) necessary conditions on the domain and the range of the function are presented.

In §1 the necessary definitions are given. Many of the original definitions for completely regular spaces such as bianalytic and Borelian as well as the notion of a complete sequence of countable covers are due to Frolik and can be found in [3]. The class of  $\sigma$ -bianalytic spaces is an enlargement of the class of bianalytic spaces. In [8] the  $\mathcal{K}_\sigma$ -fiable spaces have already been introduced. For the reader's convenience, the definitions of these concepts have been given again.

The motivation of §2 is to extend to a larger class of spaces the well-known result that if  $f(X)$  is a metrizable 1 — 1 continuous image of a  $\mathcal{K}$ -Borelian subset of a compact metric space, then  $f(X)$  is a  $\mathcal{K}$ -Borelian subset of any of its metrizable compactifications. An example is given of a Hausdorff 1 — 1 continuous image of a  $\mathcal{K}$ -Borelian subset of a compact space which is not a  $\mathcal{K}$ -Borelian subset of any Hausdorff space in which it can be embedded. However, using some of the techniques that Frolik developed in [3] for completely regular spaces, necessary and sufficient conditions in order that a 1 — 1 continuous image of  $\sigma$ -bianalytic space  $X$  be a  $\mathcal{K}$ -Borelian subset of some Hausdorff space are obtained. This result can be applied to the cases where  $X$  is a  $\mathcal{K}$ -Borelian subset of a Hausdorff space  $Y$  and  $Y$  is either a perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$ , or has property  $I$  ([10]) or is locally metrizable.

In §3 results are presented concerning the relationship between

the Borel class of a  $\mathcal{K}$ -Borelian subset of a Hausdorff  $\mathcal{K}_\sigma$  space and its complete sequence of countable covers as defined in [4]. Theorem 3.1, giving necessary and sufficient conditions for a space to be embeddable as a  $\mathcal{K}_\sigma$  subset of some Hausdorff space, is an extension of Frolik's result for completely regular spaces [4]. Theorem 3.3 gives necessary conditions for subsets of perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$  Hausdorff spaces to be  $\mathcal{K}$ -Borelian of class  $\alpha$ . The proofs depend on techniques developed by Frolik in [4].

1. Preliminaries. Throughout this article all spaces will be assumed to be Hausdorff and the notation of Frolik in [3] and [4] will be followed. If  $X \subset Y$  then  $\bar{X}^Y$  will denote the closure of  $X$  in  $Y$ .

DEFINITION 1.1.  $Y$  is said to be a  $\mathcal{K}_\sigma$ -fication of a Hausdorff space  $X$  provided that  $X$  can be embedded as a dense subspace of  $Y$  and  $Y$  is a Hausdorff  $\mathcal{K}_\sigma$  space (that is,  $Y$  is equal to a countable union of its compact subsets). A Hausdorff space for which there exists such a  $\mathcal{K}_\sigma$ -fication is said to be  $\mathcal{K}_\sigma$ -fiable.

DEFINITION 1.2.  $X$  is said to be a perfectly  $\sigma$ -normal Hausdorff  $\mathcal{K}_\sigma$  space if, in addition to being a Hausdorff  $\mathcal{K}_\sigma$  space, every closed subset is a  $\mathcal{G}_\delta$ , that is, a countable intersection of open subsets.

It should be noted that this definition does not imply that  $X$  is normal; in fact,  $X$  need not be regular. For example, let  $A, B$  be two countable disjoint dense subsets of  $[0, 1]$  in the usual topology. Let  $X = A \cup B$  endowed with the following topology  $\mathcal{T}: O \in \mathcal{T}$  if and only if  $O = O' \cup (O'' \cap B)$  where  $O', O''$  are open subsets of  $[0, 1]$  in the usual topology.  $X$  is clearly a nonregular Hausdorff perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$  space. Any perfectly normal  $\mathcal{K}_\sigma$  space is clearly a perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$  space.

DEFINITION 1.3. Let  $\{X_i\}_{i=0}^\infty$  be a sequence of Hausdorff spaces that are mutually disjoint. Then  $\sum_{i=0}^\infty X_i$  will represent the set  $X = \bigcup_{i=0}^\infty X_i$  endowed with the following topology:  $O$  is open in  $X$  if and only if  $O \cap X_i$  is open for all  $i$ .

DEFINITION 1.4. Let  $X$  be a Hausdorff space and let  $\mathcal{K}(X)$  denote the class of compact subsets of  $X$ . Then  $\mathcal{B}(\mathcal{K}(X))$ , the class of  $\mathcal{K}$ -Borelian subsets of  $X$ , is defined to be the smallest class of subsets of  $X$  containing  $\mathcal{K}(X)$  and closed under countable intersections and countable unions.

Let  $\mathcal{Z}(X)$  denote the class of zero sets of continuous functions on  $X$  and  $\mathcal{F}(X)$  the class of closed subsets of  $X$ . Let  $\mathcal{B}(\mathcal{Z}(X))$ , the class of Baire subsets of  $X$ , and  $\mathcal{B}(\mathcal{F}(X))$ , the class of  $\mathcal{F}$ -

Borelian subsets of  $X$ , be defined by analogy to the definition of  $\mathcal{B}(\mathcal{H}(X))$ .

Definitions of  $\mathcal{H}$ -Borelian or  $\mathcal{F}$ -Borelian subsets of a space  $X$  of class  $\alpha$  for each ordinal  $\alpha$  of second class (that is, for each  $\alpha$  less than the first uncountable ordinal) are given in [6] or [1].

DEFINITION 1.5. A Hausdorff space  $X$  has property I if for all  $A, B$  in  $\mathcal{H}(X)$  ( $A \setminus B$ ) is in  $\mathcal{H}_\alpha(X)$  (that is, the intersection of  $A$  and the complement of  $B$  is a  $\mathcal{H}_\alpha$  subset of  $X$ ).

The notion of a complete sequence of countable covers was introduced by Frolik in [4] and [3]. The following definitions are taken from there.

DEFINITION 1.6. A family of subsets  $\{E_r\}_{r \in \mathcal{C}}$  of a topological space  $X$  is said to have the *finite intersection property* if every finite subfamily has nonvoid intersection.

DEFINITION 1.7.  $\mu = \{\mathcal{H}_n\}_{n=1}^\infty$  is called a *sequence of countable covers of  $X$*  if, for each  $n$ ,

- (1)  $\mathcal{H}_n$  consists of a sequence of subsets  $\{H_{ni}\}_{i=1}^\infty$  of  $X$ ; and
- (2)  $X \subset \bigcup_{i=1}^\infty H_{ni}$ .

DEFINITION 1.8. A family  $\mathcal{C} = \{E_r\}_{r \in \mathcal{C}}$  of subsets of  $X$  is a  $\mu$ -*Cauchy family* if

- (1)  $\mathcal{C}$  has the finite intersection property; and
- (2) for all  $n$ , there exists a  $\gamma_n \in \mathcal{C}$  and a set  $H_{ni_n} \in \mathcal{H}_n$  such that  $E_{\gamma_n} \subset H_{ni_n}$ .

DEFINITION 1.9.  $\mu = \{\mathcal{H}_n\}_{n=1}^\infty$  is said to be a *complete sequence of countable covers of  $X$*  if, for each  $\mu$ -Cauchy family  $\mathcal{C} = \{E_r\}_{r \in \mathcal{C}}$ ,  $\bigcap_{r \in \mathcal{C}} \bar{E}_r^X \neq \emptyset$ .

DEFINITION 1.10. A Hausdorff space  $X$  is said to have a *complete sequence*  $\mu = \{\mathcal{H}_j\}_{j=1}^\infty$  of countable  $\mathcal{F}$ -Borelian covers of at most class  $\alpha$  if

- (1)  $\mu$  is complete in the sense of Definition 1.9; and
- (2) for all  $j$ ,  $\mathcal{H}_j = \{H_{ji}\}_{i=1}^\infty$  and each  $H_{ji}$  is a  $\mathcal{F}$ -Borelian subset of  $X$  of at most class  $\alpha$ .

If  $X$  is a completely regular space,  $\beta X$  will denote the Stone-Ćech compactification of  $X$ .

The following definition can be found in [3].

DEFINITION 1.11.  $\mathcal{B}(X)$  will denote the class of *Borelian spaces* which are subsets of  $X$ . A space  $Y$  is said to be a *Borelian space*

if there exists a *Borelian structure* in  $Y$ . A *Borelian structure* in  $Y$  is a complete sequence  $\{\mathcal{H}_n\}_{n=1}^\infty$  of countable disjoint coverings of  $Y$  satisfying the following two conditions:

- (1)  $\mathcal{H}_{n+1}$  refines  $\mathcal{H}_n$  for each  $n$ ,
- (2) if for each  $n$   $M_n$  and  $N_n$  are elements of  $\mathcal{H}_n$  and there exists a  $k$  such that  $M_k \neq N_k$ , then  $\bigcap_{n=1}^\infty \bar{M}_n^r \cap \bigcap_{n=1}^\infty \bar{N}_n^r = \emptyset$ .

DEFINITION 1.12. A Hausdorff space  $X$  is  $\mathcal{H}$ -analytic if it is the continuous image of a  $\mathcal{H}_\sigma$  subset of a compact space.

Let us define the following new class of spaces.

DEFINITION 1.13. A Hausdorff space  $X$  is said to be  $\sigma$ -bianalytic if there exists a sequence  $\{X_i\}_{i=1}^\infty$  of subsets of  $X$  such that

- (1)  $X = \bigcup_{i=1}^\infty X_i$ , and
- (2) each  $X_i$  is bianalytic; that is, both  $X_i$  and  $(\beta X_i \setminus X_i)$  are completely regular  $\mathcal{H}$ -analytic spaces for each  $i$ .

We should note that the union in Definition 1.13 need not be disjoint.

PROPOSITION 1.14. *Every bianalytic space is  $\sigma$ -bianalytic. However, there exist completely regular  $\sigma$ -bianalytic spaces that are not bianalytic.*

*Proof.* The first statement is obvious. If  $X = N \cup \{x\}$  where  $N$  is the space of the positive integers with the discrete topology and  $x \in (\beta N \setminus N)$ , then Frolik showed in [5] that this space was not bianalytic.

PROPOSITION 1.15. *There exists a  $\sigma$ -bianalytic space which is not a  $\mathcal{H}$ -Borelian subset of any Hausdorff space.*

*Proof.* Let  $X$  denote the Hausdorff space  $[0, 1]$  with the following topology  $\mathcal{T}$ :  $O \in \mathcal{T}$  if and only if  $O = O' \cup (O'' \cap Q)$  where  $O', O''$  are open sets in  $[0, 1]$  with respect to the usual topology and  $Q$  is the set of rationals in  $[0, 1]$ . Since  $X = I \cup Q$ , where  $I$  is the set of irrationals in  $[0, 1]$ ,  $X$  is  $\sigma$ -bianalytic. In [7] it was shown that  $X$  could not be embedded in any Hausdorff  $\mathcal{H}_\sigma$  space. Since any  $\mathcal{H}$ -Borelian subset of a Hausdorff space is necessarily contained in a  $\mathcal{H}_\sigma$ ,  $X$  is not a  $\mathcal{H}$ -Borelian subset of any Hausdorff space.

However, the following is true.

PROPOSITION 1.16. *A  $\mathcal{H}_\sigma$ -fiable (respectively regular)  $\sigma$ -bianalytic space is a  $\mathcal{H}$ -Borelian subset of any of its  $\mathcal{H}_\sigma$ -fications (respectively compactifications).*

*Proof.* Let  $X = \bigcup X_i$  where  $X_i$  is bianalytic and let  $K = \bigcup_{i=1}^{\infty} K_i$  be a  $\mathcal{K}_\sigma$ -fication of  $X$ . Now in [5] Frolik showed that closed subspaces of bianalytic spaces are bianalytic and that any bianalytic space is a Baire subset and consequently a  $\mathcal{K}$ -Borelian subset of any of its compactifications. Let  $X = \bigcup_{i,j=1}^{\infty} (X_i \cap K_j)$ . Each  $X_i \cap K_j$  is a  $\mathcal{K}$ -Borelian subset of  $K$ .

## 2. 1 – 1 continuous images of $\mathcal{K}$ -Borelian subsets of Hausdorff spaces.

**THEOREM 2.1.** *There exists a Hausdorff space  $X$  which is the 1 – 1 continuous image of a  $\mathcal{K}$ -Borelian subset  $A$  of a compact space  $Y$ , such that  $X$  is not a  $\mathcal{K}$ -Borelian subset of any Hausdorff space in which it can be embedded.*

*Proof.* Let  $X$  be the Hausdorff space  $([0, 1], \mathcal{T})$  defined in Proposition 1.15. We have shown that  $X$  is not a  $\mathcal{K}$ -Borelian subset of any Hausdorff space in which it can be embedded. It remains to show that  $X$  is the 1 – 1 continuous image of a  $\mathcal{K}$ -Borelian subset  $A$  of a compact space  $Y$ . Let  $A = \sum_{i=0}^{\infty} A_i$ ; where  $A_0$  = the irrationals in  $[0, 1]$  and  $A_i = \{q_i\}$  for each  $i \geq 1$ ,  $A_i \cap A_j = \emptyset$  ( $i \neq j$ ) and  $\bigcup_{i=1}^{\infty} \{q_i\} = Q$  = the rationals in  $[0, 1]$ .  $A$  is obviously a  $\mathcal{K}_\sigma$  subset of the one point compactification  $Y$  of the space  $\sum_{i=0}^{\infty} B_i$ ; where  $B_0 = [0, 1]$  endowed with the usual topology and  $B_i = A_i$  for all  $i \geq 1$ .  $X$  is clearly a 1 – 1 continuous image of  $A$ .

**THEOREM 2.2.** *A  $\mathcal{K}_\sigma$ -fiable 1 – 1 continuous image  $Y$  of a  $\sigma$ -bianalytic space  $X$  is a  $\mathcal{K}$ -Borelian subset of any of its  $\mathcal{K}_\sigma$ -fications. In particular, a regular 1 – 1 continuous image of a  $\sigma$ -bianalytic space is a  $\mathcal{K}$ -Borelian subset of any of its  $\mathcal{K}_\sigma$ -fications or compactifications.*

*Proof.* Let  $P$  be some Hausdorff  $\mathcal{K}_\sigma$ -fication of  $Y$ . Then  $P = \bigcup_{n=1}^{\infty} K_n$ , where  $K_n$  is a compact subset of  $P$  for each  $n$ .

Since  $X$  is  $\sigma$ -bianalytic,  $X = \bigcup_{i=1}^{\infty} X_i$  where each  $X_i$  is bianalytic.

For each  $i$  and  $n$ , let

$$Y_{i,n} = f(X_i) \cap (K_n \cap Y).$$

Now each  $Y_{i,n}$  is completely regular and  $Y = \bigcup_{i,n=1}^{\infty} Y_{i,n}$ . Since closed subspaces of bianalytic spaces are bianalytic, then each  $Y_{i,n}$  is a 1 – 1 continuous image of a bianalytic space. Therefore, from Theorem 13 [3] it follows that  $Y_{i,n} \in \mathcal{B}(P)$  for all  $i, n$ .

It follows from Theorem 11 [3] that  $Y_{i,n}$  is an element of the family consisting of countable intersections of countable disjoint unions of sets of the form  $(F \cap B)$ , where  $F$  is a closed set in  $P$  and  $B \in \mathcal{B}(\mathcal{K}(P))$ . That is, for each  $i$  and  $n$ ,

$$Y_{i,n} = \bigcap_k \bigcup_j (F_{kj}^{in} \cap B_{kj}^{in}) \quad \text{where}$$

$F_{kj}^{in}$  is a closed set in  $P$  and consequently a  $\mathcal{K}_\sigma$ ,

$B_{kj}^{in}$  is a Baire subset of  $P$  and consequently a  $\mathcal{K}$ -Borelian subset of  $P$ .

Therefore  $Y = \bigcup_{i,n=1}^\infty Y_{i,n}$  is a  $\mathcal{K}$ -Borelian subset of  $P$ .

The previous theorem can be restated as follows.

**THEOREM 2.3.** *Let  $Y$  be a 1-1 continuous of image of a  $\sigma$ -bianalytic space  $X$ . Then  $Y$  is a  $\mathcal{K}$ -Borelian subset of some Hausdorff space if and only if*

- (1)  $Y$  is Hausdorff;
- (2) there exists a sequence  $\{R_n\}_{n=1}^\infty$  of subsets  $Y$  such that
  - (i)  $Y = \bigcup_{n=1}^\infty R_n$ ,

and for each  $n$ ,

- (ii)  $R_n \subset R_{n+1}$ , and

(iii)  $R_n$  is closed and strongly regular with respect to  $Y$ ; that is,  $R_n$  is a closed subspace of  $Y$  and for each closed set  $C \subset R_n$  and for each point  $y \in (R_n \setminus C)$  there exist sets  $O, U$  open in  $Y$  such that  $y \in O, C \subset U$  and  $O \cap U = \emptyset$ .

In fact, if conditions (1) and (2) are satisfied,  $Y$  is a  $\mathcal{K}$ -Borelian subset of any of its Hausdorff  $\mathcal{K}_\sigma$ -fications.

*Proof.* Necessity of (1) and (2).

Let us suppose that  $Y$  is a  $\mathcal{K}$ -Borelian subset of some Hausdorff space. Then  $Y$  is Lindelof and  $Y$  is a subset of some Hausdorff  $\mathcal{K}_\sigma$  space. It follows from Theorem 3.1 [8] that conditions (1) and (2) are satisfied.

**COROLLARY 2.4.** *A  $\mathcal{K}_\sigma$ -fiable (respectively regular) 1-1 continuous image of a  $\mathcal{K}$ -Borelian subset of a perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$  space is a  $\mathcal{K}$ -Borelian subset of any of its  $\mathcal{K}_\sigma$ -fications (respectively compactifications).*

The following result in the case where the image was regular was first proved by Sion in [9].

**COROLLARY 2.5.** *Let  $X$  be a Hausdorff space which has property*

*I. Then a  $\mathcal{K}_\sigma$ -fiable (respectively regular)  $1 - 1$  continuous image of a  $\mathcal{K}$ -Borelian subset of  $X$  is a  $\mathcal{K}$ -Borelian subset of any of its  $\mathcal{K}_\sigma$ -fications (respectively compactifications).*

*Proof.* If  $Y$  is a  $\mathcal{K}$ -Borelian subset of  $X$  then  $Y$  is contained in a  $\mathcal{K}_\sigma$  subset  $Z$  of  $X$ . In [10] it was shown that a space  $Z$  with property I satisfies the following condition: for all  $A \in \mathcal{K}(Z)$  and  $B$  open in  $Z$ ,  $A \cap B \in \mathcal{K}_\sigma(Z)$ . Since  $Z$  is also a  $\mathcal{K}_\sigma$  then property I implies that it is perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$ .

**COROLLARY 2.6.** *A regular  $1 - 1$  continuous image  $X$  of a  $\mathcal{K}$ -Borelian subset of a metrizable  $\mathcal{K}_\sigma$  space is a  $\mathcal{K}$ -Borelian subset of any compactification of  $X$  or of any Hausdorff  $\mathcal{K}_\sigma$  space containing  $X$ .*

Theorem 1.10 can now be applied to regular locally metrizable spaces.

**COROLLARY 2.7.** *Let  $X$  be a regular locally metrizable space and let  $A$  be a  $\mathcal{K}$ -Borelian subset of  $X$ . A  $\mathcal{K}_\sigma$ -fiable (respectively regular)  $1 - 1$  continuous image of  $A$  is a  $\mathcal{K}$ -Borelian subset of any of its  $\mathcal{K}_\sigma$ -fications (respectively compactifications).*

*Proof.* If  $A$  is a  $\mathcal{K}$ -Borelian subset of  $X$ , then  $A$  is contained in a  $\mathcal{K}_\sigma$  of  $X$ . This  $\mathcal{K}_\sigma$  is Lindelof and therefore paracompact [2]. It follows from [11] that this  $\mathcal{K}_\sigma$  is metrizable. The result now follows from Corollary 2.6.

**3.  $\mathcal{K}$ -Borelian subsets of Hausdorff  $\mathcal{K}_\sigma$  spaces and their complete sequences of countable covers.** It would be interesting to characterize for each  $\alpha$  the Hausdorff spaces that can be embedded as  $\mathcal{K}$ -Borelian subsets of class  $\alpha$  of some Hausdorff  $\mathcal{K}_\sigma$  space. Theorem 3.1 answers this question in the particular case of a  $\mathcal{K}_{\sigma\sigma}$ . This is a generalization of Frolik's result in [4] for completely regular spaces.

Theorem 3.3 gives necessary conditions for a subset of a perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$  space to be a  $\mathcal{K}$ -Borelian subset of class  $\alpha$ . This theorem is proved by a method similar to that used for Theorem 9 of [4].

Both these theorems depend upon the notion of a complete sequence of countable covers defined by Frolik in [4].

**THEOREM 3.1.** *Let  $X$  be a Hausdorff space. Then  $X$  can be embedded in some Hausdorff space  $Y$  as a  $\mathcal{K}_{\sigma\sigma}$  subset if and only*



if the following two conditions are satisfied.

(1) There exists a sequence  $\{X_n\}_{n=1}^\infty$  of subspaces of  $X$  such that

(i)  $X = \bigcup_{n=1}^\infty X_n$ ,

and for each  $n$ ,

(ii)  $X_n \subset X_{n+1}$ ,

(iii)  $X_n$  is closed and strongly regular with respect to  $X$ .

(2) There exists a complete sequence  $\mu = \{\mathcal{H}_j\}_{j=1}^\infty$  of countable closed covers of  $X$  (that is, each member of each cover is closed in  $X$ ).

*Proof.* Let us suppose that  $X$  can be embedded in some Hausdorff space as a  $\mathcal{K}_o$  subset. Then  $X$  can necessarily be embedded in some Hausdorff  $\mathcal{K}_o$  space and condition (1) now follows from Proposition 1, [7]. Condition (2) follows from Proposition 4, [4].

Now, conversely, let us suppose that conditions (1) and (2) are satisfied. Condition (2) implies that for each  $n$   $\{\mathcal{H}_j \cap X_n\}_{j=1}^\infty$  is a complete sequence of countable closed covers of  $X_n$ . Now, since  $X_n$  is regular and has a complete sequence of countable closed covers, it follows from Proposition 3, [4] that  $X_n$  is Lindelof. Therefore  $X = \bigcup_{n=1}^\infty X_n$  is Lindelof and we have by Theorem 3.1 in [8] that  $X$  is embeddable in a Hausdorff  $\mathcal{K}_o$  space  $Y = (\bigcup_{n=1}^\infty K_n, \tau)$  where, for each  $n$ ,  $K_n = \beta X_n$  and  $K_n \subset K_{n+1}$ . (The topology  $\tau$  was described in [8].)

Let us consider the following subspace of  $Y$ :

$$A = \bigcap_{j=1}^\infty \bigcup_{n=1}^\infty \bigcup_{i=1}^\infty (\overline{H_{ji} \cap X_n})^{K_n},$$

where each  $\mathcal{H}_j$  in  $\mu$  consists of  $\{H_{ji}\}_{i=1}^\infty$ ; the  $X_n$  satisfy condition (1);  $K_n = \beta X_n$  for each  $n$ ; and  $(\overline{\phantom{x}})^{K_n}$  represents the closure of  $(\phantom{x})$  in  $K_n$ .

Now  $A$  is obviously a  $\mathcal{K}_o$  subset of  $Y$  and  $A$  contains  $X$ . Let us suppose that there exists  $y \in (A \setminus X)$ . Since  $y \in Y$ , then either  $y \in K_1$  or else there exists an  $n_0$  such that  $y \in (K_{n_0} \setminus K_m)$  for all  $m < n_0$ . (If  $y \in K_1$ , we shall let  $n_0 = 1$ .) Now let  $\mathcal{D}$  be the family of all closed neighborhoods of  $y$  in  $K_{n_0}$ ; let  $\mathcal{E}$  be the family of all closed subsets  $F$  of  $X$  such that  $y \in \bar{F}^{K_n}$  (the closure of  $F$  in  $K_n$ ) for some  $n \geq n_0$ . Clearly,  $\mathcal{D} \cap X = \{D \cap X : D \in \mathcal{D}\}$  is a subfamily of  $\mathcal{E}$ .

We claim that  $\mathcal{E}$  is a  $\mu$ -Cauchy family. To show that  $\mathcal{E}$  has the finite intersection property, let  $F_1, F_2$  be members of  $\mathcal{E}$ . Therefore,  $y \in (\bar{F}_1^{K_m} \cap \bar{F}_2^{K_n} \cap K_{n_0})$  for some  $m \geq n_0$  and  $n \geq n_0$ . Therefore, as it was shown in [8],  $y \in ((\bar{F}_1 \cap X_{n_0})^{K_{n_0}} \cap (\bar{F}_2 \cap X_{n_0})^{K_{n_0}})$ . From Lemma 2.1, [8] we have that  $X_{n_0}$  is strongly normal with respect to  $X$  and consequently  $X_{n_0}$  is a normal subspace of  $X$ . Since disjoint closed subsets of a normal space have disjoint closures in their Stone-Cech

compactifications, it follows that  $F_1 \cap F_2 \neq \emptyset$ . Thus  $\mathcal{E}$  has the finite intersection property.

Since  $y \in A$ , then for all  $j$  there exists an  $i_j$  and an  $n_j$  such that  $(X_{n_j} \cap H_{ji_j}) \in \mathcal{E}$ . Thus  $\mathcal{E}$  is a  $\mu$ -Cauchy family. Since  $\mu$  is complete  $\bigcap_{E \in \mathcal{E}} \bar{E}^X \neq \emptyset$ . This is impossible because  $\bigcap_{E \in \mathcal{E}} \bar{E}^X \subset \bigcap_{E \in \mathcal{E}} \bar{E}^Y \subset \bigcap_{D \in \mathcal{D}} \bar{D}^Y = \bigcap_{D \in \mathcal{D}} \bar{D}^{K_{n_0}} = \{y\}$ , and  $y \in (A \setminus X)$ . This contradiction proves that  $A = X$ .

As a corollary we have the following result due to Frolik (Theorem 7, [4]).

**COROLLARY 3.2.** *A completely regular space  $X$  is a  $\mathcal{K}_\alpha$  subset of some Hausdorff  $\mathcal{K}_\alpha$  space if and only if  $X$  has a complete sequence of countable closed covers. In particular, a completely regular space  $X$  is a  $\mathcal{K}_\alpha$  subset of its Stone-Cech compactification  $\beta X$  if and only if  $X$  has a complete sequence of countable closed covers.*

*Proof.* Let  $X_n = X$  for each  $n$  in condition (1) of Theorem 3.5 and the proof follows immediately.

**THEOREM 3.3.** *Let  $X$  be a perfectly  $\sigma$ -normal  $\mathcal{K}_\alpha$  Hausdorff space. If  $Y$  is a  $\mathcal{K}$ -Borelian subset of  $X$  of class  $\alpha$ , then  $Y$  has a complete sequence of countable  $\mathcal{F}$ -Borelian covers of at most class  $\alpha$ .*

*Proof.* We shall proceed by transfinite induction. For the case  $\alpha = 2$  the result follows from Corollary 3.6.

From the assumption the result is true for all  $\beta < \alpha$ , we must prove the result is true for  $\alpha$  in the following three cases.

*Case (1).*  $\alpha = \alpha_0$  where  $\alpha_0$  is a limit ordinal.

Since  $Y$  is a  $\mathcal{K}$ -Borelian subset of  $X$  of class  $\alpha_0$  then  $Y = \bigcap_{n=1}^{\infty} Y_n$  where each  $Y_n$  is a  $\mathcal{K}$ -Borelian subset of  $X$  of class  $\beta_n < \alpha_0$ . Thus each  $Y_n$  has a complete sequence  $\mu_n = \{\mathcal{H}_j^{n^*}\}_{j=1}^{\infty}$  of countable  $\mathcal{F}$ -Borelian covers of class at most  $\beta_n$ . Let us consider the sequence  $\mu = \{\mathcal{H}_j^{n^*} \cap Y\}_{n,j=1}^{\infty}$ . Now  $\mu$  is a sequence of countable  $\mathcal{F}$ -Borelian covers of  $Y$  of at most class  $\alpha_0$  and  $\mu$  is complete from the proof of Theorem 9, [4].

*Case (2).*  $\alpha = \alpha_0 + m$  (where  $\alpha_0$  is a limit ordinal and  $m$  is odd).

Since  $Y$  is a  $\mathcal{K}$ -Borelian subset of  $X$  of class  $\alpha = \alpha_0 + m$ , then  $Y = \bigcup_{n=1}^{\infty} Y_n$  where each  $Y_n$  is a  $\mathcal{K}$ -Borelian subset of  $X$  of class  $(\alpha - 1)$  and each  $Y_n$  has a complete sequence  $\mu_n = \{\mathcal{H}_j^{n^*}\}_{j=1}^{\infty}$  of countable  $\mathcal{F}$ -Borelian covers of at most class  $(\alpha - 1)$ .

Let us define for each  $n$  the following sequence  $\{T_n\}_{n=1}^{\infty}$  of subsets of  $X$ :  $T_0 = \emptyset$ ,

$$T_n = \bigcup_{k=1}^n Y_k \quad \text{for all } n \geq 1.$$

Let us consider the following sequence  $\mu = \{\mathcal{M}_j\}_{j=1}^\infty$  of countable covers of  $Y$  where for each  $j$ :

$$\begin{aligned} \mathcal{M}_j &= \{H_{ji}^n \cap (Y_n \setminus T_{n-1})\}_{n,i=1}^\infty, \quad \text{where for each } n \text{ and } j, \\ \mathcal{H}_j^n &= \{H_{ji}^n\}_{i=1}^\infty. \end{aligned}$$

Since  $X$  is perfectly  $\sigma$ -normal  $\mathcal{K}_\sigma$  we can easily verify that, for all  $j$ , each element of  $\mathcal{M}_j$  is a  $\mathcal{F}$ -Borelian subset of  $Y$  of at most class  $\alpha$ .

To show that  $\mu$  is complete, let  $\mathcal{E}$  be a maximal  $\mu$ -Cauchy family. Clearly there exists an  $n$  such that  $\mathcal{E}$  is a  $\mu_n$ -Cauchy family. By Proposition 2, [4],  $\emptyset \neq \bigcap_{E \in \mathcal{E}} \bar{E}^X \subset Y_n \subset Y$ . Therefore,  $\bigcap_{E \in \mathcal{E}} \bar{E}^Y \neq \emptyset$  and  $\mu$  is complete.

*Case (3).*  $\alpha = \alpha_0 + m$  (where  $\alpha_0$  is a limit ordinal and  $m$  is even). We can use the same method of proof as for Case (1).

## REFERENCES

1. G. Choquet, *Lectures on Analysis*, Volume 1, New York, 1969.
2. J. Dugundji, *Topology*, Boston, 1966.
3. Z. Frolik, *A contribution to the descriptive theory of sets and spaces*, General Topology and its Relations to Modern Analysis and Algebra. Proc. Symp. Prague Sept. (1961), 157-173 (New York, Academic Press).
4. ———, *On the descriptive theory of sets*, Czech. Math. J., **13** (1963), 335-359.
5. ———, *On bianalytic spaces*, Czech. Math. J., **13** (1963), 561-573.
6. K. Kuratowski, *Topologie I*, Warszawa, 1952.
7. B. MacGibbon, *Exemple d'espace  $K$ -analytique qui n'est  $K$ -Souslinien dans aucun espace*, Bull. Sci. Math., **94** (1970), 3-4.
8. ———, *A necessary and sufficient condition for the embedding of a Lindelof space in a Hausdorff  $K_\sigma$  space*, Pacific J. Math., **35** (1970), 459-465.
9. M. Sion, *On analytic sets in topological spaces*, Trans Amer. Math. Soc., **96** (1960), 341-354.
10. ———, *Continuous images of Borel sets*, Proc. Amer. Math. Soc., **11** (1961), 385-391.
11. Y. M. Smirnov, *On metrization of topological spaces*, Uspehi Matem. Nauk (N.S.) 6, No. 6 (46), 100-111 (1951), (Translations of AMS Series 1, Vol. **18**, 62-77).

Received January 14, 1979. The author gratefully acknowledges the support of the Canadian National Research Council during the preparation of this paper.

CONCORDIA UNIVERSITY  
MONTREAL, QUEBEC, CANADA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

DONALD BABBITT (Managing Editor)

University of California  
Los Angeles, CA 90024

HUGO ROSSI

University of Utah  
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California  
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University  
Stanford, CA 94305

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$102.00 a year (6 Vols., 12 issues). Special rate: \$51.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1981 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Thomas E. Armstrong and William David Sudderth, Nearly strategic measures .....	251
John J. Buoni, Artatrana Dash and Bhushan L. Wadhwa, Joint Browder spectrum .....	259
Jack Paul Diamond, Hypergeometric series with a $p$ -adic variable .....	265
Raymond Frank Dickman, Jack Ray Porter and Leonard Rubin, Completely regular absolutes and projective objects .....	277
James Kenneth Finch, On the local spectrum and the adjoint .....	297
Benno Fuchssteiner, An abstract disintegration theorem .....	303
Leon Gerber, The volume cut off a simplex by a half-space .....	311
Irving Leonard Glicksberg, An application of Wermer's subharmonicity theorem .....	315
William Goldman, Two examples of affine manifolds .....	327
Yukio Hirashita, On the Weierstrass points on open Riemann surfaces .....	331
Darrell Conley Kent, A note on regular Cauchy spaces .....	333
Abel Klein and Lawrence J. Landau, Periodic Gaussian Osterwalder-Schrader positive processes and the two-sided Markov property on the circle .....	341
Brenda MacGibbon, $\mathcal{H}$ -Borelian embeddings and images of Hausdorff spaces .....	369
John R. Myers, Homology 3-spheres which admit no PL involutions .....	379
Boon-Hua Ong, Invariant subspace lattices for a class of operators .....	385
Chull Park, Representations of Gaussian processes by Wiener processes .....	407
Lesley Millman Sibner and Robert Jules Sibner, A sub-elliptic estimate for a class of invariantly defined elliptic systems .....	417
Justin R. Smith, Complements of codimension-two submanifolds. III. Cobordism theory .....	423
William Albert Roderick Weiss, Small Dowker spaces .....	485
David J. Winter, Cartan subalgebras of a Lie algebra and its ideals. II .....	493