

Pacific Journal of Mathematics

**THE HOMOTOPY GROUPS OF KNOTS. I. HOW TO COMPUTE
THE ALGEBRAIC 2-TYPE**

SAMUEL JAMES LOMONACO, JR.

THE HOMOTOPY GROUPS OF KNOTS I. HOW TO COMPUTE THE ALGEBRAIC 2-TYPE

S. J. LOMONACO, JR.

Dedicated to the Memory of R. H. Fox

Let K be a CW complex with an aspherical splitting, i.e., with subcomplexes K_- and K_+ such that (a) $K = K_- \cup K_+$ and (b) $K_-, K_0 = K_- \cap K_+, K_+$ are connected and aspherical. The main theorem of this paper gives a practical procedure for computing the homology $H_*\tilde{K}$ of the universal cover \tilde{K} of K . It also provides a practical method for computing the algebraic 2-type of K , i.e., the triple consisting of the fundamental group $\pi_1 K$, the second homotopy group $\pi_2 K$ as a $\pi_1 K$ -module, and the first k -invariant kK .

The effectiveness of this procedure is demonstrated by letting K denote the complement of a smooth 2-knot (S^4, kS^2) . Then the above mentioned methods provide a way for computing the algebraic 2-type of 2-knots, thus solving problem 36 of R. H. Fox in his 1962 paper, "Some problems in knot theory." These methods can also be used to compute the algebraic 2-type of 3-manifolds from their Heegaard splittings. This approach can be applied to many other well known classes of spaces. Various examples of the computation are given.

Table of Contents

0. Introduction	349
I. Brief historical development.....	352
II. The diagram of a 2-knot	352
III. Group systems	356
IV. Chain complex basics	358
V. Free resolution systems	360
VI. Completions of chain complexes	363
VII. The main theorem (Theorem 7.1)	364
VIII. Fox-Lyndon resolutions	368
IX. A constructive form of the main theorem (Theorem 9.2).....	369
X. What is the significance of the algebraic 2-types of 2-knots?....	373
XI. Miscellaneous corollaries on 2-knots	375
XII. Conclusion: Problems and more problems	376
Appendix A. A computing manual for algebraic 2-type calculations....	378
Appendix B. Explicit calculations for 2-knots	383
References.....	387

0. Introduction. Let K be a CW complex with an aspherical splitting. An *aspherical splitting* of a CW complex K is a triad (K, K_-, K_+) of CW complexes such that (a) $K = K_- \cup K_+$ and (b)

K_- , $K_0 = K_- \cap K_+$, K_+ are connected and aspherical.¹ The main theorem (Theorem 7.1) of this paper gives an effective procedure for computing the singular homology $H_*\tilde{K}$ of the universal cover \tilde{K} of K from the groups π_1K_- , π_1K_0 , π_1K_+ and the morphisms $g_{\pm}: \pi_1K_0 \rightarrow \pi_1K_{\pm}$ induced by inclusion. It also provides a practical method for computing the *algebraic 2-type*² of K , i.e., the triple consisting of the fundamental group π_1K , the second homotopy group π_2K as a π_1K -module, and the first k -invariant kK lying in the cohomology group $H^3(\pi_1K; \pi_2K)$.

The effectiveness of this procedure is demonstrated by letting K denote the complement of a smooth (or locally flat piecewise linear (PL)) 2-knot (S^4, kS^2) . Then the above mentioned method provides a way of computing the algebraic 2-type of 2-knots from their *hyperbolic splittings* (def. 2.1). This solves problem 36 of R.H. Fox in his 1962 paper [22], and substantially more.

The effectiveness of this procedure is also demonstrated by letting K denote a closed connected 3-manifold. Then the algebraic 2-type of K can be computed from any one of its Heegaard splittings (of positive genus). The methods of this paper apply to complements of unsplitable 2-links and to many other types of spaces. Various examples of this computational procedure are given in Appendix B.

A 2-knot is said to be *quasi-aspherical* (QA) if the third homology group of the universal cover of its complement K vanishes (def. 10.1). It is demonstrated that the homotopy type of the complement K of a QA 2-knot is completely determined by the algebraic 2-type of K . This suggests a possible analogue in dimension four of Papakyriakopoulos' theorem on the asphericity of 1-knots [53]. (See Problem 1 in § XII.)

It is demonstrated that, for every homology 1-sphere K (hence, for all n -knot complements), $H_3(\pi_1K) = \pi_2K / \mathcal{F}\pi_2K = (Z \otimes_{\pi_1K} \pi_2K)$, where \mathcal{F} denotes the augmentation ideal of π_1K . (This can be thought of as a generalization of the Kervaire condition $H_2(\pi_1K) = 0$. See [36].) It follows that the first k -invariant of a homology 1-sphere K does not vanish if $H_3(\pi_1K) \neq 0$; hence, that the first k -invariant of a fibered 2-knot is nonzero if its commutator subgroup is nontrivial and finite.

NOTATION. Throughout this paper (unless stated otherwise) (S^{n+2}, kS^n) will always denote a smooth n -knot, (if $n = 2$, it can be

¹ In [43] K is called a "generalized Eilenberg-MacLane (GEM) complex. From [43], the homotopy type of the GEM complex K is completely determined by the group system $G = \{\pi_1K_- \leftarrow \pi_1K \rightarrow \pi_1K_+\}$. Hence, the notation $K = K(G, 1)$. For details, see [43].

² In [48] S. MacLane and J.H.C. Whitehead call this the algebraic 3-type.

locally flat PL). By the *exterior* of an n -knot (S^{n+2}, kS^n) is meant S^{n+2} with a small open tubular neighborhood of kS^n removed. The *complement* is S^{n+2} with kS^n removed.

Note to reader. This paper has been written to be read in any one of three modes, i.e., as

Mode 1. A mathematical treatise. Read §§ I through XII. To understand only the main theorem (Theorem 7.1), read the caveat below and §§ III through VII.

Mode 2. A “handy man’s manual” for computing algebraic 2-type. In this mode only a rudimentary understanding of [11] is needed. To compute the algebraic 2-type of:

(i) 2-knot complements, then read only § II, Theorem A 2.1, Observations 1 through 3 and the Hint (all of Appendix A), and Appendix B.

(ii) 3-manifolds, then read only Theorem A 2.1, Observations 1 through 3 and the Hint (all of Appendix A).

(iii) Arbitrary spaces with aspherical splittings, then read all of Appendices A and B.

Mode 3. A paper to browse through on a lazy Sunday afternoon. Use the table of contents and begin by browsing in any section. All terms not defined within a section are referenced back to their place of definition.

This paper started out to be a condensed and revised version of [41, 42]. To the author’s surprise, it has developed into much more. It basically contains all the material of [41, 42] with the exception of the section on identities. It also contains as its main theorem, Theorem 7.1, a distillate of the computational method hidden within the recesses of [41, 42]. This clarification makes it possible, not only to compute the entire algebraic 2-type of 2-knot complements, but of a much larger class of spaces.

Caveat. The direct limit functor \lim_{\rightarrow} defined in § V does not commute with the homology functor.

ACKNOWLEDGMENT. I would like to thank John Harper for his helpful discussions on the k -invariant. I would also like to thank John Milnor for pointing out that, as an immediate consequence of [23], all the hyperbolic points of a smooth (locally flat PL) 2-knot can be pushed into the same 3-dimensional hyperplane (see Theorem

2.1). Finally, I would like to thank the referee for his helpful comments.

I. **Brief historical development.** The development of the work on the homotopy groups of knots had its beginnings in the early 1900's when Dehn and Wirtinger found a method for computing the fundamental group of 1-knots. Later in 1957, Papakyriakopoulos [53] determined all the higher homotopy groups of 1-knots by proving his famous theorem on the asphericity of knots. This theorem essentially says that all the higher homotopy groups of 1-knots vanish, and hence, that the homotopy type of a 1-knot complement is completely determined by its algebraic 1-type (i.e., by its fundamental group).

Mathematicians then turned to higher dimensional knots, seeking methods for computing their homotopy groups and looking for a higher dimensional analogue of Papakyriakopoulos' theorem on the asphericity of knots. In 1962 Fox [21] gave a way to compute the fundamental group of 2-knots from their motion picture representations. He then asked in problem 36 of [22] for a method for computing the second homotopy group (as a π_1 -module) of locally flat PL 2-knots.

In this paper, a method is given for computing the second homotopy group as a π_1 -module of smooth (or locally flat PL) 2-knots, thus solving problem 36 of Fox. We then ask if the homotopy type, of a 2-knot complement is determined completely by its algebraic 2-type. Some evidence (Theorem 10.1) is given suggesting that this might indeed be true. A method for computing the complete algebraic 2-type of smooth (or locally flat PL) 2-knots is given.

REMARK. Please refer to the bibliography for a more complete historical development. (Also see [37] and [62].)

II. **The diagram of a 2-knot.** If we are to compute the algebraic 2-type of a 2-knot, we will first need a convenient way of representing such knots. We will utilize Fox's motion picture representation of knots, i.e., a representation in terms of a parameterized family of 3-dimensional cross-sections [21]. We will then carry this method one step further by showing that essentially only one single 3-dimensional cross-section, the so called 0-level, is needed.

As mentioned in §0, (S^4, kS^2) will denote a smooth (or locally flat PL) 2-knot. Throughout this section K will denote its *exterior*, i.e., S^4 with a small open tubular neighborhood of kS^2 removed.

For a given function $\tau: (S^4, \infty) \rightarrow (S^1, \infty)$ and for each point t in the 1-sphere S^1 , define the t -level as

$$K_t = K \cap [\tau^{-1}(t) \cup \infty].$$

If the *time function* τ is suitably chosen, then the t -levels K_t are the 3-dimensional cross-sections mentioned above.

There is no need to consider arbitrary time functions τ . More particularly, we will show that we need only consider τ of the following type.

DEFINITION 2.1. A smooth (or PL) map $\tau: (S^4, \infty) \rightarrow (S^1, \infty)$ such that $\tau^{-1}(\infty) = \infty$ is a *hyperbolic time function* of an exterior K of a 2-knot (S^4, kS^2) provided:

- (1) $\tau|_{kS^2}$ has only finitely many critical points, all of which are nondegenerate and none of which is the point at ∞ .
- (2) All elliptic points occur at levels K_{-1} and K_{+1} .
- (3) All hyperbolic points occur at level K_0 .

The spaces $K_- = \bigcup_{t \geq 0} K_t$ and $K_+ = \bigcup_{t \leq 0} K_t$ are called the *bottom* and *top halves* of K respectively. The triad (K, K_-, K_+) is called a *hyperbolic splitting* of the knot.

It now follows from the work of Fox and Milnor [23] that:

THEOREM 2.1. *Every smooth (or locally flat PL) 2-knot has a hyperbolic time function, and hence, a hyperbolic splitting.*

A hyperbolic time function of the trivial 2-knot is shown in Figure 1. One of the spun trefoil is given in Figure 2.

A quick glance at Figures 1 and 2 will convince the reader that




FIGURE 1. Hyperbolic time function of the trivial 2-knot.




FIGURE 2. Hyperbolic time function of the spun trefoil.

the above method of representation is highly redundant. The entire knot can be completely reproduced from the 0-level K_0 and a set of labels (one for each hyperbolic point) indicating how the hyperbolic points open up above, i.e., for $t > 0$. We are thus led to the following convenient and compact representation of 2-knots.

DEFINITION 2.2. Let K_0 be a 0-level of the exterior K of a smooth (or locally flat PL) 2-knot (S^4, kS^2) arising from a hyperbolic time function. Then a *diagram* of (S^4, kS^2) is a regular projection of K_0 with over and under crossings indicated in the standard way and with hyperbolic points labeled as shown in Figure 3.

By Theorem 2.1, all smooth (or locally flat PL) 2-knots have a diagram. Diagrams of the trivial 2-knot and of the spun trefoil are




FIGURE 3. Interpretation of hyperbolic point labels.




FIGURE 4. Diagram of trivial 2-knot.




FIGURE 5. Diagram of Example 1, the spun trefoil.

$$\pi_1 K = \langle a, b : abab^{-1}a^{-1}b^{-1} = 1 \rangle$$

$$\pi_1 K = (\partial U_{\square} : (1-a+ba)\partial U_{\square} = 0)$$

$$H^3(\pi_1 K; \pi_2 K) = 0 \text{ and } \mathcal{K}K = 0$$




FIGURE 6. The diagram of Example 2.

$$\pi_1 K = (a, x: a^3, ax\bar{a}^2\bar{x}) \text{ and } \pi_2 K = (\partial U_\square: (x-1)\partial U_\square = 0 = (\sum_{i=0}^{2n} a^i)\partial U_\square)$$

$$H^8(\pi_1 K; \pi_2 K) = Z_{2n+1} = (t: t^{2n+1} = 1) \text{ and } kK = t$$

shown in Figures 4 and 5 respectively. Diagrams of other 2-knots are given in Figures 6 and 7.

III. Group systems. In this section group systems and their direct limits are defined. The Van Kampen theorem is also stated. (See [10] for a more general formulation.)

A group system G is a set of three groups G_-, G_0, G_+ together with two group morphisms $g_\pm: G_0 \rightarrow G_\pm$. If g_- and g_+ are both epimorphisms (monomorphisms), then G is called an epimorphic (monomorphic) group system. (The phrase "group system" is used rather than "group triad" because all definitions and results hold for a more general class of group systems.)

If G and G' are group systems, then a morphism $f: G \rightarrow G'$




FIGURE 7. The diagram of Example 3.

$$\pi_1 K = (a, x: ax = xa^2) \text{ and } \pi_2 K = (\partial U_{\square}: (1+a-ax)\partial U_{\square} = 0)$$

$$H^3(\pi_1 K; \pi_2 K) = 0 \text{ and } kK = 0$$

from G to G' is a collection of morphisms $f_{\alpha}: G_{\alpha} \rightarrow G'_{\alpha} (\alpha = -, 0, +)$ such that the diagram

$$\begin{array}{ccccc} G_{-} & \longleftarrow & G_0 & \longrightarrow & G_{+} \\ f_{-} \downarrow & & f_0 \downarrow & & \downarrow f_{+} \\ G'_{-} & \longleftarrow & G'_0 & \longrightarrow & G'_{+} \end{array}$$

is commutative. If each morphism in f is an isomorphism onto, then f is said to be an *isomorphism* and G and G' are said to be *isomorphic*.

If G is a group system and G a group, then a *morphism* $f: G \rightarrow G$ from G into G is a collection of morphisms $f_{\alpha}: G_{\alpha} \rightarrow G$ for $\alpha = -, 0, +$ such that the diagram

$$\begin{array}{ccc} & G_0 & \\ g_{-} \swarrow & \downarrow f_0 & \searrow g_{+} \\ G_{-} & & G_{+} \\ f_{-} \swarrow & \downarrow f & \swarrow f_{+} \\ & G & \end{array}$$

is commutative. The *image* of f is the subgroup of G generated by the images of all the morphisms in f . The morphism f is said to be an *epimorphism* if its image is G itself.

DEFINITION 3.1. A *direct limit (push-out)* of a group system G is a group G together with a morphism $f: G \rightarrow G$ such that
 (1) f is an epimorphism.

(2) For every group G' and morphism $f': G \rightarrow G'$, there exists a morphism $h: G \rightarrow G'$ such that $f' = hf$, i.e., $f'_\alpha = hf_\alpha$ for $\alpha = -, 0, +$. The group G will be denoted by $\lim_{\rightarrow} G$ and the morphism f by $g^\infty = \{g_-^\infty, g_0^\infty, g_+^\infty\}$.

THEOREM 3.1 [10]. *Every group system has a direct limit unique to within isomorphism.*

The following gives some insight into the geometric significance of the above material. More will be said in later sections.

DEFINITION 3.2. Let (Q, Q_-, Q_+) be a triad of pathwise connected spaces such that $Q_0 = Q_- \cap Q_+$ is also pathwise connected. Then the group system G associated with (Q, Q_-, Q_+) is the group system consisting of the groups $\pi_1 Q_-, \pi_1 Q_0, \pi_1 Q_+$ together with the morphisms induced by inclusion.

THEOREM 3.2. (Van Kampen) (See [10] for more general version.)

Let (Q, Q_-, Q_+) be a triad of spaces such that

(1) $Q = Q_- \cup Q_+$.

(2) $Q_-, Q_0 = Q_- \cap Q_+, Q_+$ are open pathwise connected subspaces of the pathwise connected space Q .

Let G be the group system associated with (Q, Q_-, Q_+) . Then $\pi_1 Q = \lim_{\rightarrow} G$ and g^∞ is the collection of morphisms $g_\alpha: \pi_1 Q_\alpha \rightarrow \pi_1 Q$ induced by inclusion ($\alpha = -, 0, +$).

REMARK. The main theorem of this paper (Theorem 7.1) can be thought of as a π_2 -generalization of the above Van Kampen theorem.

IV. Chain complex basics. The mature mathematician probably will need only skim this section. The material in this section comes from [15, 47, 65].

If G is a group, then ZG will denote its integral group ring; and if $g: G \rightarrow G'$ is a group morphism, then the same symbol will denote the induced ring morphism $g: ZG \rightarrow ZG'$. By a left (right) G -module is meant a left (right) ZG -module. For all groups G , the infinite cyclic group Z is considered to be a left G -module under the trivial action. We write \otimes_G rather than \otimes_{ZG} . If M and M' are left G -modules, then a G -morphism $\phi: M \rightarrow M'$ means a left ZG -module morphism. If M and M' are left G - and G' -modules respectively, then a map under the group morphism $g: G \rightarrow G'$ is an additive morphism $\phi: M \rightarrow M'$ such that $\phi(xm) = g(x)\phi(m)$ for all $x \in G$ and $m \in M$. (See [65].)

A *G-chain complex* C is a sequence of left G -modules C_q for each integer q together with G -morphisms $\partial_q: C_q \rightarrow C_{q-1}$ such that (1) $C_q = 0$ for $q < 0$ and (2) $\partial_{q+1}\partial_q = 0$ for all q . C is said to be *free* if C_q is a free left G -module for all q . An *augmentation* ε of C is a G -morphism $\varepsilon: C_0 \rightarrow Z$ such that $\varepsilon\partial_0 = 0$ and $\varepsilon(xc_0) = \varepsilon(c_0)$ for all $x \in G$ and $c_0 \in C_0$. C is said to be *augmentable* if there exists an augmentation for C . An augmentable G -chain complex C together with an augmentation is called an *augmented G -chain complex* (which is also denoted by the same symbol C).

If C and C' are augmented G -chain complexes, then a *chain map* $\phi: C \rightarrow C'$ defined in dimensions up to q is a sequence of G -morphisms $\phi_i: C_i \rightarrow C'_i$ $0 \leq i \leq q$ such that

$$\partial'_i\phi_i = \phi_{i-1}\partial_i \quad (1 \leq i \leq q) \text{ and } \varepsilon'\phi_0 = \varepsilon.$$

Given two chain maps $\phi, \psi: C \rightarrow C'$ defined up to dimension q , a *chain homotopy* D from ϕ to ψ in dimensions not greater than q , written

$$D: \phi \cong \psi \quad (\dim \leq q)$$

is a sequence of G -morphisms $D_i: C_i \rightarrow C'_{i+1}$ $0 \leq i \leq q$ such that

$$\psi_i = \phi_i + \partial'_{i+1}D_i + D_{i-1}\partial_i \quad (0 \leq i \leq q),$$

where for $i = 0$, $D_{i-1}\partial_i$ is omitted. If such a chain homotopy exists, then ϕ and ψ are said to be *chain homotopic in dimensions not greater than q* , written $\phi \cong \psi$ ($\dim \leq q$). Two chain maps homotopic in dimensions not greater than q induce the same homomorphism of the integral (reduced) homology groups for $0 \leq i < q$.

A chain map $\phi: C \rightarrow C'$ defined up to dimension q is said to be a *chain equivalence in dimensions not greater than q* if there exists a chain map $\phi': C' \rightarrow C$ defined up to dimension q such that

$$\phi\phi' \cong 1' \quad (\dim \leq q) \text{ and } \phi'\phi \cong 1 \quad (\dim \leq q)$$

where 1 and $1'$ denote the identity chain maps on C and C' respectively. Such a chain equivalence induces an isomorphism of the integral (reduced) homology groups for $0 \leq i < q$.

DEFINITION 4.1. Let C' be a (augmented) free G -chain subcomplex of a (augmented) free G -chain complex C . Then C' is a *proper free subcomplex of C* or C is a *proper extension of C'* provided for all q

- (1) C'_q is a G -direct summand of C_q , and
- (2) C_q/C'_q is a free left G -module.

(Note: The boundary operator ∂ of C does not necessarily respect the direct sum decomposition $C_q = C'_q \oplus (C_q/C'_q)$.)

The following two theorems are a relativized versions of those found in [15].

THEOREM 4.1. (*Relative extension of chain maps.*) Let A, B, C be free G -chain complexes such that B is a proper subcomplex of A . Let $\phi: A \rightarrow C$ be a chain map defined up to dimension q and $\psi: B \rightarrow C$ a chain map defined up to dimension $q + 1$ and such that $\phi_i|_B = \psi_i$ for $i \leq q$. Then ϕ extends to a chain map defined up to dimension $q + 1$ such that $\phi_{q+1}|_B = \psi_{q+1}$ if and only if the $q + 1$ cocycle $\eta_q \psi_q \partial_A|_{A/B}$ lying in $Z^{q+1}(A/B, H_q C)$ vanishes, where $\eta_q: Z_q C \rightarrow H_q C$ denotes the natural morphism.

THEOREM 4.2. (*Relative extension of chain homotopies.*) Let A, B, C be free G -chain complexes such that B is a proper subcomplex of A . Let $\phi, \phi': A \rightarrow C$ be chain maps defined for all q and let $\psi, \psi': B \rightarrow C$ be chain maps defined for all q such that $\phi|_B = \psi$ and $\phi'|_B = \psi'$. If $D: B \rightarrow C$ is a chain homotopy from ψ to ψ' defined up to dimension $q + 1$ and if $D^*: A \rightarrow C$ is a chain homotopy from ϕ to ϕ' defined up to dimension q and such that $D^*|_B = D$, then D^* extends to a chain homotopy from ϕ to ϕ' defined up to dimension $q + 1$ and such that $D_{q+1}^*|_B = D_{q+1}$ if and only if the obstruction

$$\eta_{q+1}(\phi'_{q+1} - \phi_{q+1} - D_q \partial_{q+1})|_{(A/B)_{q+1}}$$

lying in $Z^{q+1}(A/B, H_{q+1} C)$ vanishes, where $\eta_{q+1}: Z_{q+1} C \rightarrow H_{q+1} C$ denotes the natural morphism.

V. Free resolution systems. A more general and slicker formulation of this material is given in [43]. The definitions of §§ III and IV are assumed.

By a *free resolution over a group G* is meant an augmented free G -chain complex C such that

$$\dots \longrightarrow C_q \xrightarrow{\partial_q} C_{q-1} \longrightarrow \dots \longrightarrow C_0 \xrightarrow{\varepsilon} Z \longrightarrow 0$$

is exact. (The infinite cyclic group Z is considered to be a left G -module under the trivial action.)

DEFINITION 5.1. Let G be a group system. (See § III.) A *free resolution system C over G* is a set of free resolutions C^-, C^0, C^+ over G_-, G_0, G_+ respectively together with maps $\gamma^\pm: C^0 \rightarrow C^\pm$ such that

(1) $ZG_\pm \otimes_{G_0} C^0$ is a proper subcomplex of C^\pm . (See Definition 4.1.)

(2) $\gamma_q^\pm: C_q^0 \rightarrow C_q^\pm$ is the map under the morphism $g_\pm: G_0 \rightarrow G_\pm$ defined as the composition of the map of C_q^0 into $ZG_\pm \otimes_{G_0} C_q^0$ given by

$$c_0 \longmapsto 1 \otimes_{G_0} c_0$$

together with the inclusion map of $ZG_\pm \otimes_{G_0} C^0$ into C^\pm .

DEFINITION 5.2. Let C and C' be free resolution systems over a group system G . A chain map $\Phi: C \rightarrow C'$ is a collection of chain maps $\phi^\alpha: C^\alpha \rightarrow C'^\alpha$ ($\alpha = -, 0, +$) such that

$$\begin{array}{ccccc} C^- & \longleftarrow & C^0 & \longrightarrow & C^+ \\ \phi^- \downarrow & & \phi^0 \downarrow & & \downarrow \phi^+ \\ C'^- & \longleftarrow & C'^0 & \longrightarrow & C'^+ \end{array}$$

is commutative. If $\Psi: C \rightarrow C'$ is also a chain map, then a chain homotopy Δ from Φ to Ψ , written $\Delta: \Phi \cong \Psi$, is a collection of chain homotopies $\Delta^\alpha: \phi^\alpha \cong \psi^\alpha$ such that

$$\begin{array}{ccccc} C^- & \longleftarrow & C^0 & \longrightarrow & C^+ \\ \Delta^- \downarrow & & \Delta^0 \downarrow & & \downarrow \Delta^+ \\ C'^- & \longleftarrow & C'^0 & \longrightarrow & C'^+ \end{array}$$


is commutative. If there is a chain equivalence from Φ to Ψ , we say that Φ is chain homotopic to Ψ , written $\Phi \cong \Psi$. A chain map $\Phi: C \rightarrow C'$ is a chain equivalence if there is a chain map $\Phi': C' \rightarrow C$ such that $\Phi\Phi' \cong 1'$ and $\Phi'\Phi \cong 1$, where 1 and $1'$ denote respectively the identity chain maps on C and C' . C and C' are said to be chain equivalent if there exists a chain equivalence from C to C' .

THEOREM 5.1. Any two free resolution systems over the same group system G are chain equivalent.

Proof. Let C and C' be any two free resolution systems over G . Since C^0 and C'^0 are free resolutions over G_0 , there exists a chain equivalence $\phi^0: C^0 \rightarrow C'^0$ which induces a chain equivalence $\bar{\phi}^0$ from $ZG_\pm \otimes_{G_0} C^0$ to $ZG_\pm \otimes_{G_0} C'^0$. Since all of the obstructions of Theorems 4.1 and 4.2 vanish, $\bar{\phi}^0$ extends to a chain equivalence from C^\pm to C'^\pm . □

DEFINITION 5.3. Let G be a group system and C be a free resolution system over G . Let $\bar{G} = \lim G$ and let \bar{C} be an augmented G -chain complex. A chain map $\bar{\Phi}: \bar{C} \rightarrow C$ is a collection of chain

maps $\phi^\alpha: C^\alpha \rightarrow C$ ($\alpha = -, 0, +$) under the respective group morphisms $g_\alpha^\infty: G_\alpha \rightarrow G$ ($\alpha = -, 0, +$) such that



is commutative. The image of Φ is the smallest G -subcomplex of C which contains the image of every morphism in Φ . The chain map Φ is an epimorphism if its image is C itself.

DEFINITION 5.4. Let C be a free resolution system over a group system G with $G = \lim G$. A direct limit (push-out) of C is an augmented G -complex \overrightarrow{C} and a chain map $\Phi': C \rightarrow \overrightarrow{C}$ such that

(1) Φ' is an epimorphism and,

(2) For every augmented G -complex C' and chain map $\Phi': C \rightarrow C'$, there exists a chain map $p: \overrightarrow{C} \rightarrow C'$ such that $\Phi' = p\Phi$.

Such an augmented G -complex \overrightarrow{C} will be denoted by $\lim C$ and Φ by Φ^∞ .

The direct limit defined universally above will now be defined constructively.

DEFINITION 5.5. Let C be a free resolution system over a group system G with $G = \lim G$. The associated triad $(\tilde{C}, \tilde{C}^-, \tilde{C}^+)$ of C is a triad of augmented \overrightarrow{G} -complexes formed as follows: Let $\tilde{C}^\pm = ZG \otimes_{G_\pm} C^\pm$. Hence, $\tilde{C}^0 = ZG \otimes_{G_0} C^0 = \tilde{C}^- \cap \tilde{C}^+$. Let $\tilde{C} = (\tilde{C}^- \oplus \tilde{C}^+) / \Delta \tilde{C}^0$, where $\Delta \tilde{C} = \{c \oplus (-c) | c \in \tilde{C}^0\}$. \tilde{C} is called the associated complex of C and \tilde{C}^0 the associated intersection. The associated morphism $\tilde{\Phi}: C \rightarrow \tilde{C}$ is defined by $\tilde{\phi}_\alpha(c) = 1_{ZG} \otimes_{G_\alpha} c$ ($\alpha = -, 0, +$), where 1_{ZG} denotes the identity of ZG .

THEOREM 5.2. Let C be a free resolution system over a group system G . Then the direct limit of C exists and is unique up to isomorphism. Moreover, the associated complex \tilde{C} together with the associated morphism $\tilde{\Phi}$ forms the direct limit of C .

THEOREM 5.3. Let C and C' be free resolution systems over the same group system G . Every chain equivalence Φ from C to C' induces a chain equivalence $\phi: \lim C \rightarrow \lim C'$ such that

$$\begin{array}{ccc}
 C & \xrightarrow{\Phi} & C' \\
 \Phi^\infty \downarrow & & \downarrow \Phi'^\infty \\
 \lim_{\rightarrow} C & \xrightarrow{\phi} & \lim_{\rightarrow} C'
 \end{array}$$

is commutative. Moreover, Φ induces a chain equivalence from the associated triad of C to the associated triad of C' . Hence, by Theorem 5.1 the direct limit of any free resolution system over G is, modulo chain equivalence, an invariant of G .

The following theorem will be of use in the calculation of the algebraic 2-type of 2-knots.

THEOREM 5.4. *Let C be a free resolution system over a group system G with $G = \lim_{\rightarrow} G$. If $\phi_\alpha: G_\alpha \rightarrow G$ is an epimorphism and if G_α is of homological dimension at most q , then*

$$ZG \otimes_{G_\alpha} \text{Ker } \partial_i^\alpha = \text{Ker } \tilde{\partial}_i^\alpha,$$

for $i \geq q + 1$, where ∂^α and $\tilde{\partial}^\alpha$ denote the boundary operators of C^α in C and of \tilde{C}^α in the associated triad of C respectively.

IV. Completions of chain complexes. The material given in this section will be used to define and to compute the first k -invariant. (See [15].)

DEFINITION 6.1. Let G be a group and \tilde{C} an augmented free G -chain complex. Let q be a nonnegative integer such that $H_i \tilde{C} = 0$ for $i < q$. Then a q -completion \bar{C} of \tilde{C} is a free resolution over G such that

- (1) \bar{C} is a proper extension of \tilde{C} (see Def. 4.1).
- (2) $\bar{C}_i = \tilde{C}_i$ and $\bar{\partial}_i = \tilde{\partial}_i$ for $i \leq q$.

DEFINITION 6.2. Let G, \bar{C}, \tilde{C} , and q be as in the above definition and let $\text{id}: \bar{C} \rightarrow \tilde{C}$ be the identity chain map defined in dimensions up to q . The obstruction $k(\bar{C}, \tilde{C})$ to extending id to \bar{C}_{q+1} is the $(q + 1)$ -cocycle

$$k(\bar{C}, \tilde{C}) = \eta \iota \bar{\partial}: \bar{C}_{q+1} \xrightarrow{\bar{\partial}} Z_q \bar{C} \stackrel{\iota}{=} Z_q \tilde{C} \xrightarrow{\eta} H_q \tilde{C}$$

lying in $Z^{q+1}(\bar{C}, H_q \tilde{C})$. (See Theorem 4.1.) Let $k(\bar{C}, \tilde{C})$ denote the cohomology class in $H^{q+1}(\bar{C}, H_q \tilde{C}) = H^{q+1}(G, H_q \tilde{C})$ represented by $k(\bar{C}, \tilde{C})$.

The invariance of the above cohomology class is given by the following theorem.

THEOREM 6.1. *Let G be a group and C and C' augmented free G -chain complexes. Let q be a nonnegative integer such that $H_i \tilde{C} = 0 = H_i \tilde{C}'$ for $i < q$ and let \bar{C} and \bar{C}' be q -completions of \tilde{C} and \tilde{C}' respectively. Then every chain equivalence from \tilde{C} to \tilde{C}' extends to a chain equivalence from (\bar{C}, \tilde{C}) to (\bar{C}', \tilde{C}') . Each such equivalence induces an isomorphism of $H^{q+1}(\bar{C}', H_q \tilde{C}')$ onto $H^{q+1}(\bar{C}, H_q \tilde{C})$ which carries $k(\bar{C}', \tilde{C}')$ onto $k(\bar{C}, \tilde{C})$. (See [15].)*

VII. The main theorem. The main theorem (Theorem 7.1) will now be stated and proven. (The direct limit functor \lim (Def. 5.4) does not commute with the homology functor. See caveat in § 0.)

DEFINITION 7.1. An *aspherical splitting* of a CW complex K is a triad (K, K_-, K_+) of CW-complexes such that

- (1) $K = K_- \cup K_+$.
- (2) $K_-, K_0 = K_- \cap K_+, K_+$ are pathwise connected.
- (3) K_-, K_0, K_+ are aspherical.³

REMARK. It follows that there exist open subsets $\hat{K}_-, \hat{K}_0 = \hat{K}_- \cap \hat{K}_+, \hat{K}_+$ of K containing: K_-, K_0, K_+ respectively as deformation retracts. Hence, the Van Kampen theorem (Theorem 3.2) can be applied. (See [10].)

THEOREM 7.1.⁴ *Let (K, K_-, K_+) be an aspherical splitting of a CW complex K . Let C be a free resolution system over the group system G associated with the triad (K, K_-, K_+) . (See Defs. 3.2 and 5.1.) Let $\tilde{K}_-, \tilde{K}_0, \tilde{K}_+$ denote the respective lifts of K_-, K_0, K_+ to the universal cover \tilde{K} of K . Then $\lim C$ is chain homotopic to the augmented chain complex $C\tilde{K}$ of singular chains of \tilde{K} . Hence,*

$$\bar{H}_* \tilde{K} \cong H_* \lim C \text{ (as left } Z\pi_1 K\text{-modules),}$$

where \bar{H} denotes the reduced singular homology. Moreover, if $(\tilde{C} = \lim C, \tilde{C}^-, \tilde{C}^+)$ denotes the associated triad of C (see Def. 5.5), then

$$\bar{H}_* \tilde{K}_\alpha = H_* \tilde{C}^\alpha = \bar{H}_*(\pi_1 K_\alpha; Z\pi_1 K),$$

for $\alpha = -, 0, +$. Finally, if \bar{C} is a 2-completion of $\lim C$ (See Def.

³ In [43], K is called a *generalized-Eilenberg MacLane (GEM) complex*. The collection $\mathbf{K} = \{K_-, K_0, K_+\}$ is called an *aspherical structure* on K . From [43], the homotopy type of a GEM complex is completely determined by the group system $G = \{\pi_1 K_- \leftarrow \pi_1 K_0 \rightarrow \pi_1 K_+\}$. Hence, the notation $K = K(G, 1)$. See [43] for details.

⁴ In [43], $\lim C$ is called the *chain complex* of the group system $G = \{\pi_1 K_0 \leftarrow \pi_1 K_- \rightarrow \pi_1 K_+\}$ and denoted by CG ; and $H_* \lim C = H_*(G; ZG)$ is the homology of the group system G with local coefficients in ZG , where $G = \lim G$. See [43] for details.

6.1), then the 3-cocycle $k(\bar{C}, \lim_{\rightarrow} C)$ given by

$$\bar{C}_3 \xrightarrow{\bar{\delta}} Z_2 \bar{C} = Z_2 (\lim_{\rightarrow} C) \xrightarrow{\eta} H_2 \lim_{\rightarrow} C$$

lying in $Z^3(C, H_2 \lim_{\rightarrow} C)$ is a representative of the k -invariant kK lying in $H^3(\bar{C}, H_2 \lim_{\rightarrow} C) = H^3(\pi_1 K; \pi_2 K)$. (See Def. 6.2)

Proof. Let $\tilde{K}_{\pm}, \tilde{K}_0$ denotes the lifts of K_{\pm}, K_0 respectively to the universal cover \tilde{K} of K and let \check{K}_{\pm} and \check{K}_0 denote respectively the universal covers of K_{\pm} and K_0 . Moreover, let σ^i denote all the i -cells of K_0 and σ^{\pm}_i all the i -cells of $K_{\pm} - K_0$. Then the i -cells of $\tilde{K}, \tilde{K}_{\pm}, \tilde{K}_0, \check{K}_{\pm}, \check{K}_0$ are respectively

$$\begin{aligned} & \{g\sigma^i \cup g\sigma^i \cup g\sigma^i \mid g \in \pi_1 K\}, \\ & \{g\sigma^i \cup g\sigma^{\pm}_i \mid g \in \pi_1 K\}, \\ & \{g\sigma^i \mid g \in \pi_1 K\}, \\ & \{g\sigma^i \cup g\sigma^{\pm}_i \mid g \in \pi_1 K_{\pm}\}, \\ & \{g\sigma^i \mid g \in \pi_1 K_0\}. \end{aligned}$$

Let $C\tilde{K}, C\tilde{K}_{\pm}, C\tilde{K}_0, C\check{K}_{\pm}, C\check{K}_0$ denote the corresponding augmented G -chain complexes and let $\gamma^{\pm}: C\check{K}_0 \rightarrow C\check{K}_{\pm}$ denote the maps induced by the covering maps. Then

$$C = \{(C\tilde{K}_-, C\tilde{K}_0, C\tilde{K}_+), \{\gamma_-, \gamma_+\}\}$$

is a free resolution system over G and $(C\tilde{K}, C\tilde{K}_-, C\tilde{K}_+)$ is the associated triad of C . Hence, by construction

$$H_q \tilde{K} = H_q C\tilde{K} = H_q \lim_{\rightarrow} C$$

and

$$H_q \tilde{K}_{\alpha} = H_q C\tilde{K}_{\alpha} = H_q(\pi_1 K_{\alpha}; Z\pi_1 K)$$

for all q and for $\alpha = -, 0, +$.

Next let \bar{C} be a 2-completion of $\lim_{\rightarrow} C$; then from § 6 an obstruction $k(\bar{C}, \lim_{\rightarrow} C)$ in $H^3(\bar{C}, H_2 \lim_{\rightarrow} C) = \bar{H}^3(\pi_1 K; \pi_2 K)$ is defined. This construction of $k(\bar{C}, \lim_{\rightarrow} C)$ is precisely the geometric definition of the k -invariant kK given in [14, 15] except that the resolution \bar{C} is used instead of the bar resolution.

This proves the theorem for a particular free resolution system over G . Let C' be any other free resolution system over G and let $(\tilde{C}' = \lim_{\rightarrow} C', \tilde{C}'^-, \tilde{C}'^+)$ be the associated triad of C' . Then from § V, the associated triad of C' is chain homotopic to $(C\tilde{K}, C\tilde{K}^-, C\tilde{K}^+)$. Finally,

let \bar{C}' be a 2-completion of \tilde{C}' . Then from § VI, (\bar{C}', \tilde{C}') is chain equivalent to $(\bar{C}, C\tilde{K})$ and the general theorem follows. □

REMARK. Theorem 7.1 holds for more general aspherical splittings and group systemt. This is why the phrases “aspherical splitting” and “group system” are used above rather than “aspherical triad” and “group triad”. (See [43]).

REMARK. *Theorem 7.1 can be thought of as a π_2 -generalization of Crowell’s version of the Van Kampen theorem [10].*

The following corollaries of Theorem 7.1 are stated more generally in [43].

COROLLARY 7.2. *Let (K, K_-, K_+) be an aspherical splitting of a CW complex K with universal cover \tilde{K} . Then*

$$\begin{aligned} \dots \longrightarrow H_{q+1}\tilde{K} &\longrightarrow H_q(\pi_1K_0; Z\pi_1K) \\ &\longrightarrow H_q(\pi_1K_-; Z\pi_1K) \oplus H_q(\pi_1K_+; Z\pi_1K) \longrightarrow H_q\tilde{K} \longrightarrow \dots \end{aligned}$$

*is a long exact sequence.*⁵

From Theorem 7.1 or Corollary 7.2 we have as a corollary the following result of J. H. C. Whitehead.

COROLLARY 7.3. (Corollary to Theorem 5 in [66].) *Let G be the group system associated with an aspherical splitting (K, K_-, K_+) of a CW-complex K , i.e., associated with the triad (K, K_-, K_+) . (See Def. 3.2.) If all the morphisms in G are monomorphisms, then K is aspherical.*

The following theorem will be of use in simplifying the calculation of the algebraic 2-type of 2-knots.

THEOREM 7.4. *Let (K, K_-, K_+) be a hyperbolic splitting of a smooth (or locally fiat PL) 2-knot. (See Def. 2.1.) Then K_{\pm} collapses to a wedge of e_{\pm} 1-spheres, where e_- and e_+ denote respectively the number of upper and lower elliptic points. Hence, π_1K_{\pm} is free of rank e_{\pm} and of homological dimension one.*

⁵ This is a special case of the long exact sequence

$$\dots \longrightarrow H_{q+1}(G; A) \longrightarrow H_q(G_0; A) \longrightarrow H_q(G_-; A) \oplus H_q(G_+; A) \longrightarrow H_q(G; A) \longrightarrow \dots$$

found in [43], where G denotes the system $\{G_{-} \leftarrow G_0 \rightarrow G_{+}\}$ with direct limit (or push-out) G and A is a right ZG -module. This is a generalization to push-outs of Swan’s [43] Mayer Vietoris sequence (with local coefficients) for free products with amalgamation. See [63] for details.

THEOREM 7.5. *Let K_0 be the 0-level of a hyperbolic splitting of a smooth (or locally flat PL) 2-knot. (See Def. 2.1.) Then the fundamental group $\pi_1 K_0$ of K_0 is of homological dimension at most two.*

Proof. $\pi_1 K_0$ is the group of an unsplitable 1-graph in S^3 . Hence, by [9, 51] it is of homological dimension at most 2. (See remark at end of § VIII.) \square

From Theorems 7.4 and 7.5 above it follows that the hyperbolic triad of a smooth (or locally flat PL) 2-knot satisfies all the hypotheses of the following theorem.

THEOREM 7.6. *Let (K, K_-, K_+) be an aspherical splitting of a CW complex K and let G be the associated group system. (See Def. 3.2.) Let \tilde{K} be the universal cover of K . If $\pi_1 K_-$, $\pi_1 K_0$, $\pi_1 K_+$ are respectively of homological dimension at most 1, 2, 1, then the exact sequence of Corollary 7.2 reduces to*

$$H_3 \tilde{K} \cong H_2(\pi_1 K_0; Z\pi_1 K)$$

and to the short exact sequence

$$\begin{aligned} 0 \longrightarrow H_2 \tilde{K} \longrightarrow H_1(\pi_1 K_0; Z\pi_1 K) \\ \longrightarrow H_1(\pi_1 K_-; Z\pi_1 K) \oplus H_1(\pi_1 K_+; Z\pi_1 K) \longrightarrow 0, \end{aligned}$$

and all the remaining reduced homology groups of \tilde{K} vanish.

The second homotopy groups of 2-knots were originally computed from the above short exact sequence [41, 42]. The method given in this paper is a more general procedure of calculation. For 2-knots and 3-manifolds, this procedure of course reduces to the use of the above exact sequence. The more general procedure is needed to compute the k -invariant.

REMARK. Heegaard splittings of positive genus of connected 3-manifolds satisfy the hypotheses of Theorem 7.6.

From Theorem 7.4 and from the asphericity of 1-knots [53], we conclude:

THEOREM 7.7. *Every hyperbolic splitting of a 2-knot is an aspherical splitting.*

It is obvious that:

THEOREM 7.8. *Every Heegaard splitting of positive genus of a closed connected 3-manifold is an aspherical splitting.*

VIII. Fox-Lyndon resolutions. We now give a method for constructing a free resolution of a group which, for our purposes, is geometrically more enlightening and computationally more economical than the bar construction. (Our construction is similar to that found in [46, 65].) This method of construction will be used to bridge the gap between the main theorem (Theorem 7.1) and its more constructive analogue (Theorem 9.2).

If C is a (augmented) G -chain complex, then by a *free basis* of C , written

$$S_0, S_1, S_2, \dots,$$

is meant a sequence $S_q (q = 0, 1, 2, \dots)$ of sets of symbols such that S_q is a free basis of the left G -module C_q for all $q \geq 0$.

Let $\mathcal{P} = (x:r)_\phi$ be a presentation [11] of a group G . A Fox-Lyndon resolution over G corresponding to \mathcal{P} is a free resolution over G constructed as follows:

Let C_0 be the free left G -module on the single symbol P and let C_1 and C_2 be the free left G -modules on the sets X and R , where X and R are sets of symbols in one-to-one correspondence with the elements of x and r respectively. Let $\varepsilon: C_0 \rightarrow Z$ denote the extension of the morphism $\varepsilon: G \rightarrow Z$ which sends each element of G to 1 and define the G -morphisms $\partial_1: C_1 \rightarrow C_0, \partial_2: C_2 \rightarrow C_1$ by

$$\begin{aligned} \partial_1 X_i &= (x_i - 1)P & \forall X_i \in X \\ \partial_2 R_j &= \sum_i (\partial r_j / \partial x_i) X_i & \forall R_j \in R, \end{aligned}$$

where $x_i - 1$ and $\partial r_j / \partial x_i$ denote respectively the image in ZG under ϕ of $x_i - 1$ and of the Fox derivative [11, 18] $\partial r_j / \partial x_i$.

Next let u be a set of generators of the left G -module $\text{Ker } \partial_2$ and U a set of symbols in one-to-one correspondence with the elements of u . Define C_3 as the free left G -module on the symbols U and $\partial_3: C_3 \rightarrow C_2$ as the G -morphism given by

$$\partial_3 U_k = u_k = \sum_j (\partial U_k / \partial R_j) R_j \quad \forall U_k \in U,$$

where $\partial U_k / \partial R_j$ denotes the left coefficient of R_j in u_k .

Similarly, let w be a set of generators of the left G -module $\text{Ker } \partial_3$ and W a set of symbols in one-to-one correspondence with the elements of w . Define C_4 as the free left G -module on the symbols W and $\partial_4: C_4 \rightarrow C_3$ as the G -morphism given by

$$\partial_4 W_l = w_l = \sum_k (\partial W_l / \partial U_k) U_k \quad (\forall W_l \in W),$$

where $\partial W_l / \partial U_k$ denotes the left coefficient of U_k in w_l .

The free resolution C over G defined inductively as above is called a *Fox-Lyndon resolution of G corresponding to \mathcal{P}* . The free basis (see definition above)

$$P, X, R, U, W, \dots$$

is called a *constructive basis of the Fox-Lyndon resolution C* .

Finally, a q -completion \bar{C} (see Def. 6.1) of an augmented free G -chain complex \tilde{C} is a *Fox-Lyndon q -completion* if \bar{C} is a Fox-Lyndon resolution over G .

REMARK 1. The symbols U correspond to identities and the W to the identities among the identities. The coefficients $\partial U_k/\partial R_j$ and $\partial W_i/\partial U_k$ are actually the images of the Fox derivatives of these identities and identities of identities. Hence, the notation. (See [41, 42, 65].)

REMARK 2. Let Q be the complement of an unsplitable graph in the 3-sphere. Then Q collapses to a 2-dimensional CW complex K containing only one 0-cell. Let $(x:r)$ be the presentation of $\pi_1 Q$ carried by the cells of K and let \tilde{K} denote the universal cover of K . Then the cells of K lift to \tilde{K} to form an augmentable $\pi_1 Q$ -chain complex $C\tilde{K}$. This chain complex, when augmented, is a Fox-Lyndon resolution of $\pi_1 Q$ corresponding to $(x:r)$. Moreover, $C_q \tilde{K} = 0$ for $q > 2$. Hence, $\pi_1 Q$ is of homological dimension at most 2.

IX. A constructive form of the main theorem. The Fox-Lyndon resolution defined in § VIII will now be used to give a more constructive form (Theorem 9.2) of the main theorem (Theorem 7.1).

DEFINITION 9.1. Let G be a group system with direct limit $G = \lim \rightarrow G$. A presentation $(x_-; x_0; x_+; r_-; r_0; r_+)$ $_{\phi_-, \phi_0, \phi_+}$ of G is a pair of morphism presentations [20] $(x_0; x_-; r_0; r_-)$ $_{\phi_0, \phi_-}$ and $(x_0; x_+; r_0; r_+)$ $_{\phi_0, \phi_+}$ of $g_-: G_0 \rightarrow G_-$ and $g_+: G_0 \rightarrow G_+$ respectively. The subscripts ϕ_0 and ϕ_{\pm} are usually omitted unless extra precision is needed. It follows from [10] that $(x_-, x_0, x_+; r_-, r_0, r_+)$ is a presentation of G .

REMARK. If $f: G \rightarrow H$ is a group morphism, then a presentation $(x; y; r; s)$ of f is a set of two presentations $(x; r)_{\phi}$ and $(x; y; r; s)_{\psi}$ of G and H respectively such that

$$\begin{array}{ccc} F(x) & \xrightarrow{\iota} & F(x \cup y) \\ \downarrow & & \downarrow \\ |x; r| & & |x, y; r, s| \\ \cong \downarrow \phi & & \downarrow \psi \cong \\ G & \xrightarrow{f} & H \end{array}$$

is commutative, i.e., such that f is the morphism induced by the inclusion ι . (See [20].)

By arguments similar to those found in [20, p. 411], it can be shown that the following transformations do not alter the isomorphism type of the presented group system.

DEFINITION. Let $\mathcal{P} = (\mathbf{x}_-; \mathbf{x}_0; \mathbf{x}_+; \mathbf{r}_-; \mathbf{r}_0; \mathbf{r}_+)$ be a presentation of a group system G . Then the following are called *Tietze transformations of \mathcal{P}* .

(I) Adjoin to r_0 any one of its consequences.

(I $_{\pm}$) Adjoin to r_{\pm} any one of the consequences of $r_0 \cup r_{\pm}$.

(II) Adjoin a new symbol x_0 to \mathbf{x}_0 and x_0u^{-1} to r_0 , where u is in the free group $F(\mathbf{x}_0)$.

(II $_{\pm}$) Adjoin a new symbol x_{\pm} to \mathbf{x}_{\pm} and $x_{\pm}u^{-1}$ to r_{\pm} , where u is in the free group $F(\mathbf{x}_0, \mathbf{x}_{\pm})$.

By a proof similar to that given in [19, p. 198], it can be shown that:

THEOREM (Tietze). *If \mathcal{P} and \mathcal{P}' are finite presentations of the same group system, then it is possible to pass from one to the other presentation by applying a finite sequence of Tietze Transformations of types*

$$(I)^{\pm 1}, (I_{-})^{\pm 1}, (I_{+})^{\pm 1}, (II)^{\pm 1}, (II_{-})^{\pm 1}, (II_{+})^{\pm 1}.$$

DEFINITION 9.2. Let $\mathcal{P} = (\mathbf{x}_-; \mathbf{x}_0; \mathbf{x}_+; \mathbf{r}_-; \mathbf{r}_0; \mathbf{r}_+)$ be a presentation of a group system G with $G = \lim_{\rightarrow} G$. Then a *Fox-Lyndon resolution system over G* is a free resolution system C constructed as follows: Let C^0 be a Fox-Lyndon resolution (see § VIII) of G_0 corresponding to $(\mathbf{x}_0; \mathbf{r}_0)$. Let C^{\pm} be a Fox-Lyndon resolution of G_{\pm} corresponding to $(\mathbf{x}_0, \mathbf{x}_{\pm}; \mathbf{r}_0, \mathbf{r}_{\pm})$ formed by extending $ZG_{\pm} \otimes_{G_0} C^0$. Then C^-, C^0, C^+ together with the obvious maps γ_-, γ_+ under the respective group morphisms g_-, g_+ is a free resolution system over G . Let $P, X_0, R_0, U_0, W_0, \dots$ be the constructive basis (see § VIII) of C^0 and let the induced free basis (see § VIII) of $ZG_{\pm} \otimes_{G_0} C^0$ also be denoted by the same symbols. Extend this free basis to a constructive basis $P, X_0 \cup X_{\pm}, R_0 \cup R_{\pm}, U_0 \cup U_{\pm}, W_0 \cup W_{\pm}, \dots$ of C^{\pm} . Then

$$P; X_-, X_0, X_+; R_-, R_0, R_+; U_-, U_0, U_+; W_-, W_0, W_+; \dots$$

is called a *constructive basis* of the Fox-Lyndon resolution system C .

THEOREM 9.1. *Let C be a Fox-Lyndon resolution system over a group system G corresponding to a presentation \mathcal{P} of G . Let $G =$*

$\lim_{\rightarrow} \mathbf{G}$ and $(\tilde{C}, \tilde{C}^-, \tilde{C}^+)$ be the associated triad (see Def. 5.5) of C . If

$$P; X_-, X_0, X_+; R_-, R_0, R_+; U_-, U_0, U_+; W_-, W_0, W_+; \dots$$

is a constructive basis of C , then

$$P, X_0, R_0, U_0, W_0, \dots$$

$$P, X_0 \cup X_{\pm}, R_0 \cup R_{\pm}, U_0 \cup U_{\pm}, W_0 \cup W_{\pm}, \dots$$

$$P, X_- \cup X_0 \cup X_+, R_- \cup R_0 \cup R_+, U_- \cup U_0 \cup U_+, W_- \cup W_0 \cup W_+, \dots$$

are respectively the induced constructive bases of $\tilde{C}_0, \tilde{C}^{\pm}, \tilde{C}$, and the boundary operators are given by

$$\begin{aligned} \tilde{\partial}^0 &= 1_G \otimes_{G_0} \partial^0 & \tilde{\partial}^{\pm} &= 1_G \otimes_{G_{\pm}} \partial^{\pm} \\ \tilde{\partial}\sigma &= \begin{cases} \tilde{\partial}^+\sigma & \text{if } \sigma \in \tilde{C}^+ \\ \tilde{\partial}^-\sigma & \text{if } \sigma \in \tilde{C}^- \end{cases}, \end{aligned}$$

where ∂_{α} is the boundary operator of C_{α} .

THEOREM 9.2. (Constructive form of main theorem.) Let C be a Fox-Lyndon resolution system over a group system \mathbf{G} corresponding to a presentation

$$\mathcal{P} = (\mathbf{x}_-; \mathbf{x}_0; \mathbf{x}_+; \mathbf{r}_-; \mathbf{r}_0; \mathbf{r}_+)$$

of \mathbf{G} . Let $G = \lim_{\rightarrow} \mathbf{G}$, let $(\tilde{C}, \tilde{C}^-, \tilde{C}^+)$ be the associated triad (see Def. 5.5) of C , and let \bar{C} be a Fox-Lyndon 2-completion of \tilde{C} (see § VIII). Let

$$P; X_-, X_0, X_+; R_-, R_0, R_+; U_-, U_0, U_+; W_-, W_0, W_+; \dots$$

be a constructive basis of C and let

$$\begin{aligned} P, X_- \cup X_0 \cup X_+, R_- \cup R_0 \cup R_+, U_- \cup U_0 \cup U_{\square} \cup U_+, \\ W_- \cup W_0 \cup W_{\square} \cup W_+, \dots \end{aligned}$$

be a constructive basis of \bar{C} formed by extending the free basis of \tilde{C} mentioned in Theorem 9.1. Then $H_2\bar{C}$ as a left G -module is generated by the elements of ∂U_{\square} and

$$(\partial U_{\square}: \sum_{v_{\square} \in v_{\square}} (\partial W_{\square} / \partial U_{\square}) \partial U_{\square} = 0, \forall W_{\square} \in W_{\square})$$

is a presentation of $H_2\bar{C}$ as a left ZG -module. Moreover, if W' is any set of generators of the left ZG -module \bar{C}' , then

$$(\partial U_{\square}: \sum_{v_{\square} \in v_{\square}} (\partial W' / \partial U_{\square}) \partial U_{\square} = 0, \forall W' \in W')$$

is also a presentation of $H_2\bar{C}$ as a left ZG -module. Moreover, the

obstruction $k(\bar{C}, \tilde{C})$ lying in $H^3(\bar{C}, H_2\tilde{C})$ is represented by the equivariant 3-cocycle $k: \bar{C}_3 \rightarrow H_2\tilde{C}$ given by

$$k(U^*) = \begin{cases} \partial U^* & \text{if } U^* \in U_\square \\ 0 & \text{if } U^* \in U_- \cup U_0 \cup U_+ . \end{cases}$$

Proof. From the long exact sequence induced by the short exact sequence

$$0 \longrightarrow \tilde{C} \longrightarrow \bar{C} \longrightarrow \bar{C}/\tilde{C} \longrightarrow 0 ,$$

we have that

$$H_2\tilde{C} \cong H_3(\bar{C}/\tilde{C})$$

as left G -modules under a isomorphism denoted by ∂_* .⁶ But

$$\phi, \phi, \phi, U_\square, W_\square, \dots$$

is a constructive basis of \bar{C}/\tilde{C} . Hence, U_\square is a set of generators of the left ZG -module $Z_3(\bar{C}/\tilde{C})$ of relative 3-cycles and

$$\partial W_\square = \sum_{v_\square \in v_\square} (\partial W_\square / \partial U_\square) U_\square$$

is a set of generators of the left ZG -module $B_3(\bar{C}/\tilde{C})$ of relative 3-boundaries. Hence,

$$(U_\square: \sum_{v_\square \in v_\square} (\partial W_\square / \partial U_\square) U_\square = 0, \forall W_\square \in W_\square)$$

is a presentation of $H_3(\bar{C}/\tilde{C})$ as a left ZG -module. Thus using the isomorphism ∂_* taking U_\square to ∂U_\square , we have the first presentation.

Next, let $\tilde{C}^{(3)}$ be a left ZG -chain complex such that $\tilde{C}^{(3)}$ and \tilde{C} agree up to and including \tilde{C}_3 and such that $\tilde{C}_q^{(3)} = 0$ for $q > 3$. Then $H_2\tilde{C} \cong H_2\tilde{C}^{(3)}$ and by the same argument, $H_2\tilde{C}^{(3)} = H_3(\bar{C}/\tilde{C}^{(3)})$, which gives the second presentation.

Finally, by Definition 6.2, the obstruction $k(\bar{C}, \tilde{C})$ is represented by

$$k = \eta id_3: \bar{C}_3 \longrightarrow Z_2\bar{C} = Z_2\tilde{C} \longrightarrow H_2\tilde{C} .$$

Hence,

$$k(U^*) = \partial U^* = \begin{cases} \partial U^* & \text{if } U^* \in U_\square \\ 0 & \text{if } U^* \in U_- \cup U_0 \cup U_+ \end{cases}$$

since $\partial U = 0$ in $H_2\tilde{C}$ for all $U \in U_- \cup U_0 \cup U_+$. □

The following theorem will be used to simplify 2-knot and 3-manifold calculations.

⁶ In the terminology of [43], $H_2(\mathbf{G}; ZG) \cong H_3(G, \mathbf{G}; ZG)$.

THEOREM 9.3. *Let (K, K_-, K_+) be an aspherical splitting of a CW complex K (see Def. 7.1) and let G be the associated group system (see Def. 3.2). Hence, by Theorem 3.2, $\pi_1 K = \lim_{\rightarrow} G$. Let C be a Fox-Lyndon resolution system over G corresponding to the presentation $(x_-; x_0; x_+; r_-; r_0; r_+)$. If $K_0 = K_- \cap K_+$ collapses to a 2-dimensional CW complex and if $(x_0; r_0)$ is aspherical, (i.e., its associated 2-complex is aspherical), then $\text{Ker } \partial_2^0 = 0$, i.e., U_0 of Theorem 9.2 is vacuous. Hence, we may also assume $\text{Ker } \partial_3^0 = 0$, i.e., W_0 of Theorem 9.2 is vacuous.*

X. What is the significance of the algebraic 2-type of 2-knots? Throughout this section (unless stated otherwise), K and K' will denote the exteriors of n -knots. (See notation in § 0.)

We now attempt to understand how significant are the remaining homotopy groups $\pi_q K$ for $q \geq 3$ for 2-knots. A complete answer to this question would indeed be a four dimensional analogue of the asphericity of 1-knots [53]. The evidence given herein (Theorem 10.1) suggests that the algebraic 2-type⁷ completely determines the homotopy type of 2-knot complements. If this is so, then the groups $\pi_q K$ for $q \geq 3$ are of little significance.

Since the asphericity of 1-knots essentially means that the algebraic 1-type (i.e., the fundamental group) completely determines the homotopy type of 1-knot complements, we now ask if the analogous property is true of 2-knots, i.e.,

QUESTION. If the complements of two smooth (or locally flat PL) 2-knots are of the same algebraic 2-type⁷, then are their complements of the same homotopy type?

In an attempt to gain some insight into the above question, we give the following definition.

DEFINITION 10.1. An n -knot is *quasi-aspherical* (QA) if the $(n + 1)$ -th homology group of the universal cover of its exterior vanishes.

THEOREM 10.1. *Let K and K' be the exteriors of two smooth (or locally flat PL) QA 2-knots. Then K and K' are of the same homotopy type if and only if they are of the same algebraic 2-type⁷.*

Proof. The exteriors K and K' are homotopic to 3-dimensional CW complexes, which we also denote by the same symbols K and K' , respectively. Hence, we need only prove that, if K and K'

⁷ In [48], S. MacLane and J. H. C. Whitehead call this the algebraic 3-type.

are two 3-dimensional CW complexes of the same algebraic 2-type, then K and K' are of the same homotopy type.

Assume that K and K' are of the same algebraic 2-type. Hence, there exists an isomorphism (f_0, h) of $T(K) = (\pi_1 K, \pi_2 K, kK)$ onto $T(K') = (\pi_1 K', \pi_2 K', kK')$ [48]; and by Theorem 3 of [48] this isomorphism has a geometric realization $\phi: K \rightarrow K'$.

Let K^i and K'^i denote the i -skeletons of K and K' , $p: \tilde{K} \rightarrow K$ and $p': \tilde{K}' \rightarrow K'$ the universal covers, and $\tilde{K}^i = p^{-1}K^i$ and $\tilde{K}'^i = p'^{-1}K'^i$.

Following [48], we define an augmentable chain complex $C = C_* K$ by $C_i K = H_i(\tilde{K}^i, \tilde{K}^{i-1})$ if $i > 0$ and by $C_0 K = H_0 \tilde{K}^0$ if $i = 0$ with boundary operator defined in the obvious way. Let $C^* = C^*_* K$ be the corresponding augmented chain complex. Define $C' = C_* K'$ and $C'^* = C^*_* K'$ in like manner. Then C^* and C'^* are acyclic in dimensions less than 2. In fact, because of quasi-asphericity $H_i C^* = H_i \tilde{K} = 0 = H_i \tilde{K}' = H_i C'^*$ if $i \neq 2$, where \tilde{K} and \tilde{K}' are the universal covers of K and K' , respectively.

By Theorem 4 of [48], (f_0, h) has a combinatorial realization $(f_0, \lambda): C^3 \rightarrow C'^3$ which induces an isomorphism of $H_i C^*$ for $i < 3$. (C^3 and C'^3 denote the 3-skeletons of C and C' respectively.) But $H_i C = 0 = H_i C'$ for $i > 2$ and $C_i K = 0 = C_i K'$ for $i > 3$. Hence, (f_0, h) has a combinatorial realization $(f_0, \lambda): C \rightarrow C'$ which induces an isomorphism of $H_i C$ onto $H_i C'$ for all i .

By Theorem 5 of [48], $C \equiv_i C'$ for all i . Thus, C is equivalent to C' . On the other hand, K and K' are J_2 -complexes of dimension 3. So by Theorem 15 of [69], K is of the same homotopy type as K' . \square

Actually, we have proven more than stated above, namely:

THEOREM 10.2. *If Q and Q' are topological spaces such that*

- (1) *Q and Q' are dominated by connected 3-dimensional CW complexes, and*
- (2) *The third homology groups of the universal covers of Q and Q' both vanish,*

then Q and Q' are of the same homotopy type if and only if they are of the same algebraic 2-type.

It is easy enough to show that the class of QA knots is immense. It is proven in [26] that:

THEOREM 10.3 [26]. *An n -knot is not QA if and only if its fundamental group G has a decomposition of the form $A^* C B$ with C finite and properly contained in A and B such that the meridian subgroup H of G lies in A .*

THEOREM 10.4. [26, 39, 41, 55, 64] *The class QA n -knots includes all spun, twist spun, fibered, nilpotent, and finitely ended knots.*

If all 2-knots were QA, then the above question would be resolved. However, this is not the case. Recently, Gonzalez-Acuna and Montesinos [26] have constructed an n -knot ($n \geq 2$) with infinitely many ends, thus solving problem 40 of R. H. Fox [22]. Ratcliffe in [55] has pointed out that this example has a group of the form given in Theorem 10.3 above, and hence, is not QA. In light of Theorem 10.4 above, the examples of [26] are indeed remarkable.

XI. Miscellaneous corollaries on 2-knots. We list below a number of miscellaneous corollaries. The following corollary is an immediate consequence of J. H. C. Whitehead's certain exact sequence [67].

COROLLARY 11.1. *Let Q be a topological space such that*

(1) *Q is homotopic to a 3-dimensional CW complex.*

(2) *The third homology group of the universal cover of Q vanishes.*

Then $\pi_3 Q$ as a $\pi_1 Q$ -module is isomorphic to $\Gamma(\pi_2 Q)$, where Γ denotes the functor defined in [67]. (See also [45].)

REMARK. If K is the exterior of a smooth (or locally flat PL) QA 2-knot or of a connected QA 3-manifold, then the above corollary can be used to compute $\pi_3 K$ from $\pi_2 K$. (For details see [45].)

From [12, 13] and from Theorem 10.1 it follows that

COROLLARY 11.2. *If K is the exterior of a smooth QA 2-knot and if $\pi_2 K = 0$, then K is a homotopy 1-sphere.*

Since all 1-knot groups are of homological and cohomological dimension at most 2 (see [9, 51] or Remark 2 in § VIII), we have:

COROLLARY 11.3. *Let K be the exterior of a spun 2-knot. Then $H^3(\pi_1 K; \pi_2 K) = 0$, and hence, the k -invariant kK vanishes.*

COROLLARY 11.4. *Let Q be a connected CW complex whose second homology group vanishes and let \mathcal{F} denote the augmentation ideal of $\pi_1 Q$. Then the k -invariant kQ lying in $H^3(\pi_1 Q; \pi_2 Q)$ is nonzero if $\pi_2 Q / \mathcal{F} \pi_2 Q \neq 0$.*

Proof. By [14], if $kQ = 0$, then $\mathcal{F} \pi_2 Q$ is then the kernel of

the natural morphism of $\pi_2 Q$ into $H_2 Q$. □

The following is a consequence of the Leray-Serre spectral sequence. (See [8, p. 356].)

COROLLARY 11.5. *Let Q be a CW-complex such that $H_2 Q = 0 = H_3 Q$. Then*

$$H_3 \pi_1 Q \cong \pi_2 Q / \mathcal{F} \pi_2 Q \quad (= Z \otimes_{\pi_1 Q} \pi_2 Q),$$

where \mathcal{F} denotes the augmentation ideal of $\pi_1 Q$.

Since every n -knot complement is by Alexander duality a homology 1-sphere, we have

COROLLARY 11.6. *Let K be the exterior (or complement) of a smooth n -knot. Then*

$$H_3 \pi_1 K \cong \pi_2 K / \mathcal{F} \pi_2 K \quad (= Z \otimes_{\pi_1 K} \pi_2 K),$$

where \mathcal{F} denotes the fundamental ideal of K . Moreover, the k -invariant kK lying in $H^3(\pi_1 K; \pi_2 K)$ is nonzero if $H_3 \pi_1 K \neq 0$.

The first part of Corollary 11.6 can be thought of as a generalization of the Kervaire condition $H_2 \pi_1 Q = 0$. (See [36].)

COROLLARY 11.7. *Let K be the exterior of a smooth (or locally flat PL) fibered 2-knot. Then $kK \neq 0$ if the commutator subgroup G' of $\pi_1 K$ is non-trivial and finite.*

Proof. If G' is finite, then by Theorem 13 of [3]

$$\pi_2 K = (\theta : (\sum_{g \in G'} g)\theta = 0 = (t - 1)\theta).$$

Hence, $\pi_2 K / \mathcal{F} \pi_2 K = Z_n \neq 0$, where n is the order of G' , and by Corollary 11.6, $kK \neq 0$. □

XII. Conclusion: Problems and more problems. We list below a number of problems which as far as I know are still unresolved.

PROBLEM 1. *Let K and K' be smooth (or locally flat PL) 2-knot exteriors. Is K of the same homotopy type as K' iff K and K' are of the same algebraic 2-type? (See § X.) If so, this is a four-dimensional analogue of the asphericity of classical knots.*

PROBLEM 2. *Let K and K' be smooth n -knot exteriors ($n \geq 3$). Is K of the same homotopy type as K' iff K and K' are of the same n -type?⁸ (See [50, 68].) (See § X.)*

PROBLEM 3. (Ratcliffe.) *Is a smooth n -knot ($n \geq 2$) quasi-aspherical if and only if it is finitely ended? (See Theorem 10.3 and last paragraph of § X. See also [39, 48, 50, 55].) (See § X.)*

PROBLEM 4. (Problem 37 of Fox [22].) *Are there any non-trivial smooth (or locally flat PL) aspherical 2-knots? This appears to be a very difficult problem. The reader should note that a smooth (or locally flat PL) 2-knot is aspherical if and only if the fundamental group $\pi_1 K$ of its exterior K is infinite cyclic. The "only if" part of this statement is proven in [12, 13]. The "if" part follows from [38, Theorem 3.4]. (See [34, 61].) Moreover, the exterior K of a smooth (or locally flat PL) aspherical 2-knot is a homotopy 1-sphere. For a map from S^1 to K can easily be constructed which induces isomorphisms $\pi_* S^1 \cong \pi_* K$. Hence, by a theorem of J. H. C. Whitehead, K is a homotopy S^1 . (See also [25].)*

PROBLEM 5. (Problem 8 of Gordon [28].) *Let K be the exterior of a smooth (or locally flat PL) 2-knot (S^4, kS^2). Is $\pi_2 K$ always finitely generated as a left $Z\pi_1 K$ -module?*

PROBLEM 6. *Let K be as in Problem 5. Is $\pi_2 K$ considered only as a group always a free abelian group? (This is true for all spun 2-knots. See [17].)*

PROBLEM 7. *Let K be as in Problem 5. Is $kK \neq 0$ if and only if $H_3 \pi_1 K \neq 0$. (See Corollaries 11.6 and 11.7.) (From [15], it follows that $H_3 \pi_1 K \neq 0$ iff the reduced K -invariant is non-zero.)*

PROBLEM 8. *Let K be as in Problem 5. What is the relationship (if any) between $H_3 \pi_1 K$, the number of ends of $\pi_1 K$, and quasi-asphericity? (See §§ 10 and 11.)*

PROBLEM 9. *When does a 2-knot diagram represent the trivial 2-knot?*

PROBLEM 10. *When do two 2-knot diagrams represent the same 2-knot?*

PROBLEM 11. *Construct a table of all 2-knot diagrams with*

⁸ This is called the $(n+1)$ -type in [68].

≤ 6 (or 7, or 8, ...) crossings and with ≤ 2 (or 3, or 4, ...) hyperbolic points. (See Problems 9 and 10 above.)

PROBLEM 12. *Is there a CW-complex K such that no cell decomposition of its underlying space $|K|$ has an aspherical splitting?* Bill Beckmann [5] has pointed out that every 2-dimensional CW complex has a subdivision which has an aspherical splitting.

PROBLEMS 13-16. These problems can be found in the addendum.

Appendix A. A computing manual for algebraic 2-type calculations. If the reader is interested only in computing the algebraic 2-type of closed connected 3-manifolds and of 2-knot complements, he or she need only read Theorem A2.1 and observations 1 through 3 and the hint, all in part A2 of this appendix. Explicit 2-knot calculations are given in Appendix B. If the reader has more general calculations in mind, then all of Appendix A should be read.

Part A1. The computation for arbitrary aspherical triads.

DEFINITION A1.1. Let (K, K_-, K_+) be an *aspherical splitting* of a CW complex K , i.e., a triad of CW-complexes such that

- (1) $K = K_- \cup K_+$.
- (2) $K_-, K_0 = K_- \cap K_+, K_+$ are pathwise connected.
- (3) K_-, K_0, K_+ are aspherical, i.e., $\pi_q K_\alpha = 0$ for $\alpha = -, 0, +$ and for $q > 1$.

Let $(x_0; r_0)$ be a presentation of $\pi_1 K_0$, let $(x_\pm, x_\pm; r_\pm)$ be a presentation of $\pi_1 K_\pm$ such that $x_{i0} \mapsto x_{i0}$ (for all $x_{i0} \in x_0$) is the morphism $g_\pm: \pi_1 K_0 \rightarrow \pi_1 K_\pm$ induced by inclusion, and let $g_\alpha^\infty: \pi_1 K_\alpha \rightarrow \pi_1 K$ denote the morphisms induced by inclusion for $\alpha = -, 0, +$. It follows from the Van Kampen theorem that $(x_-, x_0, x_+; r_-, r_0, r_+)$ is a presentation of $\pi_1 K$.

Linearly order $x_- \cup x_0 \cup x_+$ and $r_- \cup r_0 \cup r_+$ such that $x_{i-} < x_{j0} < x_{k+}$ for all i, j, k and $r_{\alpha-} < r_{\beta 0} < r_{\gamma+}$ for all α, β, γ and consider the equations:

$$\begin{aligned}
 (*0) \quad & \vec{\lambda}^0 \left(\frac{\partial r_0}{\partial x_0} \right) = \vec{0}, \\
 (*+) \quad & \vec{\lambda}^+ \left(\frac{\partial (r_+ \cup r_0)}{\partial (x_0 \cup x_+)} \right) = \vec{0}, \\
 (*-) \quad & \vec{\lambda}^- \left(\frac{\partial (r_0 \cup r_-)}{\partial (x_- \cup x_0)} \right) = \vec{0},
 \end{aligned}$$

$$(*\square) \quad \vec{\lambda}^{\square} \left(\frac{\partial(r_+ \cup r_0 \cup r_-)}{\partial(x_- \cup x_0 \cup x_+)} \right) = \vec{0},$$

where the rows and columns of the above matrices are ordered according to the above constructed linear ordering⁹ and where $\partial r_i / \partial x_j$ denotes the Fox derivative. (Note: $(\partial r_0 / \partial x_{\pm})$, $(\partial r_{\pm} / \partial x_{\mp})$ are all zero matrices.)

Let ∂U_0 denote a set of generators of the left $Z\pi_1 K_0$ -module of solutions $\vec{\lambda}^0$ of equation $(*0)$ over the ring $Z\pi_1 K_0$. Each solution of equation $(*0)$ induces a solution $\vec{\lambda}^{\pm}$ of equation $(*\pm)$ over $Z\pi_1 K_{\pm}$. Let ∂U_0 also denote the induced solutions of $(*\pm)$ over $Z\pi_1 K_{\pm}$. Let $\partial U_{\pm} \cup \partial U_0$ denote a set of generators of the left $Z\pi_1 K_{\pm}$ -module of solutions $\vec{\lambda}^{\pm}$ of equation $(*\pm)$ over $Z\pi_1 K_{\pm}$. Each solution of $(*\pm)$ over $Z\pi_1 K_{\pm}$ induces a solution of $(*\square)$ over $Z\pi_1 K$. Let $\partial U_{-} \cup \partial U_0 \cup \partial U_{+}$ also denote these induced solutions of $(*\square)$ over $Z\pi_1 K$. Finally, let $\partial U_{-} \cup \partial U_0 \cup \partial U_{\square} \cup \partial U_{+}$ denote a set of generators of the left $Z\pi_1 K$ -module of solutions $\vec{\lambda}^{\square}$ of $(*\square)$ over $Z\pi_1 K$. Then ∂U_{\square} is a set of generators of $\pi_2 K$ as a left $Z\pi_1 K$ -module.

A complete set of relations among the generators ∂U_{\square} of $\pi_2 K$ is computed as follows. Linearly order the set $\partial U_{-} \cup \partial U_0 \cup \partial U_{\square} \cup \partial U_{+}$ such that $\partial U_{\alpha} < \partial U_{\beta} < \partial U_{\gamma} < \partial U_{\delta}$ for all $\alpha, \beta, \gamma, \delta$. Let ∂R_{α} be a set of symbols in 1-1 correspondence with the relators r_{α} for $\alpha = -, 0, +$. Give $\partial R_{-} \cup \partial R_0 \cup \partial R_{+}$ the linear ordering induced by that on $r_{-} \cup r_0 \cup r_{+}$. Let $\partial U_{\sigma} / \partial R_i$ denote the component of the vector ∂U_{σ} corresponding to r_i for all ∂U_{σ} in $\partial U_{-} \cup \partial U_0 \cup \partial U_{\square} \cup \partial U_{+}$ and for all r_i in $r_{-} \cup r_0 \cup r_{+}$. And consider the equations

$$(**\square) \quad \vec{\mu}^{\square} \left(\frac{\partial U_{+} \cup \partial U_0 \cup \partial U_{\square} \cup \partial U_{-}}{\partial R_{+} \cup \partial R_0 \cup \partial R_{-}} \right) = \vec{0},$$

where the rows and columns of the above matrices are ordered according to the above constructed linear orderings.¹⁰ (Note: $(\partial U_0 / \partial R_{\pm})$, $(\partial U_{\pm} / \partial R_{\mp})$, $(\partial U_{\pm} / \partial R_{\square})$, $(\partial U_0 / \partial R_{\square})$ are all zero matrices.)

Let ∂W denote a set of generators of the left $Z\pi_1 K$ -module of solutions $\vec{\mu}^{\square}$ of $(**\square)$ over $Z\pi_1 K$. Finally, let $\partial W_{\omega} / \partial U_{\sigma}$ denote the component of ∂W_{ω} corresponding to ∂U_{σ} for all ∂W_{ω} in ∂W . Then

$$\left\{ \sum_{\sigma_{\square} \in U_{\square}} (\partial W / \partial U_{\sigma}) \partial U_{\sigma} = 0 \mid W \in W \right\}$$

is a complete set of relations in $\pi_2 K$ among the generators ∂U_{\square} . Hence,

⁹ Rows in decreasing order from top to bottom. Columns in increasing order from left to right.

¹⁰ Rows ordered as in footnote 9. Columns in decreasing order from left to right.

$$(\{\partial U_{\square} | \partial U_{\square} \in \partial U_{\square}\} : \{ \sum_{\partial U_{\square} \in \partial U_{\square}} (\partial W / \partial U_{\square}) \partial U_{\square} = 0 | \partial W \in \partial W \})$$

is a presentation of $\pi_2 K$ as a left $Z\pi_1 K$ -module.

Finally, let W, U, R be a set of symbols in 1 – 1 correspondence with the elements of $\partial W, \partial U = \partial U_{-} \cup \partial U_0 \cup \partial U_{\square} \cup \partial U_{+}, \partial R = \partial R_{-} \cup \partial R_0 \cup \partial R_{+}$, respectively and let $\bar{C}_4, \bar{C}_3, \bar{C}_2$ be the free left $Z\pi_1 K$ -modules on the sets W, U, R , respectively. Then

$$\bar{C}_4 \xrightarrow{\bar{\partial}_4} \bar{C}_3 \xrightarrow{\bar{\partial}_3} \bar{C}_2$$

is a portion of a free resolution over $\pi_1 K$, where

$$\begin{aligned} \bar{\partial}_4 W &= \sum_{U \in U} (\partial W / \partial U) U, \quad \forall W \in W \quad (\text{over } Z\pi_1 K) \\ \bar{\partial}_3 U &= \sum_{R \in R} (\partial U / \partial R) R, \quad \forall R \in R \quad (\text{over } Z\pi_1 K). \end{aligned}$$

A representative of the k -invariant kK lying in $H^3(\pi_1 K; \pi_2 K)$ is the equivariant 3-cocycle k given by

$$k(U) = \begin{cases} \partial U & \text{if } U \in U_{\square} \\ 0 & \text{if } U \in U_{-} \cup U_0 \cup U_{+}. \end{cases}$$

A2. Simplifications that occur for Heegaard and hyperbolic splittings. For the reader who is only interested in computing the algebraic 2-type (i.e., π_1, π_2 , and $k \in H^3(\pi_1; \pi_2)$) of 2-knot exteriors or of 3-manifolds, we state and prove the following theorem. (The definition of hyperbolic splitting is given in § II (Def. 2.1). A definition for Heegaard splitting can be found in any standard text on 3-manifolds. See, for example, [32, Chapter 2].)

THEOREM A2.1. *Let (K, K_{-}, K_{+}) be a hyperbolic splitting of a 2-knot (or a Heegaard splitting of positive genus of a closed connected 3-manifold). Let $(\mathbf{x}; r_0)$ be an aspherical presentation of $\pi_1 K_0$ (see Observation 1 below) and let $(\mathbf{x}; r_0 \cup r_{\pm})$ be the presentation of $\pi_1 K_{\pm}$ obtained by adjoining the hyperbolic relators r_{\pm} (see Observation 2 below) [or by adjoining the meridional handle relators r_{\pm} (see Observation 2 below)]. Let $\mathbf{x} = \mathbf{x}_0$ and $r = r_{-} \cup r_0 \cup r_{+}$ and let $(\partial r / \partial \mathbf{x}) = (\partial r_{\alpha} / \partial x_{\beta})$ denote the Jacobian matrix whose $\alpha\beta$ entry is the image of the Fox derivative [11, 18] $\partial r_{\alpha} / \partial x_{\beta}$ in the ring $Z\pi_1 K$. Let ∂U_{\pm} be a set of generators of the left $Z\pi_1 K$ -module of solutions $\bar{\lambda}$ of equation (*)*

$$(*) \quad \lambda(\partial r / \partial \mathbf{x}) = 0,$$

over $Z\pi_1K$ such that the components of $\vec{\lambda}$ corresponding to r_{\mp} all vanish. Let ∂U_{\square} be a set of solutions of (*) over $Z\pi_1K$ such that $\partial U = \partial U_{-} \cup \partial U_{\square} \cup \partial U_{+}$ is a set of generators of the left $Z\pi_1K$ -module of all solutions $\vec{\lambda}$ of (*).

Next let ∂R_{ρ} be another notation for r_{ρ} and let $\partial U_{\alpha}/\partial R_{\rho}$ denote the component of ∂U_{α} corresponding to $\partial R_{\rho} = r_{\rho}$. Now let ∂W be a set of generators of the left $Z\pi_1K$ -module of solutions $\vec{\mu}$ of equation (**)

$$(**) \quad \vec{\mu}(\partial U/\partial R) = 0 ,$$

over $Z\pi_1K$.

If $\partial W/\partial U$ denotes the component of ∂W corresponding to ∂U for all $\partial W \in \partial W$ and for all $\partial U \in \partial U$, then

$$(\partial U_{\square}: \sum_{\partial U_{\square} \in \partial U_{\square}} (\partial W/\partial U_{\square})\partial U_{\square} = 0 \text{ for all } \partial W \in \partial W)$$

is a presentation of π_2K as a left $Z\pi_1K$ -module.

Finally, let W, U, R, X be sets of symbols in 1-1 correspondence with the sets of symbols $\partial W, \partial U, r, x$, respectively; and let $\bar{C}_4, \bar{C}_3, \bar{C}_2, \bar{C}_1$, be respectively the free left $Z\pi_1k$ -modules on W, U, R, X . Let $\bar{C}^{(4)}$ denote the left $Z\pi_1K$ -chain complex

$$\bar{C}_4 \xrightarrow{\bar{\partial}_4} \bar{C}_3 \xrightarrow{\bar{\partial}_3} \bar{C}_2 \xrightarrow{\bar{\partial}_2} \bar{C}_1 \xrightarrow{\bar{\partial}_1} Z\pi_1K \xrightarrow{\varepsilon} Z \longrightarrow 0$$

given by

$$\begin{aligned} \bar{\partial}_4 W &= \sum_{U \in U} (\partial W/\partial U)U, & \bar{\partial}_3 U &= \sum_{R \in R} (\partial U/\partial R)R \\ \bar{\partial}_2 R &= \sum_{x \in X} (\partial r/\partial x)X, & \bar{\partial}_1 X &= x - 1, & \varepsilon(\sum_i n_i g_i) &= \sum_i n_i . \end{aligned}$$

Then $\bar{C}^{(4)}$ is a truncation of a free resolution of π_1K . A representative of the k -invariant kK lying in $H^3(\pi_1K; \pi_2K)$ is the equivariant 3-cocycle $k: \bar{C}_3 \rightarrow \pi_2K$ given by

$$k(U) = \begin{cases} \partial U & \text{if } U \in U_{\square} \\ 0 & \text{if } U \in U_{-} \cup U_{+} . \end{cases}$$

Proof. Theorem 9.2 gives a presentation of π_2K and a representative of the k -invariant. Since the presentation $(x_0; r_0)$ of π_1K_0 is chosen to be aspherical, it follows from Theorem 9.3 that $\partial U_0 = \emptyset = \partial W_0$.

Moreover, for both Heegaard splittings of closed connected 3-manifolds (see [32]) and hyperbolic splittings of 2-knots (see Theorem 7.4), π_1K_{\pm} is of homological dimension one. Hence, since

the morphisms $g_{\pm}^{\infty}: \pi_1 K_{\pm} \rightarrow \pi_1 K$ induced by inclusion are both epimorphisms, it follows from Theorem 5.4 that the set ∂U_{\pm} can also be obtained by finding a set of generators of the left $Z\pi_1 K$ -module of solutions $\vec{\lambda}$ of (*) over $Z\pi_1 K$ (rather than $Z\pi_1 K_{\pm}$) such that the components of $\vec{\lambda}$ corresponding to r_{\mp} all vanish. \square

OBSERVATION 1. There exist aspherical presentations $(\mathbf{x}_0; r_0)$ of $\pi_1 K_0$. (A presentation is *aspherical* if its associated 2-complex is aspherical.) Such presentations are easy enough to find. For Heegaard splittings the usual presentation of the fundamental group $\pi_1 K_0$ of the 2-manifold K_0 can be chosen. (See [7].) For hyperbolic splittings, the Wirtinger presentation of $\pi_1 K_0$ with one relator deleted is aspherical. (See [11].) Moreover, any presentation derived from an aspherical presentation by applying a finite sequence of Tietze (II)^{±1} transformations [11] is aspherical. There are many other transformations which preserve presentation asphericity. (See [41, 42].)

OBSERVATION 2. Since $g_-: \pi_1 K_0 \rightarrow \pi_1 K_-$ and $g_+: \pi_1 K_0 \rightarrow \pi_1 K_+$ are both epimorphisms, a presentation of $\pi_1 K_{\pm}$ of the form $(\mathbf{x}_0; r_0, r_{\pm})$ can be obtained from a presentation $(\mathbf{x}_0; r_0)$ by adding relators r_{\pm} . For Heegaard splittings simply add one relator for each handle, i.e., the relator carried by a meridional 2-cell of each handle. For hyperbolic splittings of 2-knot exteriors, simply adjoin one relator for each hyperbolic point, i.e., adjoin the upper (if +) or lower (if -) hyperbolic relators. (See Fig. 8.)

OBSERVATION 3. Since the groups $\pi_1 K_{\pm}$ are free for both




FIGURE 8. Hyperbolic relators.

Heegaard splittings of 3-manifolds (see [32]) and hyperbolic splittings of 2-knots (see Theorem 7.4), the reader may find it easier to find the ∂U_{\pm} as follows. First, find a set $\partial U'_{\pm}$ of generators of the left $Z\pi_1 K_{\pm}$ -module of solutions $\vec{\lambda}$ of (*) over $Z\pi_1 K_{\pm}$ for which the components of $\vec{\lambda}$ corresponding to r_{\mp} all vanish. Then ∂U_{\pm} is simply the set of solutions of (*) over $Z\pi_1 K$ induced by $\partial U'_{\pm}$.

The reader may find the following Hint of use in computing the algebraic 2-type.

HINT. (Solving equations over group rings.)

Let

$$(1) \quad \left(\sum_i x_i a_i\right)d = 0$$

be a linear equation in unknowns $\{x_i\}$ over a group ring ZG (i.e., $d \in ZG$ and $a_i \in ZG \forall i$).

Case 1. If d is not a divisor of zero, then equation (1) can be replaced by

$$\sum_i x_i a_i = 0.$$

(The element d can be shown not to be a divisor of zero by the methods of [66, § 8].)

Case 2. If d is a divisor of zero, then equation (1) can be replaced by the equation

$$\sum_i x_i a_i = \sum_j b_j d_j^+,$$

where $\{d_j^+\}$ is a set of generators of the right ideal of left annihilators of d and where the b_j 's are arbitrary.

Appendix B. Explicit calculations for 2-knots.

We now use Theorem A2.1 of Appendix A to compute the algebraic 2-type of a number of 2-knots. (The methods of [45] can be used to compute π_3 of these examples. See Cor. 11.1 above and following remark. From [49], the Euler characteristic of the embedded connected surface is the number of elliptic points minus the number of hyperbolic points.)

EXAMPLE 1. The diagram of a spun trefoil is given in Figure 5 page 355. (This is Example 9 of [21] and Example 1 of [41, 42].)

A presentation of $\pi_1 K_0$ is $(a, b, c, z: r_0)$, where

$$r_0 = caca^{-1}c^{-1}bab^{-1}a^{-1}b^{-1}.$$

The hyperbolic relators r_+ and r_- are:

$$\begin{cases} r_{+1} = bc^{-1} \\ r_{+2} = zba^{-1}b^{-1} \end{cases} \quad \begin{cases} r_{-1} = za^{-1}b^{-1}a \\ r_{-2} = bc^{-1}. \end{cases}$$

Hence, $(a, b, c, z; r_{-2}, r_{-1}, r_0, r_{+1}, r_{+2})$ which simplifies to

$$(a, b: abab^{-1}a^{-1}b^{-1})$$

is a presentation of $\pi_1 K$. Thus equation (*) of Theorem A2.1 becomes

$$\vec{0} = (\lambda_2, \lambda_1, \lambda_0, \lambda_{-1}, \lambda_{-2}) \begin{pmatrix} \text{(a)} & \text{(b)} & \text{(c)} & \text{(z)} \\ \hline -b & bab^{-1}-1 & 0 & 1 \\ 0 & 1 & -1 & 0 \\ \hline 0 & -1+a-ba & 1-a+ba & 0 \\ \hline a^{-1}-a^{-1}b & -a^{-1} & 0 & 1 \\ 0 & 1 & -1 & 0 \end{pmatrix} \begin{matrix} (r_2) \\ (r_1) \\ (r_0) \\ (r_{-1}) \\ (r_{-2}) \end{matrix}.$$

By the methods of the Hint in Appendix A, the solutions $\partial U_- \cup \partial U_\square \cup \partial U_+$ of equation (*) of Theorem A2.1 are found to be:

$$\begin{cases} \partial U_{+1} = (0, 1-a+ba, 1, 0, 0) \\ \partial U_\square = (0, 1, 0, 0, -1) \\ \partial U_{-1} = (0, 0, 1, 0, 1-a+ba). \end{cases}$$

Thus equation (**) of Theorem A2.1 becomes

$$\vec{0} = (\mu_{+1}, \mu_\square, \mu_{-1}) \begin{pmatrix} \partial U_{+1} \\ \hline \partial U_\square \\ \hline \partial U_{-1} \end{pmatrix}.$$

By the methods of the Hint in Appendix A, the solutions ∂W of equation (**) of Theorem A.1 are:

$$\partial W = (1, -1 + a - ba, -1).$$

Hence,

$$\pi_2 K = (\partial U_\square: (1 - a + ba)\partial U_\square = 0),$$

and

$$H^3(\pi_1 K; \pi_2 K) = 0, \text{ and hence } kK = 0.$$

EXAMPLE 2(n). The diagram of Example 2(n) is given in Figure

6, page 356. (This is Example 12 and 15 of [21] and Example 2 of [41, 42].)

A presentation of $\pi_1 K_0$ is

$$(a, x, z_+, z_-: r_0),$$

where

$$r_0 = u^n z_- u^{-n} x^{-1} a^{-1} z_+^{-1} x$$

and

$$u = z_-^{-1} a^{n+1} x a^{-(n+1)} z_+^{-1} x.$$

The hyperbolic relators r_+ and r_- are:

$$\begin{cases} r_{+1} = z_+ a^{n+1} x^{-1} a^{-n} \\ r_{+2} = z_- x^{-1} \end{cases} \quad \begin{cases} r_{-1} = z_+ x^{-1} \\ r_{-2} = z_- a^n x^{-1} a^{-(n+1)}. \end{cases}$$

Hence,

$$(a, x: a^{2n+1} = 1, ax = xa^{-1})$$

is a presentation of $\pi_1 K$. Thus, by the methods of Hint in Appendix A, the solutions $\partial U_- \cup \partial U_\square \cup \partial U_+$ of equation (*) of Theorem A2.1 are:

$$\begin{cases} \partial U_{+1} = (A, B, 1, 0, 0) \\ \partial U_\square = (1-a, (a-1)a^{n+1}, 0, -(a-1)a^{n+1}, -(1-a)) \\ \partial U_{-1} = (0, 0, 1, B, A) \end{cases},$$

where

$$\begin{cases} A = x^{-1} \sum_0^{n-1} a^i - \sum_1^{n+1} a^i \\ B = x^{-1} \sum_0^n a^i - \sum_2^{n+1} a^i. \end{cases}$$

The solutions ∂W of equation (***) of Theorem A2.1 are:

$$\begin{cases} \partial W_1 = (1-a, x^{-1} a^{-1} \sum_0^{n-1} a^i + \sum_1^{n+1} a^i, -(1-a)) \\ \partial W_2 = (0, \sum_0^{2n} a^i, 0) \end{cases}.$$

(See Hint in Appendix A.)

Hence,

$$\pi_2 K = (\partial U_\square: (x^{-1} a^{-1} \sum_0^{n-1} a^i + \sum_1^{n+1} a^i) \partial U_\square = 0 = (\sum_0^{2n} a^i) \partial U_\square)$$

which simplifies to

$$\pi_2 K = \left(\partial U_{\square} : (x - 1) \partial U_{\square} = 0 = \left(\sum_0^{2n} a^i \right) \partial U_{\square} \right),$$

and

$$kK = t,$$

where

$$H^3(\pi_1 K; Z\pi_2 K) = Z_{2n+1} = (t: t^{2n+1} = 1).$$

EXAMPLE 3. The diagram of Example 3 is given in Figure 7, page 9. (This is Example 10 of [21] and Example 3 of [41, 42].)

$$\begin{aligned} \pi_1 K_0 &= (a, x, y, z: r_0) \\ r_0 &= ayz^{-1}xzy^{-1}x^{-1}a^{-1}yz^{-1} \\ \begin{cases} r_{+1} = axa^{-2}y^{-1} \\ r_{+2} = yz^{-1} \end{cases} & \quad \begin{cases} r_{-1} = yz^{-1} \\ r_{-2} = xz^{-1}. \end{cases} \\ \pi_1 K &= (a, x: ax = xa^2). \end{aligned}$$

By the methods found in the Hint of Appendix A, we have:

$$\begin{cases} \partial U_{+1} = (-1 - a + ax, 0, 1, & 0, & 0) \\ \partial U_{\square} = (& 1, & 0, 0, & -1, & 0) \\ \partial U_{-1} = (& 0, & 0, 1, & -1 - a + ax, & 0). \end{cases}$$

$$\partial W = (1, 1 + a - ax, -1).$$

Hence,

$$\pi_2 K = (\partial U_{\square} : (1 + a - ax) \partial U_{\square} = 0),$$




FIGURE 9. Diagram of an embedding of RP^2 in S^4 .

$$\pi_1(S^4 - RP^2) = (a: a^2) = Z_2$$

$$\pi_2(S^4 - RP^2) = (\partial U_{\square} : (1+a)\partial U_{\square} = 0) = Z \text{ (twisted action)}$$

and

$$H^3(\pi_1 K; \pi_2 K) = 0 \text{ and } kK = 0 .$$

EXAMPLE 4. The diagram of an embedding in S^4 of the real projective plane RP^2 is given in Figure 9. Using the methods of Appendix A, we have

$$\begin{cases} \pi_1(S^4 - RP^2) = (a, b, c: ab^{-1}, ac^{-1}, ab) = (a: a^2) = Z_2 \\ \pi_2(S^4 - RP^2) = (\partial U_{\square}: (1+a)\partial U_{\square} = 0) = Z \text{ twisted action} . \end{cases}$$

(See [54].)

Addendum (Section XII continued)

PROBLEM 13. *Let K be the exterior of a smooth (or locally flat PL) 2-knot (S^4, KS^2) . Is $\pi_2 K$ always finitely related as a left $Z\pi_1 K$ -module?*

PROBLEM 14. *Let K be as in Problem 13. If $\pi_2 K$ vanishes, then is K always a spherical? (See Problem 4.)*

PROBLEM 15. *Does there exist a 2-knot whose fundamental group has no non-trivial elements of finite order but whose K -invariant is non-zero? (See Corollary 11.7.)*

PROBLEM 16. *Do there exist two 2-knots with exteriors K and K' respectively such that $\pi_1 K \cong \pi_1 K'$ and $\pi_2 K \cong \pi_2 K'$ (as left $Z\pi_1 K$ -modules) but with distinct K -invariants? (See [27].)*

REFERENCES

1. J. J. Andrews and M. L. Curtis, *Knotted 2-spheres in the 4-sphere*, Ann. of Math., **70** (1959), 565-571.
2. J. J. Andrews and S. L. Lomonaco, *The second homotopy group of spun 2-spheres in 4-space*, Ann. Math., **90**, No. 2 (1969), 199-204.
3. J. J. Andrews and D. W. Sumners, *On Higher-dimensional fibered knots*, A.M.S. Trans., **153** (1971), 415-426.
4. E. Artin, *Zur Isotopie Zweidimensionaler Flächen in R^4* , Abh. Math. Seminar Univ. Hamburg, **4** (1925), 174-177.
5. William Beckmann, Personal communication.
6. Joan S. Birman, *Braids, links, and mapping class groups*, Annals of Math. Studies, No. 82, Princeton University Press, Princeton, N. J., 1974.
7. S. S. Cairns, *Introductory Topology*, Ronald, New York, N. Y., 1962.
8. Henri Cartan and Samuel Eilenberg, *Homological Algebra*, Princeton University Press, Princeton, N. Y., 1956.
9. Richard H. Crowell, *H_2 of subgroups of knot groups*, Illinois J. Math., (1970), 665-673.
10. Richard H. Crowell, *On the Van Kampen theorem*, Pacific J. Math., **9**, No. 1 (1959), 43-50.

11. Richard H. Crowell and Ralph H. Fox, *Introduction to Knot Theory*, Springer-Verlag, New York, 1977.
12. Eldon Dyer and A. T. Vasquez, *The asphericity of higher dimensional knots*, *Canad. J. Math.*, **25**, No. 6 (1973), 1132-1136.
13. Beno Eckmann, *Aspherical manifolds and higher-dimensional knots*, *Comment. Math. Helvetici*, **51** (1976), 93-98.
14. Samuel Eilenberg and Saunders MacLane, *Determination of the second homology and cohomology groups of a space by means of homotopy invariants*, *Proc. Nat. Acad. Sci.*, (1948), 277-280.
15. ———, *Homology of spaces with operators II*, *Amer. Math. Soc. Trans.*, **65**, No. 1 (1949), 49-99.
16. Samuel Eilenberg and Norman Steenrod, *Foundations of Algebraic Topology*, Princeton University Press, Princeton, N. J., 1952.
17. D. B. A. Epstein, *Linking spheres*, *Proc. Camb. Phil. Soc.*, **56**, Part 3 (1960), 215-219.
18. Ralph H. Fox, *Free differential calculus. I Derivation in the free group ring*, *Ann. of Math.*, **57**, No. 3 (1953), 547-560.
19. ———, *Free differential calculus. II The isomorphism problem of groups*, *Ann. of Math.*, **59** No. 2 (1954), 196-210.
20. ———, *Free differential calculus. V The Alexander matrices re-examined*, *Ann. of Math.*, **71**, No. 2, (1960), 408-422.
21. Ralph H. Fox, *A quick trip through knot theory*, in "Topology of 3-Manifolds and Related Topics," edited by M. K. Fort, Jr., Prentice-Hall, Inc., Englewood Cliffs, N. J., (1962), 120-167.
22. ———, *Some problems in knot theory*, in "Topology of 3-Manifolds and Related Topics," edited by M. K. Fort, Jr., Prentice-Hall, Inc., Englewood Cliffs, N. J., (1962), 168-176.
23. R. H. Fox and J. W. Milnor, *Singularities of 2-spheres in 4-space and equivalence of knots*, (unpublished version).
24. ———, *Singularities of 2-spheres in 4-space and cobordism of knots*, *Osaka J. Math.*, **3** (1966), 257-267.
25. H. Gluck, *The embeddings of two-spheres in the four-sphere*, *Amer. Math. Soc. Trans.*, **104** (1962), 308-333.
26. F. González-Acuna and J. M. Montesinos, *Ends of knots*, *Ann. of Math.*, **108**, No. 1 (1978), 91-96.
27. C. McA Gordon, *Some higher-dimensional knots with the same homotopy groups*, *Quart. J. Math. Oxford*, **24**, No. 95 (1973), 411-422.
28. ———, *Problems*, "Knot Theory," edited by J. C. Hausmann, Springer-Verlag, New York, (1978), 309-311.
29. M. A. Gutierrez, *An exact sequence calculation for the second homotopy of a knot*, *Amer. Math. Soc. Proc.*, **32**, No. 2 (1972), 571-577.
30. M. A. Gutierrez, *An exact sequence calculation for the second homotopy group. II*, *Amer. Math. Soc. Proc.*, **40**, No. 1 (1973), 327-330.
31. John R. Harper and S. J. Lomonaco, Jr., *The k-invariant*, in preparation.
32. John Hempel, 3-Manifolds, *Annals of Math. Studies*, No. 86, Princeton Univ. Press, Princeton, N. J., 1976.
33. Sze-Tsen Hu, *Homotopy Theory*, Academic Press, New York, 1959.
34. A. Kawachi, *On partial Poincare duality and higher-dimensional knots with $\pi_1 = Z$* , preprint, Kobe University.
35. M. Kervaire, *Les noeds de dimensions supérieures*, *Bull. Soc. Math. France*, **93** (1965), 225-271.
36. ———, *On higher dimensional knots*, "Differential and Combinatorial Topology," Edited by Steward S. Cairns, Princeton Univ. Press, Princeton, N. J., (1965), 105-119.
37. M. Kervaire and Cl. Weber, *A survey of multidimensional knots*, "Knot Theory,"

- edited by J. C. Hausmann, Springer-Verlag, New York, (1978), 61-134.
38. Jerome Levine, *Knot modules I*, A.M.S. Trans., **229** (1977), 1-50.
 39. S. J. Lomonaco, Jr., *Finitely ended knots are quasi-aspherical*, submitted.
 40. ———, *The fundamental ideal and π_2 of higher dimensional knots*, Amer. Math. Soc. Proc., **38**, No. 2, (1973), 431-433.
 41. ———, *The homotopy groups of knots. I. The four dimensional analogue of the asphericity of knots*, (manuscript), version 9-14-77.
 42. ———, *The homotopy group of knots. II. A solution to problem 36 of R.H. Fox*, (manuscript), version 9-14-77.
 43. ———, *The homology of group systems with applications to low dimensional topology*, A.M.S. Bull., **3**, No. 3 (1980), 1049-1052.
 44. ———, *The second homotopy group of a spun knot*, Topology, **8** (1969), 95-98.
 45. ———, *The third homotopy group of some higher dimensional knots*, in "Knots, Groups and 3-Manifolds" edited by L.P. Neuwirth, Annals of Math. Studies, Princeton Univ. Press, Princeton, N. J., (1975), 35-43.
 46. Roger C. Lyndon, *Cohomology theory of groups with a single defining relation*, Ann. Math., **52**, No. 3 (1950), 650-665.
 47. Saunders MacLane, *Homology*, Academic Press, New York, N. Y., 1963.
 48. Saunders MacLane and J. H. C. Whitehead, *On the 3-type of a complex*, Proc. Nat. Acad. Sci., **36** (1950), 41-48.
 49. J. Milnor, *Morse Theory*, Annals of Math. Studies, No. 51, Princeton Univ. Press, Princeton, N. J., 1963.
 50. Robert E. Mosher and Martin C. Tangora, *Cohomology Operations and Applications in Homotopy Theory*, Harper & Row, N. Y., 1968.
 51. L. P. Neuwirth, *Knot Groups*, Annals of Math. Studies, No. 56, Princeton Univ. Press, Princeton, N. J., 1965.
 52. D. G. Northcott, *An Introduction to Homological Algebra*, Cambridge Univ. Press, New York, N. Y., 1960.
 53. C. D. Papakyriakopoulos, *On Dehn's lemma and the asphericity of knots*, Ann. Math., **66**, No. 1 (1957), 1-26.
 54. T. M. Price and D. M. Roseman, *Embeddings of the projective plane in four space*, manuscript.
 55. John Ratcliffe, *On the ends of higher dimensional knot groups*, Quart. J. Oxford (to appear).
 56. Kurt Reidemeister, *Complexes and homotopy chains*, Amer. Math. Soc. Bull., (1950), 297-307.
 57. ———, *Homotopiegruppen von Komplexen*, Hamburg. Abh., **10** (1934), 211-215.
 58. ———, *Überdeckungen von Komplexen*, J. reine angew. Math., **173** (1935), 164-173.
 59. Dale Rolfsen, *Knots and Links*, Publish or Perish, Berkeley, CA., 1976.
 60. Edwin H. Spanier, *Algebraic Topology*, McGraw-Hill, New York, N. Y., 1966.
 61. D. W. Sumners, *On asphericity of knots S^2 in S^4* , preprint.
 62. Shinichi Suzuki, *Knotting problems of 2-spheres in 4-sphere*, Mathematics Seminar Notes, Kobe Univ., **4**, No. 3, Nada, Kobe, Japan, (1976).
 63. Richard G. Swan, *Groups of cohomological dimension one*, J. Algebra, **12**, (1969), 585-601.
 64. G. A. Swarup, *An unknotting criterion*, J. Pure and Appl. Algebra, **6** (1975), 291-296.
 65. H. F. Trotter, *Homology of group systems with applications to knot theory*, Ann. Math., **76**, No. 3, (1962), 464-498.
 66. J. H. C. Whitehead, *On the asphericity of regions in a 3-sphere*, Fund. Math., **32** (1939), 149-166.
 67. ———, *A certain exact sequence*, Ann. Math., **52** (1950), 51-110.
 68. ———, *Combinatorial homotopy I*, Amer. Math. Soc. Bull., **55** (1949), 213-245.

69. J. H. C. Whitehead, *Combinatorial homotopy II*, Amer. Math. Soc. Bull., **55** (1949), 453-496.
70. ———, *The second boundary operator*, Proc. Nat. Acad. Sci., **36** (1950), 55-60.
71. E. C. Zeeman, *Twisting spun knots*, Amer. Math. Soc. Trans., **115** (1965), 471-495.

Received May 21, 1979 and in revised form June 10, 1980. Supported in part by State University of New York Summer Research Grant and also by the L-O-O-P Fund.

STATE UNIVERSITY OF NEW YORK
ALBANY, NY 12222
AND
INSTITUTE FOR DEFENSE ANALYSES
PRINCETON, NJ 08540

Current Address: 7004 South Ridge Drive
Dallas, TX 75214

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ANDREW OGG

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$102.00 a year (6 Vols., 12 issues). Special rate: \$51.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1981 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

George E. Andrews, The Rogers-Ramanujan reciprocal and Minc's partition function	251
Allan Calder, William H. Julian, Ray Mines, III and Fred Richman, ε -covering dimension	257
Thomas Curtis Craven and George Leslie Csordas, An inequality for the distribution of zeros of polynomials and entire functions	263
Thomas Jones Enright and R. Parthasarathy, The transfer of invariant pairings to lattices	281
Allen Roy Freedman and John Joseph Sember, Densities and summability	293
Robert Heller and Francis Aubra Roach, A generalization of a classical necessary condition for convergence of continued fractions	307
Peter Wilcox Jones, Ratios of interpolating Blaschke products	311
V. J. Joseph, Smooth actions of the circle group on exotic spheres	323
Mohd Saeed Khan, Common fixed point theorems for multivalued mappings	337
Samuel James Lomonaco, Jr., The homotopy groups of knots. I. How to compute the algebraic 2-type	349
Louis Magnin, Some remarks about C^∞ vectors in representations of connected locally compact groups	391
Mark Mandelker, Located sets on the line	401
Murray Angus Marshall and Joseph Lewis Yucas, Linked quaternionic mappings and their associated Witt rings	411
William Lindall Paschke, K -theory for commutants in the Calkin algebra	427
W. J. Phillips, On the relation $PQ - QP = -iI$	435
Ellen Elizabeth Reed, A class of Wallman-type extensions	443
Sungwoo Suh, The space of real parts of algebras of Fourier transforms	461
Antonius Johannes Van Haagen, Finite signed measures on function spaces	467
Richard Hawks Warren, Identification spaces and unique uniformity	483