

Pacific Journal of Mathematics

CERTAIN TRANSFORMATIONS OF BASIC HYPERGEOMETRIC SERIES AND THEIR APPLICATIONS

V. K. JAIN

CERTAIN TRANSFORMATIONS OF BASIC HYPERGEOMETRIC SERIES AND THEIR APPLICATIONS

V. K. JAIN

We obtain identities of Rogers-Ramanujan type related to the modulus 13. We also obtain the q -analogues of the nearly-poised summation theorems and use them for obtaining q -analogues of general transformations of nearly-poised hypergeometric series. We also discuss some important applications of the transformations obtained in this note.

Recently, Askey and Wilson [4] derived the transformation

$$(1.1) \quad {}_4\phi_3 \left[\begin{matrix} a^2, b^2, c, d; q; q \\ ab\sqrt{q}, -ab\sqrt{q}, -cd \end{matrix} \right] = {}_4\phi_3 \left[\begin{matrix} a^2, b^2, c^2, d^2; q^2; q^2 \\ a^2b^2q, -cd, -cdq \end{matrix} \right],$$

(provided a, b, c , or d is of the form q^{-N} , N a nonnegative integer). In an earlier paper [11] we have an alternative proof of (1.1). We begin this note by showing in §3 that all the transformations proved by Singh [13], for obtaining the q -analogues of identities of the Cayley-Orr type, can be deduced from (1.1). We also show that (1.1) may be used effectively to prove the following transformation:

$$(1.2) \quad {}_8\phi_6 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, iq^{-n}, -iq^{-n}, -q^{-n}, q^{-n}, o; q; -aq^{1+4n} \\ \sqrt{a}, -\sqrt{a}, -iaq^{1+n}, iaq^{1+n}, -aq^{1+n}, aq^{1+n} \end{matrix} \right] \\ = \frac{[aq; q]_{4n}}{[a^4q^{4+4n}; q^4]_n} {}_3\phi_2 \left[\begin{matrix} -q^{-2n}, q^{-2n}, o; q^2; q^2 \\ q^{-4n}/a, q^{1-4n}/a \end{matrix} \right],$$

due to Andrews [2] which is his key result for obtaining the identities of the Rogers-Ramanujan type of modulus 11. In fact, we shall prove the transformation:

$$(1.3) \quad {}_8\phi_7 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, c, e, -e, -q^{-n}, q^{-n}; q; \frac{a^2q^{2+2n}}{ce^2} \\ \sqrt{a}, -\sqrt{a}, aq/c, aq/e, -aq/e, -aq^{1+n}, aq^{1+n} \end{matrix} \right] \\ = \frac{[a^2q^2; q^2]_n [-aq/e^2; q]_{2n}}{[a^2q^2/c^2; q^2]_n [-aq; q]_{2n}} \\ \times \frac{[a^2q^2/c^2e^2; q^2]_ne^{2n}}{[a^2q^2/e^2; q^2]_n} {}_4\phi_3 \left[\begin{matrix} \frac{ce^2}{a^2}q^{-2n-1}, \frac{ce^2}{a^2}q^{-2n}, e^2, q^{-2n}; q^2; q^2 \\ \frac{e^2}{a^2}c^2q^{-2n}, -\frac{e^2}{a}q^{-2n}, -\frac{e^2}{a}q^{1-2n} \end{matrix} \right]$$

which is a generalization of (1.2) and to which it reduces for $e = iq^{-n}$, $c \rightarrow 0$. (1.3) can be used with advantage for obtaining the identities of Rogers-Ramanujan type related to the modulus 13, not given, thus far. In the sequel, we also present a generalization of (1.1) along the lines of a similar result of Burchall and Chaundy [9].

In § 4, we prove the q -analogue of the summation theorem for the nearly-poised ${}_4F_3(1)$:

$$(1.4) \quad {}_4F_3 \left[\begin{matrix} 2a, 1+a, c, -N \\ a, 1+2a-c, 1+2c-N \end{matrix} ; - \right] = \frac{(2a-2c)_N(-c)_N}{(1+2a-c)_N(-2c)_N},$$

in the form

$$(1.5) \quad {}_4\phi_3 \left[\begin{matrix} a^2, aq, c, q^{-N}; q; q \\ a, a^2q/c, c^2q^{1-N} \end{matrix} \right] = \frac{[a^2/c^2; q]_N [c^{-1}; q]_N [-aq/c; q]_N}{[a^2q/c; q]_N [c^{-2}; q]_N [-a/c; q]_N}.$$

This result also gives the q -analogue of the summation theorem for nearly-poised ${}_3F_2$, viz.

$$(1.6) \quad {}_3F_2 \left[\begin{matrix} 2a, c, -N \\ 1+2a-c, 1+2c-N \end{matrix} ; - \right] = \frac{(2a-2c)_N(1+a-c)_N(-c)_N}{(1+2a-c)_N(a-c)_N(-2c)_N},$$

on replacing ' a ' by ' $-a$ ' and then proceeding to the limits in the usual way.

In this connection it may be of interest to note that Andrews had obtained a q -analogue of (1.6) in the form

$$(1.7) \quad {}_4\phi_3 \left[\begin{matrix} a^2, c, q^{-N}, -a^2q^2/c; q; q \\ a^2q/c, c^2q^{1-N}, -a^2q/c \end{matrix} \right] = \frac{[c^{-1}; q]_N [a^2c^{-2}; q]_N}{[a^2qc^{-1}; q]_N [c^{-2}; q]_N} \\ \times \frac{\{(1+a^2c^{-1})(1-a^2c^{-2}q^{1+N}) + a^2qc^{-1}(1-q^{N-1})(1+c^{-1})\}}{(1-a^2c^{-2})(1+a^2c^{-1}q)}.$$

However, in view of the identity

$$(1.8) \quad {}_4\phi_3 \left[\begin{matrix} a^2, c, q^{-N}, -a^2q^2/c; q; q \\ a^2q/c, c^2q^{1-N}, -a^2q/c \end{matrix} \right] = \frac{(1+a)}{(1+a^2qc^{-1})} {}_4\phi_3 \left[\begin{matrix} a^2, -aq, c, q^{-N}; q; q \\ -a, a^2q/c, c^2q^{1-N} \end{matrix} \right] \\ - \frac{a(1-aqc^{-1})}{(1+a^2qc^{-1})} {}_3\phi_2 \left[\begin{matrix} a^2, c, q^{-N}; q; q^2 \\ a^2q/c, c^2q^{1-N} \end{matrix} \right] \\ = \frac{(1+a)}{(1+a^2qc^{-1})} {}_4\phi_3 \left[\begin{matrix} a^2, -aq, c, q^{-N}; q; q \\ -a, a^2q/c, c^2q^{1-N} \end{matrix} \right] \\ - \frac{a(1-aqc^{-1})}{(1+a^2qc^{-1})} {}_3\phi_2 \left[\begin{matrix} a^2, cq, q^{-N}; q; q \\ a^2q/c, c^2q^{1-N} \end{matrix} \right]$$

$$+ \frac{aq(1-aqc^{-1})(1-a^2)(1-q^{-N})}{(1+a^2qc^{-1})(1-a^2qc^{-1})(1-c^2q^{1-N})} {}_3\phi_2 \left[\begin{matrix} a^2q, cq, q^{1-N}; q; q \\ a^2q^2/c, c^2q^{2-N} \end{matrix} \right],$$

(the last two series may be summed by q -analogue of Saalschütz summation theorem), it is not difficult to show the equivalence of the summation theorems (1.5) and (1.7). However, we prefer to stick to the form (1.5) as it has the added advantage that it gives the q -analogue of (1.4) as well as of (1.6) in an straight forward form.

The summation (1.5) has been further employed for obtaining two transformations connecting a terminating nearly-poised Saalschützian ${}_6\phi_5$ into a terminating well-poised ${}_{12}\phi_{11}$. It may be remarked that Bailey in his book [7] has mentioned four known transformations of nearly-poised hypergeometric series [7; 4.5 (3-6)]. The q -analogues of two of these [7; 4.5(3) and 4.5(6)] only were obtained by Bailey [8]. The above two transformations deduced by us are q -analogues of the remaining two transformations 4.5(4) and 4.5(5) given in Bailey's Tract [7]. We conclude the paper by obtaining the summation formula

$$(1.9) \quad {}_{10}\phi_9 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, \frac{a}{d}, \frac{a^2}{d}q^{1+N}, \sqrt{d}, -\sqrt{d}, \sqrt{dq}, -\sqrt{dq}, q^{-N}; q; q \\ \sqrt{a}, -\sqrt{a}, dq, \frac{d}{a}q^{-N}, \frac{aq}{\sqrt{d}}, -\frac{aq}{\sqrt{d}}, a\sqrt{\frac{q}{d}}, -a\sqrt{\frac{q}{d}}, aq^{1+N} \end{matrix} \right] \\ = \frac{[aq; q]_N [a^2qd^{-2}; q]_N}{[aqd^{-1}; q]_N [a^2qd^{-1}; q]_N},$$

which is a q -analogue of a summation theorem for well-poised ${}_7F_6(1)$ (different from the Dougall's theorem) due to Bailey [7; Ex. 8, p. 98] (see also [6]).

2. Definitions and notations. If we let,

$$|q| < 1, [a; q]_n = (1-a)(1-aq) \cdots (1-aq^{n-1}), [a; q]_0 = 1$$

and

$$[a; q]_\infty = \prod_{r=0}^{\infty} (1-aq^r),$$

then we may define the basic hypergeometric series as

$${}_{p+1}\phi_{p+r} \left[\begin{matrix} a_1, a_2, \dots, a_{p+1}; q; x \\ b_1, b_2, \dots, b_{p+r} \end{matrix} \right] \\ = \sum_{n=0}^{\infty} \frac{[a_1; q]_n \cdots [a_{p+1}; q]_n (-)^{r^n} x^n q^{(r/2)n(n-1)}}{[q; q]_n [b_1; q]_n \cdots [b_{p+r}; q]_n},$$

where the series ${}_{p+1}\phi_{p+r}(x)$ converges for all positive integral values of r and for all x , except when $r = 0$, it converges only for $|x| < 1$.

Further, we shall denote by

$$\Pi \left[\begin{matrix} a_1, a_2, \dots, a_r; q \\ b_1, b_2, \dots, b_s \end{matrix} \right], \text{ the infinite product } \prod_{j=0}^{\infty} \frac{(1 - a_1 q^j) \cdots (1 - a_r q^j)}{(1 - b_1 q^j) \cdots (1 - b_s q^j)}.$$

3. For obtaining the transformations used by Singh [13] to obtain the q -analogues of identities of Cayley-Orr, we begin by setting $b = ak$, $c = -a^2 b^2 k^2 q^N$ and $d = q^{-N}$ in (1.1) to obtain

$$(3.1) \quad {}_4\phi_3 \left[\begin{matrix} a^2, a^2 k^2, -a^2 b^2 k^2 q^N, q^{-N}; q; q \\ a^2 k \sqrt{q}, -a^2 k \sqrt{q}, a^2 b^2 k^2 \end{matrix} \right] = {}_4\phi_3 \left[\begin{matrix} a^2, a^2 k^2, a^4 b^4 k^4 q^{2N}, q^{-2N}; q^2; q^2 \\ (abk)^2 q, (abk)^2, a^4 k^2 q \end{matrix} \right].$$

Using the transformation [12; 8.3]

$$(3.2) \quad {}_4\phi_3 \left[\begin{matrix} a, b, c, q^{-N}; q; q \\ e, g, h \end{matrix} \right] = \frac{\left[\frac{g}{c}; q \right]_N \left[\frac{eg}{ab}; q \right]_N}{[g; q]_N \left[\frac{eg}{cab}; q \right]_N} {}_4\phi_3 \left[\begin{matrix} \frac{e}{a}, \frac{e}{b}, c, q^{-N}; q; q \\ e, \frac{c}{g} q^{1-N}, \frac{c}{h} q^{1-N} \end{matrix} \right],$$

(where $abc = eghq^{N-1}$) on both the sides of (3.2) (in the left hand side with $a \rightarrow a^2 k^2$, $b \rightarrow -a^2 b^2 k^2 q^N$, $c \rightarrow a^2$, $e \rightarrow (abk)^2$, $g \rightarrow a^2 k \sqrt{q}$, $h \rightarrow -a^2 k \sqrt{q}$ and on the right hand side with $q \rightarrow q^2$, $a \rightarrow a^2$, $b \rightarrow (abk)^4 q^{2N}$, $c \rightarrow a^2 k^2$, $e \rightarrow (abk)^2 q$, $g \rightarrow (abk)^2$, $h \rightarrow a^4 k^2 q$), we get

$$(3.3) \quad {}_4\phi_3 \left[\begin{matrix} a^2, b^2, -q^{-N}, q^{-N}; q; q \\ (abk)^2, \frac{1}{k} q^{-N+1}, -\frac{1}{k} q^{-N+1} \end{matrix} \right] = \frac{[a^2 q; q^2]_N [b^2; q^2]_N k^{2N}}{[a^2 b^2 k^2; q^2]_N [k^2 q; q^2]_N} \\ \times {}_4\phi_3 \left[\begin{matrix} a^2 k^2, b^2 k^2 q, (abk)^{-2} q^{1-2N}, q^{-2N}; q^2; q^2 \\ a^2 b^2 k^2 q, \frac{1}{b^2} q^{1-2N}, \frac{1}{a^2} q^{1-2N} \end{matrix} \right].$$

Again, using the transformation (3.2) on the right hand side of (3.3) (with $q \rightarrow q^2$, $a \rightarrow a^2 k^2$, $b \rightarrow 1/(abk)^2 q^{1-2N}$, $c \rightarrow b^2 k^2 q$, $e \rightarrow b^{-2} q^{2-2N}$, $g \rightarrow a^{-2} q^{1-2N}$, $h \rightarrow a^2 b^2 k^2 q$), we have

$$(3.4) \quad {}_4\phi_3 \left[\begin{matrix} a^2, b^2, -q^{-N}, q^{-N}; q; q \\ a^2 b^2 k^2, k^{-1} q^{-N+1}, -k^{-1} q^{-N+1} \end{matrix} \right] = \frac{[a^2; q^2]_N [b^2; q^2]_N [(abk q)^2; q^2]_N k^{2N}}{[a^2 b^2 k^2; q^2]_N [k^2 q; q^2]_N} \\ \times {}_4\phi_3 \left[\begin{matrix} b^2 k^2 q, a^2 k^2 q, (abk)^{-2} q^{2-2N}, q^{-2N}; q^2; q^2 \\ b^{-2} q^{2-2N}, a^{-2} q^{2-2N}, (abk q)^2 \end{matrix} \right].$$

Once again using the transformation (3.2) on the right hand side of (3.4) (with $q \rightarrow q^2$, $a \rightarrow b^2 k^2 q$, $b \rightarrow a^2 k^2 q$, $c \rightarrow (abk)^2 q^{-2-2N}$, $e \rightarrow (abkq)^2$, $g \rightarrow b^{-2} q^{2-2N}$, $h \rightarrow a^{-2} q^{2-2N}$), we get

$$\begin{aligned}
 (3.5) \quad & \left[\begin{matrix} a^2, b^2, -q^{-N}, q^{-N}; q; q \\ a^2 b^2 k^2, k^{-1} q^{-N} + \frac{1}{2}, -k^{-1} q^{-N} + \frac{1}{2} \end{matrix} \right] \\
 &= \frac{[a^2 k^2; q^2]_N [b^2 k^2; q^2]_N [(abkq)^2; q^2]_N}{[(abk)^2; q]_{2N} [k^2 q; q^2]_N} \\
 & \quad {}_4\phi_3 \left[\begin{matrix} a^2 q, b^2 q, (abk)^{-2} q^{2-2N}, q^{-2N}; q^2; q^2 \\ (abkq)^2, b^{-2} k^{-2} q^{2-2N}, a^{-2} k^{-2} q^{2-2N} \end{matrix} \right].
 \end{aligned}$$

(3.5) is one of the results proved by Singh [13]. All the other results due to Singh [13] may be deduced by applying the transformation (3.2) to (3.1) and (3.3) (see [1] for details).

Next, for proving (1.3), we start with the Watson's transformation [14; 3.4.1.5]:

$$\begin{aligned}
 (3.6) \quad & {}_8\phi_7 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, c, d, e, f, q^{-n}; q; \frac{a^2 q^{2+n}}{cdef} \\ \sqrt{a}, -\sqrt{a}, \frac{aq}{c}, \frac{aq}{d}, \frac{aq}{e}, \frac{aq}{f}, aq^{1+n} \end{matrix} \right] \\
 &= \frac{[-aq; q]_n \left[\frac{aq}{ef}; q \right]_n}{\left[\frac{aq}{e}; q \right]_n \left[\frac{aq}{f}; q \right]_n} {}_4\phi_3 \left[\begin{matrix} \frac{aq}{cd}, e, f, q^{-n}; q; q \\ \frac{ef}{a} q^{-n}, \frac{aq}{c}, \frac{aq}{d} \end{matrix} \right].
 \end{aligned}$$

Reversing the order of the series on the right hand side of (3.6), we obtain (on setting $f = -e$, $d = -q^{-n}$):

$$\begin{aligned}
 & {}_8\phi_7 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, c, e, -e, -q^{-n}, q^{-n}; q; \frac{a^2 q^{2+2n}}{ce^2} \\ \sqrt{a}, -\sqrt{a}, \frac{aq}{c}, \frac{aq}{e}, -\frac{aq}{e}, -aq^{1+n}, aq^{1+n} \end{matrix} \right] \\
 &= \frac{(-aq)^n [aq; q]_n [e^2; q^2]_n \left[-\frac{a}{c} q^{1+n}; q \right]_n}{e^{2n} [-aq^{1+n}; q]_n \left[\frac{aq}{c}; q \right]_n \left[\frac{a^2 q^2}{e^2}; q^2 \right]_n} \\
 & \quad \times {}_4\phi_3 \left[\begin{matrix} -\frac{aq}{e^2}, -\frac{q^{-2n}}{a}, \frac{cq^{-n}}{a}, q^{-n}; q; q \\ \frac{q^{1-n}}{e}, -\frac{q^{1-n}}{e}, -\frac{cq^{-2n}}{a} \end{matrix} \right]
 \end{aligned}$$

$$\begin{aligned}
&= \frac{[e^2; q^2]_n [a^2 q^2; q^2]_n \left[-\frac{aq}{c}; q \right]_{2n} (-aq)^n}{[-aq; q]_{2n} \left[\frac{a^2 q^2}{c^2}; q^2 \right]_n \left[\frac{a^2 q^2}{e^2}; q^2 \right]_n e^{2n}} \\
&\quad \times {}_4\phi_3 \left[\begin{matrix} -\frac{aq}{e^2}, -\frac{q^{-2n}}{a}, -\frac{c^2}{a^2} q^{-2n}, q^{-2n}; q^2; q^2 \\ \frac{q^{2-2n}}{e^2}, -\frac{c}{a} q^{-2n}, -\frac{c}{a} q^{1-2n} \end{matrix} \right],
\end{aligned}$$

(using (1.1) with $a^2 \rightarrow -aq/e^2$, $b^2 \rightarrow -q^{-2n}/a$, $c \rightarrow (c/a)q^{-n}$, $d \rightarrow q^{-n}$)

$$\begin{aligned}
&= \frac{[e^2; q^2]_n [a^2 q^2; q^2]_n \left[\frac{a^2 q^2}{e^2 c}; q \right]_{2n}}{\left[\frac{a^2 q^2}{c^2}; q^2 \right]_n \left[\frac{a^2 q^2}{e^2}; q^2 \right]_n [-aq; q]_{2n}} \\
&\quad \times {}_4\phi_3 \left[\begin{matrix} -\frac{aq}{e^2}, -\frac{aq^2}{e^2}, \frac{a^2 q^2}{e^2 c^2}, q^{-2n}; q^2; q^2 \\ \frac{q^{2-2n}}{e^2}, \frac{a^2 q^3}{e^2 c}, \frac{a^2 q^2}{e^2 c} \end{matrix} \right],
\end{aligned}$$

(using (3.2) with $q \rightarrow q^2$, $a \rightarrow -a^{-1}q^{-2n}$, $b \rightarrow c^2 a^{-2} q^{-2n}$, $c \rightarrow -aqe^{-2}$, $e \rightarrow e^{-2}q^{2-2n}$, $g \rightarrow -ca^{-1}q^{-2n}$, $h \rightarrow -ca^{-1}q^{1-2n}$).

Reversing the order of the series ${}_4\phi_3$ in the right hand side of the above expression, we get (1.3).

Furthermore, using (1.3) we prove the following three transformations. These transformations on specialization yield identities of Rogers-Ramanujan type related to the moduli 11 and 13:

$$\begin{aligned}
(3.7) \quad & [a^4 q^4; q^4]_\infty \sum_{n=0}^{\infty} \frac{[aq; q]_{4n} [-a^2 q^2 c^{-2}; q^2]_{2n} (-)^n a^{4n} q^{2n(n-2p)}}{[q^4; q^4]_n [a^4 q^4; q^4]_{2n} [a^4 q^4 c^{-4}; q^4]_n} \\
& \quad \times {}_4\phi_3 \left[\begin{matrix} -ca^{-2} q^{-4n-1}, -ca^{-2} q^{-4n}, -q^{-2n}, q^{-2n}; q^2; q^2 \\ -c^2 a^{-2} q^{-4n}, a^{-1} q^{-4n}, a^{-1} q^{1-4n} \end{matrix} \right] \\
&= \sum_{s=0}^p \frac{[q^{-4p}; q^4]_s (-)^s a^{4s} q^{2s(s+1)}}{[q^4; q^4]_s} \\
& \quad \times \sum_{n=0}^{\infty} \frac{[a; q]_n (1 - aq^{2n}) [c; q]_n a^{6n} q^{2n(3n-2p+4s)}}{[q; q]_n (1 - a) \left[\frac{aq}{c}; q \right]_n c^n},
\end{aligned}$$

$$\begin{aligned}
(3.8) \quad & [a^4 q^4; q^4]_\infty \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[aq; q]_{4n+2r} (-)^n a^{4n+4r} q^{2n^2+3r^2+4nr+4n+3r}}{[q^2; q^2]_r [q^4; q^4]_n [a^4 q^4; q^4]_{2n+2r} (1 + a^2 q^{4n+4r+2})} \\
&= \sum_{n=0}^{\infty} \frac{[aq; q]_n (1 - a^2 q^{4n+2}) (-)^n a^{6n} q^{(1/2)n(13n+9)}}{[q; q]_n}
\end{aligned}$$

and

$$\begin{aligned}
 (3.9) \quad [a^4q^4; q^4]_\infty & \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[aq; q]_{4n+2r}(-)^r a^{4n+6r} q^{4n^2+8r^2+12nr+4n+5r}}{[q^2; q^2]_r [q^4; q^4]_n [a^4q^4; q^4]_{2n+2r} (1 + a^2 q^{4n+4r+2})} \\
 & = \sum_{n=0}^{\infty} \frac{[aq; q]_n (1 - a^2 q^{4n+2}) (-)^n a^{5n} q^{(1/2)n(11n+7)}}{[q; q]_n}.
 \end{aligned}$$

Proof of (3.7). Setting $e = iq^{-n}$ in (1.3), we get

$$\begin{aligned}
 (3.10) \quad & \sum_{r=0}^n \frac{[a; q]_r (1 - aq^{2r}) [c; q]_r a^{2r} q^{2r^2}}{[q; q]_r (1 - a) \left[\frac{aq}{c}; q \right]_r [a^4q^4; q^4]_{n+r} [q^4; q^4]_{n-r} c^r} \\
 & = \frac{[aq; q]_{4n} [-a^2q^2c^{-2}; q^2]_{2n} (-)^n q^{-2n^2}}{[a^4q^4; q^4]_{2n} [a^4q^4c^{-4}; q^4]_n [q^4; q^4]_n} \\
 & \quad \times {}_4\phi_3 \left[\begin{matrix} -ca^{-2}q^{-4n-1}, -ca^{-2}q^{-4n}, -q^{-2n}, q^{-2n}; q^2; q^2 \end{matrix} \right].
 \end{aligned}$$

Now, in Bailey's transformation [14] choosing

$$u_s = \frac{1}{[q^4; q^4]_s}, \quad v_s = \frac{1}{[a^4q^4; q^4]_s}, \quad \alpha_s = \frac{[a; q]_s (1 - aq^{2s}) [c; q]_s a^{2s} q^{2s^2}}{[q; q]_s (1 - a) \left[\frac{aq}{c}; q \right]_s c^s}$$

and

$$\delta_s = \frac{[x; q^4]_s [y; q^4]_s a^{4s} q^{4s(1-p)}}{x^s y^s}$$

and evaluating $\langle \beta_n \rangle$, $\langle \gamma_n \rangle$ by using (3.10) and following formula [15]

$$(3.11) \quad {}_2\phi_1 \left[\begin{matrix} a, b; q; \frac{ec}{ab} \\ e \end{matrix} \right] = \Pi \left[\begin{matrix} \frac{e}{a}, \frac{e}{b}; q \\ e, \frac{e}{ab} \end{matrix} \right] {}_3\phi_2 \left[\begin{matrix} a, b, c; q; q \\ \frac{abq}{e}, 0 \end{matrix} \right],$$

(where, either a, b , or c is of the form q^{-p} , p a nonnegative integer. In case only c is of the form q^{-p} then (3.8) is valid only if $|ec/ab| < 1$), we get (3.7) on letting $x, y \rightarrow \infty$.

Proof of (3.8). In (3.10), letting $c \rightarrow \infty$, we have

$$\begin{aligned}
 & \frac{[aq; q]_{4n} (-)^n q^{-2n^2}}{[a^4q^4; q^4]_{2n} [q^4; q^4]_n} \sum_{r=0}^n \frac{[q^{-4n}; q^4]_r q^{r(r-4n)}}{[q^2; q^2]_r [a^{-1}q^{-4n}; q]_{2r} a^{2r}} \\
 & = \sum_{r=0}^n \frac{[a; q]_r (1 - aq^{2r}) (-)^r a^{2r} q^{(1/2)r(5r-1)}}{[q; q]_r (1 - a) [a^4q^4; q^4]_{n+r} [q^4; q^4]_{n-r}} \\
 & = \sum_{r=0}^n \frac{[aq; q]_r (-)^r a^{2r} q^{(1/2)r(5r+1)}}{[q; q]_r [a^4q^4; q^4]_{n+r} [q^4; q^4]_{n-r}} \\
 & \quad + \sum_{r=1}^n \frac{[aq; q]_{r-1} (1 - q^r) (-)^r a^{2r} q^{(1/2)r(5r-1)}}{[q; q]_r [a^4q^4; q^4]_{n+r} [q^4; q^4]_{n-r}}
 \end{aligned}$$

$$= \sum_{r=0}^n \frac{[aq; q]_r (-)^r a^{2r} q^{(1/2)r(5r+1)} \{(1 - a^4 q^{4n+4r+4}) - a^2 q^{4r+2} (1 - q^{4n-4r})\}}{[q; q]_r [a^4 q^4; q^4]_{n+r+1} [q^4; q^4]_{n-r}}$$

or,

$$(3.12) \quad \sum_{r=0}^n \frac{[aq; q]_r (1 - a^2 q^{4r+2}) (-)^r a^{2r} q^{(1/2)r(5r+1)}}{[q; q]_r [a^4 q^8; q^4]_{n+r} [q^4; q^4]_{n-r}} \\ = \frac{(1 - a^4 q^4) (-)^n q^{-2n^2}}{(1 + a^2 q^{4n+2}) [a^4 q^4; q^4]_{2n}} \sum_{r=0}^n \frac{[aq; q]_{4n-2r} (-)^r q^{r(r-1)}}{[q^2; q^2]_r [q^4; q^4]_{n-r}}.$$

Next, in Bailey's transformation [14] choosing

$$u_s = \frac{1}{[q^4; q^4]_s}, \quad v_s = \frac{1}{[a^4 q^8; q^4]_s}, \quad \alpha_s = \frac{[aq; q]_s (1 - a^2 q^{4s+2}) (-)^s q^{\frac{1}{2}s(5s+1)} a^{2s}}{[q; q]_s}$$

$\delta_s = [x; q^4]_s [y; q^4]_s a^{4s} q^{8s} / x^s y^s$ and evaluating $\langle \beta_n \rangle$, $\langle \gamma_n \rangle$ by using (3.12) and the q -analogue of Gauss' summation theorem [14; 3.3.2.5], we get (3.8) on letting $x, y \rightarrow \infty$.

Proof of (3.9). In (3.10), setting $c \rightarrow 0$, we get

$$(3.13) \quad \frac{[aq; q]_{4n}}{[q^4; q^4]_n [a^4 q^4; q^4]_{2n}} \sum_{r=0}^n \frac{[q^{-4n}; q^4]_r q^{2r}}{[q^2; q^2]_r [a^{-1} q^{-4n}; q]_{2r}} \\ = \sum_{r=0}^n \frac{[a; q]_r (1 - a q^{2r}) (-)^r a^r q^{(1/2)r(3r-1)}}{[q; q]_r (1 - a) [a^4 q^4; q^4]_{n+r} [q^4; q^4]_{n-r}}$$

(3.13) may be rewritten in the following form (its proof follows on the lines of the proof of (3.12))

$$(3.14) \quad \sum_{r=0}^n \frac{[aq; q]_r (1 - a^2 q^{4r+2}) (-a)^r q^{(1/2)r(3r-1)}}{[q; q]_r [a^4 q^8; q^4]_{n+r} [q^4; q^4]_{n-r}} \\ = \frac{(1 - a^4 q^4)}{(1 + a^2 q^{4n+2}) [a^4 q^4; q^4]_{2n}} \sum_{r=0}^n \frac{[aq; q]_{4n-2r} (-)^r a^{2r} q^{r(4n+1)}}{[q^2; q^2]_r [q^4; q^4]_{n-r}}.$$

However, in Bailey's transformation, choosing

$$u_s = \frac{1}{[q^4; q^4]_s}, \quad v_s = \frac{1}{[a^4 q^8; q^4]_s}, \quad \alpha_s = \frac{[aq; q]_s (1 - a^2 q^{4s+2}) (-a)^s q^{(1/2)s(3s-1)}}{[q; q]_s}, \\ \delta_s = \frac{[x; q^4]_s [y; q^4]_s a^{4s} q^{8s}}{x^s y^s}$$

and evaluating $\langle \beta_n \rangle$, $\langle \gamma_n \rangle$ by using (3.14) and the q -analogue of Gauss' summation theorem [14; 3.3.2.5], we get (3.9) on letting $x, y \rightarrow \infty$.

Identities of Rogers-Ramanujan type related to the modulus 13.
(3.7) for $c \rightarrow \infty$, $a = 1$ and $p = 0$ yields

$$\begin{aligned}
 (3.15) \quad & \frac{[q^4; q^4]_\infty}{[q; q]_\infty} \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[q; q]_{4n+2r} (-)^n q^{2n^2+3r^2+4nr-r}}{[q^2; q^2]_r [q^4; q^4]_n [q^4; q^4]_{2n+2r}} \\
 &= \prod_{n \neq 0, 6, 7 \pmod{13}} (1 - q^n)^{-1}.
 \end{aligned}$$

But, (3.7) for $c \rightarrow \infty$, $a = 1$ and $p = 1$, gives

$$\begin{aligned}
 (3.16) \quad & \frac{[q^4; q^4]_\infty}{[q; q]_\infty} \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[q; q]_{4n+2r} (-)^n q^{2n^2+3r^2+4nr-4n-5r}}{[q^2; q^2]_r [q^4; q^4]_n [q^4; q^4]_{2n+2r}} \\
 &= \prod_{n \neq 0, 2, 11 \pmod{13}} (1 - q^n)^{-1} + \prod_{n \neq 0, 3, 10 \pmod{13}} (1 - q^n)^{-1}.
 \end{aligned}$$

On the other hand, (3.7) for $c \rightarrow \infty$, $a = q$ and $p = 0$, reduces to

$$\begin{aligned}
 (3.17) \quad & \frac{[q^4; q^4]_\infty}{[q; q]_\infty} \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[q; q]_{4n+2r+1} (-)^n q^{2n^2+3r^2+4nr+4n+3r}}{[q^2; q^2]_r [q^4; q^4]_n [q^4; q^4]_{2n+2r+1}} \\
 &= \prod_{n \neq 0, 1, 12 \pmod{13}} (1 - q^n)^{-1}.
 \end{aligned}$$

Next, on setting $a = 1$, (3.8) yields

$$\begin{aligned}
 (3.18) \quad & \frac{[q^4; q^4]_\infty}{[q; q]_\infty} \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[q; q]_{4n+2r} (-)^n q^{2n^2+3r^2+4nr+4n+3r}}{[q^2; q^2]_r [q^4; q^4]_n [q^4; q^4]_{2n+2r} (1 + q^{4n+4r+2})} \\
 &= \prod_{n \neq 0, 2, 11 \pmod{13}} (1 - q^n)^{-1}
 \end{aligned}$$

Whereas, in (3.8) setting $a = q^{-1}$ and using (3.15), we get

$$\begin{aligned}
 (3.19) \quad & \frac{[q^4; q^4]_\infty}{[q; q]_\infty} \left\{ 1 + \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[q; q]_{4n+2r+1} (-)^n q^{2n^2+3r^2+4nr+8n+7r+4}}{[q^2; q^2]_r [q^4; q^4]_n [q^4; q^4]_{2n+2r+2}} \right\} \\
 &= \prod_{n \neq 0, 5, 8 \pmod{13}} (1 - q^n)^{-1}.
 \end{aligned}$$

Lastly, in (3.8) setting $a = q$ and using (3.19), we have

$$\begin{aligned}
 (3.20) \quad & \frac{[q^4; q^4]_\infty}{[q; q]_\infty} \left\{ 1 + \sum_{r=0}^{\infty} \sum_{n=0}^{\infty} \frac{[q; q]_{4n+2r+1} (-)^n q^{2n^2+3r^2+4nr+12n+11r+8}}{[q^2; q^2]_r [q^4; q^4]_n [q^4; q^4]_{2n+2r+2}} \right\} \\
 &= \prod_{n \neq 0, 4, 9 \pmod{13}} (1 - q^n)^{-1}.
 \end{aligned}$$

Similarly the five identities of Rogers-Ramanujan type related to the modulus 11 due to Andrews [2] may be obtain from (3.7) and (3.9).

In view of the above applications of (1.1), it may be of interest to record a generalization of (1.1). In fact we prove that if a, b, e, f is of the form q^{-N} , then

$$\begin{aligned}
 (3.21) \quad & {}_4\phi_3 \left[\begin{matrix} a^2, b^2, e^2, f^2; q^2; \frac{c^2 q}{a^2 b^2} \end{matrix} \right] = \sum_{n \geq 0} \frac{[a^2; q^2]_n [b^2; q^2]_n \left[\frac{a^2 b^2 q}{c^2}; q^2 \right]_n}{[q^2; q^2]_n [c^2; q^2]_{2n}} \\
 & \times \frac{[e; q]_{2n} [f; q]_{2n} c^{4n} q^{n(1-2n)}}{[-ef; q]_{2n} (ab)^{4n}} {}_4\phi_3 \left[\begin{matrix} a^2 q^{2n}, b^2 q^{2n}, eq^{2n}, fq^{2n}; q; \frac{c^2 q^{-2n}}{a^2 b^2} \end{matrix} \right].
 \end{aligned}$$

(3.21) reduces to (1.1) for $c = ab\sqrt{q}$.

We complete the proof of (3.21) by evaluating

$$(3.22) \quad S = \sum_{r \geq 0} \frac{[a^2; q^2]_r [b^2; q^2]_r [e; q]_r [f; q]_r c^{2r} q^{-(1/2)r(r-1)}}{[q; q]_r [c^2; q^2]_r [-ef; q]_r (ab)^{2r}} \\ \times {}_3\phi_2 \left[\begin{matrix} q^{-r}, q^{1-r}, \frac{q^{2-2r}}{c^2}; q^2; q^2 \end{matrix} \right]_{a^{-2}q^{2-2r}, b^{-2}q^{2-2r}},$$

in two different ways. Firstly, if we substitute the series definition of ${}_3\phi_2$, change the order of summations and then diagonalize the two series, we get

$$S = \sum_{r \geq 0} \frac{[a^2; q^2]_r [b^2; q^2]_r [e; q]_r [f; q]_r c^{2r} q^{-(1/2)r(r-1)}}{[q; q]_r [c^2; q^2]_r [-ef; q]_r (ab)^{2r}} {}_3\phi_2 \left[\begin{matrix} q^{-r}, eq^r, fq^r; q; q \end{matrix} \right]_{-q, -efq^r}.$$

Summing the inner ${}_3\phi_2$ by the q -analogue of Saalschütz summation theorem [14; 3.3.2.2], we get the left hand side of (3.21).

Secondly, we may rewrite (3.22) as

$$(3.23) \quad S = \sum_{r \geq 0} \frac{[a^2; q^2]_{2r} [b^2; q^2]_{2r} [e; q]_{2r} [f; q]_{2r} c^{4r} q^{r(1-2r)}}{[q; q]_{2r} [c^2; q^2]_{2r} [-ef; q]_{2r} (ab)^{4r}} \\ \times {}_3\phi_2 \left[\begin{matrix} q^{-2r}, q^{1-2r}, \frac{q^{2-4r}}{c^2}; q^2; q^2 \end{matrix} \right]_{\frac{q^{2-4r}}{a^2}, \frac{q^{2-4r}}{b^2}} \\ + \sum_{r \geq 0} \frac{[a^2; q^2]_{2r+1} [b^2; q^2]_{2r+1} [e; q]_{2r+1} [f; q]_{2r+1} c^{4r+2} q^{-r(1+2r)}}{[q; q]_{2r+1} [c^2; q^2]_{2r+1} [-ef; q]_{2r+1} (ab)^{4r+2}} \\ \times {}_3\phi_2 \left[\begin{matrix} q^{-2r}, q^{-2r-1}, \frac{q^{-4r}}{c^2}; q^2; q^2 \end{matrix} \right]_{\frac{q^{-4r}}{a^2}, \frac{q^{-4r}}{b^2}}.$$

In the transformation

$$(3.24) \quad {}_3\phi_2 \left[\begin{matrix} b, c, q^{-N}; q; \frac{egq^N}{bc} \\ e, g \end{matrix} \right] = \frac{\left[\frac{g}{c}; q \right]_N}{[g; q]_N} {}_3\phi_2 \left[\begin{matrix} \frac{e}{b}, c, q^{-N}; q; q \\ e, \frac{c}{g} q^{1-N} \end{matrix} \right],$$

(which is obtained from (3.2) by substituting for h and then letting $a \rightarrow \infty$), transforming the ${}_3\phi_2$ on the left hand side by the same formula (3.24) (with e replaced by g), we get

$$(3.25) \quad {}_3\phi_2 \left[\begin{matrix} \frac{e}{b}, c, q^{-N}; q; q \\ e, \frac{c}{g} q^{1-N} \end{matrix} \right] = \frac{[g; q]_N \left[\frac{e}{c}; q \right]_N}{\left[\frac{g}{c}; q \right]_N [e; q]_N} {}_3\phi_2 \left[\begin{matrix} \frac{g}{b}, c, q^{-N}; q; q \\ g, \frac{c}{e} q^{1-N} \end{matrix} \right].$$

Now, using (3.25) for transforming the two ${}_3\phi_2$ series in (3.23) [to transform the first of the two ${}_3\phi_2$ in (3.23), we use (3.25) with $q \rightarrow q^2$, $N = r$, $e \rightarrow q^{2-4r}/a^2$, $e/b \rightarrow q^{2-4r}/c^2$, $c \rightarrow q^{1-2r}$, $c/g \rightarrow q^{-2r}/b^2$ and for transforming the second ${}_3\phi_2$ in (3.23), we use (3.25) with $q \rightarrow q^2$, $N = r$, $e \rightarrow q^{-4r}/a^2$, $e/b \rightarrow q^{-4r}/c^2$, $c \rightarrow q^{-2r-1}$, $c/g \rightarrow q^{-2r-2}/b^2$], we get

$$\begin{aligned} S &= \sum_{r \geq 0} \frac{[a^2; q]_{2r}[b^2; q]_{2r}[e; q]_{2r}[f; q]_{2r}c^{4r}}{[q; q]_{2r}[c^2; q^2]_{2r}[-ef; q]_{2r}(ab)^{4r}} {}_3\phi_2 \left[\begin{matrix} \frac{a^2b^2q}{c^2}, q^{1-2r}, q^{-2r}; q^2; q^2 \end{matrix} \right] \\ &\quad + \sum_{r \geq 0} \frac{[a^2; q]_{2r+1}[b^2; q]_{2r+1}[e; q]_{2r+1}[f; q]_{2r+1}c^{4r+2}}{[q; q]_{2r+1}[c^2; q^2]_{2r+1}[-ef; q]_{2r+1}(ab)^{4r+2}} \\ &\quad \times {}_3\phi_2 \left[\begin{matrix} \frac{a^2b^2q}{c^2}, q^{-2r-1}, q^{-2r}; q^2; q^2 \end{matrix} \right] \\ &= \sum_{r \geq 0} \frac{[a^2; q]_r[b^2; q]_r[e; q]_r[f; q]_rc^{2r}}{[q; q]_r[c^2; q^2]_r[-ef; q]_r(ab)^{2r}} {}_3\phi_2 \left[\begin{matrix} \frac{a^2b^2q}{c^2}, q^{1-r}, q^{-r}; q^2; q^2 \end{matrix} \right]. \end{aligned}$$

Writing the series definition for inner ${}_3\phi_2$ and then interchanging the order of summations of the two series, we get the right hand side of (3.21).

If a or b is of the form q^{-N} , $e = x$, $f = o$, then (3.21) yields

$$\begin{aligned} {}_3\phi_2 \left[\begin{matrix} a^2, b^2, x^2; q^2; \frac{c^2q}{a^2b^2} \end{matrix} \right] &= \sum_{n \geq 0} \frac{[a^2; q^2]_n[b^2; q^2]_n \left[\frac{a^2b^2q}{c^2}; q^2 \right]_n [x; q]_{2n}c^{4n}}{[q^2; q^2]_n[c^2; q^2]_{2n}(ab)^{4n}q^{n(2n-1)}} \\ &\quad \times {}_3\phi_2 \left[\begin{matrix} a^2q^{2n}, b^2q^{2n}, xq^{2n}; q; \frac{c^2q^{-2n}}{a^2b^2} \end{matrix} \right]. \end{aligned}$$

In which replacing a, b, c , by q^{-N}, q^b, q^c respectively and letting $q \rightarrow 1$, we only get a terminating version of the following formula of Burchnell and Chaundy [9; 5.7] (with x replaced by $1 - 2x$):

$$\begin{aligned} (3.26) \quad {}_2F_1 \left[\begin{matrix} a, b; 4x(1-x) \\ c \end{matrix} \right] &= \sum_{n=0}^{\infty} \frac{(a)_n(b)_n \left(a + b - c + \frac{1}{2} \right)_n 4^n x^{2n}}{(1)_n(c)_{2n}} \\ &\quad \times {}_2F_1 \left[\begin{matrix} 2a + 2n, 2b + 2n; x \\ c + 2n \end{matrix} \right]. \end{aligned}$$

On the other hand to obtain the non-terminating version of (3.26), we start (3.21) by replacing e by $-e$, $f = q^{-N}$ and then

replace a, b, c, e by q^a, q^b, q^c, q^e respectively and let $q \rightarrow 1$ to obtain

$$\begin{aligned}
 & {}_4F_3 \left[\begin{matrix} a, b, e, -N; \\ c, \frac{1}{2}(e-N), \frac{1}{2}(e-N+1) \end{matrix} \right] \\
 (3.27) \quad &= \sum_{n=0}^{[N/2]} \frac{(a)_n (b)_n \left(a + b - c + \frac{1}{2} \right)_n (-N)_{2n} 4n}{(1)_n (c)_{2n} (e-N)_{2n}} \\
 &\quad \times {}_3F_2 \left[\begin{matrix} 2a + 2n, 2b + 2n, -N + 2n; \\ c + 2n, e - N + 2n \end{matrix} \right].
 \end{aligned}$$

In (3.27) on replacing e by $N(1 - 1/x)$ and letting $N \rightarrow \infty$, we get the non-terminating version of (3.26).

4. We begin this section by proving a q -analogue of the transformation due to Bailey [5; 2.5] in the form:

$$\begin{aligned}
 & {}_4\phi_3 \left[\begin{matrix} \frac{a^2}{q}, a\sqrt{q}, -a\sqrt{q}, \frac{a}{b}\sqrt{q}; q; b^2z \end{matrix} \right] \\
 (4.1) \quad & \left[\begin{matrix} \frac{a}{\sqrt{q}}, -\frac{a}{\sqrt{q}}, ab\sqrt{q} \end{matrix} \right] \\
 &= \frac{[a^2z; q]_\infty}{[b^2z; q]_\infty} {}_4\phi_3 \left[\begin{matrix} \frac{b^2}{q}, b\sqrt{q}, -b\sqrt{q}, \frac{b}{a\sqrt{q}}; q; a^2z \end{matrix} \right] \\
 & \left[\begin{matrix} \frac{b}{\sqrt{q}}, -\frac{b}{\sqrt{q}}, ab\sqrt{q} \end{matrix} \right]
 \end{aligned}$$

provided $|a^2z| < 1$, $|b^2z| < 1$.

Proof of (4.1). Using the q -analogue of a nearly-poised summation theorem due to Bailey [8; (3)] in the form

$$\begin{aligned}
 & {}_5\phi_4 \left[\begin{matrix} \frac{b^2}{q}, b\sqrt{q}, -b\sqrt{q}, \frac{b}{a\sqrt{q}}, q^{-n}; q; q \end{matrix} \right] \\
 & \left[\begin{matrix} \frac{b}{\sqrt{q}}, -\frac{b}{\sqrt{q}}, ab\sqrt{q}, \frac{b^2}{a^2}q^{1-n} \end{matrix} \right] \\
 &= \frac{[a^2q^{-1}; q]_n [a^2q; q^2]_n \left[\frac{a}{b\sqrt{q}}; q \right]_n}{[a^2q^{-1}; q^2]_n [ab\sqrt{q}; q]_n [a^{+2}b^{-2}; q]_n},
 \end{aligned}$$

the left hand side of (4.1) may be rewritten as:

$$\sum_{n=0}^{\infty} \frac{[a^2b^{-2}; q]_n b^{2n} z^n}{[q; q]_n} {}_5\phi_4 \left[\begin{matrix} \frac{b^2}{q}, b\sqrt{q}, -b\sqrt{q}, \frac{b}{a\sqrt{q}}, q^{-n}; q; q \end{matrix} \right]$$

$$= \sum_{r=0}^{\infty} \frac{[b^2 q^{-1}; q]_r [b^2 q; q^2]_r \left[\frac{b}{a\sqrt{q}}; q \right]_r a^{2r} z^r}{[q; q]_r [b^2 q^{-1}; q^2]_r [ab\sqrt{q}; q]_r} {}_1\phi_0 \left[\begin{matrix} \frac{a^2}{b^2}; q; b^2 z \\ - \end{matrix} \right].$$

Summing the ${}_1\phi_0$, we get the right hand side of (4.1).

Augmenting parameters on both sides of ϕ -series (4.1) by using q -beta transform [10], we get

$$(4.2) \quad \begin{aligned} & {}_6\phi_5 \left[\begin{matrix} \frac{a^2}{q}, a\sqrt{q}, -a\sqrt{q}, \frac{a}{b\sqrt{q}}, c, d; q; b^2 z \\ \frac{a}{\sqrt{q}}, -\frac{a}{\sqrt{q}}, ab\sqrt{q}, e, f \end{matrix} \right] \\ &= \sum_{n=0}^{\infty} \frac{[b^2 q^{-1}; q]_n [b^2 q; q^2]_n}{[q; q]_n [b^2 q^{-1}; q^2]_n} \\ & \quad \times \frac{\left[\frac{b}{a\sqrt{q}}; q \right]_n [c; q]_n [d; q]_n a^{2n} z^n}{[ab\sqrt{q}; q]_n [e; q]_n [f; q]_n} {}_3\phi_2 \left[\begin{matrix} \frac{a^2}{b^2}, cq^n, dq^n; q; b^2 z \\ eq^n, fq^n \end{matrix} \right]. \end{aligned}$$

In (4.2) setting $d = q^{-N}$, $z = qb^{-2}$, $f = a^2 cb^{-2} e^{-1} q^{1-N}$ and summing the inner ${}_3\phi_2$ on the right hand side by the q -analogue of Saalschütz summation theorem, we get

$$(4.3) \quad \begin{aligned} & {}_6\phi_5 \left[\begin{matrix} \frac{a^2}{q}, a\sqrt{q}, -a\sqrt{q}, \frac{a}{b\sqrt{q}}, c, q^{-N}; q; q \\ \frac{a}{\sqrt{q}}, -\frac{a}{\sqrt{q}}, ab\sqrt{q}, e, \frac{a^2 cq^{1-N}}{eb^2} \end{matrix} \right] = \frac{[ec^{-1}; q]_N [eb^2 a^{-2}; q]_N}{[e; q]_N [eb^2 c^{-1} a^{-2}; q]_N} \\ & \quad \times {}_6\phi_5 \left[\begin{matrix} \frac{b^2}{q}, b\sqrt{q}, -b\sqrt{q}, \frac{b}{a\sqrt{q}}, c, q^{-N}; q; q \\ \frac{b}{\sqrt{q}}, -\frac{b}{\sqrt{q}}, ab\sqrt{q}, \frac{eb^2}{a^2}, \frac{c}{e} q^{1-N} \end{matrix} \right]. \end{aligned}$$

(4.3) for $N \rightarrow \infty$ yields the q -analogue of a non-terminating version of a transformation due to Bailey [5; 2.51] in the form (with e replaced by $a^2 e$):

$$(4.4) \quad \begin{aligned} & \frac{[a^2 e; q]_{\infty}}{[a^2 ec^{-1}; q]_{\infty}} {}_6\phi_4 \left[\begin{matrix} \frac{a^2}{q}, a\sqrt{q}, -a\sqrt{q}, \frac{a}{b\sqrt{q}}, c; q; \frac{b^2 e}{c} \\ \frac{a}{\sqrt{q}}, -\frac{a}{\sqrt{q}}, ab\sqrt{q}, a^2 e \end{matrix} \right] \\ &= \frac{[b^2 e; q]_{\infty}}{[b^2 ec^{-1}; q]_{\infty}} {}_6\phi_4 \left[\begin{matrix} \frac{b^2}{q}, b\sqrt{q}, -b\sqrt{q}, \frac{b}{a\sqrt{q}}, c; q; \frac{a^2 e}{c} \\ \frac{b}{\sqrt{q}}, -\frac{b}{\sqrt{q}}, ab\sqrt{q}, b^2 e \end{matrix} \right]. \end{aligned}$$

On the other hand (4.3), for $b = -1$, reduces to the summation theorem:

$$(4.5) \quad {}_4\phi_3 \left[\begin{matrix} \frac{a^2}{q}, a\sqrt{q}, c, q^{-N}; q; q \\ \frac{a}{\sqrt{q}}, e, \frac{a^2 c}{e} q^{1-N} \end{matrix} \right] = \frac{\left[\frac{e}{c}; q \right]_N [ea^{-2}; q]_N}{[e; q]_N [ec^{-1}a^{-2}; q]_N} \left\{ 1 + \frac{(1-c)(1-q^{-N})a\sqrt{q}}{(a^2 - e)\left(1 - \frac{c}{e}q^{1-N}\right)} \right\}.$$

It may be worthwhile to remark that (4.5) could have been obtained directly by transforming the Saalschützian ${}_4\phi_3$ in (4.5) by using (3.2) with $a \rightarrow a^2q^{-1}$, $b \rightarrow a\sqrt{q}$, $e \rightarrow a/\sqrt{q}$, $g \rightarrow e$ and $h \rightarrow (c/e)a^2q^{1-N}$.

Now, if we specialize $e = a^2/c$ in (4.5), we get (1.5) (on replacing a by $a\sqrt{q}$).

Next, using the summation theorem (1.5), we can prove the q -analogue of a transformation of Bailey [7; 4.5 (4)] in the form if $k = a^2q/bcd$ then

$$(4.6) \quad {}_6\phi_5 \left[\begin{matrix} a, q\sqrt{a}, b, c, d, q^{-N}; q; q \\ \sqrt{a}, \frac{aq}{b}, \frac{aq}{c}, \frac{aq}{d}, \frac{a^2}{k^2} q^{1-N} \end{matrix} \right] = \frac{\left[\frac{k}{a}; q \right]_N \left[\frac{k^2}{a}; q \right]_N \left[-\frac{kq}{\sqrt{a}}; q \right]_N}{[kq; q]_N [k^2a^{-2}; q]_N \left[-\frac{k}{\sqrt{a}}; q \right]_N} \\ \times {}_{12}\phi_{11} \left[\begin{matrix} k, q\sqrt{k}, -q\sqrt{k}, \frac{kb}{a}, \frac{kc}{a}, \frac{kd}{a}, q\sqrt{a}, -\sqrt{a}, \sqrt{aq}, \\ -\sqrt{aq}, \frac{k^2}{a} q^N, q^{-N}; q; q \\ \sqrt{k}, -\sqrt{k}, \frac{aq}{b}, \frac{aq}{c}, \frac{aq}{d}, \frac{k}{\sqrt{a}}, -\frac{kq}{\sqrt{a}}, k\sqrt{\frac{q}{a}}, \\ -k\sqrt{\frac{q}{a}}, \frac{a}{k} q^{1-N}, kq^{1+N} \end{matrix} \right].$$

Further replacing " \sqrt{a} " by " $-\sqrt{a}$ " in (4.6), we get the q -analogue of another result of Bailey [7; 4.5(5)].

Proof of (4.6). Using the q -analogue of Dougall's theorem [14; 3.3.1.1] in the form

$${}_8\phi_7 \left[\begin{matrix} k, q\sqrt{k}, -q\sqrt{k}, \frac{kb}{a}, \frac{kc}{a}, \frac{kd}{a}, aq^n, q^{-n}; q; q \\ \sqrt{k}, -\sqrt{k}, \frac{aq}{b}, \frac{aq}{c}, \frac{aq}{d}, \frac{k}{a} q^{1-n}, kq^{1+n} \end{matrix} \right]$$

$$= \frac{[kq; q]_n [b; q]_n [c; q]_n [d; q]_n}{\left[\frac{a}{k}; q \right]_n \left[\frac{aq}{b}; q \right]_n \left[\frac{aq}{c}; q \right]_n \left[\frac{aq}{d}; q \right]_n},$$

we may rewrite the left hand side of (4.6) (denoted by S) in the form:

$$\begin{aligned} S &= \sum_{n=0}^N \frac{[a; q]_n [q\sqrt{a}; q]_n \left[\frac{a}{k}; q \right]_n [q^{-N}; q]_n q^n}{[q; q]_n [\sqrt{a}; q]_n [a^2 k^{-2} q^{1-N}; q]_n [kq; q]_n} \\ &\quad \times {}_8\phi_7 \left[\begin{matrix} k, q\sqrt{k}, -q\sqrt{k}, \frac{kb}{a}, \frac{kc}{a}, \frac{kd}{a} aq^n, q^{-n}; q; q \\ \sqrt{k}, -\sqrt{k}, \frac{aq}{b}, \frac{aq}{c}, \frac{aq}{d}, \frac{k}{a}, q^{1-n}, kq^{1+n} \end{matrix} \right] \\ &= \sum_{r=0}^N \frac{[k; q]_r [kq^2; q^2]_r \left[\frac{kb}{a}; q \right]_r \left[\frac{kc}{a}; q \right]_r \left[\frac{kd}{a}; q \right]_r [a; q]_{2r} [q\sqrt{a}; q]_r}{[q; q]_r [k; q^2]_r \left[\frac{aq}{b}; q \right]_r \left[\frac{aq}{c}; q \right]_r \left[\frac{aq}{d}; q \right]_r [kq; q]_{2r} [\sqrt{a}; q]_r} \\ &\quad \times \frac{[q^{-N}q]_r a^r q^r}{\left[\frac{a^2}{k^2} q^{1-N}; q \right]_r k^r} \cdot {}_4\phi_3 \left[\begin{matrix} aq^{2r}, \sqrt{a} q^{1+r}, \frac{a}{k}, q^{-N+r}; q; q \\ \sqrt{a} q^r, kq^{1+2r}, a^2 k^{-2} q^{1-N+r} \end{matrix} \right], \end{aligned}$$

summing the inner ${}_4\phi_3$ by (1.5), we get the desired result.

Lastly, we prove the formula (1.9).

Proof of (1.9). In view of the q -analogue of Dougall's theorem in the form:

$$\begin{aligned} &{}_8\phi_7 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, c, dq^n, eq^{-n}, kq^n, q^{-n}; q; q \\ \sqrt{a}, -\sqrt{a}, \frac{aq}{c}, \frac{a}{d} q^{1-n}, \frac{a}{e} q^{1+n}, \frac{a}{k} q^{1-n}, aq^{1+n} \end{matrix} \right] \\ &= \frac{[aq; q]_n \left[\frac{aq}{ce}; q \right]_{2n} \left[\frac{aq}{de}; q \right]_n \left[\frac{aq}{e}; q \right]_n \left[\frac{cd}{a}; q \right]_n}{\left[\frac{aq}{c}; q \right]_n \left[\frac{aq}{e}; q \right]_{2n} \left[\frac{aq}{cde}; q \right]_n \left[\frac{aq}{ce}; q \right]_n \left[\frac{d}{a}; q \right]_n c^n} \end{aligned}$$

(where $k = a^2 q(cde)^{-1}$), we have

$$\sum_{n=0}^N \frac{\left[\frac{cd}{a}; q \right]_n \left[\frac{aq}{de}; q \right]_n \left[\frac{aq}{ce}; q \right]_{2n} [d; q]_n [k; q]_n [q^{-N}; q]_n q^n}{[q; q]_n \left[\frac{q}{e}; q \right]_n \left[\frac{aq}{c}; q \right]_n \left[\frac{aq}{ce}; q \right]_n [d^2 a^{-2} q^{-N}; q]_n \left[\frac{aq}{e}; q \right]_{2n} c^n}$$

$$\begin{aligned}
&= \sum_{n=0}^N \frac{[d; q]_n [k; q]_n \left[\frac{aq}{cde}; q \right]_n \left[\frac{d}{a}; q \right]_n [q^{-N}; q]_n q^n}{[q; q]_n \left[\frac{q}{e}; q \right]_n [aq; q]_n \left[\frac{aq}{e}; q \right]_n [d^2 a^{-2} q^{-N}; q]_n} \\
&\quad \times {}_8\phi_7 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, c, dq^n, eq^{-n}, kq^n, q^{-n}; q; q \\ \sqrt{a}, -\sqrt{a}, \frac{aq}{c}, \frac{a}{d} q^{1-n}, \frac{a}{e} q^{1+n}, \frac{a}{k} q^{1-n}, aq^{1+n} \end{matrix} \right] \\
&= \sum_{r=0}^N \frac{[a; q]_r [aq^2; q^2]_r [c; q]_r [d; q]_{2r} [k; q]_{2r} [q^{-N}; q]_r q^r}{[q; q]_r [a; q^2]_r \left[\frac{aq}{c}; q \right]_r \left[\frac{aq}{e}; q \right]_{2r} [d^2 a^{-2} q^{-N}; q]_{2r} [aq; q]_{2r} c^r} \\
(4.7) \quad &\quad \times {}_5\phi_4 \left[\begin{matrix} dq^{2r}, kq^{2r}, \frac{d}{a}, \frac{k}{a}, q^{-N+r}; q; q \\ \frac{a}{e} q^{1+2r}, aq^{1+2r}, \frac{q}{e}, d^2 a^{-2} q^{-N+r} \end{matrix} \right].
\end{aligned}$$

In (4.7) taking $c = a/d$, $k = aq/e$ and then summing the inner ${}_8\phi_7$ on the right hand side by the q -analogue of Saalschütz summation theorem [14; 3.3.2.2], we get (1.9).

I am grateful to Dr. Arun Verma for suggesting the problem and for his helpful discussions during the preparation of this paper.

REFERENCES

1. R. P. Agarwal, *Generalized Hypergeometric Series*, (1963) (Monograph, Council of Science and Industrial Research U. P.) Asia Publishing Co.
2. G. E. Andrews, *On Rogers-Ramanujan type identities related to the modulus 11*, Proc. London Math. Soc., (3) **30** (1975), 330-346.
3. ———, *On q -analogues of the Watson and Whipple summations*, SIAM J. Math. Anal., **7** (1976), 332-336.
4. R. Askey and J. Wilson, (Private communication).
5. W. N. Bailey, *Transformations of generalized hypergeometric series*, Proc. London Math. Soc., (2) **29** (1929), 495-502.
6. ———, *Some identities involving generalized hypergeometric series*, Proc. London Math. Soc., (2) **29** (1929), 503-516.
7. ———, *Generalized Hypergeometric Series*, (Cambridge Tract 1935).
8. ———, *A transformation of nearly-poised basic hypergeometric series*, J. London Math. Soc., **22** (1947), 237-240.
9. J. L. Burchall and T. W. Chaundy, *The hypergeometric identities of Cayley, Orr and Bailey*, Proc. London Math. Soc., (2) **50** (1949), 56-74.
10. F. H. Jackson, *q -difference equations*, Amer. J. Math., **32** (1910), 305-314.
11. V. K. Jain, *Some transformations of basic hypergeometric functions*, SIAM J. Math. Anal., **12** (6), (1981).
12. D. B. Sears, *On the transformation theory of basic hypergeometric functions*, Proc. London Math. Soc., (2) **53** (1951), 158-180.
13. V. N. Singh, *Basic analogue of identities of the Cayley-Orr type*, J. London Math. Soc., **34** (1959), 15-22.
14. L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, 1966.

15. A. Verma, *Certain summation formulae for basic hypergeometric series*, Canad. Math. Bull., **20** (1977), 369-375.

Received August 6, 1980.

BAREILLY COLLEGE
BAREILLY (U. P.) INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR AGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90007

R. FINN and J. MILGRAM

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF AAWAH

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies,

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

50 reprints to each author are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966, Regular subscription rate: \$114.00 a year (6 Vol., 12 issues). Special rate: \$57.00 a year to individual members of supporting institution.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.).
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1982 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Jean Bourgain , A Hausdorff-Young inequality for B -convex Banach spaces	255
J. L. Brenner and Lorraine L. Foster , Exponential Diophantine equations	263
Henry H. Glover and William Duncan Homer, II , Fixed points on flag manifolds	303
Lothar Hahn , A note on stochastic methods in connection with approximation theorems for positive linear operators	307
James P. Henderson , Approximating cellular maps between low-dimensional polyhedra	321
V. K. Jain , Certain transformations of basic hypergeometric series and their applications	333
Charles David Keys , On the decomposition of reducible principal series representations of p -adic Chevalley groups	351
M. S. Klamkin and A. Meir , Ptolemy's inequality, chordal metric, multiplicative metric	389
Robert F. Lax , Independence of normal Weierstrass points under deformation	393
Leonid A. Luxemburg , On compactifications of metric spaces with transfinite dimensions	399
Carlton James Maxson, Martin Ross Pettet and Kirby C. Smith , On semisimple rings that are centralizer near-rings	451
Teodor C. Przymusiński , Extending functions from products with a metric factor and absolutes	463
Giorgio Talenti , A note on the Gauss curvature of harmonic and minimal surfaces	477
D. M. Terlinden , A spectral containment theorem analogous to the semigroup theory result $e^{t\sigma(A)} \subseteq \sigma(e^{tA})$	493