

Pacific Journal of Mathematics

**ALTMAN'S CONTRACTORS AND FIXED POINTS OF
MULTIVALUED MAPPINGS**

K. BALAKRISHNA REDDY AND P. V. SUBRAHMANYAM

ALTMAN'S CONTRACTORS AND FIXED POINTS OF MULTIVALUED MAPPINGS

K. BALAKRISHNA REDDY AND P. V. SUBRAHMANYAM

Let $P_i: D \subset X_1 \times \cdots \times X_n \rightarrow \text{CL}(Y_i)$ be multivalued mappings where X_i , Y_i are Banach spaces and $\text{CL}(Y_i)$ is the set of all nonempty closed subsets of Y_i , $i = 1, \dots, n$. We prove a theorem ensuring that $\theta_i \in P_i(x_1, \dots, x_n)$ for some $(x_1, \dots, x_n) \in D$ and deduce the fixed point theorems for multivalued mappings proved earlier by Czerwinski, Nadler and Reich as corollaries. Besides, generalizations for multivalued mappings of the existence theorems proved by Altman using his theory of contractors are also obtained.

1. Introduction. In [2] we showed how the fixed point theorems of Altman [1] and Matkowski ([5], [6]) can be unified in the set-up of Banach spaces. The present paper studies further the relationship between Altman's theory of contractors and Matkowski's fixed point theorem and offers an existence theorem for the multivalued operator equation $\theta \in Px$ on subsets of a Banach space. We deduce as a corollary a comprehensive fixed point theorem proved by Czerwinski [4] for multivalued mappings. Czerwinski's fixed point theorem generalized the earlier fixed point theorems for set valued transformations on metric spaces obtained by Nadler [7], Covitz and Nadler [3] and Reich [9]. Apart from Czerwinski's theorem, our main result obtains as corollaries generalizations to multivalued mappings of Altman's existence theorems and Matkowski's fixed point theorem. Section 3 gives the main result of the paper, while § 2 provides preliminaries basic to § 3.

2. Let X be a Banach space. We employ the following notation of [7] and [8]:

$$(2.1) \quad \text{CL}(X) = \{C: C \text{ is a nonempty closed subset of } X\}.$$

$$(2.2) \quad N(\varepsilon, C) = \{x \in X: \|x - c\| < \varepsilon \text{ for some } c \in C\}, \\ \varepsilon > 0, \quad C \in \text{CL}(X).$$

$$(2.3) \quad H(A, B) = \begin{cases} \inf \varepsilon > 0, & A \subset N(\varepsilon, B) \text{ and } B \subset N(\varepsilon, A), \\ & \text{if the infimum exists,} \\ \infty, & \text{otherwise,} \end{cases} \\ A, B \in \text{CL}(X).$$

The function H is called the generalized Hausdorff distance for

$\text{CL}(X)$ induced by the norm of X .

$$(2.4) \quad D(x, A) = \inf \{\|x - a\| : a \in A\}.$$

The lemma given below is well-known and is used in § 3.

LEMMA 2.1. Suppose $A, B \in \text{CL}(X)$ and $a \in A$. Then, for $q > 0$, there exists an element $b \in B$ such that

$$d(a, b) \leq H(A, B) + q.$$

A point $x \in X$ is said to be a fixed point for the multivalued mapping $f: X \rightarrow \text{CL}(X)$ if $x \in f(x)$.

We follow the notation of [6].

Let (a_{ik}) be an $n \times n$ nonnegative matrix. Define

$$(2.5) \quad a_{ik}^l = \begin{cases} a_{ik}, & i \neq k, \\ 1 - a_{ik}, & i = k, \end{cases} \quad i, k = 1, \dots, n.$$

$$(2.6) \quad a_{ik}^{l+1} = \begin{cases} a_{11}^l a_{i+1k+1}^l + a_{i+11}^l a_{1k+1}^l, & i \neq k \\ a_{11}^l a_{i+1k+1}^l - a_{i+11}^l a_{1k+1}^l, & i = k \end{cases} \quad l = 1, \dots, n-1; \quad i, k = 1, \dots, n-l$$

Matkowski ([5], [6]) proved the following

LEMMA 2.2. Let $a_{ik}^l > 0$, $i, k = 1, \dots, n$. The system of inequalities

$$(2.7) \quad \sum_{k=1}^n a_{ik} r_k < r_i, \quad i = 1, \dots, n,$$

has a solution $r_i > 0$, $i = 1, \dots, n$ if and only if the following inequalities hold:

$$(2.8) \quad a_{ii}^l > 0, \quad l = 1, \dots, n; \quad i = 1, \dots, n+1-l.$$

Using this lemma he obtained the following fixed point theorem. (Actually Matkowski proved this theorem in the setting of complete metric spaces.)

THEOREM 2.1. Let X_i be Banach spaces and $T_i: X_1 \times \dots \times X_n \rightarrow X_i$, $i = 1, \dots, n$ be mappings such that

$$(2.9) \quad \|T_i(x_1, \dots, x_n) - T_i(y_1, \dots, y_n)\| \leq \sum_{k=1}^n a_{ik} \|x_k - y_k\|, \\ i = 1, \dots, n, \quad x_k, y_k \in X_k, \quad k = 1, \dots, n,$$

where $a_{ik} > 0$, $i, k = 1, \dots, n$. If the numbers a_{ik}^l , $l = 1, \dots, n$;

$i, k = 1, \dots, n + 1 - l$ defined by (2.5) and (2.6) fulfill (2.8), then the system of equations

$$x_i = T_i(x_1, \dots, x_n), \quad i = 1, \dots, n,$$

has exactly one solution $x_i \in X_i$, $i = 1, \dots, n$. Moreover,

$$x_i = \lim_{m \rightarrow \infty} x_i^m, \quad i = 1, \dots, n,$$

where $x_i^{m+1} = T_i(x_1^m, \dots, x_n^m)$, $x_i^m \in X_i$, $i = 1, \dots, n$ is arbitrarily chosen.

Considerations of some of the fundamental problems of numerical analysis and operator theory led Altman [1] to the concept of contractors.

DEFINITION 2.1 [p. 8, [1]]. Let $P: D(P) \subset X \rightarrow Y$, $D(P)$ being the domain of P , be a nonlinear operator from a Banach space X to a Banach space Y and $\Gamma(x): Y \rightarrow X$ be a bounded linear operator associated with $x \in X$. We say that P has a contractor $\Gamma(x)$ if there is a positive number $q < 1$ such that

$$(2.10) \quad \|P(x + \Gamma(x)y) - Px - y\| \leq q\|y\|,$$

where $x \in D(P)$, $y \in Y$ are defined by the particular problem.

DEFINITION 2.2 [p. 10, [1]]. A contractor $\Gamma(x)$ is called regular if (2.10) is satisfied for all $y \in Y$ and $D(P) = \Gamma(x)(Y)$.

DEFINITION 2.3 [p. 6, [1]]. An operator $P: D(P) \subset X \rightarrow Y$ is closed on $D(P)$ if $x_n \in D(P)$, $x_n \rightarrow x$ and $Px_n \rightarrow y$ imply $x \in D(P)$ and $Px = y$.

Altman proved the following theorem:

THEOREM 2.2 [p. 13, Theorem 5.1, [1]]. Suppose that the closed nonlinear operator $P: D(P) \subset X \rightarrow Y$ has a bounded contractor Γ such that

$$(2.11) \quad (a) \quad x + \Gamma(x)y \in D(P), \quad \text{whenever } x \in D(P), \quad y \in Y;$$

$$(2.11) \quad (b) \quad \|P(x + \Gamma(x)y) - Px - y\| \leq q\|y\|, \quad y \in Y, \quad 0 < q < 1;$$

$$(2.12) \quad \|\Gamma(x)\| \leq B \quad \text{for all } x \in D(P).$$

Then the equation $Px = y$ has a solution for $y \in Y$. When Γ is regular, the assumption (2.11)(a) is readily satisfied and further, the solution is unique.

3. We generalize the notion of a closed operator to multivalued mappings as follows:

DEFINITION 3.1. A nonlinear multivalued operator $P: D(P) \subset X \rightarrow \text{CL}(Y)$, X, Y being Banach spaces, is closed on $D(P)$, if $x_n \rightarrow x$, $y_n \in Px_n$ and $y_n \rightarrow y$ imply that $x \in D(P)$ and $y \in Px$.

Let X_i, Y_i , $i = 1, \dots, n$ be Banach spaces and $b_{ik}, c_{ik} \geq 0$ $i, k = 1, \dots, n$. Let $a_{ik} = b_{ik} + c_{ik}$, $i, k = 1, \dots, n$, be positive and the numbers a'_{ik} defined by (2.5) and (2.6) fulfill (2.8). Then, by Lemma 2.2, there exists a positive solution r_1, \dots, r_n of the system of inequalities (2.7). We define

$$(3.1) \quad q = \max_i \left(r_i^{-1} \sum_{k=1}^n a_{ik} r_k \right).$$

Clearly, $0 < q < 1$, and

$$(3.2) \quad \sum_{k=1}^n a_{ik} r_k \leq q r_i, \quad i = 1, \dots, n.$$

THEOREM 3.1. Suppose that the closed nonlinear transformations $P_i: D \subset X_1 \times \dots \times X_n \rightarrow \text{CL}(Y_i)$, $i = 1, \dots, n$ fulfill the following:

(3.3) $\begin{aligned} &\text{there exist bounded linear operators } \Gamma_i(x_i): Y_i \rightarrow X_i, \quad x_i \in X_i, \\ &i = 1, \dots, n \text{ such that} \end{aligned}$

$$\|\Gamma_i(x_i)\| \leq B, \quad (x_1, \dots, x_n) \in D, \quad i = 1, \dots, n;$$

(3.4) (a) $(x_1 + \Gamma_1(x_1)y_1, \dots, x_n + \Gamma_n(x_n)y_n) \in D$ whenever
 $(x_1, \dots, x_n) \in D$ and $y_i \in Y_i$, $i = 1, \dots, n$;

(3.4) (b) $H_i[P_i(x_1 + \Gamma_1(x_1)y_1, \dots, x_n + \Gamma_n(x_n)y_n), P_i(x_1, \dots, x_n) + y_i]$
 $\leq \sum_{k=1}^n b_{ik} \|y_k\| + \sum_{k=1}^n c_{ik} D_k[y_k, y_k - P_k(x_1, \dots, x_n)]$
 $+ c D_i[y_i, y_i]$
 $- \Gamma_i(x_i) P_i(x_1 + \Gamma_1(x_1)y_1, \dots, x_n + \Gamma_n(x_n)y_n)]$
 $\text{for } (x_1, \dots, x_n) \in D, \quad y_i \in Y_i, \quad i = 1, \dots, n,$

where $a_{ik} = b_{ik} + c_{ik}$, $i, k = 1, \dots, n$ are positive and the numbers a'_{ik} defined by (2.5) and (2.6) satisfy (2.8);

(3.5) c is a constant such that $0 \leq cB < 1 - q$.

Then there exists $(x_1, \dots, x_n) \in D$ such that $\theta_i \in P_i(x_1, \dots, x_n)$ $i = 1, \dots, n$, where θ_i is the zero element of the Banach space Y_i , $i = 1, \dots, n$.

Proof. Let $(\dot{x}_1, \dots, \dot{x}_n) \in D$ be an arbitrary element. Choose

$\dot{y}_i \in P_i(\dot{x}_1, \dots, \dot{x}_n)$, $i = 1, \dots, n$. We can assume without loss of generality that $\|\dot{y}\| \leq r_i$, $r_i \geq 1$, $i = 1, \dots, n$, since the set of solutions to the system (2.7) is closed with respect to multiplication by positive scalars. Define

$$(3.6) \quad x_i^1 = \dot{x}_i - \Gamma_i(\dot{x}_i)\dot{y}_i, \quad i = 1, \dots, n.$$

Replacing x_i by \dot{x}_i and y_i by $-\dot{y}_i$ in (3.4)(b) and using (3.6) we get,

$$\begin{aligned} (3.7) \quad & H_i[P_i(x_1^1, \dots, x_n^1), P_i(\dot{x}_1, \dots, \dot{x}_n) - \dot{y}_i] \\ & \leq \sum_{k=1}^n b_{ik} \|\dot{y}_k\| + \sum_{k=1}^n c_{ik} D_k[-\dot{y}_k, -\dot{y}_k - P_k(\dot{x}_1, \dots, \dot{x}_n)] \\ & \quad + c D_i[-\dot{y}_i, -\dot{y}_i - \Gamma_i(\dot{x}_i)P_i(x_1^1, \dots, x_n^1)] \\ & \leq \sum_{k=1}^n b_{ik} \|\dot{y}_k\| + \sum_{k=1}^n c_{ik} \|\dot{y}_k\| + c D_i[\theta_i, \Gamma_i(\dot{x}_i)P_i(x_1^1, \dots, x_n^1)]. \end{aligned}$$

As $\dot{y}_i \in P_i(\dot{x}_1, \dots, \dot{x}_n)$, $\theta_i \in P_i(\dot{x}_1, \dots, \dot{x}_n) - \dot{y}_i$, $i = 1, \dots, n$. So, for $q > 0$, by Lemma 2.1, there exists an element $y_i^1 \in P_i(x_1^1, \dots, x_n^1)$, $i = 1, \dots, n$ such that

$$(3.8) \quad \|y_i^1 - \theta_i\| \leq H_i[P_i(x_1^1, \dots, x_n^1), P_i(\dot{x}_1, \dots, \dot{x}_n) - \dot{y}_i] + q.$$

From (3.7) and (3.8) we have

$$\begin{aligned} \|y_i^1\| & \leq \sum_{k=1}^n (b_{ik} + c_{ik}) \|\dot{y}_k\| + c \|\Gamma_i(x_i) y_i^1\| + q \\ & \leq \sum_{k=1}^n a_{ik} \|\dot{y}_k\| + cB \|y_i^1\| + q, \quad \text{by (3.3)} \end{aligned}$$

$$\begin{aligned} (1 - cB) \|y_i^1\| & \leq \sum_{k=1}^n a_{ik} r_k + q, \quad \text{by our assumption} \\ & \leq qr_i + q, \quad \text{by (3.2)} \\ & \leq 2qr_i, \quad \text{as } r_i \geq 1, \quad i = 1, \dots, n \quad \text{and } 0 < q < 1. \end{aligned}$$

Hence

$$(3.9) \quad \|y_i^1\| \leq \frac{2q}{(1 - cB)} r_i, \quad i = 1, \dots, n.$$

We shall now construct inductively sequences $\{x_i^m\}$ and $\{y_i^m\}$ $i = 1, \dots, n$ such that

$$(3.10) \quad (a) \quad (x_1^m, \dots, x_n^m) \in D, \quad y_i^m \in P_i(x_1^m, \dots, x_n^m)$$

$$(3.10) \quad (b) \quad \|y_i^m\| \leq (m+1) \left(\frac{q}{1 - cB} \right)^m r_i, \quad i = 1, \dots, n.$$

For $m = 1$, the above hypotheses are true, in view of (3.4)(a), (3.6) and (3.9). Assume the truth of (3.10)(a), (3.10)(b) for $m - 1 \in N$, i.e.,

$$(3.11) \quad (\text{a}) \quad (x_1^{m-1}, \dots, x_n^{m-1}) \in D, \quad y_i^{m-1} \in P_i(x_1^{m-1}, \dots, x_n^{m-1}),$$

$$(3.11) \quad (\text{b}) \quad \|y_i^{m-1}\| \leq m \left(\frac{q}{1 - cB} \right)^{m-1} r_i, \quad i = 1, \dots, n.$$

Define

$$(3.12) \quad x_i^m = x_i^{m-1} - \Gamma_i(x_i^m) y_i^{m-1}, \quad i = 1, \dots, n.$$

In (3.4)(b), replacing x_i by x_i^{m-1} , y_i by $-y_i^{m-1}$ and using (3.12) we have

$$\begin{aligned} (3.13) \quad & H_i[P_i(x_1^m, \dots, x_n^m), P_i(x_1^{m-1}, \dots, x_n^{m-1}) - y_i^{m-1}] \\ & \leq \sum_{k=1}^n b_{ik} \|y_k^{m-1}\| + \sum_{k=1}^n c_{ik} D_k[-y_k^{m-1}, -y_k^{m-1} - P_k(x_1^{m-1}, \dots, x_n^{m-1})] \\ & \quad + c D_i[-y_i^{m-1}, -y_i^{m-1} - \Gamma_i(x_i^{m-1}) P_i(x_1^m, \dots, x_n^m)] \\ & \leq \sum_{k=1}^n b_{ik} \|y_k^{m-1}\| + \sum_{k=1}^n c_{ik} \|y_k^{m-1}\| \\ & \quad + c D_i[\theta_i, \Gamma_i(x_i^{m-1}) P_i(x_1^m, \dots, x_n^m)]. \end{aligned}$$

Since $\theta_i \in P_i(x_1^{m-1}, \dots, x_n^{m-1}) - y_i^{m-1}$, given $q/(1 - cB)^{m-1} > 0$, there exists $y_i^m \in P_i(x_1^m, \dots, x_n^m)$ such that

$$\begin{aligned} (3.14) \quad & \|y_i^m - \theta_i\| \leq H_i[P_i(x_1^m, \dots, x_n^m), P_i(x_1^{m-1}, \dots, x_n^{m-1}) - y_i^{m-1}] \\ & \quad + \frac{q^m}{(1 - cB)^{m-1}}. \end{aligned}$$

From (3.13) and (3.14) we get

$$\begin{aligned} \|y_i^m\| & \leq \sum_{k=1}^n (b_{ik} + c_{ik}) \|y_k^{m-1}\| + cB \|y_i^m\| + \frac{q^m}{(1 - cB)^{m-1}} \\ (1 - cB) \|y_i^m\| & \leq m \left(\frac{q}{1 - cB} \right)^{m-1} \sum_{k=1}^n a_{ik} r_k + \frac{q^m}{(1 - cB)^{m-1}} \\ & \leq m a r_i \left(\frac{q}{1 - cB} \right)^{m-1} + \frac{q^m}{(1 - cB)^{m-1}}, \quad \text{by (3.2)} \end{aligned}$$

i.e.,

$$\begin{aligned} \|y_i^m\| & \leq m \left(\frac{q}{1 - cB} \right)^m r_i + \left(\frac{q}{1 - cB} \right)^m r_i \\ & \leq (m+1) \left(\frac{q}{1 - cB} \right)^m r_i, \quad \text{as } r_i \geq 1, \quad i = 1, \dots, n. \end{aligned}$$

Hence by induction (3.10)(b) holds for all $m = 0, 1, 2, \dots$. From (3.12), and (3.4)(a), it follows that $(x_1^m, \dots, x_n^m) \in D$. By construction, $y_i^m \in P_i(x_1^m, \dots, x_n^m)$. Hence by induction (3.10)(a) holds. By (3.5), $0 \leq q/(1 - cB) < 1$ and it follows from (3.10)(b) that $y_i^m \rightarrow \theta_i$, as $m \rightarrow \infty$, $i = 1, \dots, n$. From (3.12),

$$\begin{aligned} \|x_i^{m+1} - x_i^m\| &\leq \|\Gamma_i(x_i^m)y_i^m\| \\ &\leq B(m+1)\left(\frac{q}{1-cB}\right)^m r_i. \end{aligned}$$

Hence $\{x_i^m\}$ is a Cauchy sequence in X_i , $i = 1, \dots, n$. Therefore, $x_i^m \rightarrow x_i$, $i = 1, \dots, n$. As the operator P_i is closed, $y_i^m \in P_i(x_1^m, \dots, x_n^m)$, $y_i^m \rightarrow \theta_i$, $x_i^m \rightarrow x_i$, $i = 1, \dots, n$, imply that $(\theta_1, \dots, \theta_n) \in D$ and $\theta_i \in P(x_1, \dots, x_n)$, $i = 1, \dots, n$. \square

We now deduce Czerwinski's Theorem [4] as a corollary in the set-up of Banach spaces. Czerwinski proved his theorem for multivalued mappings on complete metric spaces.

THEOREM 3.2 [Theorem, [4]]. *Let X_i , $i = 1, \dots, n$ be Banach spaces and $b_{ik}, c_{ik} \geq 0$ for $i, k = 1, \dots, n$. Let $a_{ik} = b_{ik} + c_{ik}$, $i, k = 1, \dots, n$ be positive and let the numbers a_{ik}^l defined by (2.5) and (2.6) fulfill (2.8). Suppose that the transformations $F_i: X_1 \times \dots \times X_n \rightarrow \text{CL}(X_i)$, $i = 1, \dots, n$ fulfill*

$$\begin{aligned} (3.15) \quad H_i[F_i(x_1, \dots, x_n), F_i(z_1, \dots, z_n)] \\ &\leq \sum_{k=1}^n b_{ik} \|x_k - z_k\| + \sum_{k=1}^n c_{ik} D_k[x_k, F_k(x_1, \dots, x_n)] \\ &\quad + c D_i[z_i, F_i(z_1, \dots, z_n)], \quad i = 1, \dots, n; \end{aligned}$$

for $x_j, z_j \in X_j$, $j = 1, \dots, n$, where c fulfills the condition $0 \leq c < 1 - q$, q being defined by

$$q = \max_i \left(r_i^{-1} \sum_{k=1}^n a_{ik} r_k \right).$$

Then the system (F_1, \dots, F_n) has a fixed point, i.e., there exist points $x_i \in X_i$, $i = 1, \dots, n$ such that $x_i \in F_i(x_1, \dots, x_n)$, $i = 1, \dots, n$.

Proof. That the above theorem follows as a corollary for the Theorem 3.1 can be seen if we put $\Gamma_i(x_i) = I$, $(x_i \in X_i)$, the identity operator on X_i and $P_i(x_1, \dots, x_n) = x_i - F_i(x_1, \dots, x_n)$ $i = 1, \dots, n$ in (3.4)(b) and observe that it reduces to

$$\begin{aligned} H_i[x_i + y_i - F_i(x_1 + y_1, \dots, x_n + y_n), x_i + y_i - F_i(x_1, \dots, x_n)] \\ &\leq \sum_{k=1}^n b_{ik} \|y_k\| + \sum_{k=1}^n c_{ik} D_k[y_k, y_k - x_k + F_k(x_1, \dots, x_n)] \\ &\quad + c D_i[x_i + y_i, x_i + y_i - (x_i + y_i) + F_i(x_1 + y_1, \dots, x_n + y_n)] \end{aligned}$$

i.e.,

$$\begin{aligned} H_i[F_i(x_1 + y_1, \dots, x_n + y_n), F_i(x_1, \dots, x_n)] \\ &\leq \sum_{k=1}^n b_{ik} \|y_k\| + \sum_{k=1}^n c_{ik} [y_k - x_k + F_k(x_1, \dots, x_n)] \\ &\quad + c D_i[x_i + y_i, F_i(x_1 + y_1, \dots, x_n + y_n)]. \end{aligned}$$

Taking $x_i + y_i = z_i$, $i = 1, \dots, n$, we have

$$\begin{aligned} H_i[F_i(x_1, \dots, x_n), F_i(z_1, \dots, z_n)] \\ \leq \sum_{k=1}^n b_{ik} \|x_k - z_k\| + \sum_{k=1}^n c_{ik} D_k[x_k, F_k(x_1, \dots, x_n)] \\ + c D_i[z_i, F_i(z_1, \dots, z_n)], \end{aligned}$$

which is nothing but condition (3.15). It can be similarly shown that (3.15) implies (3.4)(b) in this case. To prove that the operator F_i , $i = 1, \dots, n$ is closed in the sense of Definition 3.1, observe that we have shown in the proof of Theorem 3.1 that $x_i^m \rightarrow x_i$ and $y_i^m \in P_i(x_1^m, \dots, x_n^m)$, i.e., $y_i^m \in x_i^m - F_i(x_1^m, \dots, x_n^m)$ $i = 1, \dots, n$, and $y_i^m \rightarrow 0$ as $m \rightarrow \infty$, i.e., $x_i^m - y_i^m \rightarrow x_i$, $i = 1, \dots, n$. It remains to show that $x_i \in F_i(x_1, \dots, x_n)$, $i = 1, \dots, n$.

$$\begin{aligned} D_i[x_i, F_i(x_1, \dots, x_n)] \\ \leq \|x_i - x_i^m\| + D_i[x_i^m, F_i(x_1, \dots, x_n)] \\ \leq \|x_i - x_i^m\| + H_i[F_i(x_1^{m-1}, \dots, x_n^{m-1}), F_i(x_1, \dots, x_n)] \\ \leq \|x_i - x_i^m\| + \sum_{k=1}^n b_{ik} \|x_k^{m-1} - x_k\| \\ + \sum_{k=1}^n c_{ik} D_k[x_k^{m-1}, F_k(x_1^{m-1}, \dots, x_n^{m-1})] + c D_i[x_i, F_i(x_1, \dots, x_n)] \\ \leq \|x_i - x_i^m\| + \sum_{k=1}^n b_{ik} \|x_k^{m-1} - x_k\| \\ + \sum_{k=1}^n c_{ik} \|x_k^{m-1} - x_k^m\| + c D_i[x_i, F_i(x_1, \dots, x_n)]. \end{aligned}$$

As $0 < c < 1$ and

$$\begin{aligned} D_i[x_i, F_i(x_1, \dots, x_n)] \\ \leq \frac{1}{1-c} \left[\|x_i - x_i^m\| + \sum_{k=1}^n b_{ik} \|x_k^{m-1} - x_k\| + \sum_{k=1}^n c_{ik} \|x_k^{m-1} - x_k^m\| \right] \end{aligned}$$

it follows that $D_i[x_i, F_i(x_1, \dots, x_n)] = 0$. Since $F_i(x_1, \dots, x_n)$ is a closed set, $x_i \in F_i(x_1, \dots, x_n)$. \square

Theorem 3.3 below is a generalization of Matkowski's Theorem 2.1 to multivalued mappings.

THEOREM 3.3. *Let X_i , $i = 1, \dots, n$ be Banach spaces and a_{ik} , $i, k = 1, \dots, n$ be positive and a'_{ik} be defined by (2.5) and (2.6) and fulfill (2.8). Suppose that the transformations $F_i: X_1 \times \dots \times X_n \rightarrow \text{CL}(X_i)$, $i = 1, \dots, n$ satisfy*

$$H_i[F_i(x_1, \dots, x_n), F_i(z_1, \dots, z_n)] \leq \sum_{k=1}^n a_{ik} \|x_k - z_k\|$$

for all $x_j, z_j \in X_j$, $j = 1, \dots, n$. Then the operator $F = (F_1, \dots, F_n)$ has a fixed point, i.e., there exist points $x_i \in X_i$, such that $x_i \in F_i(x_1, \dots, x_n)$ for all $i = 1, \dots, n$.

Proof. In Theorem 3.2, let $c_{ik} = 0$, $b_{ik} = a_{ik}$, $i, k = 1, \dots, n$ and $c = 0$. \square

The following is the multivalued version of Altman's Theorem 2.2.

THEOREM 3.4. Suppose a nonlinear closed operator $P: D(P) \subset X \rightarrow \text{CL}(Y)$ has a bounded contractor Γ satisfying

$$(3.16) \quad \begin{aligned} \text{(a)} \quad & x + \Gamma(x)y \in D(P), \text{ whenever } x \in D(P), y \in Y, \\ & D(P) \text{ being the domain of } P; \end{aligned}$$

$$(3.16) \quad \begin{aligned} \text{(b)} \quad & H[P(x + \Gamma(x)y), Px + y] \leq q\|y\|, \\ & x \in D(P), y \in Y, 0 < q < 1; \end{aligned}$$

$$(3.17) \quad \|\Gamma(x)\| \leq B, \quad x \in D(P).$$

Then there exists $x \in D(P)$ such that $\theta \in Px$, where θ is the zero element of Y .

Proof. For $n = 1$, Theorem 3.1 reduces to the above theorem for the choice $c_{ik} = 0$, $b_{ik} = a_{ik}$, $i, k = 1, \dots, n$ i.e., $b_{11} = a_{11} = q < 1$, and $c = 0$. \square

Besides, Theorem 3.1 yields as corollaries several fixed point theorems for single-valued mappings including the following theorem proved elsewhere (Theorem 2.1, [2]).

THEOREM 3.5. Let X_i, Y_i , $i = 1, \dots, n$ be Banach spaces and $T_i: D \subset X_1 \times \dots \times X_n \rightarrow Y_i$, $i = 1, \dots, n$ be closed non-linear operators. Suppose that there exist bounded linear operators $\Gamma_i(x_i): Y_i \rightarrow X_i$, $i = 1, \dots, n$ such that

$$(3.18) \quad \begin{aligned} \text{(a)} \quad & (x_1 + \Gamma_1(x_1)y_1, \dots, x_n + \Gamma_n(x_n)y_n) \in D \\ & \text{whenever } (x_1, \dots, x_n) \in D, y_i \in Y_i, i = 1, \dots, n; \end{aligned}$$

$$(3.18) \quad \begin{aligned} \text{(b)} \quad & \|T_i(x_1 + \Gamma_1(x_1)y_1, \dots, x_n + \Gamma_n(x_n)y_n) - T_i(x_1, \dots, x_n) - y_i\| \\ & \leq \sum_{k=1}^n a_{ik}\|y_k\|, \end{aligned}$$

where the nonnegative numbers a_{ik} , $i, k = 1, \dots, n$ are defined by (2.5) and (2.6) and fulfill (2.8);

$$(3.19) \quad \|\Gamma_i(x_i)\| \leq B, \quad i = 1, \dots, n, \quad (x_1, \dots, x_n) \in D.$$

Then the system of operator equations

$$(3.20) \quad T_i(x_1, \dots, x_n) = y_i, \quad i = 1, \dots, n$$

has a solution in D for arbitrary $y_i \in Y_i$, $i = 1, \dots, n$.

Proof. In Theorem 3.1, let $c_{ik} = 0$, $b_{ik} = a_{ik}$, $i, k = 1, \dots, n$ and $c = 0$. Define $P_i(x_1, \dots, x_n) = \{T_i(x_1, \dots, x_n)\}$ for $(x_1, \dots, x_n) \in D$, $i = 1, \dots, n$. Clearly the assumptions of Theorem 3.1 are satisfied and hence the system of equations (3.20) has a solution in D . \square

The above Theorem proved in [2] unified, in the setting of Banach spaces, Altman's extension of the contraction principle and Matkowski's fixed point theorem.

REFERENCES

1. M. Altman, *Contractors and contractor directions, Theory and Applications*, Marcel Dekker, New York, 1977.
2. K. Balakrishna Reddy and P. V. Subrahmanyam, *Altman's contractors and Matkowski's fixed point theorem*, to appear in J. Nonlinear Analysis, Theory methods and applications, 1981.
3. H. Covitz and S. B. Nadler, Jr., *Multivalued contraction mappings in generalized metric spaces*, Israel J. Math., **8** (1970), 5-11.
4. S. Czerwinski, *A fixed point theorem for a system of multivalued transformations*, Proc. Amer. Math. Soc., **55** (1976), 136-139.
5. J. Matkowski, *Some inequalities and generalization of Banach's principle*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys., **21** (1973), 323-324.
6. ———, *Integrable solutions of functional equations*, Dissert. Math. (Rozprawy Mat.), CXXVII (1975).
7. S. B. Nadler, Jr., *Multivalued contraction mappings*, Pacific J. Math., **30** (1969), 475-488.
8. ———, *Some results on multivalued contraction mappings*, Lecture Notes in Math., Vol. 171, Springer-Verlag, Berlin 1970, 64-69.
9. S. Reich, *Kannan's fixed point theorem*, Boll. Un. Mat. Ital., **4** (1971), 1-11.

Received September 3, 1980.

THE RAMANUJAN INSTITUTE
UNIVERSITY OF MADRAS
MADRAS 600 005
INDIA
AND
INDIAN INSTITUTE OF TECHNOLOGY
MADRAS 600036
INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, California 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR AGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. FINN and J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 99, No. 1

May, 1982

Mariano Giaquinta, Jindrich Nečas, O. John and J. Stará, On the regularity up to the boundary for second order nonlinear elliptic systems	1
Siegfried Graf, Realizing automorphisms of quotients of product σ -fields	19
Alfred Washington Hales and Ernst Gabor Straus, Projective colorings	31
Sandra Hayes, The weak Nullstellensatz for finite-dimensional complex spaces	45
Gerald Norman Hile and Murray Harold Protter, The Cauchy problem and asymptotic decay for solutions of differential inequalities in Hilbert space	57
Robert D. Little, Projective space as a branched covering with orientable branch set	89
Jaroslav Mach, On the proximinality of Stone-Weierstrass subspaces	97
John C. Morgan, II, On product bases	105
K. Balakrishna Reddy and P. V. Subrahmanyam, Altman's contractors and fixed points of multivalued mappings	127
James Ted Rogers Jr., Decompositions of homogeneous continua	137
Ahmed Ramzy Sourour, Characterization and order properties of pseudo-integral operators	145
Robert Moffatt Stephenson Jr., Pseudocompact and Stone-Weierstrass product spaces	159
Bruce Stewart Trace, On attaching 3-handles to a 1-connected 4-manifold	175
Akihito Uchiyama, The construction of certain BMO functions and the corona problem	183
Thomas Alva Whitehurst, An application of orthogonal polynomials to random walks	205
David J. Winter, Root locologies and idempotents of Lie and nonassociative algebras	215
William Robin Zame, The classification of uniform algebras on plane domains	231