

Pacific Journal of Mathematics

ON THE APPROXIMATION OF SINGULARITY SETS BY ANALYTIC VARIETIES

HERBERT JAMES ALEXANDER AND JOHN WERMER

ON THE APPROXIMATION OF SINGULARITY SETS BY ANALYTIC VARIETIES

HERBERT ALEXANDER AND JOHN WERMER

We study the problem of approximating the singularity set of an analytic function of two complex variables, lying in a product domain in \mathbb{C}^2 , by analytic varieties.

Let D denote the open unit disk and let

$$D \times \mathbb{C} = \{(z, w) \mid z \in D, w \in \mathbb{C}\} \subset \mathbb{C}^2.$$

We consider a compact set X contained in the unit bidisk $|z| \leq 1, |w| \leq 1$. Let X^0 denote $X \cap (D \times \mathbb{C})$. We assume that there exists a function ϕ which is analytic on $D \times \mathbb{C} \setminus X^0$ and singular at each point of X^0 . If there exists such a ϕ we call X a *singularity set*.

For each λ in \bar{D} we put

$$X_\lambda = \{w \in \mathbb{C} \mid (\lambda, w) \in X\}.$$

Each X_λ is then a compact subset of $|w| \leq 1$. We assume $X_\lambda \neq \emptyset$, for each λ .

Singularity sets were first studied by Hartogs, in [3]. Hartogs showed that if for some integer p X_λ contains at most p points for each λ , then X^0 is an analytic subvariety of $D \times \mathbb{C}$. Further results on singularity sets were given by Oka, [5], and Nishino, [4].

Recently one of us in [7] and Slodkowski in [6] studied general singularity sets. In particular, Theorem 1 in [7] gives that the maximum principle holds on X^0 for restrictions to X^0 of polynomials in z and w , in the sense that for each compact subset N of X^0 and each $(z_0, w_0) \in N$,

$$|P(z_0, w_0)| \leq \max_{\partial N} |P|,$$

for each polynomial P .

(See also [6], Theorem II, (vi).) Here ∂N denotes the boundary of N relative to X^0 . In particular, fix $R < 1$. Put $N = \{(\lambda, w) \in X \mid |\lambda| \leq R\}$. Then $\partial N = \{(\lambda, w) \in X \mid |\lambda| = R\}$. Hence, for $(z_0, w_0) \in X$, $|z_0| < R$, we have for each polynomial P ,

$$|P(z_0, w_0)| \leq \max_{X \cap \{|z|=R\}} |P|.$$

Letting $R \rightarrow 1$, we get

$$(1) \quad |P(z_0, w_0)| \leq \max_{X \cap \{|z|=1\}} |P|, \quad (z_0, w_0) \in X.$$

In order to account for the inequality (1), one of us, (Alexander), suggested that it may be possible to approximate an arbitrary singularity set by one-dimensional analytic subvarieties of $D \times \mathbb{C}$.

Question 1. Given a singularity set X and an open neighborhood U of X in \mathbb{C}^2 . Can we find a one-dimensional analytic subvariety of $D \times \mathbb{C}$ which is contained in U ?

In this paper, we look at a special case of this question. For each λ in $|\lambda| < 1$, we fix an open disk D_λ with center $a(\lambda)$ and radius r (r fixed and > 0). Consider the tube

$$T = \{(\lambda, w) \mid |\lambda| < 1, w \in D_\lambda\}.$$

Question 2. Suppose X is a singularity set contained in the tube. Then for each λ , $X_\lambda \subset D_\lambda$. Can we find a function f analytic on $|\lambda| < 1$ such that the graph of f lies in T , i.e. such that, for every λ , $f(\lambda) \in D_\lambda$, or, in other words,

$$|f(\lambda) - a(\lambda)| < r?$$

We have the following partial answer.

THEOREM. *Let $\lambda \mapsto a(\lambda)$ be a continuous function defined for $|\lambda| \leq 1$ with $|a(\lambda)| \leq 1$ for all λ . Fix $r > 0$. Suppose that there exists a singularity set X such that for each λ , $|\lambda| \leq 1$, $X_\lambda \subset \{w \mid |w - a(\lambda)| < r\}$. Then there exists an analytic function $\lambda \mapsto f(\lambda)$ such that for each λ in the unit disk*

$$(2) \quad |f(\lambda) - a(\lambda)| \leq 4r.$$

Let $C(\Gamma)$ denote the Banach space of functions g which are continuous on $\Gamma = \{|z| = 1\}$, with

$$\|g\| = \max_{\Gamma} |g|.$$

Let $A(\Gamma)$ denote the closed subspace of $C(\Gamma)$ consisting of those functions in $C(\Gamma)$ which have an analytic extension to $|\lambda| < 1$. We write a for the element of $C(\Gamma)$ obtained from our function a by restricting it to Γ . The distance from a to $A(\Gamma)$, in the norm of $C(\Gamma)$, equals

$$\sup |L(a)|$$

taken over all bounded linear functionals L on $C(\Gamma)$ with $\|L\| = 1$ and $L(g) = 0$ for every $g \in A(\Gamma)$. Each such functional L is given, in view of the Theorem of F. and M. Riesz, by a summable function ϕ in the Hardy space H^1 such that

$$(3a) \quad L(g) = \frac{1}{2\pi} \int_0^{2\pi} g(e^{i\theta}) \phi(e^{i\theta}) d\theta,$$

$$(3b) \quad \phi(0) = 0,$$

and

$$(3c) \quad \frac{1}{2\pi} \int_0^{2\pi} |\phi(e^{i\theta})| d\theta = 1.$$

LEMMA. Let $\phi \in H^1$ and assume (3b), (3c). Then

$$(4) \quad \left| \frac{1}{2\pi} \int_0^{2\pi} a(e^{i\theta}) \phi(e^{i\theta}) d\theta \right| \leq r.$$

Proof. Since polynomials in λ are dense in H^1 in L^1 -sense, it suffices to prove (4) where ϕ is a polynomial.

Fix w_0 so that $(O, w_0) \in X$. It follows from the inequality (1) that there exists a representing measure μ for (O, w_0) relative to polynomials which is carried on $X \cap \{|\lambda| = 1\}$. (See e.g. [2], pp. 79–81.) μ is a probability measure such that

$$(5) \quad Q(O, w_0) = \int Q d\mu$$

for every polynomial Q .

We denote by μ^* the projection of μ under the map λ . Then μ^* is a probability measure on the circle $|\lambda| = 1$ so that

$$(6) \quad \int_{X \cap \{|\lambda|=1\}} F(\lambda) d\mu(\lambda, w) = \int_{|\lambda|=1} F(\lambda) d\mu^*(\lambda)$$

for every F in $C(\Gamma)$.

Then for $n = 1, 2, \dots$

$$\int_{|\lambda|=1} \lambda^n d\mu^*(\lambda) = \int \lambda^n d\mu = \lambda^n(O, w_0) = 0.$$

Conjugating, we get

$$\int_{|\lambda|=1} \overline{\lambda^n} d\mu^*(\lambda) = 0, \quad n = 1, 2, \dots$$

Finally, μ^* has total mass 1. So $\mu^* = d\theta/2\pi$ on Γ . Thus for all $F \in C(\Gamma)$:

$$(7) \quad \int F(\lambda) d\mu(\lambda, w) = \frac{1}{2\pi} \int_{|\lambda|=1} F(\lambda) d\theta.$$

Fix now a polynomial ϕ satisfying (3b), (3c). Then

$$\begin{aligned} \frac{1}{2\pi} \int_{|\lambda|=1} a(\lambda)\phi(\lambda) d\theta &= \int_{X \cap \{|\lambda|=1\}} a(\lambda)\phi(\lambda) d\mu(\lambda, w) \\ &= \int (a(\lambda) - w)\phi(\lambda) d\mu(\lambda, w), \end{aligned}$$

since

$$\int w\phi(\lambda) d\mu(\lambda, w) = w_0\phi(0) = 0.$$

Hence

$$\left| \frac{1}{2\pi} \int_{|\lambda|=1} a(\lambda)\phi(\lambda) d\theta \right| \leq \int |a(\lambda) - w| |\phi(\lambda)| d\mu(\lambda, w).$$

For each $(\lambda, w) \in \text{supp } \mu$, $|a(\lambda) - w| \leq r$. Hence the right hand side

$$\leq \int r |\phi(\lambda)| d\mu(\lambda, w) = \int_{|\lambda|=1} r |\phi(\lambda)| \frac{d\theta}{2\pi} = r.$$

Thus (4) holds and we are done.

Proof of Theorem. In view of the Lemma, $\sup |L(a)|$, taken over all functionals L with $\|L\| = 1$ and $L = 0$ on $A(\Gamma)$, $\leq r$, and so the distance from a to $A(\Gamma) \leq r$. Thus $\exists f_0 \in A(\Gamma)$ with

$$(8) \quad |f_0(\lambda) - a(\lambda)| \leq 2r, \quad |\lambda| = 1.$$

Without loss of generality, we take f_0 to be a polynomial in λ .

In view of (8), for each λ on $|\lambda| = 1$ $|w - f_0(\lambda)| \leq 3r$ for every $w \in X_\lambda$.

Denote by Q the polynomial in λ and w ,

$$Q(\lambda, w) = w - f_0(\lambda).$$

We just saw that

$$|Q(\lambda, w)| \leq 3r$$

for each $(\lambda, w) \in X \cap (|\lambda| = 1)$. Because of (1), it follows that for every (λ_0, w_0) in X ,

$$|Q(\lambda_0, w_0)| \leq 3r.$$

Thus

$$|f_0(\lambda_0) - w_0| \leq 3r, \quad (\lambda_0, w_0) \in X.$$

Hence

$$|f_0(\lambda_0) - a(\lambda_0)| \leq 4r, \quad |\lambda_0| \leq 1.$$

This was to be proved.

Note. A natural extension of the situation treated in the Theorem is the following: let X be a singularity set. Fix $r > 0$.

Assume there exist continuous functions $a_1(\lambda), a_2(\lambda), \dots, a_n(\lambda)$ defined on $|\lambda| \leq 1$ such that the roots $w_1(\lambda), w_2(\lambda), \dots, w_n(\lambda)$ of the equation

$$(9) \quad w^n + a_1(\lambda)w^{n-1} + \dots + a_n(\lambda) = 0$$

give an approximation to X_λ in the sense that for each λ in $|\lambda| \leq 1$, X_λ is contained in the union of the disks

$$|w - w_j(\lambda)| < r, \quad j = 1, 2, \dots, n.$$

Can we find good approximations $a_1^0(\lambda), \dots, a_n^0(\lambda)$ to the $a_j(\lambda)$, where the a_j^0 are analytic on $|\lambda| < 1$? In this case, the goodness of approximation should be expressed in terms of r and n . The analytic variety defined by

$$(10) \quad w^n + a_1^0(\lambda)w^{n-1} + \dots + a_n^0(\lambda) = 0, \quad |\lambda| < 1,$$

will then provide an approximation to X .

Note. Singularity sets occur in the spectral theory of linear operators. See Aupetit, [1], and Slodkowski, [6].

REFERENCES

1. B. Aupetit, *Analytic multivalued functions in Banach Algebras and Uniform Algebras*, Department of Mathematics, Université Laval, Quebec (1981) (multigraphed).
2. A. Browder, *Introduction to Function Algebras*, W. A. Benjamin, Inc. (1979).
3. F. Hartogs, *Über die aus den singulären Stellen einer analytischer Funktion mehrerer Veränderlichen bestehenden Gebilde*, Acta Math., **32** (1909), 57–79.
4. T. Nishino, *Sur les ensembles pseudo-concaves*, J. Math. Kyoto Univ., **1** (1962), 225–245.
5. K. Oka, *Notes sur les familles des fonctions analytiques multiforme etc.*, J. Sci. Hiroshima Univ., Series A, **4** (1934), 93–98.
6. Z. Slodkowski, *Analytic Set-Valued Functions and Spectra*, Math. Annalen, **256** (3), (1981).

7. J. Wermer, *Maximum modulus algebras and singularity sets*, Proc. Royal Soc. of Edinburgh, **86A** (1980), 327–331.

Received July 1, 1981 and in revised form November 1, 1981.

UNIVERSITY OF ILLINOIS AT CHICAGO CIRCLE
CHICAGO, IL 60680

AND

BROWN UNIVERSITY
PROVIDENCE, RI 02912

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P. O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1983 by Pacific Journal of Mathematics

Leo James Alex, Simple groups and a Diophantine equation	257
Herbert James Alexander and John Wermer, On the approximation of singularity sets by analytic varieties	263
Waleed A. Al-Salam and Mourad Ismail, Orthogonal polynomials associated with the Rogers-Ramanujan continued fraction	269
J. L. Brenner and Roger Conant Lyndon, Permutations and cubic graphs	285
Ian George Craw and Susan Ross, Separable algebras over a commutative Banach algebra	317
Jesus M. Dominguez, Non-Archimedean Gel'fand theory	337
David Downing and Barry Turett, Some properties of the characteristic of convexity relating to fixed point theory	343
James Arthur Gerhard and Mario Petrich, Word problems for free objects in certain varieties of completely regular semigroups	351
Moses Glasner and Mitsuru Nakai, Surjective extension of the reduction operator	361
Takesi Isiwata, Ultrafilters and mappings	371
Lowell Duane Loveland, Double tangent ball embeddings of curves in E^3	391
Douglas C. McMahon and Ta-Sun Wu, Homomorphisms of minimal flows and generalizations of weak mixing	401
P. H. Maserick, Applications of differentiation of \mathcal{L}_p -functions to semilattices	417
Wayne Bruce Powell and Constantine Tsinakis, Free products in the class of abelian l -groups	429
Bruce Reznick, Some inequalities for products of power sums	443
C. Ray Rosentrater, Compact operators and derivations induced by weighted shifts	465
Edward Silverman, Basic calculus of variations	471
Charles Andrew Swanson, Criteria for oscillatory sublinear Schrödinger equations	483
David J. Winter, The Jacobson descent theorem	495