

Pacific Journal of Mathematics

AN INTERPOLATION THEOREM FOR H_E^∞

KNUT ØYMA

AN INTERPOLATION THEOREM FOR H_E^∞

KNUT ØYMA

We prove a synthesis of Carleson's interpolation theorem, the Rudin-Carleson theorem and an interpolation theorem of S. A. Vinogradov.

Let D be the open unit disc in \mathbb{C} and let T be its boundary. By $A(D)$ we mean the set of functions continuous on \bar{D} analytic on D . H^∞ is the set of bounded analytic functions on D , and if E is a subset of T , H_E^∞ is the set of functions continuous on $D \cup E$ bounded and analytic on D .

The Rudin-Carleson theorem states that if K is a closed subset of T of measure zero, then $A(D)|_K = C(K)$. This was proved independently by W. Rudin and L. Carleson [8], [3].

A sequence $\{z_n\} \subset D$ is said to be uniformly separated if

$$\inf_n \prod_{m \neq n} \left| \frac{z_n - z_m}{1 - \bar{z}_n z_m} \right| = \delta > 0.$$

Carleson's interpolation theorem states that $H^\infty|_{\{z_n\}} = l^\infty$ if and only if $\{z_n\}$ is uniformly separated. This was first proved in [2]. Other proofs can be found in [5] and [10].

Let $F \subset \mathbb{N} \cup \{0\}$. A function $f(z) = \sum a_n z^n \in H^1$ is said to be an F function if $a_n = 0$ for $n \notin F$. For a definition and properties of the H^p spaces see [4]. F is said to be of type $\Lambda(s)$ if for every $r < s$ there is a constant K depending on F , r and s only such that $\|f\|_s \leq K\|f\|_r$ for every F function. If $F = \{n_k\}$ satisfies $n_{k+1}/n_k > \lambda > 1$, then F is of type $\Lambda(s)$ for every $s \in \langle 0, \infty \rangle$. Other sets of type $\Lambda(s)$ exist. See [7]. Let $\{n_k\}$ be of type $\Lambda(2)$ and let R be the operator from $A(D) \rightarrow l^2$ defined by $R(\sum a_n z^n) = \{a_{n_k}\}$. S. A. Vinogradov proved that R is onto. In fact he proved much more. See [11].

These results do not live their own lives separate from each other. In [6] E. A. Heard and J. H. Wells proved that if E is an open subset of T and S is a relatively closed subset of $D \cup E$ such that $S \cap E$ has measure zero and $S \cap D$ is uniformly separated, then $H_E^\infty|_S = C_b(S)$, the space of all bounded continuous functions on S . Vinogradov proved in [11] that if K is a closed subset of T of measure zero, $g \in C(K)$ and $\{b_k\} \in l^2$, then

there is an $f \in A(D)$ such that $f|_K = g$ and $R(f) = \{b_k\}$. We intend to prove:

THEOREM. *Let E be an open subset of T and assume that S is a relatively closed subset of $D \cup E$ such that $S \cap E$ has measure zero, $S \cap D$ is uniformly separated and $0 \notin S$. Assume $F = \{n_k\}$ is an increasing sequence of integers of type $\Lambda(2)$ such that $\lim_{k \rightarrow \infty} (n_{k+1} - n_k) = \infty$. If $\beta(S) \in C_b(S)$ and $\{b_k\} \in l^2$, there is a function $f(z) = \sum a_n z^n \in H_E^\infty$ such that $f|_S = \beta$ and $a_{n_k} = b_k$ for all k .*

REMARK. $0 \notin S$ represents no loss of generality since we may have $0 \in \{n_k\}$.

Before proving the theorem, we are going to develop some background material. Let $S \cap D = \{z_n\}$ and let

$$\inf_n \prod_{m \neq n} \left| \frac{z_m - z_n}{1 - \bar{z}_m z_n} \right| = \delta > 0.$$

Then there exists a real number M with the following property: Given $\{w_n\} \in \text{ball } l^\infty$, we can find a real number α and a Blaschke product $B(z)$ such that $Me^{i\alpha} B(z_n) = w_n$ for all n . The zeros $\{\xi_n\}$ of $B(z)$ can be chosen to satisfy $\psi(z_n, \xi_n) < \delta$ where $\psi(a, b) = |(a - b)/(1 - \bar{a}b)|$ is the pseudo-hyperbolic metric on D . This shows that $B(z)$ has analytic continuation across $T \setminus \{z_n\}$. The result is due to J. Earl [5]. We want to prove that the mass of the Taylor coefficients of $B(z)$ regarded as an element of H^2 is concentrated on the first coefficients.

LEMMA 1. *Let $B(z) = \sum a_n z^n$ be as above. If $\varepsilon > 0$ then there is an integer $N = N(\varepsilon)$ independent of $\{\xi_n\}$ such that $\sum_{n=N}^\infty |a_n|^2 < \varepsilon$.*

Proof. ε is now fixed. Let

$$B_K(z) = \prod_{n=K}^\infty \frac{|\xi_n|}{\xi_n} \cdot \frac{\xi_n - z}{1 - \bar{\xi}_n z}.$$

Since $\psi(\xi_n, z_n) < \delta$, a calculation shows that

$$1 - |\xi_n| \leq (2/(1 - \delta))(1 - |z_n|).$$

Hence $\lim_{K \rightarrow \infty} \sum_{n=K}^\infty (1 - |\xi_n|) = 0$ uniformly in $\{\xi_n\}$. This shows that $B_K(0) \rightarrow 1$. Since $\|B_K\|_2 = 1$, $B_K(z) = \sum_{n=0}^\infty a_{n,K} z^n$ satisfies $\sum_{n=N_K}^\infty |a_{n,K}|^2 < \varepsilon/2$ for $N_K = 1$ if K is chosen large.

$$B_{K-1}(z) = B_K(z) \cdot \frac{|\xi_{K-1}|}{\xi_{K-1}} \cdot \frac{\xi_{K-1} - z}{1 - \bar{\xi}_{K-1} z}.$$

We have

$$B_K(z) = \sum_{n=0}^{N_K} a_{n,K} z^n + \sum_{n=N_K+1}^{\infty} a_{n,K} z^n = p(z) + \varepsilon_p(z)$$

where $\|\varepsilon_p\|_2^2 < \varepsilon/2$ and $\|p\|_2 \leq 1$.

$$\frac{|\xi_{K-1}|}{\xi_{K-1}} \cdot \frac{\xi_{K-1} - z}{1 - \xi_{K-1}z} = \sum_{n=0}^{\infty} b_n(\xi_{K-1}) z^n.$$

Since $\psi(z_{K-1}, \xi_{K-1}) < \delta$ this converges uniformly on D independent of ξ_{K-1} . Choose R such that

$$\sum_{n=0}^R b_n(\xi_{K-1}) z^n + \sum_{n=R+1}^{\infty} b_n(\xi_{K-1}) z^n = q(z) + \varepsilon_q(z)$$

satisfies $\|\varepsilon_q\|_\infty < \eta$, $\|q\|_\infty < 1 + \eta$ where η is to be chosen below. We have

$$B_{K-1} = (p + \varepsilon_p)(q + \varepsilon_q) = pq + \varepsilon_p q + p \varepsilon_q + \varepsilon_p \varepsilon_q.$$

pq is a polynomial of degree $N_K + R$. It is not the $(N_K + R)$ -partial sum of the Taylor series of B_{K-1} , but deleting coefficients decreases the $\|\cdot\|_2$ norm. For $B_{K-1}(z) = \sum C_n z^n$ we therefore have

$$\begin{aligned} \left(\sum_{n=R+N_K+1}^{\infty} |C_n|^2 \right)^{1/2} &= \|B_{K-1}(z) - \sum_{n=0}^{R+N_K} C_n z^n\|_2 \\ &\leq \|\varepsilon_p \cdot q\|_2 + \|p \varepsilon_q\|_2 + \|\varepsilon_p \varepsilon_q\|_2 \\ &\leq \|\varepsilon_p\|_2 \cdot \|q\|_\infty + \|p\|_2 \cdot \|\varepsilon_q\|_\infty + \|\varepsilon_p\|_2 \cdot \|\varepsilon_q\|_\infty \\ &\leq \sqrt{\varepsilon/2} (1 + \eta) + \eta + \sqrt{\varepsilon/2} \cdot \eta < \sqrt{3\varepsilon/4} \end{aligned}$$

if η is chosen small. Continuing in the same way, the lemma is proved in a finite number of steps.

We are now going to take a look at Vinogradov's theorem. If $F = \{n_k\}$ is of type $\Lambda(2)$, the mapping $R: A(D) \rightarrow l^2: \sum a_n z^n \rightarrow \{a_{n_k}\}$ is onto. The open mapping theorem gives that $R(\text{ball } A(D)) \supseteq c \text{ ball } l^2$ for some $c > 0$. To obtain an estimate for c we need a result of Smirnov. Let $f(\xi)$ be integrable over the unit circle and let

$$h(z) = \frac{1}{2\pi} \int_T \frac{f(\xi)}{\xi - z} d\xi.$$

Then $h \in H^{1/2}$ and $\|h\|_{1/2} \leq K_1 \|f\|_1$. For a proof see p. 35 of [4] or [11]. Since F is of type $\Lambda(2)$, we have $\|f\|_2 \leq K_2 \|f\|_{1/2}$ for every F function in

H^2 . Vinogradov proves his theorem by showing that the adjoint mapping $R^*: (l^2)^* = l^2 \rightarrow A(D)^*$ satisfies

$$\|R^*(x)\| \geq (1/2\pi K_1 K_2) \|x\|.$$

This is proved more generally on the first seven pages of [11]. Using a result of Banach, Lemma 4.13 of [9], we get $R(\text{ball } A(D)) \supseteq (1/2\pi K_1 K_2) \text{ball } l^2$ if by ball we mean open ball. Our balls are open from now on.

If $F = \{n_k\}_{k=1}^\infty$, consider the set $F' = \{n_k - n_K\}_{k=K+1}^\infty$. F' is also of type $\Lambda(2)$, and it is not difficult to see that the associated constant $K'_2 \leq K_2$. If R' is the operator from $A(D)$ to l^2 associated with F' we see that $R'(\text{ball } A(D)) \supseteq (1/2\pi K_1 K_2) \text{ball } l^2$.

The proof of the theorem will also make use of

LEMMA 2. *Let $T: X \rightarrow Y$ be a continuous linear mapping between Banach spaces. Assume there are constants $\varepsilon < 1$ and M such that for all $y \in \text{ball } B$ there is $x \in X$ such that $\|x\| < M$ and $\|Tx - y\| < \varepsilon$. Then T is onto.*

For a proof see [1]. We now prove the theorem. Assume first that $S \cap E = \emptyset$. Choose an integer K such that $f_K(z) = B_K(z)/B_K(0) = 1 + \varepsilon(z)$ satisfies $\|f_K\|_\infty < 2$ and $\|\varepsilon\|_2 < 1/4\pi K_1 K_2$. Let $B_K \cdot H_E^\infty$ be the subspace of H_E^∞ consisting of the functions that vanish at z_n for $n \geq K$. Given $\{b_k\} \in \text{ball } l^2$, choose $g(z) = \sum a_n z^n \in A(D)$ such that $a_{n_k} = b_k$ for all k and $\|g(z)\|_\infty \leq 2\pi K_1 K_2$. Let

$$g_K(z) = g(z)f_K(z) = \sum c_n z^n \in B_K H_E^\infty,$$

$$\|g_K(z)\|_\infty \leq 4\pi K_1 K_2$$

and

$$\|\{b_k - c_{n_k}\}\|_2 \leq \|\varepsilon(z)g(z)\|_2 \leq \|\varepsilon(z)\|_2 \cdot \|g(z)\|_\infty < 1/2.$$

Lemma 2 now proves that $R(B_K H_E^\infty) = l^2$. Let $\{w_n\}_{n=K}^\infty \in l^\infty$ and $\{b_k\} \in l^2$ be given. Choose $h(z) = \sum d_n z^n \in H_E^\infty$ such that $h(z_n) = w_n$ for $n \geq K$ and choose $j(z) = \sum l_n z^n \in B_K H_E^\infty$ such that $l_{n_k} = b_k - d_{n_k}$ for all k . The function $r(z) = h(z) + j(z) = \sum t_n z^n$ satisfies $r(z_n) = w_n$ for $n \geq K$ and $t_{n_k} = b_k$ for all k . This proves the theorem for $\{z_n\}_{n=1}^\infty$ replaced by $\{z_n\}_{n=K}^\infty$. The proof will be complete if we can prove that K can be replaced by $K-1$. To obtain this it is enough to find a function $f(z) = \sum a_n z^n \in B_K \cdot H_E^\infty$ such that $a_{n_k} = 0$ for all k and $f(z_{K-1}) = 1$.

Such a function is likely to exist because it is easy to prove that there are many functions in $B_K H_E^\infty$ with F coefficients zero. All these functions could, however, vanish at z_{K-1} (a black hole). In that case, then for every $f(z) = \sum a_n z^n \in B_K \cdot H_E^\infty$, $f(z_{K-1})$ would be a function of $\{a_{n_k}\}$ alone.

Let $f(z) = \sum a_n z^n \in B_K H_E^\infty$. Look at $f(z_{K-1}) \leftarrow f(z) \xrightarrow{R} \{a_{n_k}\} \in l^2$. $\{a_{n_k}\} \rightarrow f(z_{K-1})$ is now seen to be a well-defined linear functional on l^2 since R is onto. This functional is continuous since every $x \in \text{ball } l^2$ comes from a function of norm $< C$ as an application of the open mapping theorem shows. Therefore there exists a unique $\{\lambda_k\} \in l^2$ such that

$$(*) \quad f(z_{K-1}) = \sum_k a_{n_k} \lambda_k \quad \text{for every } f(z) = \sum a_n z^n \in B_K \cdot H_E^\infty.$$

Infinitely many $\lambda_k \neq 0$. If this were not so, let λ_M be the largest. If $f(z) \in B_K H_E^\infty$. Then $z^{n_{M+1}} f(z)$ would vanish at z_{K-1} . This is clearly impossible. Since $\{\lambda_k\}$ is unique, the relation $(*)$ is impossible if we delete some n_N from F for which $\lambda_N \neq 0$. If we do so, K can be replaced by $K-1$. We may choose n_N arbitrary large. Doing so we have pushed the problem from $\{z_n\}$ to F . We now prove that n_N can be replaced.

Let $\{z_n^*\} = \{z_n\}_{n=K-1}^\infty \cup \{0\}$. Every sequence $\{w_n\} \in \text{ball } l^\infty$ can be interpolated at $\{z_n^*\}$ by a function of the form $Me^{i\alpha} B(z) = \sum l_n z^n$ as pointed out above. Choose an integer Q independent of $\{w_n\}$ such that

$$(**) \quad \left(\sum_{n=Q}^\infty |l_n|^2 \right)^{1/2} < \frac{1}{10\pi K_1 K_2}.$$

This is possible by Lemma 1.

Choose n_N such that $\lambda_N \neq 0$ and $n_{N+1} - n_N > Q$. Let $F' = \{n_k - n_N\}_{k=N+1}^\infty$ and let

$$\mathfrak{B} = \{f(z) = \sum a_n z^n \in H_E^\infty : a_n = 0 \text{ for } n \in F'\}.$$

We want to prove that $\mathfrak{B} | \{z_n^*\} = l^\infty$. Let $\{w_n\} \in \text{ball } l^\infty$ be given. Choose α and $B(z)$ as above such that $Me^{i\alpha} B(z_n^*) = w_n$ for all n . Choose $h(z) = \sum b_n z^n \in A(D)$ such that $b_n = l_n$ for $n \in F'$ and such that $\|h(z)\| \leq \frac{1}{5}$. This is possible by $(**)$ and the remark following Vinogradov's theorem. $f(z) = Me^{i\alpha} B(z) - h(z)$ has the following properties: $f \in \mathfrak{B}$, $\|f\| \leq M + \frac{1}{5}$, $|f(z_n^*) - w_n| < \frac{1}{5}$. Lemma 2 now proves that $\mathfrak{B} | \{z_n^*\} = l^\infty$. n_N can now be replaced: Let $\{w_n\} \in l^\infty$, $\{b_k\} \in l^2$. Take $f(z) = \sum a_n z^n \in H_E^\infty$ such that $f(z_n) = w_n$ for $n \geq K-1$ and $a_{n_k} = b_k$ for $n_k \in F \setminus \{n_N\}$.

Choose $g(z) \in \mathfrak{B}$ such that $g(0) = 1$, $g(z^*) = 0$ for $z_n^* \neq 0$. Let $r(z) = z^n g(z) = \sum t_n z^n$. We have: $r(z_n) = 0$ for $n \geq K - 1$, $t_n = 0$ for $n \in F \setminus \{n_N\}$, $t_{n_N} = 1$. Our interpolation problem is now solved by the function $f(z) + \lambda r(z)$ for a proper choice of λ .

The proof is now complete except we assumed $S \cap E = \emptyset$. Using the Heard and Wells result, we may assume $\beta|_S = 0$. Let $E' = E \setminus S$, $\mathcal{C} = \{f \in H_E^\infty: f|_S = 0\}$, $\mathcal{C}' = \{f \in H_{E'}^\infty: f|_S = 0\}$. The proof will be complete if we can prove $R(\mathcal{C}) = l^2$. By what we have just proved and the open mapping theorem, $R(k \cdot \text{ball } \mathcal{C}') \supseteq \text{ball } l^2$ for some constant k . Now choose $g \in H_E^\infty$ such that $g = 0$ on $S \cap T$, $\|g\| \leq 1$ and $g(z) = 1 + \varepsilon(z)$ satisfies $\|\varepsilon(z)\|_2 < 1/2k$. This is possible by Lemma 4 of [6]. Let $\{b_k\} \in \text{ball } l^2$. Take $f(z) = \sum a_n z^n \in \mathcal{C}'$ such that $\|f\| \leq k$ and $a_{n_k} = b_k$ for all k . $h(z) = f(z)g(z) = \sum c_n z^n$ satisfies: $h \in \mathcal{C}$, $\|h\| \leq k$,

$$\| \{c_{n_k} - b_k\}_k \|_2 \leq \|\varepsilon(z)\|_2 \cdot \|f(z)\|_\infty < 1/2.$$

Lemma 2 now proves $R(\mathcal{C}) = l^2$ and the proof is complete.

REFERENCES

- [1] W. G. Bade and P. C. Curtis, Jr. *Embedding theorems for commutative Banach algebras*, Pacific J. Math., **18** (1966), 391–409.
- [2] L. Carleson, *An interpolation problem for bounded analytic functions*, Amer. J. Math., **80** (1958), 921–930.
- [3] ———, *Representations of continuous functions*, Math. Z., **66** (1957), 447–451.
- [4] P. L. Duren, *Theory of H^p Spaces*, Academic Press.
- [5] J. Earl, *On the interpolation of bounded sequences by bounded functions*, J. London Math. Soc., **2** (1970), 544–548.
- [6] E. A. Heard and J. H. Wells, *An interpolation problem for subalgebras of H^∞* , Pacific J. Math., **28** No. 3 (1969), 543–553.
- [7] W. Rudin, *Trigonometric series with gaps*, J. Math. Mech., **9** (1960), 203–228.
- [8] ———, *Boundary values of continuous functions*, P.A.M.S., **7** (1956).
- [9] ———, *Functional Analysis*, McGraw-Hill Book Company.
- [10] H. S. Shapiro and A. L. Shields, *On some interpolation problems for analytic functions*, Amer. J. Math., **83** (1961), 513–532.
- [11] S. A. Vinogradov, *The Banach-Rudin-Carleson Theorems and Embedding Operators*, Seminars in Math. V. A. Steklov Math. Inst. Leningrad, Vol. **19**, 1–28. Consultants Bureau New York–London (1972).

Received February 25, 1981.

AGDER DISTRIKTSKOLE
4600 KRISTIANSAND, NORWAY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* ISSN 0030-8730 is published monthly by the *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1983 by Pacific Journal of Mathematics

Tibor Bisztriczky , On the singularities of almost-simple plane curves	257
Peter B. Borwein , On Sylvester's problem and Haar spaces	275
Emilio Bujalance , Cyclic groups of automorphisms of compact nonorientable Klein surfaces without boundary	279
Robert Jay Daverman and John J. Walsh , Acyclic decompositions of manifolds	291
Lester Eli Dubins , Bernstein-like polynomial approximation in higher dimensions	305
Allan L. Edelson and Jerry Dee Schuur , Nonoscillatory solutions of $(rx^n)^n \pm f(t, x)x = 0$	313
Akira Endô , On units of pure quartic number fields	327
Hector O. Fattorini , A note on fractional derivatives of semigroups and cosine functions	335
Ronald Fintushel and Peter Sie Pao , Circle actions on homotopy spheres with codimension 4 fixed point set	349
Stephen Michael Gagola, Jr. , Characters vanishing on all but two conjugacy classes	363
Saverio Giulini , Singular characters and their L^p norms on classical Lie groups	387
Willy Govaerts , Locally convex spaces of non-Archimedean valued continuous functions	399
Wu-Chung Hsiang and Bjørn Jahren , A remark on the isotopy classes of diffeomorphisms of lens spaces	411
Hae Soo Oh , Compact connected Lie groups acting on simply connected 4-manifolds	425
Frank Okoh and Frank A. Zorzitto , Subsystems of the polynomial system	437
Knut Øyma , An interpolation theorem for H_E^∞	457
Nikolaos S. Papageorgiou , Nonsmooth analysis on partially ordered vector spaces. II. Nonconvex case, Clarke's theory	463