Pacific Journal of Mathematics

A REMARK ON A PAPER: "COMMON FIXED POINTS OF NONEXPANSIVE MAPPINGS BY ITERATION"

GIOVANNI EMMANUELE

Vol. 110, No. 2

October 1984

A REMARK ON A PAPER OF KUHFITTIG

GIOVANNI EMMANUELE

It is observed that an iterative scheme introduced by Kuhfittig satisfies a property which allows us to obtain sufficient conditions for its strong and weak convergence to a common fixed point of a finite family of nonexpansive mappings, via other known results. In particular we generalize the results obtained by Kuhfittig.

Let *E* be a strictly convex Banach space and let *C* be a closed, convex subset of *E*. We consider a finite family of nonexpansive mappings $\{T_i: i = 1, 2, ..., k\}$, with $\bigcap_{i=1}^k \mathfrak{F}(T_i) \neq \emptyset$, where $\mathfrak{F}(T_i)$ is the fixed point set of T_i . In [5] Kuhfittig introduced the following functions U_i by putting

$$U_{\iota}(x) = [(1-\alpha)I + \alpha T_{\iota}U_{\iota-1}](x), \qquad x \in C,$$

where $\alpha \in [0, 1[, U_0 = I, I]$ identity on *E*, and he studied convergence of the sequence $\{U_k^n(x_0)\}, x_0 \in C$, to a point of $\bigcap_{i=1}^k \mathcal{F}(T_i)$.

Purpose of this note is to obtain convergence results for such a sequence.

Our results are based on the following property of $\{U_k^n(x_0)\}$: the sequence $\{x_n\}$, $x_n = U_k^n(x_0)$, $x_0 \in C$, $n \in N$, is a particular case of the following iterative procedure considered by Ishikawa (see [4]) for a nonexpansive mapping F

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n F(x_n), \qquad n \in N,$$

 $\alpha_n \in [0, b] \subset [0, 1[$. In our case $F = T_k U_{k-1}$. Then, following Ishikawa, we have

(1)
$$\lim_{n} ||x_{n} - T_{k}U_{k-1}(x_{n})|| = 0, \quad x_{0} \in C,$$

(2)
$$\lim_{n} \left\| U_{k}^{n}(x_{0}) - U_{k}^{n+1}(x_{0}) \right\| = 0, \quad x_{0} \in C.$$

Moreover, like in [5], Theorem 1, we have the following

LEMMA. Let E, C, T_i , U_i , α be as above. If $y \in C$ is such that $y = T_k U_{k-1}(y)$, then y is a common fixed point of $\{T_i: i = 1, 2, ..., k\}$.

Now, we are ready to show our theorems. First is an improvement of Theorem 1 of [5]

THEOREM 1. Let E, C, T_i , U_i , α be as above. We suppose that $j \in \{1, 2, ..., k\}$ exists for which T_j is α -condensing, i.e. $\alpha(T(A)) < \alpha(A)$, for each bounded $A \subset C$, $\alpha(A) > 0$, where α denotes the Kuratowski measure of non compactness.

Then, $\{x_n\}$ converges strongly to a point of $\bigcap_{i=1}^k \mathfrak{F}(T_i)$.

Proof. It is easy to show that $T_k U_{k-1}$ is α -condensing and so $\{x_n\}$ converges strongly to a fixed point of $T_k U_{k-1}$, by a result of [3]. The Lemma completes the proof.

THEOREM 2. Let E, C, T_i , U_i , α be as above. If E is uniformly convex and C = -C, then $\{U_k^n(x_0)\}$ converges strongly to a common fixed point of $\{T_i: i = 1, 2, ..., k\}$, if any T_i is odd.

Proof. We observe that U_k is odd and asymptotically regular, by (2). Baillon, Bruck and Reich [1] have shown that $\{U_k^n(x_0)\}$ converges strongly to a fixed point of U_k , i.e. a fixed point of $T_k U_{k-1}$, under these assumptions. Like in Theorem 1, we conclude the proof.

The following Theorem 3 generalizes a result by Kuhfittig ([5], Theorem 2) if (i) below holds.

THEOREM 3. Let E, C, T_i , U_i , α be as above. We suppose that C is boundedly weakly compact and one of the following conditions is satisfied:

(i) E satisfies Opial's condition.

(ii) E is uniformly convex and it has a Frechét differentiable norm.

Then, $\{U_k^n(x_0)\}$ converges weakly to a point of $\bigcap_{i=1}^k \mathfrak{F}(T_i)$.

Proof. If (i) is true, using (1) we show that $\{x_n\}$ converges weakly to a fixed point of $T_k U_{k-1}$, by a result of [3]. Then, the Lemma concludes the proof.

Moreover, if (ii) is satisfied, we have to use results by Bruck [2] in order to show that $\{U_k^n(x_0)\}$ converges weakly to a fixed point of U_k , i.e. a fixed point of $T_k U_{k-1}$. Using the Lemma, we have our thesis.

References

- [1] J. B. Baillon, R. E. Bruck and S. Reich, On the asymptotic behavior of nonexpansive mappings and semigroups in Banach spaces, Houston J. Math., 4, no. 1, (1978), 1–9.
- [2] R. E. Bruck, A simple proof of the mean ergodic theorem for nonlinear contractions in Banach spaces, Israel J. Math., 32, no. 2-3, (1979), 107-116.
- [3] G. Emmanuele, Convergence of the Mann-Ishikawa iterative process for nonexpansive mappings, to appear in J. Nonlinear Analysis, TMA.
- [4] S. Ishikawa, Fixed points and iteration of a nonexpansive mapping in a Banach space, Proc. Amer. Math. Soc., **59** (1976), 65–71.
- P. K. F. Kuhfittig, Common fixed points of nonexpansive mappings by iteration, Pacific J. Math., 97, (1981), 137–139.

Received April 1, 1982.

Università di Catania Viale A. Doria 6, 95125 Catania, Italy

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor) University of California Los Angeles, CA 90024

Hugo Rossi University of Utah Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS University of California Berkeley, CA 94720 J. DUGUNDJI Department of Mathematics University of Southern California Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON Stanford University Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH B. I (1906–1982)

B. H. NEUMANN F.

F. WOLF K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONAUNIVERUNIVERSITY OF BRITISH COLUMBIAUNIVERCALIFORNIA INSTITUTE OF TECHNOLOGYSTANFCUNIVERSITY OF CALIFORNIAUNIVERMONTANA STATE UNIVERSITYUNIVERUNIVERSITY OF NEVADA, RENOUNIVERNEW MEXICO STATE UNIVERSITYWASHINOREGON STATE UNIVERSITYUNIVER

UNIVERSITY OF OREGON UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF HAWAII UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION Copyright © 1984 by Pacific Journal of Mathematics

Pacific Journal of Mathematics Vol. 110, No. 2 October, 1984

Robert A. Bekes, The range of convolution operators	57
Dennis K. Burke and Sheldon Davis, Subsets of ${}^{\omega}\omega$ and generalized metric	73
Giovanni Emmanuele. A remark on a paper: "Common fixed points of	15
nonexpansive mappings by iteration"	33
I. Erdélyi and Sheng-Wang Wang, On strongly decomposable operators 28	37
Gerhard Gierz, Injective Banach lattices with strong order units	9 7
Maurizio Letizia, Quotients by complex conjugation of nonsingular	
quadrics and cubics in $\mathbf{P}_{\mathbf{C}}^3$ defined over \mathbf{R})7
P. H. Maserick and Franciszek Hugon Szafraniec, Equivalent definitions	
of positive definiteness	15
Costel Peligrad and S. Rubinstein, Maximal subalgebras of C*-crossed	
products	25
Derek W. Robinson and Sadayuki Yamamuro, Hereditary cones, order	
ideals and half-norms	35
Derek W. Robinson and Sadayuki Yamamuro, The Jordan decomposition	
and half-norms	15
Richard Rochberg, Interpolation of Banach spaces and negatively curved	
vector bundles	55
Dale Rolfsen, Rational surgery calculus: extension of Kirby's theorem	17
Walter Iaan Seaman, Helicoids of constant mean curvature and their Gauss	~ -
maps	57
Diana Shelstad, Endoscopic groups and base change C/R	¥7
Jerrold Norman Siegel and Frank Williams, Numerical invariants of	17
nomotopies into spheres	17
Alladi Sitaram, Some remarks on measures on noncompact semisimple Lie	20
groups	29
Teruhiko Soma, Atoroidal, irreducible 3-manifolds and 3-told branched	75
Lon de Viries On the C composition of and dusts	22
Jan de vries, On the G-compactification of products	+/
Hans weber, lopological Boolean rings. Decomposition of finitely additive	71
set functions	/1