

Pacific Journal of Mathematics

NUMERICAL INVARIANTS OF HOMOTOPIES INTO SPHERES

JERROLD NORMAN SIEGEL AND FRANK WILLIAMS

NUMERICAL INVARIANTS OF HOMOTOPIES INTO SPHERES

JERROLD SIEGEL AND FRANK WILLIAMS

A classical problem in the calculus of variations may be expressed: *Given two points in the same component of a Riemannian manifold M , what is the length of the shortest path connecting them?* In this paper we discuss parametric analogs to this problem. The *width* of a homotopy is the supremum of the lengths of the paths traced by the points of X . Work of Allan Calder and the first author on the topology of Stone-Cech compactifications led to the study of the question: *Given a space X and two homotopic maps $f, g: X \rightarrow M$, what is the width of the shortest homotopy between them?*

In this paper we obtain bounds $b_q(M)$ that depend only on the dimension q of X for the answer to this question in case M is a sphere or a projective space (with the standard metric). We also introduce a related sequence of invariants $B_q(M)$ and compute these numbers for spheres and projective spaces. Of particular interest is the fact that $b_{2n-2}(S^n)$ detects elements of Hopf invariant one while $B_{2n-2}(S^n)$ does not.

One may regard this question as a parametric variational problem with parameters in the space X , or, equivalently, as a brachistochrone problem in the appropriately metrized function space M^X .

Of course in general the answer to the parametric problem may be infinity: If $X = \mathbf{R}$ and $M = S^1$, the complex numbers of unit norm, then the exponential map $\exp(t) = e^{2\pi it}$ is homotopic to the constant map, but not by a homotopy of finite width. However it was proved in [2] that S^1 is the exceptional case: *If X is a finite-dimensional normal space and M is a compact Riemannian manifold with $\pi_1(M)$ finite, then any two homotopic maps f and g are connected by a homotopy of finite width.*

Of more interest is the fact that these widths are bounded with respect to the dimension of X . In the classical problem above, if M is compact then there is a bound $b_0(M)$ such that any two points in the same component are connected by a path of length $\leq b_0(M)$. (Simply take $b_0(M)$ to be the greatest of the diameters of the components of M under the Riemannian distance.) The main theorem of [3] extended this result to homotopies: *If $q \geq 0$ is an integer and $\pi_1(M)$ is finite, then there exists a bound $\tilde{b}_q(M)$ such that any two homotopic maps $f, g: X \rightarrow M$, $\dim(M) \leq q$, are homotopic via a homotopy of width $\leq \tilde{b}_q(M)$.*

Consequently there is a sequence of invariants of M defined by letting $b_q(M) = \infimum \text{ of such } \tilde{b}_q(M)$. (Clearly this $b_0(M)$ coincides with the one defined above.) It was also shown in [3] that if $\dim(M) > 0$ then $\lim_{q \rightarrow \infty} b_q(M) = \infty$, so that these invariants are non-trivial.

As a first step toward understanding these numbers, we began to calculate the $b_q(S^n)$, where S^n is the standard unit sphere, $n \geq 2$. By a simplicial approximation argument one sees that $b_q(S^n) = \pi$ for $0 \leq q < n$, and some elementary algebraic topology shows that $b_n(S^n) \geq 2\pi$, [3]. In [4] we computed all of the $b_q(S^n)$ with one exception (see Theorem 1.2 below). These numbers follow a somewhat regular pattern with exceptions when $q = n - 1$ and when $q = 2n - 2$ and $n = 2, 4$, or 8 .

In the present paper we obtain a better understanding of these invariants by introducing a new and independently interesting sequence of numbers $B_q(M)$ that do follow a regular pattern. $B_q(M)$ is the infimum of the numbers $\tilde{B}_q(M)$ such that any $H: X \times I \rightarrow M$, $\dim(X) \leq q$, can be deformed *keeping the endmaps fixed* into one of width less than $\tilde{B}_q(M)$. The proof in [3] of the finiteness of $b_q(M)$ actually shows that under the same hypotheses the numbers $B_q(M)$ are finite. (Variants of $B_q(M)$ have also been considered by M. Gromov in [5].)

Clearly $b_q(M) \leq B_q(M)$. One would expect from the definitions that the inequality would usually be strict, but for spheres that turns out to be more the exception than the rule. In this paper we compute the $B_q(S^n)$, $q \geq 0$, $n \geq 2$, then use the results of this calculation to obtain the missing value, $b_{14}(S^8)$, via the solution of a differential equation in the Cayley bundle $S^{15} \rightarrow S^8$. Comparison of these results with those of [4] yields:

0.1. THEOREM. *If $n \geq 2$, then $b_q(S^n) = B_q(S^n)$ unless $q = n - 1$ or $n = 2, 4$, or 8 and $q = 2n - 2$, in which cases $b_q(S^n) = B_{q-1}(S^n) < B_q(S^n)$.*

Since the numbers $B_q(S^n)$ are easy to compute and describe (see Table I below), it would be desirable to have a direct proof of this theorem and to understand the relationship between $b_q(M)$ and $B_q(M)$ in general. We hope to deal with this question in the future. For the present, we do observe that $b_{2n-2}(S^n)$ is sensitive to the presence of elements of Hopf invariant one in $\pi_{2n-1}(S^{n-1})$ and that $b_q(S^n) = B_q(S^n)$ except when the Hurewicz morphism $\pi_q(\Omega S^n) \rightarrow H_q(\Omega S^n)$ is a non-zero epimorphism. We conjecture:

0.2. Conjecture. *Let M be a compact Riemannian manifold with $\pi_1(M)$ finite. Then $b_q(M)$ is equal to either $B_q(M)$ or $B_{q-1}(M)$. It is equal to the*

former except when the Hurewicz morphism $\pi_q(\Omega M) \rightarrow H_q(\Omega M)$ is a non-zero epimorphism. In these cases $b_q(M) = B_{q-1}(M)$.

Further support for this conjecture comes through calculation of $B_q(P_n)$ and $b_q(P_n)$ for P_n a real, complex, or quaternionic projective. We include in §4 a computation of $B_q(CP_n)$ and $b_q(CP_n)$ as a further example of our methods. The results appear in Table II.

q	$b_q(S^n)$	$B_q(S^n)$
$0 \leq q < n - 1$	π	π
$q = n - 1$	π	2π
$n \leq q < 2n - 2$	2π	2π
$q = 2n - 2 \begin{cases} n = 2, 4, 8 \\ n \neq 2, 4, 8 \end{cases}$	2π 3π	3π 3π
$2n - 2 < q < 3n - 3$	3π	3π
$(k - 1)(n - 1) \leq q < k(n - 1), k \geq 4$	$k\pi$	$k\pi$

TABLE I

Thus, for example, since $b_{22}(S^7) = 4\pi = B_{22}(S^7)$ any homotopy from a 22-dimensional complex X to S^7 may be deformed (rel $X \times \{0, 1\}$) into a homotopy of width arbitrarily close to 4π . Furthermore, there exists a 22-complex X and maps $f, g: X \rightarrow S^7$ such that f and g are homotopic but not by a homotopy of width less than 4π .

q	$b_q(CP_n)$	$B_q(CP_n)$
0	$\pi/2$	$\pi/2$
1	$\pi/2$	π
$2 \leq q \leq 2n - 1$	π	π
$2n$	π	$3\pi/2$
$2(k - 1)n < q < 2kn, k \geq 2$	$k\pi$	$k\pi$
$2kn, k \geq 2$	$(2k + 1)\pi/2$	$(2k + 1)\pi/2$

TABLE II

1. Definitions and statement of results. Throughout this paper the domain space X will be a finite-dimensional polyhedron. By standard “bridge” techniques [9] our results will also hold if X is a finite-dimensional normal space.

1.1. DEFINITIONS. Let (M, d) be a metric space.

(a) The *length* $|\sigma|$ of a path $\sigma: I \rightarrow M$ is given by the formula

$$|\sigma| = \sup \left\{ \sum_{i=1}^k d(\sigma(t_i), \sigma(t_{i-1})) \right\}$$

taken over all partitions $0 = t_0 < t_1 < \cdots < t_k = 1$, for all k .

(b) The *width* $|H|$ of a homotopy $H: X \times I \rightarrow M$ is given by the formula $|H| = \sup\{|H_x| : x \in X\}$.

For maps $f, g: X \rightarrow M$ and a homotopy H from f to g , we consider the two extended real numbers:

(c) $W(H) = \inf\{|H'| : H' \sim H \text{ rel } X \times \{0, 1\}\}$.

(d) $w(f, g) = \inf\{|H'| : H'_0 = f, H'_1 = g\}$.

The subjects of this paper are the following sequences:

(e) $B_q(M) = \sup\{W(H) : H: X \times I \rightarrow M, \dim(X) \leq q\}$.

(f) $b_q(M) = \sup\{w(f, g) : f \sim g: X \rightarrow M, \dim(X) \leq q\}$.

Let S^n denote the unit sphere in R^{n+1} . If $x, x' \in S^n$, let $\text{dist}(x, x')$ be the length of a minimum geodesic from x to x' . For the numbers $b_q(S^n)$, $n \geq 2$, we have the following result.

1.2. THEOREM [4]. *If (n, q) is a pair of integers such that $n \geq 2$, $q \geq 0$, and (n, q) is not $(2, 2)$, $(4, 6)$, or $(8, 14)$, then*

$$b_q(S^n) = \begin{cases} \pi, & 0 \leq q \leq n-1, \\ 2\pi, & n \leq q < 2n-2, \\ k\pi, & (k-1)(n-1) \leq q < k(n-1), k \geq 3. \end{cases}$$

Moreover, $b_2(S^2) = 2\pi = b_6(S^4)$ and $2\pi \leq b_{14}(S^8) \leq 3\pi$.

In §3 we shall show that:

1.3. THEOREM. *The value of $b_{14}(S^8)$ is 2π .*

The numbers $B_q(S^n)$ are relatively easy to compute. The following theorem will be proved in §2.

1.4. THEOREM. *If n, q , and k are integers such that $n \geq 2$, $k \geq 1$, and $(k-1)(n-1) \leq q < k(n-1)$, then $B_q(S^n) = k\pi$.*

We shall use the following kind of approximate fibration in the sequel:

1.5. DEFINITION. Let (M, d) be a metric space. A map $p: E \rightarrow M$ has the *approximate covering homotopy property through dimension k* , $\text{XCHP}(k)$, provided that if X is a normal space of dimension $\leq k$, and $G: X \times I \rightarrow M$ and $g: X \rightarrow E$ are such that $pg(x) = G(x, 0)$, then for any $\varepsilon > 0$ there exists $H: X \times I \rightarrow E$ such that $H(x, 0) = g(x)$ and $d(pH(x, t), G(x, t)) < \varepsilon$ for all $(x, t) \in X \times I$.

Let $E = \{\sigma: I \rightarrow S^n: \sigma \text{ is piecewise } C^\infty \text{ with constant speed}\}$. We topologize E as in [8, p. 88]. For $0 < c \leq \infty$ we define $E_c = \{\sigma \in E: |\sigma| < c\}$ and let $p_c: E_c \rightarrow S^n \times S^n$ be given by $p_c(\sigma) = (\sigma(0), \sigma(1))$.

1.6. THEOREM [4, 1.4]. *If $k \geq 0$ is an integer and $c > k\pi$ then $p_c: E_c \rightarrow S^n \times S^n$ has the $\text{XCHP}(k(n-1)-1)$.*

2. Proof of 1.4. The following theorem will be used in this section to obtain the upper bounds for $B_q(S^n)$ and in §3 to compute $b_{14}(S^8)$.

2.1. THEOREM. *Let X be a complex of dimension less than $k(n-1)$, where $k \geq 1$ and $n \geq 2$. Suppose $H: X \times I \rightarrow S^n$ is a homotopy and $\varepsilon > 0$. Then there exists $\bar{H}: X \times I \rightarrow S^n$ such that $\bar{H} \sim H(\text{rel } X \times \{0, 1\})$ and $|\bar{H}| < k\pi + \varepsilon$. Furthermore \bar{H} may be chosen so that \bar{H}_x is piecewise C^∞ of constant speed for all $x \in X$.*

Proof. Choose $\delta, \tau > 0$ such that $\rho < \pi$ and $\delta + 2\rho < \varepsilon$. Define $G: X \times I \rightarrow S^n \times S^n$ by $G(x, t) = (H(x, 0), H(x, t))$. Let $g: X \rightarrow E_{k\pi+\delta}$ be given by $g(x) = C[H(x, 0)]$, the constant path at $H(x, 0)$. By 1.6 there exists $\bar{G}: X \times I \rightarrow E_{k\pi+\delta}$ such that $\bar{G}(x, 0) = g(x)$ and $\text{dist}[(H(x, 0), H(x, t)), p\bar{G}(x, t)] < \rho$. Since $\rho < \pi$, any two points $y, y' \in S^n$ such that $\text{dist}(y, y') < \rho$ may be joined by a unique minimal geodesic $m(y, y')$. Thus we may define $\hat{G}: X \times I \times I \rightarrow S^n$ by the formula

$$\hat{G}(x, t, s) = \begin{cases} m(H(x, 0), \bar{G}(x, t)[0])[3s], & 0 \leq s \leq \frac{1}{3}, \\ \bar{G}(x, t)[3s-1], & \frac{1}{3} \leq s \leq \frac{2}{3}, \\ m(\bar{G}(x, t)[1], H(x, t))[3s-2], & \frac{2}{3} \leq s \leq 1. \end{cases}$$

Note that for each $(x, t) \in X \times I$, $\hat{G}_{(x,t)}$ is a piecewise C^∞ path from $H(x, 0)$ to $H(x, t)$ of length less than $k\pi + \delta + 2\rho$, which is less than

$k\pi + \varepsilon$. Let \tilde{G} be \hat{G} reparametrized so as to have constant speed in the s parameter for each $(x, t) \in X \times I$.

We define $\bar{H}(x, s)$ to be $\tilde{G}(x, 1, s)$. A homotopy K from H to \bar{H} is given by

$$K(x, t, r) = \begin{cases} \tilde{G}(x, t, 1 - 2r(1 - t)), & 0 \leq r \leq \frac{1}{2}, \\ \tilde{G}(x, t + (2r - 1)(1 - t), t), & \frac{1}{2} \leq r \leq 1. \end{cases}$$

It is immediate that K stays fixed on $X \times \{0, 1\}$. \square

If we denote the dimension of X by q , we see that a consequence of 2.1 is that $q < k(n - 1)$ implies $B_q(S^n) \leq k\pi$.

The proof of 1.4 will be completed by showing that $B_q(M) \geq k\pi$ if $q \geq (k - 1)(n - 1)$. By 1.2, unless $k = 2$, or $n = 2, 4$, or 8 and $k = 3$, we have $b_q(S^n) \geq k\pi$, whence $B_q(S^n) \geq k\pi$. Although we could dispose of the remaining numbers on a case-by-case basis, the following theorem gives us an easy proof of the desired inequality for all values of q and n at once, independent of any appeal to 1.2.

2.2. THEOREM. *For all $n \geq 2$, $k \geq 1$, if $q \geq (k - 1)(n - 1)$, then $B_q(S^n) \geq k\pi$.*

Proof. Let $(S^{n-1})_r$ denote the r th James reduced product space of S^{n-1} , [6], and let $E_\lambda(p, q)$ be the subspace of E_λ consisting of paths from p to q , for $p, q \in S^n$. Let k be odd. In [4, 1.5] we noted that if $\lambda \leq k\pi$ then $E_\lambda(p, -p) \sim (S^{n-1})_r$ for some $r < k - 1$. Let $X = (S^{n-1})_{k-1}$ and let $\eta: (S^{n-1})_{k-1} \rightarrow E(p, -p)$ be any map that is essential on homology in dimension $(k - 1)(n - 1)$. We consider η as a homotopy between the constant maps of X to p and $-p$, respectively. If $\lambda = W(\eta) \leq k\pi$, then η would factor up to homotopy through $E_\lambda(p, -p)$, contradicting the fact that $H_{(k-1)(n-1)}(E_\lambda(p, -p)) = 0$. Thus $W(\eta) > k\pi$ and hence $B_{(k-1)(n-1)}(S^n) \geq k\pi$.

If k is even we apply a similar argument to $E_\lambda(p, p)$. \square

3. The Cayley bundle. In this section we show that $b_{14}(S^8) = 2\pi$. Our proof involves a detailed study of the geometry of the Hopf-Cayley bundle. In fact our method also works for the Hopf complex and quaternionic bundles.

3.1. Notation. Let K and R denote the Cayley numbers and the real numbers. Let S^{15} and S^8 be the unit spheres in $R^{16} = K \times K$ and

$R^9 = K \times R$. The Cayley bundle is given by the map $h: S^{15} \rightarrow S^8$, where

$$h(c, d) = (2c\bar{d}, |c|^2 - |d|^2).$$

We use juxtaposition to denote Cayley multiplication and “bar” to denote conjugation of Cayley numbers. We shall also use the vector dot product $c \cdot d$.

3.2. THEOREM. *Suppose $F: X \times I \rightarrow S^8$ is a piecewise C^∞ homotopy. Let $g: X \rightarrow S^{15}$ be such that $hg = F_0$. There exists $G: X \times I \rightarrow S^{15}$ such that $G_0 = g$, $hG = F$; and for each x , G_x is piecewise C^∞ and $|G_x| = \frac{1}{2} |F_x|$ on common points of smoothness. Thus*

$$|G| = \frac{1}{2} |F|.$$

Proof. Our proof amounts to showing that h is a Riemannian submersion, [10]. This would be easy if h were a principal bundle. The difficulty arises from the non-associativity of the Cayley numbers. We represent points of S^{15} as $(u \cos \theta, v \sin \theta)$, $u, v \in S^7$. The points of S^8 are represented as $(w \sin \Phi, \cos \Phi)$, $w \in S^7$. With respect to this representation h becomes

$$h(u \cos \theta, v \sin \theta) = (u\bar{v} \sin 2\theta, \cos 2\theta).$$

For each point $x \in X$ we may write

$$F(x, t) = (u(x, t) \sin[\psi(x, t)], \cos[\psi(x, t)]).$$

Locally

$$g(x) = (u(x, 0)v(x, 0) \cos[\frac{1}{2}\psi(x, 0)], v(x, 0) \sin[\frac{1}{2}\psi(x, 0)]).$$

Moreover, any path covering F is locally of the form

$$G(x, t) = (uv \cos(\frac{1}{2}\psi), v \sin(\frac{1}{2}\psi))$$

where $v(x, t)$ is a path in S^7 starting at $v(x, 0)$. Thus to find a $g(x, t)$ that satisfies the requirements of 3.2 amounts to finding a suitable $v(x, t)$ such that

$$(3.3) \quad |\dot{G}(x, t)| = \frac{1}{2} |\dot{F}(x, t)|.$$

Firstly $\dot{F} = (u\dot{\psi} \cos(\psi) + \dot{u} \sin(\psi), -\dot{\psi} \sin(\psi))$. Using the fact that $u \cdot \dot{u} = 0$ (since $|u| = 1$), we have

$$|\dot{F}|^2 = \frac{1}{4} (\dot{\psi})^2 + |\dot{u}|^2 \sin^2 \psi.$$

Similarly,

$$|\dot{G}|^2 = \frac{1}{4}(\dot{\psi})^2 + |\dot{u}|^2 \cos^2(\psi/2) + 2(\dot{u}v) \cdot (u\dot{v}) \cos^2(\psi/2) + |\dot{v}|^2.$$

By combining these last two relations, we see that 3.3 is equivalent to

$$(3.4) \quad |\dot{u}|^2 \cos^2(\psi/2) + 2(\dot{u}v) \cdot (u\dot{v}) \cos^2(\psi/2) + |\dot{v}|^2 = \frac{1}{4} \sin^2(\psi).$$

Using the identity $\cos^2(\psi/2) - \frac{1}{2} \sin^2(\psi) = \cos^4(\psi/2)$, we reduce 3.4 to $|u\dot{v} + \dot{u}v \cos^2(\psi/2)|^2 = 0$. Hence it suffices to find v such that

$$(3.5) \quad u\dot{v} + \cos^2(\psi/2)\dot{u}v = 0.$$

Since multiplication by u is invertible as a linear operator, 3.5 may be considered to be a differential equation of the form $\dot{v} + Av = 0$, where A is a continuous family of linear operators. It is a standard result that such a system with continuous initial conditions $v(x, 0)$ has a solution $v(x, t)$ that is continuous in x . (See for example [1; Theorem 3.8].)

Finally we observe that dotting 3.5 with uv yields $v \cdot \dot{v} = 0$ so $v(x, t) \in S^7$. For this last observation we use the relation $(a \cdot b)|c|^2 = ac \cdot bc$. \square

3.6. Notation. In order to define our homotopies it will be convenient to have a radial coordinate for our standard 14-cell. To this end let S^{13} be a 13-sphere and D^{14} be the cone on S^{13} , $D^{14} = (I \times S^{13})/(\{0\} \times S^{13})$. Write $x = s \wedge y$ for points of D^{14} , where $(s, y) \in I \times S^{13}$.

3.7. THEOREM. *Let $F: D^{14} \times I \rightarrow S^8$ be such that $F|S^{13} \times I$ is piecewise C^∞ and of width less than 2π . Let $\varepsilon > 0$. Then there exists $\bar{F}: D^{14} \times I \rightarrow S^8$ such that*

- (a) $|\bar{F}| < 2\pi$,
- (b) $\bar{F} = F$ on $S^{13} \times I$,
- (c) $\text{dist}(\bar{F}(x, i), F(x, i)) < \varepsilon$, $x \in D^{14}$, $i = 0, 1$.

Proof. Since D^{14} is contractible, $F|D^{14} \times \{0\}$ lifts to $G: D^{14} \times \{0\} \rightarrow S^{15}$. By 3.2 we may extend G to a lift of $F|((S^{13} \times I) \cup (D^{14} \times \{0\}))$ such that $|G_y| = \frac{1}{2}|F_y| < \pi$ for every $y \in S^{13}$. Now apply the covering homotopy extension property to lift F to all of $D^{14} \times I$.

If $y \in S^{13}$ then $G(y, 0) \neq -G(y, 1)$. Thus by dimensionality we may deform G to \tilde{G} so that $G = \tilde{G}$ on $S^{13} \times I$, $\text{dist}(G(x, i), \tilde{G}(x, i)) < \varepsilon/4$ for all $(x, i) \in D^{14} \times \{0, 1\}$ and $\tilde{G}(x, 0) \neq -\tilde{G}(x, 1)$ for all x .

Let $0 < s_0 < 1$ be small enough so that

- (a) $\text{dist}(\tilde{G}(s \wedge y, i), \tilde{G}(1 \wedge y, i)) < \varepsilon/4$; $y \in S^{13}$, $s_0 \leq s \leq 1$, $i = 0, 1$,

and

(b) $\text{dist}(\tilde{G}(s \wedge y, i), \tilde{G}(s/s_0 \wedge y, i)) < \varepsilon/4$; $y \in S^{13}$, $0 \leq s \leq s_0$, $i = 0, 1$.

Define

$$\hat{G}(s \wedge y, i) = \begin{cases} \tilde{G}(1 \wedge y, i), & s_0 \leq s \leq 1, \\ \tilde{G}(s/s_0 \wedge y, i), & 0 \leq s \leq s_0. \end{cases}$$

For $s \wedge y \in D^{14}$ and $t \in I$, let

$$\bar{G}(s \wedge y, t)$$

$$= \begin{cases} m(\hat{G}(s \wedge y, 0), \hat{G}(s \wedge y, 1))[t], & 0 \leq s \leq s_0, 0 \leq t \leq 1, \\ m\left(G(1 \wedge y, 0), G\left(1 \wedge y, \frac{1-s}{1-s_0}\right)\right)\left[\frac{1-s_0}{1-s}t\right], & s_0 \leq s \leq 1, 0 \leq t \leq \frac{1-s}{1-s_0}, \\ G(1 \wedge y, t), & s_0 \leq s \leq 1, \frac{1-s}{1-s_0} \leq t \leq 1. \end{cases}$$




FIGURE 1

Note that $\bar{G} = G$ on $S^{13} \times I$ and $\text{dist}(\bar{G}(x, i), G(x, i)) < \varepsilon/2$ for $x \in D^{14}$, $i = 0, 1$. Moreover if $0 \leq s \leq s_0$, then

$$|\bar{G}_{s \wedge y}| = \text{dist}(\hat{G}(s \wedge y, 0), \hat{G}(s \wedge y, 1)) < \pi.$$

Finally, if $s_0 \leq s \leq 1$, then

$$\begin{aligned} |\bar{G}_{s \wedge y}| &= \left| m \left(G(1 \wedge y, 0), G \left(1 \wedge y, \frac{1-s}{1-s_0} \right) \right) \right| + \left| G_{1 \wedge y} \left[\frac{1-s}{1-s_0}, 1 \right] \right| \\ &\leq \left| G_{1 \wedge y} \left[0, \frac{1-s}{1-s_0} \right] \right| + \left| G_{1 \wedge y} \left[\frac{1-s}{1-s_0}, 1 \right] \right| = |G_{1 \wedge y}| < \pi, \end{aligned}$$

where $G_{1 \wedge y}[s_1, s_2]$ denotes the restriction of the path $G_{1 \wedge y}$ to the subinterval $[s_1, s_2]$.

Set $\bar{F} = h\bar{G}$. Since the dilatation of h is 2, $|\bar{F}| < 2\pi$ and $\text{dist}(F(x, i), \bar{F}(x, i)) < \varepsilon, i = 0, 1$. By definition, $F = \bar{F}$ on $S^{13} \times I$. \square

3.8. THEOREM. *Let X be a 14-dimensional complex, $f, f': X \rightarrow S^8$ be homotopic maps, and $\varepsilon > 0$. Then f and f' are homotopic by a homotopy of width less than $2\pi + \varepsilon$. Hence $b_{14}(S^8) \leq 2\pi$.*

Proof. Let $F: X \times I \rightarrow S^8$ be a homotopy of f to f' and let $0 < \varepsilon_1 < \varepsilon/8$. If we let $X^{(13)}$ denote the 13-skeleton of X , we may apply 3.1 to obtain $\bar{F}: X^{(13)} \times I \rightarrow S^8$ such that \bar{F} is piecewise C^∞ of constant speed, of width less than $2\pi + \varepsilon_1$, and homotopic to $F(\text{rel } X^{(13)} \times \{0, 1\})$. We now use the homotopy extension property to extend \bar{F} to a homotopy from f to f' .

We define $G: X^{(13)} \times I \rightarrow S^8$ by

$$G(x, t) = \bar{F} \left(x, \frac{2\pi - \varepsilon_1}{2\pi + \varepsilon_1} t \right).$$

Observe that $|G| \leq 2\pi - \varepsilon_1 < 2\pi$ and $\text{dist}(G(x, 1), f'(x)) < \varepsilon/4$. By a patching process similar to that used in the proof of 3.7 we extend G to $K \times I$ such that $G(x, 0) = f(x)$ and $\text{dist}(g(x, 1), f'(x)) < \varepsilon/4$ for all $x \in X$.

We next apply 3.7 to G on the individual 14-cells of X , using $\varepsilon/4$ in place of ε . Since 3.7 leaves G unchanged on the boundary of each 14-cell, the resulting homotopy $\bar{G}: X \times I \rightarrow S^8$ is such that $\bar{G} = G$ on $X^{(13)} \times I$, $|\bar{G}| \leq 2\pi$, and $\text{dist}(\bar{G}(x, i)) < \varepsilon/4$, for all $x \in X$ and $i = 0, 1$.

Now $\text{dist}(\bar{G}(x, 0), f(x)) < \varepsilon/4$ and

$$\begin{aligned} \text{dist}(\bar{G}(x, 1), f'(x)) &\leq \text{dist}(\bar{G}(x, 1), G(x, 1)) + \text{dist}(G(x, 1), f'(x)) \\ &< \varepsilon/4 + \varepsilon/4 = \varepsilon/2. \end{aligned}$$

If we compose \bar{G} with the minimal geodesics on each end from $f(x)$ to $\bar{G}(x, 0)$ and from $\bar{G}(x, 1)$ to $g(x)$, as in the proof of 3.1, we obtain a homotopy \hat{G} from f to f' such that $|G| \leq 2\pi + 3\varepsilon/4 < 2\pi + \varepsilon$. \square

Since 1.2 tells us that $b_{14}(S^8) \geq 2\pi$, the proof of 1.3 is accomplished.

4. Projective spaces. In this section we shall compute the invariants for complex projective spaces (with the metric induced from the Hopf map $S^{2n+1} \rightarrow \mathbf{C}P_n$). We shall proceed as for spheres, in that we shall calculate upper and lower bounds for $B_q(\mathbf{C}P_n)$ and lower bounds for $b_q(\mathbf{C}P_n)$. When these do not coincide we shall use special arguments.

The geodesic structure of $\mathbf{C}P_n$ is well known, see e.g. [7]. Let E be the space of piecewise C^∞ paths of constant speed and $E \rightarrow \mathbf{C}P_n \times \mathbf{C}P_n$ be the projection, as in the paragraph preceding 1.6. Since $\Omega \mathbf{C}P_n \sim S^1 \times \Omega S^{2n+1}$, it follows that for $p, q \in \mathbf{C}P_n$, $E_\lambda(p, q)$ is homotopic to a skeleton $K^{(m)}$ of $K = S^1 \times (S^{2n})_\infty$. Indeed, if $\lambda \leq k\pi$, then $E_\lambda(p, p) \sim K^{(m)}$ for some $m \leq 2(k-1)n$. Furthermore, regarding p as a complex line in \mathbf{C}^n , if p' is an orthogonal line and $\lambda \leq (2k+1)\pi/2$, then $E_\lambda(p, p') \sim K^{(m)}$ for some $m \leq 2kn-1$. Thus, as in 2.2, we have:

4.1. THEOREM. (a) *If $q \geq 2(k-1)n+1$, then $B_q(\mathbf{C}P_n) \geq k\pi$.*

(b) *If $q \geq 2kn$, then $B_q(\mathbf{C}P_n) \geq (2k+1)\pi/2$.* □

Moving to upper bounds on $B_q(\mathbf{C}P_n)$, we have from Morse theory:

4.2. (a) *If $\lambda > k\pi$, then for all p, p' , $E_\lambda(p, p') \sim K^{(m)}$ for some $m \geq 2kn-1$.*

(b) *If $\lambda > (2k+1)\pi/2$, then for all p, p' , $E_\lambda(p, p') \sim K^{(m)}$ for some $m \geq 2kn$.*

Thus in these cases, as in [4, 1.4], $E_\lambda \rightarrow \mathbf{C}P_n \times \mathbf{C}P_n$ has the XCHP(m) for the appropriate m . Hence:

4.3. THEOREM. (a) *If $q \leq 2kn-1$, then $B_q(\mathbf{C}P_n) \leq k\pi$.*

(b) *If $q \leq 2kn$, then $B_q(\mathbf{C}P_n) \leq (2k+1)\pi/2$.*

Combining 4.1 and 4.3, we obtain the values for $B_q(\mathbf{C}P_n)$ listed in Table II.

The lower bounds for $b_q(\mathbf{C}P_n)$ are somewhat more complicated. We content ourselves here by referring to [4, pp. 27–289], since the method of computation used there applies in our present situation. The results here are somewhat simpler since for all n , the Hurewicz morphism $\pi_r(\Omega \mathbf{C}P_n) \rightarrow H_r(\Omega \mathbf{C}P_n)$ is zero except when $r = 1$ or $2n$. We obtain:

4.4. THEOREM. *If $q \neq 1$ or $2n$, then $b_q(\mathbf{C}P_n) = B_q(\mathbf{C}P_n)$.* □

If $q = 1$, the fact that $\pi_1(\mathbf{C}P_n) = 0$ yields

$$b_1(\mathbf{C}P_n) = \text{diam}(\mathbf{C}P_n) = \pi/2.$$

If $q = 2n$, we consider the bundle $S^1 \rightarrow S^{2n+1} \rightarrow \mathbf{C}P_n$. Since this bundle is principal and S^1 operates by isometries, the argument of [4, Theorem 2.3] shows that $b_{2n}(\mathbf{C}P_n) \leq b_{2n}(S^{2n+1}) = \pi$. Hence:

4.5. THEOREM. $b_{2n}(\mathbf{C}P_n) = \pi$. □

We conclude by observing that the same method also yields the values of b_q and B_q for the real and quaternionic projective spaces. In all these examples, as in the case of spheres, we are able to compute the values of B_q since the upper and lower bounds we obtain for B_q coincide with each other. A goal for future work is a characterization of those spaces for which this phenomenon occurs and, if possible, to discover the relationship between b_q , B_q , and the topology of the space of loops on the manifold.

REFERENCES

- [1] Brauer and Nohel, *Qualitative Theory of Differential Equations*, Benjamin 1969.
- [2] A. Calder and J. Siegel, *Homotopy and uniform homotopy*, Trans. Amer. Math. Soc., **235** (1978), 245–270.
- [3] ———, *On the width of homotopies*, Topology **19** (1980), 209–220.
- [4] A. Calder, J. Siegel and F. Williams, *The width of homotopies into spheres*, Topology, **21** (1982), 281–290.
- [5] M. Gromov, *Homotopical effects of dilatation*, J. Differential Geom., **13** (1978), 303–310.
- [6] I. M. James, *Reduced product spaces*, Ann. Math., **62** (2) (1955), 170–197.
- [7] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Vol. 2, Interscience, New York, 1969.
- [8] J. Milnor, *Morse Theory*, Annals of Mathematics Studies Number 51, Princeton University, 1963.
- [9] K. Morita, *Cech cohomology and covering dimension for topological spaces*, Adv. in Math., **15** (1975), 198–206.
- [10] B. O'Neill, *The fundamental equations of a submersion*, Michigan Math. J., **13** (1966), 459–469.

Received December 7, 1982.

UNIVERSITY OF MISSOURI
ST. LOUIS, MO 63121

AND

NEW MEXICO STATE UNIVERSITY
LAS CRUCES, NM 88003

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)

University of California
Los Angeles, CA 90024

HUGO ROSSI

University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS

University of California
Berkeley, CA 94720

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON

Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

Robert A. Bekes, <i>The range of convolution operators</i>	257
Dennis K. Burke and Sheldon Davis, <i>Subsets of ${}^\omega\omega$ and generalized metric spaces</i>	273
Giovanni Emmanuele, <i>A remark on a paper: "Common fixed points of nonexpansive mappings by iteration"</i>	283
I. Erdélyi and Sheng-Wang Wang, <i>On strongly decomposable operators</i> ...	287
Gerhard Gierz, <i>Injective Banach lattices with strong order units</i>	297
Maurizio Letizia, <i>Quotients by complex conjugation of nonsingular quadrics and cubics in \mathbf{P}_C^3 defined over \mathbf{R}</i>	307
P. H. Maserick and Franciszek Hugon Szafraniec, <i>Equivalent definitions of positive definiteness</i>	315
Costel Peligrad and S. Rubinstein, <i>Maximal subalgebras of C^*-crossed products</i>	325
Derek W. Robinson and Sadayuki Yamamuro, <i>Hereditary cones, order ideals and half-norms</i>	335
Derek W. Robinson and Sadayuki Yamamuro, <i>The Jordan decomposition and half-norms</i>	345
Richard Rochberg, <i>Interpolation of Banach spaces and negatively curved vector bundles</i>	355
Dale Rolfsen, <i>Rational surgery calculus: extension of Kirby's theorem</i>	377
Walter Iaan Seaman, <i>Helicoids of constant mean curvature and their Gauss maps</i>	387
Diana Shelstad, <i>Endoscopic groups and base change \mathbf{C}/\mathbf{R}</i>	397
Jerrold Norman Siegel and Frank Williams, <i>Numerical invariants of homotopies into spheres</i>	417
Alladi Sitaram, <i>Some remarks on measures on noncompact semisimple Lie groups</i>	429
Teruhiko Soma, <i>Atoroidal, irreducible 3-manifolds and 3-fold branched coverings of S^3</i>	435
Jan de Vries, <i>On the G-compactification of products</i>	447
Hans Weber, <i>Topological Boolean rings. Decomposition of finitely additive set functions</i>	471