

Pacific Journal of Mathematics

**ON THE REVERSE WEAK TYPE INEQUALITY FOR THE
HARDY MAXIMAL FUNCTION AND THE WEIGHTED
CLASSES $L(\log L)^k$**

KENNETH F. ANDERSEN AND WO-SANG YOUNG

ON THE REVERSE WEAK TYPE INEQUALITY FOR THE HARDY MAXIMAL FUNCTION AND THE WEIGHTED CLASSES $L(\log L)^k$

KENNETH F. ANDERSEN AND WO-SANG YOUNG

Muckenhoupt has given a necessary and sufficient condition to be satisfied by the weight functions U and V in order that the Hardy-Littlewood maximal function Mf should satisfy a weighted weak type $(1, 1)$ inequality. In this note, conditions on the weight functions U and V are given in order that the sense of this inequality may be reversed. This is then applied to give conditions which ensure that the integrability of Mf with respect to a weight implies that f belongs to a weighted Zygmund class $L \log L$, thus extending a result of Stein. Analogous results related to the strong maximal function and the classes $L(\log L)^k$ are also given. These extend certain results of Fava, Gatto and Gutiérrez.

If f is locally integrable on R^n , the Hardy-Littlewood maximal function Mf is defined by

$$(Mf)(x) = \sup |Q|^{-1} \int_Q |f|$$

where the supremum is taken over all cubes Q containing x . Here and henceforth, by "cube" we shall always mean "cube with sides parallel to the co-ordinate axis". As usual, $|E|$ denotes the Lebesgue measure of the measurable set E , and more generally if $U(x) \geq 0$ is defined on E we write $|E|_U = \int_E U(x) dx$. If Q is a given cube, RQ denotes the cube concentric with Q but with side R times as long. Q_0 will denote a fixed but arbitrary cube in R^n . Our first result is the following theorem.

THEOREM 1. *Suppose the non-negative weight functions U and V are defined on Q_0 . If there is a constant C depending only on U and V such that*

$$(1) \quad \frac{|Q|_U}{|Q|} \geq C \operatorname{ess\,sup}_{x \in Q} V(x) \quad \text{for all cubes } Q \subset Q_0,$$

then

$$(2) \quad |\{x \in Q_0 : (Mf)(x) > \lambda\}|_U \geq C 2^{-n} \lambda^{-1} \int_{\{x: f(x) > \lambda\}} f(x) V(x) dx$$

holds for all non-negative f supported on Q_0 with $\int_{Q_0} f < \infty$ and all $\lambda \geq \lambda_0 = |Q_0|^{-1} \int_{Q_0} f$. Conversely, if (2) holds for some constant C independent of all f which are characteristic functions of measurable sets $E \subset Q_0$ with $|E| > 0$ and for all $\lambda \geq \lambda_0 = |Q_0|^{-1} |E|$, then

$$(3) \quad \frac{|(2Q) \cap Q_0|_U}{|Q|} \geq C 4^{-n} \operatorname{ess\,sup}_{x \in Q} V(x)$$

for all cubes $Q \subset Q_0$.

COROLLARY 1. Let $U(x) \geq 0$, $V(x) \geq 0$ be defined on R^n . If there is a constant C depending only on U , V such that

$$(4) \quad \frac{|Q|_U}{|Q|} \geq C \operatorname{ess\,sup}_{x \in Q} V(x) \quad \text{for all cubes } Q \subset R^n$$

then

$$(5) \quad |\{x \in R^n: (Mf)(x) > \lambda\}|_U \geq C 2^{-n} \lambda^{-1} \int_{\{x: f(x) > \lambda\}} f(x) V(x) dx$$

holds for all non-negative f and all $\lambda > 0$. Conversely, if (5) holds for all f which are characteristic functions of measurable sets $E \subset R^n$ with $0 < |E| < \infty$ and all $\lambda > 0$, then

$$(6) \quad \frac{|2Q|_U}{|Q|} \geq C 4^{-n} \operatorname{ess\,sup}_{x \in Q} V(x) \quad \text{for all cubes } Q \subset R^n.$$

If U satisfies the doubling condition

$$|2Q|_U \leq C |Q|_U \quad \text{for all cubes } Q$$

then (6) shows that (4) is both necessary and sufficient for (5). Observe that the example $U(x) = 1$ on $R^n - Q_0$, $U(x) = \infty$ otherwise, satisfies (4) with $U = V$ but does not satisfy the doubling condition. On the other hand, if U is locally integrable and satisfies (4) with $U = V$ then U necessarily satisfies the doubling condition. To see this, note that (4) is equivalent to

$$\frac{|Q|_U}{|Q|} \geq C \frac{|E|_V}{|E|} \quad \text{for all measurable } E \subset Q \text{ with } |E| > 0$$

so that with $U = V$ it follows that U satisfies the A_∞ condition of Muckenhoupt [2] and hence also the doubling condition.

The inequality (5) may be viewed as a reverse of the weak type (1, 1) inequality

$$|\{x \in R^n: (Mf)(x) > \lambda\}|_U \leq C\lambda^{-1} \int_{\{x: f(x) > \lambda/2\}} f(x)V(x) dx$$

which holds if and only if U, V satisfy the A_1 condition of Muckenhoupt [3], namely, $|Q|_U/|Q| \leq C \operatorname{ess\,inf}_{x \in Q} V(x)$ for all cubes Q . Restricting to the case $U = V$, the A_1 condition implies that U satisfies the doubling condition, and hence the weak type inequality and its reverse both hold in this case if and only if there are positive constants c_1 and c_2 such that $c_1 \leq U(x) \leq c_2$ for almost all $x \in R^n$.

If $U(x)^{-1}$ satisfies the A_1 condition, Hölder's inequality shows that (4) holds with $U = V$. The functions $U(x) = V(x) = |x|^\alpha$, $\alpha \geq 0$, provide further examples that satisfy (4).

Theorem 1 may be used to prove the following result.

THEOREM 2. *Suppose $f(x) \geq 0$ is supported on Q_0 and that there is a constant $C > 0$ such that the weight functions $U(x) \geq 0$, $V(x) \geq 0$ satisfy*

$$(7) \quad \frac{|Q|_U}{|Q|} \geq C \operatorname{ess\,sup}_{x \in Q} V(x) \quad \text{for all cubes } Q \subset Q_0$$

Then

$$\int_{Q_0} (Mf)(x)U(x) dx < \infty$$

implies

$$\int_{Q_0} [f(x) \log^+ f(x)]V(x) dx < \infty.$$

Corollary 1 and Theorem 2 generalize some results of Stein [4] who considered the unweighted case, $U(x) \equiv V(x) \equiv 1$.

Let $1 \leq i \leq n$ and let M_i denote the Hardy-Littlewood maximal function in the i th variable, that is

$$(M_i f)(x) = \operatorname{ess\,sup}_{a, b > 0} (a + b)^{-1} \int_{-a}^b |f(x_1, \dots, x_{i-1}, x_i + t, x_{i+1}, \dots, x_n)| dt$$

where $x = (x_1, \dots, x_n)$. The following generalize certain results of Fava, Gatto and Gutiérrez [1] who considered the unweighted case, $U(x) \equiv 1$.

THEOREM 3. *Let $U(x) \geq 0$ be defined on R^n and let k be a fixed integer, $1 \leq k \leq n$. If there is a constant C depending only on U and k such that for each i , $1 \leq i \leq k$,*

$$(8) \quad (a+b)^{-1} \int_{-a}^b U(x_1, \dots, x_{i-1}, x_i + t, x_{i+1}, \dots, x_n) dt \\ \geq C \operatorname{ess\,sup}_{-a < t < b} U(x_1, \dots, x_{i-1}, x_i + t, x_{i+1}, \dots, x_n)$$

for all $a, b > 0$ and almost all $x \in R^n$, then

$$(9) \quad |\{x \in R^n: (M_k \cdots M_1 f)(x) > \lambda\}|_U \\ \geq \frac{2^{-k} C^k}{(k-1)! \lambda} \int_{\{x: f(x) > \lambda\}} f(x) \left[\log^+ \left(\frac{f(x)}{\lambda} \right) \right]^{k-1} U(x) dx$$

holds for all $\lambda > 0$ and all $f \geq 0$.

COROLLARY 2. *With the same hypothesis as Theorem 3 and $C > 0$, if $(M_k \cdots M_1 f)(x) \rightarrow 0$ as $|x| \rightarrow \infty$ and $\int_E (M_k \cdots M_1 f)(x) U(x) dx < \infty$ for every bounded set $E \subset R^n$, then*

$$\int_{R^n} |f(x)| [\log^+ |f(x)|]^k U(x) dx < \infty.$$

Proof of Theorem 1. First we will prove that (1) implies (2). Since $\lambda \geq \lambda_0$ and f is supported on Q_0 , $\lambda \geq |Q_0|^{-1} \int_{R^n} f$ so the Calderón-Zygmund decomposition [5, Theorem 4, p. 17] shows that there are pairwise disjoint cubes $\{Q_j\}$ satisfying

$$(10) \quad \bigcup Q_j \subset Q_0,$$

$$(11) \quad \lambda < |Q_j|^{-1} \int_{Q_j} f \leq 2^n \lambda$$

and

$$(12) \quad f(x) \leq \lambda \quad \text{if } x \notin \bigcup Q_j.$$

The definition of Mf and (11) shows that $(Mf)(x) > \lambda$ if $x \in Q_j$ and then (10) yields $\bigcup Q_j \subset \{x \in Q_0: (Mf)(x) > \lambda\}$ so that

$$(13) \quad |\{x \in Q_0: (Mf)(x) > \lambda\}|_U \geq \sum_j |Q_j|_U.$$

Now from (11) and hypothesis (1) it follows that

$$\begin{aligned}
 \sum_j |Q_j|_U &\geq 2^{-n} \lambda^{-1} \sum_j |Q_j|_U \left(|Q_j|^{-1} \int_{Q_j} f(x) dx \right) \\
 &\geq 2^{-n} \lambda^{-1} C \sum_j \left(\operatorname{ess\,sup}_{x \in Q_j} V(x) \right) \left(\int_{Q_j} f(x) dx \right) \\
 &\geq 2^{-n} \lambda^{-1} C \sum_j \int_{Q_j} f(x) V(x) dx = 2^{-n} \lambda^{-1} C \int_{\cup Q_j} f(x) V(x) dx,
 \end{aligned}$$

while from (12) $\cup Q_j \supset \{x \in R^n: f(x) > \lambda\}$ so that

$$\int_{\cup Q_j} f(x) V(x) dx \geq \int_{\{x: f(x) > \lambda\}} f(x) V(x) dx.$$

Thus (1) implies (2).

Conversely, let $Q \subset Q_0$ be given. Let $\varepsilon > 0$ and select $E_\varepsilon \subset Q$ with $0 < |E_\varepsilon| < 2^{-n} |Q|$ so that $V(x) > \operatorname{ess\,sup}_{t \in Q} V(t) - \varepsilon$ for $x \in E_\varepsilon$. Then with $f(x) = 1$ if $x \in E_\varepsilon$, $f(x) = 0$ otherwise, and $\lambda = 2^n |E_\varepsilon| / |Q|$, elementary geometry shows that $(Mf)(x) \leq \lambda$ if $x \notin 2Q$. Thus (2) yields

$$\begin{aligned}
 |(2Q) \cap Q_0|_U &\geq |\{x \in Q_0: (Mf)(x) > \lambda\}|_U \\
 &\geq C 4^{-n} \frac{|Q|}{|E_\varepsilon|} \int_{E_\varepsilon} V(x) dx \geq C 4^{-n} |Q| \left(\operatorname{ess\,sup}_{t \in Q} V(t) - \varepsilon \right).
 \end{aligned}$$

Since ε is arbitrary we obtain (3). This proves Theorem 1.

Proof of Corollary 1. The proof that (5) implies (6) is similar to that used to prove that (2) implies (3) and is therefore omitted. Now if (4) holds and $f(x) \geq 0$, let Q_0 be a fixed cube and let $f_t(x) = f(x)$ if $f(x) \leq t$ and $x \in tQ_0$, $f_t(x) = 0$ otherwise. For $\lambda > 0$, Theorem 1 yields

$$(14) \quad |\{x \in RQ_0: (Mf_t)(x) > \lambda\}|_U \geq C 2^{-n} \lambda^{-1} \int_{\{x: f_t(x) > \lambda\}} f_t(x) V(x) dx$$

provided R satisfies $|RQ_0|^{-1} \int_{R^n} f_t \leq \lambda$ and $R \geq t$. Note that $f_t(x) \uparrow f(x)$ and $(Mf_t)(x) \uparrow (Mf)(x)$ as $t \rightarrow \infty$. Hence (5) follows from (14) by the monotone convergence theorem, letting $R \rightarrow \infty$ first, then $t \rightarrow \infty$. This proves the Corollary.

Proof of Theorem 2. Assume that $V(x) > 0$ on a set of positive measure in Q_0 for otherwise there is nothing to prove. Then (7) shows that $|Q_0|_U > 0$. Hence, if $f(x) \not\equiv 0$ a.e. then $(Mf)(x) \geq |Q_0|^{-1} \int_{Q_0} f > 0$ for

$x \in Q_0$ and the hypothesis $\int_{Q_0} (Mf)(x)U(x) dx < \infty$ implies that f and U are integrable on Q_0 , and in view of (7), V is also. Thus it suffices to show that

$$(15) \quad \int_{\{x: f(x) > \lambda_0\}} f(x) \log \left(\frac{f(x)}{\lambda_0} \right) V(x) dx < \infty$$

where $\lambda_0 = |Q_0|^{-1} \int_{Q_0} f$.

Fubini's Theorem shows that the left side of (15) is equal to

$$\int_{\lambda_0}^{\infty} \frac{d\lambda}{\lambda} \int_{\{x: f(x) > \lambda\}} f(x) V(x) dx$$

and Theorem 1 shows that this is bounded above by

$$C^{-1} 2^n \int_{\lambda_0}^{\infty} |\{x \in Q_0: (Mf)(x) > \lambda\}|_U d\lambda.$$

This integral is bounded by

$$\int_0^{\infty} |\{x \in Q_0: (Mf)(x) > \lambda\}|_U d\lambda = \int_{Q_0} (Mf)(x)U(x) dx$$

so we obtain (15) and the theorem is proved.

Proof of Theorem 3. Observe first that for any fixed i , $1 \leq i \leq k$, Corollary 1 and (8) show that

$$\int_{\{x_i \in R^1: (M_i f)(x) > \lambda\}} U(x) dx_i \geq \frac{2^{-1}C}{\lambda} \int_{\{x_i: f(x) > \lambda\}} f(x)U(x) dx_i.$$

Integrating this inequality over the remaining variables yields

$$(16) \quad |\{x \in R^n: (M_i f)(x) > \lambda\}|_U \geq \frac{2^{-1}C}{\lambda} \int_{\{x: f(x) > \lambda\}} f(x)U(x) dx.$$

Now the proof proceeds by induction. As we have just proved, (9) holds with $k = 1$. Assume that (9) holds for some k , $k \leq n - 1$. Then (16) yields

$$\begin{aligned} & |\{x \in R^n: (M_{k+1} \cdots M_1 f)(x) > \lambda\}|_U \\ & \geq \frac{2^{-1}C}{\lambda} \int_{\{x: (M_k \cdots M_1 f)(x) > \lambda\}} (M_k \cdots M_1 f)(x)U(x) dx \\ & \geq 2^{-1}C \int_{\{x: (M_k \cdots M_1 g)(x) > 1\}} (M_k \cdots M_1 g)(x)U(x) dx \end{aligned}$$

where we have set $g(x) = f(x)/\lambda$ if $f(x) > \lambda$ and $g(x) = 0$ otherwise. Now

$$\begin{aligned} \int_{\{x: (M_k \cdots M_1 g)(x) > 1\}} (M_k \cdots M_1 g)(x) U(x) dx \\ = \int_0^\infty |\{x: (M_k \cdots M_1 g)(x) > \max(1, \alpha)\}|_U d\alpha \\ \geq \int_1^\infty |\{x: (M_k \cdots M_1 g)(x) > \alpha\}|_U d\alpha \end{aligned}$$

so that the inductive hypothesis shows this is bounded below by

$$\begin{aligned} \frac{2^{-k} C^k}{(k-1)!} \int_1^\infty \frac{d\alpha}{\alpha} \int_{\{x: g(x) > \alpha\}} g(x) \left[\log \left(\frac{g(x)}{\alpha} \right) \right]^{k-1} U(x) dx \\ = \frac{2^{-k} C^k}{k!} \int_{\{x: g(x) > 1\}} g(x) [\log g(x)]^k U(x) dx \\ = \frac{2^{-k} C^k}{k! \lambda} \int_{\{x: f(x) > \lambda\}} f(x) \left[\log \left(\frac{f(x)}{\lambda} \right) \right]^k U(x) dx. \end{aligned}$$

Thus we have (9) for $k+1$ and the proof is complete.

Proof of Corollary 2. Since $(M_k \cdots M_1 f)(x) \rightarrow 0$ as $|x| \rightarrow \infty$ the set $E = \{x: (M_k \cdots M_1 f)(x) \geq 1\}$ is bounded. Thus, Theorem 3 shows

$$\begin{aligned} \infty > \int_E (M_k \cdots M_1 f)(x) U(x) dx \geq \int_1^\infty |\{x: (M_k \cdots M_1 f)(x) > \lambda\}|_U d\lambda \\ \geq \frac{2^{-k} C^k}{(k-1)!} \int_1^\infty \frac{d\lambda}{\lambda} \int_{\{x: f(x) > \lambda\}} f(x) \left[\log \left(\frac{f(x)}{\lambda} \right) \right]^{k-1} U(x) dx \\ = \frac{2^{-k} C^k}{k!} \int_{R^n} f(x) [\log^+ f(x)]^k U(x) dx \end{aligned}$$

where we have used Fubini's Theorem to obtain the last equality. This proves the Corollary.

REFERENCES

- [1] N. A. Favo, E. A. Gatto and C. Gutiérrez, *On the strong maximal function and Zygmund's class $L(\log^+ L)^n$* , Studia Math., **69** (1980), 155–158.
- [2] B. Muckenhoupt, *The equivalence of two conditions for weight functions*, Studia Math., **49** (1974), 101–106.
- [3] ———, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc., **167** (1972), 207–226.

- [4] E. M. Stein, *Note on class $L \log L$* , *Studia Math.*, **32** (1969), 305–310.
- [5] ———, *Singular Integrals and the Differentiability Properties of Functions*, Princeton University Press, 1970.

Received June 11, 1982. Research of the first author was supported in part by NSERC grant A-8185. The second author's research was supported in part by NSERC grant A-5165.

UNIVERSITY OF ALBERTA
EDMONTON, T6G 2G1 CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)
University of California
Los Angeles, CA 90024

HUGO ROSSI
University of Utah
Salt Lake City, UT 84112

C. C. MOORE and ARTHUR OGUS
University of California
Berkeley, CA 94720

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, CA 90089-1113

R. FINN and H. SAMELSON
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$132.00 a year (6 Vol., 12 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* ISSN 0030-8730 is published monthly by the Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P. O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS. A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

Kenneth F. Andersen and Wo-Sang Young , On the reverse weak type inequality for the Hardy maximal function and the weighted classes $L(\log L)^k$	257
Richard Eugene Bedient , Double branched covers and pretzel knots	265
Harold Philip Boas , Holomorphic reproducing kernels in Reinhardt domains	273
Janey Antonio Daccach and Arthur Gabriel Wasserman , Stiefel's theorem and toral actions	293
Michael Fried , The nonregular analogue of Tchebotarev's theorem	303
Stanley Joseph Gurak , Minimal polynomials for circular numbers	313
Norimichi Hirano and Wataru Takahashi , Nonlinear ergodic theorems for an amenable semigroup of nonexpansive mappings in a Banach space ...	333
Jim Hoste , Sewn-up r -link exteriors	347
Mohammad Ahmad Khan , The existence of totally dense subgroups in LCA groups	383
Mieczysław Kula, Murray Angus Marshall and Andrzej Śladek , Direct limits of finite spaces of orderings	391
Luis Montejano Peimbert , Flat Hilbert cube manifold pairs	407
Steven C. Pinault , An a priori estimate in the calculus of variations	427
McKenzie Y. K. Wang , Some remarks on the calculation of Stiefel-Whitney classes and a paper of Wu-Yi Hsiang's	431
Brian Donald Wick , The calculation of an invariant for Tor	445
Wolfgang Wollny , Contributions to Hilbert's eighteenth problem	451