

Pacific Journal of Mathematics

GYSIN HOMOMORPHISM AND SCHUBERT CALCULUS

ERSAN AKYILDIZ

GYSIN HOMOMORPHISM AND SCHUBERT CALCULUS

ERSAN AKYILDIZ

Let G be a connected reductive linear algebraic group over the field of complex numbers, and B a fixed Borel subgroup of G . The study of the homological properties of G/B can be carried out by two well-known methods. The first of these methods is due to A. Borel and involves the identification of the cohomology ring of G/B with the quotient ring of the ring of polynomials on the Lie algebra \mathfrak{h} of the Cartan subgroup $H \subset G$ by the ideal generated by the W -invariant polynomials (where W is the Weyl group of G). The second method is classical, and based on the calculation of the homology with the aid of the partition of G/B into cells, the so-called Schubert cells. The correspondence between these approaches has been studied in the paper by Bernstein, Gel'fand and Gel'fand, where in the quotient ring of the polynomial ring figuring in Borel's model of the cohomology, the authors have found a symmetrical basis dual to the Schubert cells. Moreover, they have given a formula (Intersection formula) which expresses the intersection of any Schubert cell with a cell of codimension one. In the same paper, the authors have also generalized these results, except the intersection formula, to the case when B is replaced by an arbitrary parabolic subgroup $P \subset G$.

On the other hand, for any parabolic subgroup $P \subset G$ containing B , the cohomology Gysin homomorphism of $\pi: G/B \rightarrow G/P$ has been studied by Akyildiz and Carrell, where an explicit formula has been obtained between the rings figuring in Borel's model of the cohomologies. This formula also enables one to obtain some of the results mentioned above for G/P from the corresponding results on G/B . In this note, we consider the case where $G = \mathrm{GL}(n+1)$ and G/P is the Grassmann manifold. By using the explicit description of the Gysin homomorphism and the intersection formula given in the cohomology ring of G/B we obtain three main theorems of the symbolic formalism, known as Schubert Calculus, concerning the cohomology ring structure of the Grassmann manifold. Although there are several different approaches for proving these theorems (see Kleiman and Laksov), it seems that none of them uses the cohomology ring structure of $\mathrm{GL}(n+1)/B$, where B is the group of upper triangular matrices in $\mathrm{GL}(n+1)$. We thus hope that this alternative point of view may be used to understand the generalized Schubert Calculus.

1. Gysin homomorphism of $\pi: G/B \rightarrow G/P$. In this section we first give the rings figuring in Borel's model of the cohomologies of G/B , G/P , and then we write the formula for the cohomology Gysin homomorphism of $\pi: G/B \rightarrow G/P$ between these rings. Here G is a connected

reductive linear algebraic group over the field of complex numbers, B is a fixed Borel subgroup of G , and P is a parabolic subgroup of G containing B . We adopt the following notation: B_u the unipotent radical of B , H a fixed maximal torus contained in B , \mathfrak{g} the Lie algebra of G , \mathfrak{h} and \mathfrak{b}_u the Lie algebras of H and B_u respectively, $\Delta \subset \mathfrak{h}^*$ the root system of \mathfrak{h} in \mathfrak{g} , Δ_+ the set of positive roots, namely the set of roots of \mathfrak{h} in \mathfrak{b}_u , $\Sigma \subset \Delta_+$ the set of simple roots, W the Weyl group of G , Θ is the subset of Σ such that the parabolic subgroup P_Θ corresponding to Θ is equal to P , W_Θ the subgroup of W generated by the reflections σ_α , $\alpha \in \Theta$, Δ_Θ the subset of Δ_+ consisting of linear combinations of the elements of Θ , W_Θ^\perp the set of $w \in W$ such that $w\Theta \subset \Delta_+$.

Let $X(H)$ be the group of characters of H , and $R = \text{Sym}(C \otimes_{\mathbb{Z}} X(H))$, the symmetric algebra of $C \otimes_{\mathbb{Z}} X(H)$. We denote by the same symbol an element of $X(H)$ and the corresponding element of Δ when it can be done without any ambiguity. For the basic facts about algebraic groups and the Borel model of the cohomologies of homogeneous spaces, the reader is referred to [6] and [1], [3], [5] respectively. Since W acts on $X(H)$ we get an action of W on R in the usual way. Let R^W be the ring of invariants of W , and $I = \{f \in R^W: f(0) = 0\}$. For any character α of H , let L_α be the associated homogeneous line bundle on G/B : $L_\alpha = G \times C / \sim$, $(g, z) \sim (g', z')$ if and only if $g' = gb$ for some b in B and $z' = \alpha(b^{-1})z$, where α is extended on B with $\alpha(u) = 1$ for u in B_u . It is shown in [1] that the characteristic homomorphism $c: R \rightarrow H^*(G/B, C)$ determined by $c(\alpha) = c_1(L_\alpha)$, the first Chern class of L_α for any $\alpha \in X(H)$, induces an isomorphism of graded algebras $\bar{c}: R/IR \xrightarrow{\sim} H^*(G/B, C)$ (see also [5]). Moreover, if R^{W_Θ} is the ring of invariants of W_Θ , then one also has the following commutative diagram between the graded algebras (cf. [1]):

$$\begin{array}{ccc} \bar{c}: R/IR & \xrightarrow{\sim} & H^*(G/B, C) \\ \uparrow & & \uparrow \pi^* \\ \bar{c}: R^{W_\Theta}/IR^{W_\Theta} & \xrightarrow{\sim} & H^*(G/P, C), \end{array}$$

where π^* is the cohomology map of $\pi: G/B \rightarrow G/P$. This is the Borel model of the cohomologies of G/B and G/P .

For the cohomology Gysin homomorphism of $\pi: G/B \rightarrow G/P$, we first recall the operators $A_\omega: R \rightarrow R$ for $\omega \in W$ (cf. [3], [4], [5]). For each $\alpha \in \Delta$ the element $f - \sigma_\alpha \cdot f$ is divisible by α for any $f \in R$, where σ_α is the reflection corresponding to α . Thus $A_\alpha: R \rightarrow R$, $A_\alpha f = (f - \sigma_\alpha \cdot f)/\alpha$, is a well-defined R^W -linear operator on R . Let $\alpha_1, \dots, \alpha_l \in \Sigma$, and let $\omega = \sigma_{\alpha_1} \cdots \sigma_{\alpha_l}$ be any element of W . Then

- (i) if the length $l(\omega)$ of ω is less than l , then $A_{\alpha_1} \cdots A_{\alpha_l} = 0$,
 (ii) if $l(\omega) = l$, then the operator $A_{\alpha_1} \cdots A_{\alpha_l}$ depends only on ω and not on the representation of ω in the form $\omega = \sigma_{\alpha_1} \cdots \sigma_{\alpha_l}$. In this case we put $A_\omega = A_{\alpha_1} \cdots A_{\alpha_l}$.

We note that the operator $A_\omega: R \rightarrow R$ preserves the ideal IR , and thus induces an operator $\bar{A}_\omega: R/IR \rightarrow R/IR$ of homogeneous degree $-l(\omega)$. Moreover for $\omega_1, \omega_2 \in W$, one has $\bar{A}_{\omega_1} \bar{A}_{\omega_2} = \bar{A}_{\omega_1 \omega_2}$ if $l(\omega_1 \omega_2) = l(\omega_1) + l(\omega_2)$, $\bar{A}_{\omega_1} \bar{A}_{\omega_2} = 0$ otherwise (cf. [4], [5]). It is shown in [2] that the cohomology Gysin homomorphism $\pi_*: H^*(G/B, C) \rightarrow H^*(G/P, C)$ of the natural map $\pi: G/B \rightarrow G/P$ is given by the following commutative diagram:

$$(1) \quad \begin{array}{ccc} \bar{c}: R/IR & \xrightarrow{\sim} & H^*(G/B, C) \\ \bar{A}_{\tau_\Theta} \downarrow & & \downarrow \pi_* \\ \tilde{c}: R^{W_\Theta}/IR^{W_\Theta} & \xrightarrow{\sim} & H^*(G/P, C), \end{array}$$

where τ_Θ is the unique element of W_Θ of maximal length. Moreover, one has the following formula for \bar{A}_{τ_Θ} (cf. [2]):

$$(2) \quad A_{\tau_\Theta}(f) = \sum_{\tau \in W_\Theta} \det(\tau) \tau \cdot f / \prod_{\alpha \in \Delta_\Theta} \alpha.$$

This is the explicit description of the cohomology Gysin homomorphism of $\pi: G/B \rightarrow G/P$.

2. Schubert calculus. The correspondence between the Borel's model of the cohomology of G/P and the calculation of the homology with the aid of partition of G/P into Schubert cells has been studied in [3]. In the case of G/B , the authors have also given a formula (intersection formula) in R/IR which expresses the intersection of any Schubert cell with a cell of codimension one. In this section, we first give this correspondence. Then we write the intersection formula only for the case $G = \text{GL}(n+1)$. Later, by using this formula and the results of §1. We obtain three main theorems, known as Schubert Calculus, concerning the cohomology ring structure of Grassmannians.

We keep the notation of §1. Let ω_0 be the unique element of W of maximal length, and $P_{\omega_0} = (1/|W|) \prod_{\alpha \in \Delta_+} \alpha \pmod{IR}$, where $|W|$ is the order of W . For each $\omega \in W$, let $P_\omega = \bar{A}_{\omega^{-1}\omega_0}(P_{\omega_0})$. It is shown in [3] that $\{P_\omega: \omega \in W\}$ is a free \mathbb{Z} -basis of R/IR which is dual to the Schubert cells. Namely, for each $\omega \in W$, let X_ω denote the cycle class of the closure of $B(\omega_0\omega)x_0$ in $H_*(G/B, \mathbb{Z})$, $x_0 = B \in G/B$. Then under the Poincare

duality map \mathcal{P} , $\mathcal{P}(X_\omega) = \bar{c}(P_\omega)$. Similarly, for the parabolic subgroup $P = P_\Theta$ of G containing B , one has the following: For each $\sigma \in W_\Theta^\perp$ let Y_σ be the cycle class of the closure of $B(\omega_0\sigma)\pi(x_0)$ in $H_*(G/P, \mathbb{Z})$, where $\pi: G/B \rightarrow G/P$ is the natural map. Then the Poincare dual $\mathcal{P}(Y_\sigma)$ of Y_σ , $\sigma \in W_\Theta^\perp$, is equal to $\bar{c}(P_\sigma \pmod{IR^{W_\Theta}})$. Moreover $\{P_\sigma \pmod{IR^{W_\Theta}}: \sigma \in W_\Theta^\perp\}$ is a free \mathbb{Z} -basis of $R^{W_\Theta}/IR^{W_\Theta}$. Although the proof of this fact is given in [3], for the sake of completeness of our approach we will give a different proof by using the Gysin homomorphism (cf. [2]). Let τ_Θ be the unique element of W_Θ of maximal length. Since $l(\sigma\tau) = l(\sigma) + l(\tau)$ for any $\sigma \in W_\Theta^\perp$ and $\tau \in W_\Theta$, we have

$$\bar{A}_{\tau_\Theta}(P_{\sigma\tau}) = \bar{A}_{\tau_\Theta}\bar{A}_{\tau^{-1}\sigma^{-1}\omega_0}(P_{\omega_0}) = \delta_{\tau_\Theta, \tau}A_{\sigma^{-1}\omega_0}(P_{\omega_0}) = \delta_{\tau_\Theta, \tau}P_\sigma.$$

This shows that $\{P_\sigma \pmod{IR^{W_\Theta}}: \sigma \in W_\Theta^\perp\}$ is a basis of $R^{W_\Theta}/IR^{W_\Theta}$, because the Gysin homomorphism \bar{A}_{τ_Θ} is surjective and any element ω of W can be written uniquely in the form $\omega = \sigma\tau$ for some $\sigma \in W_\Theta^\perp$ and $\tau \in W_\Theta$. On the other hand, it is clear that for $\sigma \in W_\Theta^\perp$, the cycle class $X_{\sigma\tau_\Theta}$ goes under the homology Gysin homomorphism of $\pi: G/B \rightarrow G/P$ to the cycle class Y_σ in $H_*(G/P, \mathbb{Z})$. Thus from the commutativity of the diagram (1) and $\bar{c}(P_{\sigma\tau_\Theta}) = \mathcal{P}(X_{\sigma\tau_\Theta})$, we get $\bar{c}(P_\sigma \pmod{IR^{W_\Theta}}) = \mathcal{P}(Y_\sigma)$ for any $\sigma \in W_\Theta^\perp$.

There is a formula (intersection formula) given in R/IR which expresses the product $P_{\sigma_\alpha} \cdot P_\sigma$ in terms of P_ω , where σ_α is the reflection corresponding to the simple root $\alpha \in \Sigma$ (cf. [3]). We will now write this formula for the group of invertible matrices $GL(n+1)$. Let G be $GL(n+1)$, B the group of upper triangular matrices in G , and H the group of diagonal matrices in B . Then

$$R = \text{Sym}(C \otimes_{\mathbb{Z}} X(H)) = C[x_0, \dots, x_n],$$

$$\Delta_+ = \{\alpha_{i,j} = x_i - x_j; 0 \leq i < j \leq n\},$$

$$\Sigma = \{\alpha_{i,i+1}; 0 \leq i \leq n-1\}.$$

Let S_{n+1} be the symmetric group in $0, 1, \dots, n$, and for $\sigma \in S_{n+1}$ let $\sigma(I)$ be the permutation matrix obtained from $(n+1) \times (n+1)$ identity matrix I . The homomorphism $\sigma^{-1} \rightarrow \sigma(I)$ gives an isomorphism between S_{n+1} and the Weyl group W of G . Moreover it can be checked that the action of W on R is given by

$$\begin{aligned} \sigma(I) \cdot f(x_0, \dots, x_n) &= \sigma f(x_0, \dots, x_n) \\ &= f(x_{\sigma_0}, \dots, x_{\sigma_n}), \quad \text{for any } \sigma \in S_{n+1}. \end{aligned}$$

Thus IR is the ideal generated by the elementary symmetric functions of x_0, \dots, x_n .

We will now write the operators A_ω , and the elements P_ω of R/IR in terms of the elements of S_{n+1} . Let $\sigma = (a_0 \cdots a_n)$ be any element of S_{n+1} , namely $\sigma i = a_i$ for $i = 0, 1, \dots, n$. Then the reflection $\sigma_{\alpha_{i,j}}$ corresponding to the root $\alpha_{i,j}$ is given by $\sigma_{\alpha_{i,j}} = (i, j)(I)$, where $(i, j) \in S_{n+1}$ is the transposition obtained by changing i with j . Let $A_{(i,i+1)} = A_{\alpha_{i,i+1}}$, and $P_\sigma = P_{\sigma(I)}$ for $\sigma \in S_{n+1}$. Then for any $\sigma \in S_{n+1}$, we get

$$A_{\sigma^{-1}} = A_{\sigma(I)}, \quad \text{and} \quad P_\sigma = \bar{A}_{\sigma\omega_0}(P_{\omega_0}),$$

where $\omega_0 = (n \ n - 1 \ \cdots \ 1 \ 0)$, and

$$P_{\omega_0} = \frac{1}{(n+1)!} \prod_{0 \leq i < j \leq n} (x_i - x_j) \pmod{IR}.$$

The intersection formula given in [3] can now be written as follows:

THEOREM (*Intersection formula*, cf. [3, p. 17]). *For any $r = 0, 1, \dots, n - 1$, and $\sigma = (a_0 \cdots a_n) \in S_{n+1}$, we have*

$$P_{(r,r+1)} P_\sigma = \sum P_{(i,j)\sigma},$$

where the summation is over all $0 \leq i < j \leq n$ such that

- (a) $i \leq r < j$,
- (b) $a_i < a_j$,
- (c) if $i < k < j$, then either $a_k < a_i$ or $a_k > a_j$.

We are now ready to discuss the cohomology ring structure of Grassmannians. For any $q = 0, 1, \dots, n - 1$, let $\Theta = \Sigma \setminus \{\alpha_{q,q+1}\}$. Then the parabolic subgroup $P = P_\Theta$ of G corresponding to Θ consists of all matrices of the form $\begin{pmatrix} A & * \\ 0 & B \end{pmatrix}$, where $A \in GL(q+1)$, and $B \in GL(n-q)$. Thus G/P is the Grassmann manifold $G_{q,n}$ of $(q+1)$ -planes in C^{n+1} . Moreover it is easy to see that corresponding to Θ , we have

$$W_\Theta = \{(a_0 \cdots a_q a_{q+1} \cdots a_n) \in S_{n+1}; 0 \leq a_0, \dots, a_q \leq q, \\ q+1 \leq a_{q+1}, \dots, a_n \leq n\},$$

and

$$W_\Theta^\perp = \{(a_0 \cdots a_q \cdots a_n) \in S_{n+1}; a_0 < \cdots < a_q, a_{q+1} < \cdots < a_n\}.$$

A sequence of integers $0 \leq a_0 < \cdots < a_q \leq n$ corresponds to a unique element $(a_0 \cdots a_q a_{q+1} \cdots a_n)$ of W_Θ^\perp , and we denote this element by $\sigma(a_0, \dots, a_q)$. For any sequence of integers $0 \leq a_0 < \cdots < a_q \leq n$, let $\Omega(a_0, \dots, a_q)$ denote the Poincare dual of $Y_{\sigma(a_0, \dots, a_q)}$ in $H^*(G_{q,n}, Z) \subset H^*(G_{q,n}, C)$. $\Omega(a_0, \dots, a_q)$ is called the Schubert cycle corresponding to

$0 \leq a_0 < \cdots < a_q \leq n$. The Schubert cycle corresponding to $0 < 1 < \cdots < q-1 < q+k \leq n$ for $k = 0, 1, \dots, n-q$ is called the special Schubert cycle, and it is denoted by $\Omega(k)$.

THEOREM (The basis theorem). *The Schubert cycles $\Omega(a_0, \dots, a_q)$, $0 \leq a_0 < \cdots < a_q \leq n$, form a basis of $H^*(G_{q,n}, \mathbb{Z})$.*

Proof. It follows from the above observations because $\Omega(a_0, \dots, a_q) = P_{\sigma(a_0, \dots, a_q)} \pmod{IR^{W_\Theta}}$ for any sequence of integers $0 \leq a_0 < \cdots < a_q \leq n$.

We will now compute $\Omega(a_0, \dots, a_q) = P_{\sigma(a_0, \dots, a_q)} \pmod{IR^{W_\Theta}}$. Let $\Theta_1 = \{\alpha_{i, i+1} : 0 \leq i \leq q-1\}$, and $P_1 = P_{\Theta_1}$ be the corresponding parabolic subgroup of G . Then we have

$$W_{\Theta_1} = \{(a_0 \cdots a_q q+1 \cdots n) \in S_{n+1} : 0 \leq a_0, \dots, a_q \leq q\},$$

$$W_{\Theta_1}^\perp = \{(a_0 \cdots a_q \cdots a_n) \in S_{n+1} : a_0 < \cdots < a_q\},$$

and $\tau_0 = (q \ q-1 \ \cdots \ 0 \ q+1 \ \cdots \ n)$ is the unique element of W_{Θ_1} of maximal length. Since $W_\Theta^\perp \subset W_{\Theta_1}^\perp$, we can also use the Gysin homomorphism \bar{A}_{τ_0} of $\pi_1: G/B \rightarrow G/P_1$ to compute P_σ for any $\sigma \in W_\Theta^\perp$. Namely, for $\sigma \in W_\Theta^\perp$ we have $P_\sigma = \bar{A}_{\tau_0}(P_{\tau_0\sigma})$. Although that it is possible to compute P_σ , $\sigma \in W_\Theta^\perp$, by using the Gysin homomorphism \bar{A}_{τ_Θ} , $P_\sigma = \bar{A}_{\tau_\Theta}(P_{\tau_\Theta\sigma})$, it is much simpler to do this by using \bar{A}_{τ_0} . For that, we need to compute $P_{\tau_0\sigma}$ for any $\sigma \in W_\Theta^\perp$.

PROPOSITION. *Let $\sigma = (a_0 \cdots a_q \cdots a_n) \in W_\Theta^\perp$. Then we have $P_{\tau_0\sigma} = x_0^{a_q} \cdots x_q^{a_0} \pmod{IR}$.*

Proof. Let $\tau_0\sigma = (a_q \cdots a_0 \ a_{q+1} \cdots a_n) = (b_0 \cdots b_n) \in S_{n+1}$. Since $b_0 > b_1 > \cdots > b_q$ and $b_{q+1} < \cdots < b_n$, it is easy to see that from the intersection formula one gets the following:

(i) $P_{(q, q+1)} P_{\tau_0\sigma} = \sum P_{(i, q+k)\tau_0\sigma}$, where the summation is all over $0 \leq i \leq q$ and $1 \leq k \leq n-q$ such that $b_{q+k-1} \leq b_i < b_{q+k}$,

(ii) $P_{(q-1, q)} P_{\tau_0\sigma} = \sum P_{(j, q+r)\tau_0\sigma}$, where the summation is all over $0 \leq j \leq q-1$ and $1 \leq r \leq n-q$ such that $b_{q+r-1} \leq b_j < b_{q+r}$. Thus

$$\begin{aligned} & (P_{(q, q+1)} - P_{(q-1, q)}) P_{\tau_0\sigma} \\ &= \begin{cases} P_{(q, q+k)\tau_0\sigma}, & \text{if there exists } 1 \leq k \leq n-q \\ & \text{such that } b_{q+k-1} \leq b_q < b_{q+k}, \\ 0 \pmod{IR}, & \text{otherwise.} \end{cases} \end{aligned}$$

We note that the fundamental dominant weight χ_r corresponding to the simple root $\alpha_{r,r+1}$ is given by $\chi_r = x_0 + \cdots + x_r$. By [3, p. 18], since $P_{(r,r+1)} = \chi_r \pmod{IR}$, we have $P_{(r,r+1)} - P_{(r-1,r)} = x_r \pmod{IR}$ for $r = 0, 1, \dots, n-1$.

We prove the proposition by induction on q . If $q = 0$, then $\tau_0\sigma = (b_0 \cdots b_n)$, $b_1 < \cdots < b_n$. In this case we can have either $\tau_0\sigma = e$ (the identity) or $\tau_0\sigma = \omega(p+1) = (p+1 \ 0 \ \cdots \ p \ p+2 \ \cdots \ n)$ for some $p = 0, 1, \dots, n-1$. If $\tau_0\sigma = e$, then the claim follows from the formula (2), $P_e = \bar{A}_{\omega_0}(P_{\omega_0}) = 1 \pmod{IR}$. If $\tau_0\sigma = \omega(p+1)$ for some $p = 0, 1, \dots, n-1$, then we claim that $P_{\omega(p+1)} = x_0^{p+1} \pmod{IR}$. We prove this by induction on p . If $p = 0$, then $\omega(1) = (0, 1)$. Since $P_{(0,1)} = \chi_0 = x_0 \pmod{IR}$, we have the claim for $p = 0$. We now assume $P_{\omega(p)} = x_0^p \pmod{IR}$. But by the intersection formula we have $P_{(0,1)}P_{\omega(p)} = P_{(0,p+1)\omega(p)}$. Since $(0, p+1)\omega(p) = \omega(p+1)$, by the induction hypothesis we have $P_{\omega(p+1)} = x_0x_0^p = x_0^{p+1} \pmod{IR}$. This finishes the induction argument for $q = 0$. We now assume the claim for $q-1$. Let $\tau_0\sigma = (a_q \cdots a_0 a_{q+1} \cdots a_n) = (b_0 \cdots b_n)$. If $a_0 = b_q = 0$, then the claim follows from the induction hypothesis for $q-1$ because $b_0 > \cdots > b_{q-1}$, $0 = b_q < \cdots < b_n$. If $b_q > 0$, then we claim that $P_{\tau_0\sigma} = x_0^{b_0} \cdots x_q^{b_q} \pmod{IR}$. This will be proved by induction on b_q . If $b_q = 1$, then $b_{q+1} = 0$. In this case $\omega = (q, q+1)\tau_0\sigma = (b_0 \cdots b_{q-1} \ 0 \ 1 \ b_{q+2} \cdots b_n)$, and thus (by the induction hypothesis for $q-1$) $P_\omega = x_0^{b_0} \cdots x_{q-1}^{b_{q-1}} \pmod{IR}$. Since $P_{(q,q+1)} - P_{(q-1,q)} = x_q \pmod{IR}$, we have from above

$$(P_{(q,q+1)} - P_{(q-1,q)})P_\omega = P_{(q,q+1)\omega} = P_{\tau_0\sigma} = x_0^{b_0} \cdots x_{q-1}^{b_{q-1}}x_q \pmod{IR}.$$

We now assume the claim for $b_q - 1 \geq 0$. If $b_q > 1$, then $b_{q+1} = 0$ and $b_q - 1 \in \{b_{q+2}, \dots, b_n\}$. Let $b_q - 1 = b_{q+2+s}$ for some $s = 0, 1, \dots, n-q-2$. Consider

$$\begin{aligned} \omega &= (q, q+2+s)\tau_0\sigma \\ &= (b_0 \cdots b_{q-1} \ b_q - 1 \ b_{q+1} \cdots b_{q+1+s} \ b_q \ b_{q+3+s} \cdots b_n). \end{aligned}$$

Since $b_0 > \cdots > b_q - 1$, $b_{q+1} < \cdots < b_{q+1+s} < b_q < b_{q+3+s} < \cdots < b_n$, by the induction hypothesis we have $P_\omega = x_0^{b_0} \cdots x_q^{b_q-1} \pmod{IR}$. But from above and the induction hypothesis, we get

$$\begin{aligned} (P_{(q,q+1)} - P_{(q-1,q)})P_\omega &= P_{(q,q+2+s)\omega} = P_{\tau_0\sigma} \\ &= x_qx_0^{b_0} \cdots x_q^{b_q-1} = x_0^{b_0} \cdots x_q^{b_q} \pmod{IR} \end{aligned}$$

which is the claim. This completes the proof of the proposition.

COROLLARY. For any sequence of integers $0 \leq a_0 < \cdots < a_q \leq n$, we have

$$P_{\sigma(a_0, \dots, a_q)} = \begin{vmatrix} h(a_0) & \cdots \cdots \cdots & h(a_0 - q) \\ \vdots & & \vdots \\ \vdots & & \vdots \\ \vdots & & \vdots \\ h(a_q) & \cdots \cdots \cdots & h(a_q - q) \end{vmatrix} \pmod{IR},$$

where $h(m)$ is the m th complete homogeneous symmetric function of x_0, x_1, \dots, x_q . In particular $P_{\sigma(0, 1, \dots, q-1, q+k)} = h(k) \pmod{IR}$ for any $k = 0, 1, \dots, n - q$.

Proof. By the proposition we have $P_{\tau_0 \sigma(a_0, \dots, a_q)} = x_0^{a_q} \cdots x_q^{a_0} \pmod{IR}$. On the other hand, from formula (2) we get

$$\begin{aligned} P_{\sigma(a_0, \dots, a_q)} &= \overline{A}_{\tau_0} \left(P_{\tau_0 \sigma(a_0, \dots, a_q)} \right) \\ &= \frac{\sum_{\tau \in W_{\Theta_1}} \det(\tau) x_{\tau_0}^{a_q} \cdots x_{\tau q}^{a_0}}{\prod_{0 \leq i < j \leq q} (x_i - x_j)} \pmod{IR}. \end{aligned}$$

But this bialternant is equal to (cf. [8, p. 92])

$$\begin{vmatrix} h(a_q - q) & \cdots \cdots \cdots & h(a_q) \\ \vdots & & \vdots \\ h(a_0 - q) & \cdots \cdots \cdots & h(a_0) \end{vmatrix} = \begin{vmatrix} h(a_0) & \cdots \cdots \cdots & h(a_0 - q) \\ \vdots & & \vdots \\ h(a_q) & \cdots \cdots \cdots & h(a_q - q) \end{vmatrix},$$

which gives the claim.

THEOREM (The determinantal formula). For all sequence of integers $0 \leq a_0 < \cdots < a_q \leq n$ the following formula holds in the cohomology ring $H^*(G_{q,n}, \mathbb{Z})$:

$$\Omega(a_0, \dots, a_q) = \begin{vmatrix} \Omega(a_0) & \cdots \cdots \cdots & \Omega(a_0 - q) \\ \vdots & & \vdots \\ \Omega(a_q) & \cdots \cdots \cdots & \Omega(a_q - q) \end{vmatrix},$$

where we agree to put $\Omega(k) = 0$ for $k > n - q$ or $k < 0$.

Proof. It follows from the Corollary above the fact that; if $k > n - q$, then the k th complete homogeneous symmetric function of x_0, \dots, x_q

(being equal to \pm the k th elementary symmetric function of x_{q+1}, \dots, x_n (mod IR)) is equal to zero (mod IR^{w_Θ}), (cf. [8]).

This theorem, together with the basis theorem, implies that the special Schubert cycles generate the cohomology ring $H^*(G_{q,n}, \mathbb{Z})$ as a \mathbb{Z} -algebra. Moreover, it reduces the problem of determining the product of two arbitrary Schubert cycles to the case where one (or for that matter, each) is a special Schubert cycle. This case is handled by the following theorem.

THEOREM (Pieri's formula). *For all sequences of integers $0 \leq a_0 < \dots < a_q \leq n$ and $k = 0, 1, \dots, n - q$, the following formula holds in the cohomology ring $H^*(G_{q,n}, \mathbb{Z})$:*

$$\Omega(k)\Omega(a_0, \dots, a_q) = \sum \Omega(b_0, \dots, b_q),$$

where the sum ranges over all sequences of integers $b_0 < \dots < b_q$ satisfying $0 \leq a_0 \leq b_0 < a_1 \leq b_1 < \dots < a_q \leq b_q \leq n$ and $\sum_{i=0}^q (b_i - a_i) = k$.

Proof. For any sequence of integers $0 \leq a_0 < \dots < a_q \leq n$ and for $k = 0, 1, \dots, n - q$, by the proposition and it's corollary we have

$$P_{\tau_0\sigma(a_0, \dots, a_q)} = x_0^{a_q} \dots x_q^{a_0} \pmod{IR}$$

and $P_{\sigma(0, 1, \dots, q-1, q+k)} = h(k) = \sum x_0^{l_q} \dots x_q^{l_0} \pmod{IR}$, where the sum ranges over all integers $l_i \geq 0$ such that $\sum_{i=0}^q l_i = k$. Thus we have

$$(3) \quad P_{\sigma(0, 1, \dots, q-1, q+k)} P_{\tau_0\sigma(a_0, \dots, a_q)} = \sum x_0^{a_q + l_q} \dots x_q^{a_0 + l_0} \\ = f(x_0, \dots, x_q) \pmod{IR}$$

where the sum ranges over all integers $l_i \geq 0$ such that $\sum_{i=0}^q l_i = k$. We will now apply the operator \bar{A}_{τ_0} to equation (3). Since \bar{A}_{τ_0} is $R^{w_{\Theta_1}}$ -linear and $h(k) \in R^{w_{\Theta_1}}$ (where $R^{w_{\Theta_1}}$ is the ring of invariants of the group W_{Θ_1}), we have

$$(4) \quad \bar{A}_{\tau_0} \left(P_{\sigma(0, 1, \dots, q-1, q+k)} P_{\tau_0\sigma(a_0, \dots, a_q)} \right) = P_{\sigma(0, 1, \dots, q-1, q+k)} \bar{A}_{\tau_0} \left(P_{\tau_0\sigma(a_0, \dots, a_q)} \right) \\ = P_{\sigma(0, 1, \dots, q-1, q+k)} P_{\sigma(a_0, \dots, a_q)}.$$

Let $J_{\Theta_1}(g) = \sum_{\tau \in W_{\Theta_1}} \det(\tau) \tau g$ for $g \in R$. Since $J_{\Theta_1}(\tau g) = \det(\tau) J_{\Theta_1}(g)$ for any $\tau \in W_{\Theta_1}$, we have in particular

$$(i) \quad J_{\Theta_1}(X_0^{m_0} \dots X_q^{m_q}) = \det(\tau) J_{\Theta_1}(X_0^{m_0} \dots X_q^{m_q}) \text{ for any } \tau \in W_{\Theta_1},$$

$$(ii) \quad J_{\Theta_1}(X_0^{m_0} \dots X_q^{m_q}) = 0 \text{ if } m_i = m_j \text{ for some } i \neq j.$$

By using (i), (ii), and the fact that $X_0^{n+1} = 0 \pmod{IR}$, it is to see that

$$J_{\Theta_1}(f(x_0, \dots, x_q)) = \sum J_{\Theta_1}(x_0^{a_q + \lambda_q} \dots x_q^{a_0 + \lambda_0}),$$

where the sum ranges over all integers $\lambda_i \geq 0$ satisfying $a_0 + \lambda_0 < \cdots < a_q + \lambda_q \leq n$ and $\sum_{i=0}^q \lambda_i = k$. This gives by the proposition

$$\bar{J}_{\Theta_1}(\bar{f}) = \sum \bar{J}_{\Theta_1}(P_{\tau_0 \sigma(b_0, \dots, b_q)}),$$

where $\bar{f} = f(x_0, \dots, x_q) \pmod{IR}$, and the sum ranges over all sequences of integers $b_0 < b_1 < \cdots < b_q$ satisfying $0 \leq a_0 \leq b_0 < a_1 \leq b_1 < \cdots < a_q \leq b_q \leq n$ and $\sum_{i=0}^q (b_i - a_i) = k$. Thus from formula (2) we get

$$(5) \quad \bar{A}_{\tau_0}(\bar{f}) = \sum \bar{A}_{\tau_0}(P_{\tau_0 \sigma(b_0, \dots, b_q)}) = \sum P_{\sigma(b_0, \dots, b_q)},$$

where the sum is as above. By comparing (3), (4) and (5) we obtain

$$(6) \quad P_{\sigma(0, 1, \dots, q-1, q+k)} P_{\sigma(a_0, \dots, a_q)} = \sum P_{\sigma(b_0, \dots, b_q)},$$

where the sum ranges over all sequences of integers $b_0 < b_1 < \cdots < b_q$ satisfying $0 \leq a_0 \leq b_0 < a_1 \leq b_1 < \cdots < a_q \leq b_q \leq n$ and $\sum_{i=0}^q (b_i - a_i) = k$. By considering equation (6) $\pmod{IR^{W_\Theta}}$, we get the claim, because \tilde{c} is a graded algebra isomorphism and $\tilde{c}(P_{\sigma(a_0, \dots, a_q)} \pmod{IR^{W_\Theta}}) = \Omega(a_0, \dots, a_q)$.

REFERENCES

- [1] E. Akyildiz, *Vector Fields and Cohomology of G/P* , Lecture Notes in Math. **956** (1982), Springer-Verlag, pp. 1–9.
- [2] E. Akyildiz and J. B. Carrell, *An algebraic formula for the Gysin homomorphism from G/B to G/P* , to appear.
- [3] I. N. Bernstein, I. M. Gel'fand and S. I. Gel'fand, *Schubert cells and cohomology of the space G/P* , Russian Math. Surveys, **28** (1973), 1–26.
- [4] M. Demazure, *Invariant symétriques entiers des groupes de Weyl et torsion*, Inventiones Math., **21** (1973), 287–301.
- [5] ———, *Désingularisation des variétés de Schubert généralisées*, Ann. Scient. Éc. Norm. Sup., 4^e serie, t.7 (1974), 53–88.
- [6] J. E. Humphreys, *Linear Algebraic Groups*, New York-Heidelberg-Berlin: Springer-Verlag, GTM 21, 1975.
- [7] S. Kleiman and D. Laksov, *Schubert calculus*, Amer. Math. Monthly, **79** (1972), 1061–1082.
- [8] D. E. Littlewood, *A University Algebra*, second edition, London: Heinemann 1958.

Received April 18, 1983.

On leave from the Middle East
Technical University, Ankara, Turkey

UNIVERSITY OF PETROLEUM AND MINERALS
P.O. Box 144, UPM #841
DHAHRAN, SAUDIA ARABIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
University of Southern California
Los Angeles, CA 90089-1113

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

C. C. MOORE
University of California
Berkeley, CA 94720

ARTHUR OGUS
University of California
Berkeley, CA 94720

HUGO ROSSI
University of Utah
Salt Lake City, UT 84112

H. SAMELSON
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

Ersan Akyildiz, Gysin homomorphism and Schubert calculus	257
Marilyn Breen, Clear visibility and unions of two starshaped sets in the plane	267
Robert F. Brown, Retraction methods in Nielsen fixed point theory	277
Herbert Busemann and Bhalchandra B. Phadke, A general version of Beltrami's theorem in the large	299
Gerald Arthur Edgar and Robert Francis Wheeler, Topological properties of Banach spaces	317
Yaakov Friedman and Bernard Russo, Conditional expectation without order	351
Robert Allen Goggins, Cobordism of manifolds with strong almost tangent structures	361
Mike Hoffman, Noncoincidence index of manifolds	373
William H. Julian, ε -continuity and monotone operations	385
Gerasimos E. Ladas, Y. G. Sficas and I. P. Stavroulakis, Nonoscillatory functional-differential equations	391
Arnold William Miller and Karel Libor Prikry, When the continuum has cofinality ω_1	399
Jean-Leah Mohrherr, Density of a final segment of the truth-table degrees	409
Carl Norman Mutchler, The flat Cauchy problem for radially hyperbolic operators from a characteristic manifold of high codimension	421
Kenji Nakagawa, On the orders of automorphisms of a closed Riemann surface	435
W. Ricker, Representation of vector-valued functions by Laplace transforms	445
Jorge Donato Samur, On semigroups of convolution operators in Hilbert space	463
Joseph Gail Stampfli, One-dimensional perturbations of operators	481
Andrew George Earnest and John Sollion Hsia, Correction to: "Spinor norms of local integral rotations. II"	493